



US008273958B2

(12) **United States Patent**  
**Napier et al.**(10) **Patent No.:** **US 8,273,958 B2**  
(45) **Date of Patent:** **Sep. 25, 2012**(54) **PROCESS FOR THE PRODUCTION OF**  
**ARACHIDONIC ACID AND/OR**  
**EICOSAPENTAENOIC ACID**WO WO-2004/057001 A2 7/2004  
WO WO-2005/012316 A2 2/2005  
WO WO-2005/047479 A2 5/2005(75) Inventors: **Johnathan A. Napier**, Preston (GB);  
**Olga Sayanova**, Harpenden (GB)

## OTHER PUBLICATIONS

(73) Assignee: **BASF Plant Science GmbH** (DE)Wang, X.M., et al., "Biosynthesis and regulation of linolenic acid in higher plants", *Plant Physiol. Biochem.*, 1988, vol. 26, No. 6, pp. 777-792.

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

McKeon, T., et al., "Stearoyl-acyl carrier protein desaturase from safflower seeds", *Methods in Enzymol.*, 1981, vol. 71, 275-281.(21) Appl. No.: **13/186,030**Huang, Y.-S., et al., "Cloning of  $\Delta 12$ - and  $\Delta 6$ -desaturases from *Mortierella alpina* and recombinant production of  $\gamma$ -linolenic acid in *Saccharomyces cerevisiae*", *Lipids*, 1999, vol. 34, No. 7, pp. 649-659.(22) Filed: **Jul. 19, 2011**Wada, H., et al., "Enhancement of chilling tolerance of a cyanobacterium by genetic manipulation of fatty acid desaturation", *Nature*, 1990, vol. 347, pp. 200-203.(65) **Prior Publication Data**

US 2012/0011618 A1 Jan. 12, 2012

Stukey, J.E., et al., "The OLE1 gene of *Saccharomyces cerevisiae* encodes the  $\Delta 9$  fatty acid desaturase and can be functionally replaced by the rat stearoyl-CoA desaturase gene", *J. Biol. Chem.*, 1990, vol. 265, No. 33, pp. 20144-20149.**Related U.S. Application Data**

(62) Division of application No. 12/083,300, filed as application No. PCT/EP2006/067223 on Oct. 10, 2006, now Pat. No. 8,017,839.

Sayanova, O., et al., "A bifunctional  $\Delta 12$ ,  $\Delta 15$ -desaturase from *Acanthamoeba castellanii* directs the synthesis of highly unusual *n*-1 series unsaturated fatty acids", *J. Biol. Chem.*, 2006, vol. 281, No. 48, pp. 36533-36541.(30) **Foreign Application Priority Data**

(Continued)

Oct. 13, 2005 (GB) ..... 0520843.4

(51) **Int. Cl.****A01H 5/00** (2006.01)  
**C12N 15/82** (2006.01)  
**C07H 21/04** (2006.01)*Primary Examiner* — Elizabeth McElwain(52) **U.S. Cl.** ..... **800/298**; 800/281; 536/23.2(74) *Attorney, Agent, or Firm* — Connolly Bove Lodge & Hutz LLP(58) **Field of Classification Search** ..... None  
See application file for complete search history.(57) **ABSTRACT**(56) **References Cited**

## U.S. PATENT DOCUMENTS

5,614,393 A 3/1997 Thomas et al.  
2005/0132442 A1 6/2005 Yadav et al.

## FOREIGN PATENT DOCUMENTS

CA 2180154 C 7/1995  
CA 2533613 A1 2/2005  
EP 0550162 A1 7/1993  
EP 0794250 A1 9/1997  
EP 0472722 B1 5/2003  
WO WO-91/13972 A1 9/1991  
WO WO-93/06712 A1 4/1993  
WO WO-93/11245 A1 6/1993  
WO WO-94/11516 A1 5/1994  
WO WO-94/18337 A1 8/1994  
WO WO-95/18222 A1 7/1995  
WO WO-96/21022 A2 7/1996  
WO WO-97/21340 A1 6/1997  
WO WO-97/30582 A1 8/1997  
WO WO-98/46763 A1 10/1998  
WO WO-98/46764 A1 10/1998  
WO WO-98/46765 A1 10/1998  
WO WO-98/46776 A2 10/1998  
WO WO-99/27111 A1 6/1999  
WO WO-99/33958 A2 7/1999  
WO WO-99/64616 A2 12/1999  
WO WO-00/21557 A1 4/2000  
WO WO-00/34439 A1 6/2000

The present invention relates to a new process for the production of arachidonic acid and/or eicosapentaenoic acid in plants through the co-expression of a  $\Delta$ -12-/ $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase and a  $\Delta$ -5-desaturase and a process for the production of lipids or oils having an increased content of unsaturated fatty acids, in particular  $\omega$ -3 and  $\omega$ -6 fatty acids having at least two double bonds and a 18 or 20 carbon atom chain length. Preferably the arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio. The invention furthermore relates to the production of a transgenic plants, preferably a transgenic crop plant, having an increased content of arachidonic acid and/or eicosapentaenoic acid, oils or lipids containing  $C_{18}$ - or  $C_{20}$ -fatty acids with a double bond in position  $\Delta$ 5, 8, 9, 11, 12, 14, 15 or 17 of the fatty acid produced, respectively due to the expression of the  $\Delta$ -12-/ $\Delta$ -15-desaturase, of the  $\Delta$ -9-elongase, of the  $\Delta$ -8-desaturase and of the  $\Delta$ -5-desaturase in the plant. The expression of the inventive  $\Delta$ -12-/ $\Delta$ -15-desaturase leads preferably to linoleic acid and linolenic acid as products having a double bond in the position  $\Delta$ 9, 12 and 15 of the fatty acid. The invention additionally relates to specific nucleic acid sequences encoding for proteins with  $\Delta$ -12-/ $\Delta$ -15-desaturase-,  $\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- or  $\Delta$ -5-desaturase-activity, nucleic acid constructs, vectors and transgenic plants containing said nucleic acid sequences.

**22 Claims, 13 Drawing Sheets**

OTHER PUBLICATIONS

- "Mucor circinelloides D12d mRNA for delta-12 fatty acid desaturase complete cds", Database NCBI, Accession No. AB052087, Oct. 2, 2001.
- Qi, B., et al. "Identification of a cDNA encoding a novel C18- $\Delta^9$  polyunsaturated fatty acid-specific elongating activity from the docosahexaenoic acid (DHA)-producing microalga, *IsochrYSIS galbana*", FEBS Letters, 2002, vol. 510, pp. 159-165.
- Wallis, J.G., et al., "The  $\Delta^8$ -desaturase of *Euglena gracilis*: an alternate pathway for synthesis of 20-carbon polyunsaturated fatty acids", Archives of Biochemistry and Biophysics, 1999, vol. 365, No. 2, pp. 307-316.
- Napier, J.A., et al., "Progress toward the production of long-chain polyunsaturated fatty acids in transgenic plants", Lipids, 2004, vol. 39, No. 11, pp. 1067-1075.
- Qi, B., et al., "Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants", Nature Biotechnology, 2004, vol. 22, No. 6, pp. 739-745.

Figure 1: Biosynthesis pathway to ARA and/or EPA




Figure 2 A: Comparison of the fatty acid profile of yeast transformed with the constructs pYES2 (2A) as control and construct pYES2-12Ac (2B). The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3.




Figure 2B: Comparison of the fatty acid profile of yeast transformed with the constructs pYES2 (2A) as control and construct pYES2-12Ac (2B). The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3.




Figure 3 A: Fatty acid profile of yeasts transformed with the construct pYES2 as control (Figure 3 A) and pYES2-8Ac (Figure 3 B) and fed with the fatty acid C20:2 $\Delta^{11,14}$ . The respective fatty acids are marked.




Figure 3 B: Fatty acid profile of yeasts transformed with the construct pYES2 as control (Figure 3 A) and pYES2-8Ac (Figure 3 B) and fed with the fatty acid C20:2<sup>Δ11,14</sup>. The respective fatty acids are marked.




Figure 4 A: Fatty acid profile of yeast transformed with the construct pYES2 (Figure 4 A) as control and pYES2-8Ac (Figure 4 B) and fed with the fatty acid C20:3 $\Delta^{11,14,17}$ . The respective fatty acids are marked.






Figure 4 B: Fatty acid profile of yeast transformed with the construct pYES2 (Figure 4 A) as control and pYES2-8Ac (Figure 4 B) and fed with the fatty acid C20:3<sup>A11,14,17</sup>. The respective fatty acids are marked.




Figure 5 A: Comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (Figure 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (Figure 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).




Figure 5 B: Comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (Figure 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (Figure 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).




Figure 6: Expression of AcD8 in double transgenic Arabidopsis




Figure 7 A: Expression of the  $\Delta$ -9-elongase or  $\Delta$ -9-elongase and  $\Delta$ -8-desaturase in transgenic Arabidopsis

7 A) Arabidopsis Wild




Figure 7 B: Expression of the  $\Delta$ -9-elongase or  $\Delta$ -9-elongase and  $\Delta$ -8-desaturase in transgenic Arabidopsis

7 B) Arabidopsis  $\Delta$ 9elo





Figure 7 C: Expression of the  $\Delta$ -9-elongase or  $\Delta$ -9-elongase and  $\Delta$ -8-desaturase in transgenic Arabidopsis

7 C) Arabidopsis  $\Delta 9elo\Delta 8des$



**PROCESS FOR THE PRODUCTION OF  
ARACHIDONIC ACID AND/OR  
EICOSAPENTAENOIC ACID**

CROSS REFERENCE TO RELATED  
APPLICATIONS

This application is a divisional application of U.S. application Ser. No. 12/083,300 filed Apr. 9, 2008, now U.S. Pat. No. 8,017,839 which is a national stage application (under 35 U.S.C. §371) of PCT/EP2006/067223 filed Oct. 10, 2006, which claims benefit to United Kingdom application 0 520 843.4 filed Oct. 13, 2005. The entire contents of each of these applications are hereby incorporated by reference herein in their entirety.

SUBMISSION OF SEQUENCE LISTING

The Sequence Listing associated with this application is filed in electronic format via EFS-Web and hereby incorporated by reference into the specification in its entirety. The name of the text file containing the Sequence Listing is Revised\_Sequence\_List\_17418\_00071\_US. The size of the text file is 120 KB, and the text file was created on Sep. 2, 2011.

DESCRIPTION

The present invention relates to a new process for the production of arachidonic acid and/or eicosapentaenoic acid in plants through the co-expression of a  $\Delta$ -12/ $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase and a  $\Delta$ -5-desaturase and a process for the production of lipids or oils having an increased content of unsaturated fatty acids, in particular  $\omega$ -3 and  $\omega$ -6 fatty acids having at least two double bonds and a 18 or 20 carbon atom chain length. Preferably the arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio.

The invention furthermore relates to the production of a transgenic plants, preferably a transgenic crop plant, having an increased content of arachidonic acid and/or eicosapentaenoic acid, oils or lipids containing  $C_{18}$ - or  $C_{20}$ -fatty acids with a double bond in position  $\Delta$ 5, 8, 9, 11, 12, 14, 15 or 17 of the fatty acid produced, respectively due to the expression of the  $\Delta$ -12/ $\Delta$ -15-desaturase, of the  $\Delta$ -9-elongase, of the  $\Delta$ -8-desaturase and of the  $\Delta$ -5-desaturase in the plant. The expression of the inventive  $\Delta$ -12/ $\Delta$ -15-desaturase leads preferably to linoleic acid and  $\alpha$ -linolenic acid as products having a double bond in the position  $\Delta$ 9, 12 and 15 of the fatty acid.

The invention additionally relates to specific nucleic acid sequences encoding for proteins with  $\Delta$ -12/ $\Delta$ -15-desaturase-,  $\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- or  $\Delta$ -5-desaturase-activity, nucleic acid constructs, vectors and transgenic plants containing said nucleic acid sequences.

Plants and especially oil crops have been used for centuries as sources for edible and non-edible products. There are written records and archaeological excavations that oil crops such as linseed, olive and sesame were widespread use at least six thousand years ago.

Non-edible products of oilseed crops such as rapeseed were used and included in lubricants, oil lamps, and cosmetics such as soaps. Oil crops differ in their cultural, economic and utilization characteristics, for example rapeseed and linseed are adapted to relatively cool climates, whereas oil palm and coconut are adapted to warm and damp climates. Some plants are a real oilseed plant that means the main product of such plants is the oil, whereas in case of others such as cotton

or soybean the oil is more or less a side product. The oils of different plants are basically characterized by their individual fatty acid pattern.

Fatty acids and triglycerides have numerous applications in the food industry, animal nutrition, cosmetics and in the drug sector. Depending on whether they are free saturated or unsaturated fatty acids or triglycerides with an increased content of saturated or unsaturated fatty acids, they are suitable for the most varied applications; thus, for example, long chain polyunsaturated fatty acids (=LCPUFAs) are added to infant formula to increase its nutritional value. The various fatty acids and triglycerides are mainly obtained from microorganisms such as *Mortierella* or from oil-producing plants such as soybean, oilseed rape, sunflower and others, where they are usually obtained in the form of their triacylglycerides. Alternatively, they are obtained advantageously from animals, such as fish. The free fatty acids are prepared advantageously by hydrolysis.

Whether oils with unsaturated or with saturated fatty acids are preferred depends on the intended purpose; thus, for example, lipids with unsaturated fatty acids, specifically polyunsaturated fatty acids, are preferred in human nutrition since they have a positive effect on the cholesterol level in the blood and thus on the possibility of heart disease. They are used in a variety of dietetic foodstuffs or medicaments. In addition PUFAs are commonly used in food, feed and in the cosmetic industry. Poly unsaturated  $\omega$ -3- and/or  $\omega$ -6-fatty acids are an important part of animal feed and human food. Because of the common composition of human food polyunsaturated  $\omega$ -3-fatty acids, which are an essential component of fish oil, should be added to the food to increase the nutritional value of the food; thus, for example, polyunsaturated fatty acids such as Docosahexaenoic acid (=DHA,  $C_{22:6}^{\Delta 4,7,10,13,16,19}$ ) or Eicosapentaenoic acid (=EPA,  $C_{20:5}^{\Delta 5,8,11,14,17}$ ) are added as mentioned above to infant formula to increase its nutritional value. Whereas DHA has a positive effect on the brain development of babies. The addition of poly unsaturated  $\omega$ -3-fatty acids is preferred as the addition of poly unsaturated  $\omega$ -6-fatty acids like Arachidonic acid (=ARA,  $C_{20:4}^{\Delta 5,8,11,14}$ ) to common food have an undesired effect for example on rheumatic diseases such as rheumatoid arthritis. Poly unsaturated  $\omega$ -3- and  $\omega$ -6-fatty acids are precursor of a family of paracrine hormones called eicosanoids such as prostaglandins which are products of the metabolism of Dihomo- $\gamma$ -linoleic acid, ARA or EPA. Eicosanoids are involved in the regulation of lipolysis, the initiation of inflammatory responses, the regulation of blood circulation and pressure and other central functions of the body. Eicosanoids comprise prostaglandins, leukotrienes, thromboxanes, and prostacyclins.  $\omega$ -3-fatty acids seem to prevent arteriosclerosis and cardiovascular diseases primarily by regulating the levels of different eicosanoids. Other Eicosanoids are the thromboxanes and leukotrienes, which are products of the metabolism of ARA or EPA.

Principally microorganisms such as *Mortierella* or oil producing plants such as soybean, rapeseed or sunflower or algae such as *Cryptocodinium* or *Phaeodactylum* are a common source for oils containing PUFAs, where they are usually obtained in the form of their triacyl glycerides. Alternatively, they are obtained advantageously from animals, such as fish. The free fatty acids are prepared advantageously by hydrolysis with a strong base such as potassium or sodium hydroxide.

Plant oils are in general rich in fatty acids such as monounsaturated fatty acids like oleic acid or poly unsaturated fatty acids (=PUFA) like linoleic or linolenic acid. LCPUFAs like arachidonic acid or eicosapentaenoic acid are rarely found in plants exceptions are some *Nephelium* and *Salvia* species in



which arachidonic acid is found and some *Santalum* species in which eicosapentaenoic acid is found. The LCPUFA Docosahexaenoic acid is not found in plants. LCPUFAs such as DHA, EPA, ARA, Dihomo- $\gamma$ -linoleic acid ( $C_{20:3}^{\Delta 8,11,14}$ ) or Docosapentaenoic acid (=DPA,  $C_{22:5}^{\Delta 7,10,13,16,19}$ ) are not produced by oil producing plants such as soybean, rapeseed, safflower or sunflower. A natural sources for said fatty acids are fish for example herring, salmon, sardine, redfish, eel, carp, trout, halibut, mackerel, pikeperch, tuna or algae.

Approximately 80% of the oils and fats are used in the food industry. Nearly about 84% of all world wide used vegetable oils are stemming from only six crops/oil crops, which are soybean, oil palm, rapeseed, sunflower, cottonseed, and groundnut.

On account of their positive properties there has been no shortage of attempts in the past to make available genes which participate in the synthesis of fatty acids or triglycerides for the production of oils in various organisms having a modified content of unsaturated fatty acids. Thus, in WO 91/13972 and its US equivalent a  $\Delta$ -9-desaturase is described. In WO 93/11245 a  $\Delta$ -15-desaturase and in WO 94/11516 a  $\Delta$ -12-desaturase is claimed. WO 00/34439 discloses a  $\Delta$ -5- and a  $\Delta$ -8-desaturase. Other desaturases are described, for example, in EP-A-0 550 162, WO 94/18337, WO 97/30582, WO 97/21340, WO 95/18222, EP-A-0 794 250, Stukey et al., J. Biol. Chem., 265, 1990: 20144-20149, Wada et al., Nature 347, 1990: 200-203 or Huang et al., Lipids 34, 1999: 649-659. To date, however, the various desaturases have been only inadequately characterized biochemically since the enzymes in the form of membrane-bound proteins are isolable and characterizable only with very great difficulty (McKeon et al., Methods in Enzymol. 71, 1981: 275-277, Wang et al., Plant Physiol. Biochem., 26, 1988: 777-792). Generally, membrane-bound desaturases are characterized by introduction into a suitable organism, which is then investigated for enzyme activity by means of analysis of starting materials and products.  $\Delta$ -6-Desaturases are described in WO 93/06712, U.S. Pat. No. 5,614,393, U.S. Pat. No. 5,614,393, WO 96/21022, WO0021557 and WO 99/27111 and their application to production in transgenic organisms is also described, e.g. in WO 9846763, WO 9846764 and WO 9846765. At the same time the expression of various fatty acid biosynthesis genes, as in WO 9964616 or WO 9846776, and the formation of poly-unsaturated fatty acids is also described and claimed. With regard to the effectiveness of the expression of desaturases and their effect on the formation of polyunsaturated fatty acids it may be noted that through expression of a desaturases and elongases as described to date only low contents of poly-unsaturated fatty acids/lipids, such as by way of example eicosapentaenoic or arachidonic acid, have been achieved. Therefore, an alternative and more effective pathway with higher product yield is desirable.

Accordingly, there is still a great demand for new and more suitable genes, which encode enzymes, which participate in the biosynthesis of unsaturated fatty acids and make it possible to produce certain fatty acids specifically on an industrial scale without unwanted byproducts forming. In the selection of genes for biosynthesis two characteristics above all are particularly important. On the one hand, there is as ever a need for improved processes for obtaining the highest possible contents of polyunsaturated fatty acids. Advantageously genes should be as selective as possible and should if possible have more than one activity in the fatty acid biosynthesis chain.

Accordingly, it is an object of the present invention to provide further genes of desaturase and elongase enzymes for the synthesis of polyunsaturated fatty acids in plants prefer-

ably in oilseed plants and to use them in a commercial process for the production of PUFAs especially LCPUFAs. Said process should increase LCPUFA content in plants as much as possible preferably in seeds of an oil producing plant.

#### BRIEF SUMMARY OF THE INVENTION

We have found that a process for the production of arachidonic acid or eicosapentaenoic acid achieves this object or arachidonic acid and eicosapentaenoic acid in transgenic plants that produces mature seeds with a content of at least 1% by weight of said compounds referred to the total lipid content of said organism, which comprises the following steps:

- a) introduction of at least one nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity, and
- b) introduction of at least one second nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -9-elongase activity, and
- c) introduction of at least one third nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -8-desaturase activity, and
- d) introduction of at least a one fourth nucleic acid sequence, which encodes a polypeptide having a  $\Delta$ -5-desaturase activity, and
- e) cultivating and harvesting of said transgenic plant.

According to the invention the used nucleic acid sequences are isolated nucleic sequences coding for polypeptides having a  $\Delta$ -12-desaturase- and  $\Delta$ -15-desaturase-,  $\Delta$ 9-elongase-,  $\Delta$ -8 desaturase- or  $\Delta$ 5-desaturase-activity.

Advantageously nucleic acid sequences are used in the abovementioned process of the invention, which encode polypeptides having  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity,  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase or  $\Delta$ -5-desaturase activity and which are selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO:13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23, and
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 according to the degeneracy of the genetic code,
- c) derivatives of the nucleic acid sequences depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which encode polypeptides having at least 50% homology to the sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity,  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase or  $\Delta$ -5-desaturase activity.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1: Biosynthesis pathway to ARA and/or EPA.  
 FIGS. 2A and 2B: Comparison of the fatty acid profile of yeast transformed with the constructs pYES2 (FIG. 2A) as control and construct pYES2-12Ac (FIG. 2B). The fatty acids

are marked. The new fatty acids synthesized are, in the case of construct pYES2-12Ac (FIG. 2B), the fatty acids C16:2, C16:3, C18:2 and C18:3.

FIGS. 3A and 3B: Fatty acid profile of yeasts transformed with the constructs pYES2 as control (FIG. 3A) and construct pYES2-8Ac (FIG. 3B) and fed with the fatty acid C20:2<sup>011,14</sup>. The respective fatty acids are marked.

FIGS. 4A and 4B: Fatty acid profile of yeast transformed with the constructs pYES2 as control (FIG. 4A) and pYES2-8Ac (FIG. 4B) and fed with the fatty acid C20:3<sup>011,14,17</sup>. The respective fatty acids are marked.

FIGS. 5A and 5B: Comparison of the fatty acid profile of yeasts transformed with the constructs pYES2 as control (FIG. 5A) and pYES2-5Pm (FIG. 5B) and fed with the fatty acid C20:3n-6. The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).

FIG. 6: Expression of AcD8 in double transgenic *Arabidopsis*.

FIGS. 7A-7C: Expression of the  $\Delta$ -9-elongase or  $\Delta$ -9-elongase and  $\Delta$ -8-desaturase in transgenic *Arabidopsis*.

#### DETAILED DESCRIPTION OF THE INVENTION

In the inventive process the nucleic acid sequence encoding the bifunctional  $\Delta$ -12-desaturase- and  $\Delta$ -15-desaturase-enzyme leads to an increased flux from oleic acid (C18:1 $\Delta$ 9) to linolenic acid (C18:3<sup>A9,12,15</sup>) and thereby to an increase of  $\omega$ -3-fatty acids in comparison to the  $\omega$ -6-fatty acids. Furthermore this bifunctional enzyme acts on C16-fatty acids having one double bond in the fatty acid molecule as well as on C18-fatty acids having one double bond in the fatty acid molecule. This leads to a further increase in flux from precursor fatty acids such as C18 fatty acids such as oleic acid towards C18 fatty acids such as linoleic and linolenic acid. This is especially of advantage in plants such as oilseed plants having a high content of oleic acid like such as those from the family of the Brassicaceae, such as the genus *Brassica*, for example oilseed rape or canola; the family of the Elaeagnaceae, such as the genus *Elaeagnus*, for example the genus and species *Olea europaea*, or the family Fabaceae, such as the genus *Glycine*, for example the genus and species *Glycine max*, which are high in oleic acid. But also in other plants such oilseed plants like *Brassica juncea*, *Camelina sativa*, sunflower or safflower and all other plants mentioned herein this leads to a higher amount of  $\omega$ -3-fatty acids. By using said inventive nucleic acid sequence and the activity of its gene product  $\omega$ -3-fatty acids to the  $\omega$ -6-fatty acids are produced in at least a 1:2 ratio, preferably in at least a 1:3 or 1:4 ratio, more preferably in at least a 1:5 or 1:6 ratio. That means especially arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio, preferably in at least a 1:3 or 1:4 ratio, more preferably in at least a 1:5 or 1:6 ratio.

In particular  $\omega$ -3-fatty acids or  $\omega$ -6-fatty acids molecules are produced in the inventive process, arachidonic acid and eicosapentaenoic acid are most preferred produced. We have found that this object is advantageously achieved by the combined expression of four isolated nucleic acid sequences according to the invention which encode for polypeptides having the following activities: a polypeptide with  $\Delta$ -12-desaturase- and  $\Delta$ -15-desaturase-activity, a polypeptide with a C18- $\Delta$ -9-elongase-activity, a polypeptide with C20- $\Delta$ -8-desaturase-activity and a C20- $\Delta$ -5-desaturase-activity. This objective was achieved in particular by the co-expression of the isolated nucleic acid sequences according to the invention. C18 fatty acids with a single double bond in  $\Delta$ -9-position are desaturated a first time to linoleic acid by the  $\Delta$ -12-de-

saturase and  $\Delta$ -15-desaturase and thereafter a second time to linolenic acid by the same enzyme advantageously used in the inventive process. The produced C18 fatty acids linoleic and linolenic acid both having a double bond in  $\Delta$ -9-position are then elongated by the  $\Delta$ -9-elongase, which is advantageously used in the inventive process. By the  $\Delta$ -8-desaturase used in the process a double bond in  $\Delta$ -8-position is introduced into C20 fatty acids. In addition a double bond is introduced into the produced fatty acid molecules in  $\Delta$ -5-position by the  $\Delta$ -5-desaturase. The end products of the whole enzymatic reaction are arachidonic acid and eicosapentaenoic acid.

The  $\omega$ -3-fatty acids or  $\omega$ -6-fatty acids, preferably  $\omega$ -3-fatty acids produced in the process are advantageously bound in membrane lipids and/or triacylglycerides or mixtures of different glycerides, but may also occur in the plants as free fatty acids or else bound in the form of other fatty acid esters.

The fatty acid esters with  $\omega$ -3-fatty acids or  $\omega$ -6-fatty acids especially arachidonic acid and eicosapentaenoic acid molecules can be isolated in the form of an oil or lipid, for example in the form of compounds such as sphingolipids, phosphoglycerides, lipids, glycolipids such as glycosphingolipids, phospholipids such as phosphatidylethanolamine, phosphatidylcholine, phosphatidylserine, phosphatidylglycerol, phosphatidylinositol or diphosphatidylglycerol, monoacylglycerides, diacylglycerides, triacylglycerides or other fatty acid esters such as the acetyl-coenzyme A esters from the plants which have been used for the preparation of the fatty acid esters; preferably, they are isolated in the form of their diacylglycerides, triacylglycerides and/or in the form of phosphatidylcholine, especially preferably in the form of the triacylglycerides. In addition to these esters, the LCPUFAs are also present in the plants, advantageously in the oilseed plants as free fatty acids or bound in other compounds. As a rule, the various abovementioned compounds (fatty acid esters and free fatty acids) are present in the plants with an approximate distribution of 80 to 90% by weight of triglycerides, 2 to 5% by weight of diglycerides, 5 to 10% by weight of monoglycerides, 1 to 5% by weight of free fatty acids, 2 to 8% by weight of phospholipids, the total of the various compounds amounting to 100% by weight.

In the inventive process(es) [the singular shall include the plural and vice versa] the LCPUFAs are produced in a content of at least 1% by weight, preferably at least 2, 3, 4 or 5% by weight, more preferably at least 6, 7, 8, or 9% by weight, most preferably 10, 20 or 30% by weight referred to the total lipid content of the plant used in the process. That means Arachidonic acid and eicosapentaenoic acid are produced in a content of at least 1% by weight, preferably at least 2, 3, 4 or 5% by weight, more preferably at least 6, 7, 8, or 9% by weight, most preferably 10, 20 or 30% by weight referred to the total lipid content. Preferred starting material for the inventive process is oleic acid (C18:1), which is transformed to the preferred end products ARA or EPA. As for the inventive process plants are used the product of the process is not a product of one pure substance per se. It is a mixture of different substances where one or more compounds are the major product and others are only contained as side products. Advantageously the side products shall not exceed 20% by weight referred to the total lipid content of the plant, preferably the side products shall not exceed 15% by weight, more preferably they shall not exceed 10% by weight, most preferably they shall not exceed 5% by weight. In the event that a mixture of different fatty acids such as ARA and EPA are the product of the inventive process said fatty acids can be further purified by method known by a person skilled in the art such as distillation, extraction, crystallization at low temperatures, chromatography or a combination of said methods. These

chemically pure fatty acids or fatty acid compositions are advantageous for applications in the food industry sector, the cosmetic sector and especially the pharmacological industry sector.

Fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise 6 to 15% of palmitic acid, 1 to 6% of stearic acid, 7 to 85% of oleic acid, 0.5 to 8% of vaccenic acid, 0.1 to 1% of arachic acid, 7 to 25% of saturated fatty acids, 8 to 85% of monounsaturated fatty acids and 60 to 85% of polyunsaturated fatty acids including LCPUFAs, in each case based on 100% and on the total fatty acid content of the organisms. Advantageous LCPUFAs, which are present in the fatty acid esters or fatty acid mixtures are preferably at least 1%, 2%, 3%, 4% or 5% by weight of arachidonic acid and/or preferably at least 5%, 6%, 7%, 8%, 9% or 10% by weight of eicosapentaenoic acid, based on the total fatty acid content.

Moreover, the fatty acid esters or fatty acid mixtures which have been produced by the process of the invention advantageously comprise fatty acids selected from the group of the fatty acids erucic acid (13-docosaenoic acid), stercularic acid (9,10-methyleneoctadec-9-enoic acid), malvalic acid (8,9-methyleneheptadec-8-enoic acid), chaulmoogric acid (cyclopentenedodecanoic acid), furan fatty acid (9,12-epoxyoctadeca-9,11-dienoic acid), vernolic acid (9,10-epoxyoctadec-12-enoic acid), tariric acid (6-octadecynoic acid), 6-nonadecynoic acid, santalbic acid (11-octadecen-9-ynoic acid), 6,9-octadecenynoic acid, pyrulic acid (10-heptadecen-8-ynoic acid), crepenynic acid (9-octadecen-12-ynoic acid), 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diynoic acid, petroselenic acid (cis-6-octadecenoic acid), 9c,12t-octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid), catalpic acid (9t11t3c-octadecatrienoic acid), eleostearic acid (9c11t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), puniolic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid), pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or coriolic acid (13-hydroxy-9c,11t-octadecadienoic acid). The abovementioned fatty acids are, as a rule, advantageously only found in traces in the fatty acid esters or fatty acid mixtures produced by the process according to the invention, that is to say that, based on the total fatty acids, they occur to less than 30%, preferably to less than 25%, 24%, 23%, 22% or 21%, especially preferably to less than 20%, 15%, 10%, 9%, 8%, 7%, 6% or 5%, very especially preferably to less than 4%, 3%, 2% or 1%. In a further preferred form of the invention, these abovementioned fatty acids occur to less than 0.9%, 0.8%, 0.7%, 0.6% or 0.5%, especially preferably to less than 0.4%, 0.3%, 0.2%, 0.1%, based on the total fatty acids. The fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise less than 0.1%, based on the total fatty acids, and/or no butyric acid, no cholesterol, no clupanodonic acid (=docosapentaenoic acid, C22:5<sup>A4,8,12,15,21</sup>) and no nisinic acid (tetracosahexaenoic acid, C23:6<sup>A3,8,12,15,21</sup>).

The isolated nucleic acid sequences used in the process according to the invention encode proteins or parts of these, where the proteins or the individual protein or parts thereof comprise(s) an amino acid sequence with sufficient homology to an amino acid sequence which is shown in the sequences SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 and SEQ ID NO: 24 so that the proteins or parts thereof retain a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase-,

$\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- and/or  $\Delta$ -5-desaturase activity. The proteins or parts thereof which is/are encoded by the nucleic acid molecule(s) preferably retains their essential enzymatic activity and the ability of participating in the metabolism of compounds required for the synthesis of cell membranes or lipid bodies in organisms, advantageously in plants, or in the transport of molecules across these membranes. Advantageously, the proteins encoded by the nucleic acid molecules have at least approximately 50%, preferably at least approximately 60% and more preferably at least approximately 70%, 80% or 90% and most preferably at least approximately 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity with the amino acid sequences shown in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 and SEQ ID NO: 24. For the purposes of the invention, homology or homologous is understood as meaning identity or identical, respectively.

The homology was calculated over the entire amino acid or nucleic acid sequence region. The skilled worker has available a series of programs which are based on various algorithms for the comparison of various sequences. Here, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981)), which are part of the GCG software packet [Genetics Computer Group, 575 Science Drive, Madison, Wis., USA 53711 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program GAP and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for the sequence alignments.

Moreover, in the process of the invention advantageously nucleic acid sequences are used which differ from one of the nucleotide sequences shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23 (and parts thereof) owing to the degeneracy of the genetic code and which thus encode the same  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase as those encoded by the nucleotide sequences shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23.

Suitable plants for the production in the process according to the invention are, in principle all plants that produces mature seeds especially crop plants such as oilseed plants.

Plants which are suitable are, in principle, all those plants which are capable of synthesizing fatty acids and that produce mature seeds, such as all dicotyledonous or monocotyledonous plants. Advantageous plants are selected from the group consisting of the plant families Anacardiaceae, Asteraceae, Apiaceae, Boraginaceae, Brassicaceae, Cannabaceae, Elaeagnaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Leguminosae, Linaceae, Lythraeae, Malvaceae, Onagraceae, Palmae, Poaceae, Rubiaceae, Scrophulariaceae, Solanaceae, Sterculiaceae and Theaceae or vegetable plants or ornamentals. More preferred plants are selected from the group consisting of the plant genera of

*Pistacia*, *Mangifera*, *Anacardium*, *Calendula*, *Carthamus*, *Centaurea*, *Cichorium*, *Cynara*, *Helianthus*, *Lactuca*, *Locusta*, *Tagetes*, *Valeriana*, *Borago*, *Daucus*, *Brassica*, *Camelina*, *Melanosinapis*, *Sinapis*, *Arabadopsis*, *Orychophragmus*, *Cannabis*, *Elaeagnus*, *Manihot*, *Janipha*, *Jatropha*, *Ricinus*, *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicajo*, *Glycine*, *Dolichos*, *Phaseolus*, *Pelargonium*, *Cocos*, *Oleum*, *Juglans*, *Wallia*, *Arachis*, *Linum*, *Punica*, *Gossypium*, *Camissonia*, *Oenothera*, *Elaeis*, *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea*, *Triticum*, *Coffea*, *Verbascum*, *Capsicum*, *Nicotiana*, *Solanum*, *Lycopersicon*, *Theobroma* and *Camellia*.

Examples which may be mentioned are the following plants selected from the group consisting of Anacardiaceae such as the genera *Pistacia*, *Mangifera*, *Anacardium*, for example the genus and species *Pistacia vera* [pistachio], *Mangifer indica* [mango] or *Anacardium occidentale* [cashew], Asteraceae, such as the genera *Calendula*, *Carthamus*, *Centaurea*, *Cichorium*, *Cynara*, *Helianthus*, *Lactuca*, *Locusta*, *Tagetes*, *Valeriana*, for example the genus and species *Calendula officinalis* [common marigold], *Carthamus tinctorius* [safflower], *Centaurea cyanus* [cornflower], *Cichorium intybus* [chicory], *Cynara scolymus* [artichoke], *Helianthus annuus* [sunflower], *Lactuca sativa*, *Lactuca crispa*, *Lactuca esculenta*, *Lactuca scariola* L. ssp. *sativa*, *Lactuca scariola* L. var. *integrata*, *Lactuca scariola* L. var. *integrifolia*, *Lactuca sativa* subsp. *romana*, *Locusta communis*, *Valeriana locusta* [salad vegetables], *Tagetes lucida*, *Tagetes erecta* or *Tagetes tenuifolia* [african or french marigold], Apiaceae, such as the genus *Daucus*, for example the genus and species *Daucus carota* [carrot], Boraginaceae, such as the genus *Borago*, for example the genus and species *Borago officinalis* [borage], Brassicaceae, such as the genera *Brassica*, *Camelina*, *Melanosinapis*, *Sinapis*, *Arabadopsis*, for example the genera and species *Brassica napus*, *Brassica rapa* ssp. [oilseed rape], *Sinapis arvensis* *Brassica juncea*, *Brassica juncea* var. *juncea*, *Brassica juncea* var. *crispifolia*, *Brassica juncea* var. *foliosa*, *Brassica nigra*, *Brassica sinapioides*, *Camelina sativa*, *Melanosinapis communis* [mustard], *Brassica oleracea* [fodder beet] or *Arabidopsis thaliana*, Cannabaceae, such as the genus *Cannabis*, such as the genus and species *Cannabis sativa* [hemp], Elaeagnaceae, such as the genus *Elaeagnus*, for example the genus and species *Olea europaea* [olive], Euphorbiaceae, such as the genera *Manihot*, *Janipha*, *Jatropha*, *Ricinus*, for example the genera and species *Manihot utilissima*, *Janipha manihot*, *Jatropha manihot*, *Manihot aipil*, *Manihot dulcis*, *Manihot manihot*, *Manihot melanobasis*, *Manihot esculenta* [cassava] or *Ricinus communis* [castor-oil plant], Fabaceae, such as the genera *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicajo*, *Glycine*, *Dolichos*, *Phaseolus*, soybean, for example the genera and species *Pisum sativum*, *Pisum arvense*, *Pisum humile* [pea], *Albizia berteriana*, *Albizia julibrissin*, *Albizia lebeck*, *Acacia berteriana*, *Acacia littoralis*, *Albizia berteriana*, *Albizia berteriana*, *Cathormion berteriana*, *Feuillea berteriana*, *Inga fragrans*, *Pithecolobium berterianum*, *Pithecolobium fragrans*, *Pithecolobium berterianum*, *Pseudalbizia berteriana*, *Acacia julibrissin*, *Acacia nemu*, *Albizia nemu*, *Feuillea julibrissin*, *Mimosa julibrissin*, *Mimosa speciosa*, *Sericandra julibrissin*, *Acacia lebeck*, *Acacia macrophylla*, *Albizia lebeck*, *Feuillea lebeck*, *Mimosa lebeck*, *Mimosa speciosa*, *Medicago sativa*, *Medicago falcata*, *Medicago varia* [alfalfa] *Glycine max* *Dolichos soja*, *Glycine gracilis*, *Glycine hispida*, *Phaseolus max*, *Soja hispida* or *Soja max* [soybean], Geraniaceae, such as the genera *Pelargonium*, *Cocos*, *Oleum*,

for example the genera and species *Cocos nucifera*, *Pelargonium grossularioides* or *Oleum cocois* [coconut], Gramineae, such as the genus *Saccharum*, for example the genus and species *Saccharum officinarum*, Juglandaceae, such as the genera *Juglans*, *Wallia*, for example the genera and species *Juglans regia*, *Juglans ailanthifolia*, *Juglans sieboldiana*, *Juglans cinerea*, *Wallia cinerea*, *Juglans bixbyi*, *Juglans californica*, *Juglans hindsii*, *Juglans intermedia*, *Juglans jamaicensis*, *Juglans major*, *Juglans microcarpa*, *Juglans nigra* or *Wallia nigra* [walnut], Leguminosae, such as the genus *Arachis*, for example the genus and species *Arachis hypogaea* [peanut], Linaceae, such as the genera *Adenolinum*, for example the genera and species *Linum usitatissimum*, *Linum humile*, *Linum austriacum*, *Linum bienne*, *Linum angustifolium*, *Linum catharticum*, *Linum flavum*, *Linum grandiflorum*, *Adenolinum grandiflorum*, *Linum lewisii*, *Linum narbonense*, *Linum perenne*, *Linum perenne* var. *lewisii*, *Linum pratense* or *Linum trigynum* [linseed], Lythraeae, such as the genus *Punica*, for example the genus and species *Punica granatum* [pomegranate], Malvaceae, such as the genus *Gossypium*, for example the genera and species *Gossypium hirsutum*, *Gossypium arboreum*, *Gossypium barbadense*, *Gossypium herbaceum* or *Gossypium thurberi* [cotton], Onagraceae, such as the genera *Camissonia*, *Oenothera*, for example the genera and species *Oenothera biennis* or *Camissonia brevipes* [evening primrose], Palmae, such as the genus *Elaeis*, for example the genus and species *Elaeis guineensis* [oil palm], Poaceae, such as the genera *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea* (maize), *Triticum*, for example the genera and species *Hordeum vulgare*, *Hordeum jubatum*, *Hordeum murinum*, *Hordeum secalinum*, *Hordeum distichon* *Hordeum aegiceras*, *Hordeum hexastichon*, *Hordeum hexastichum*, *Hordeum irregulare*, *Hordeum sativum*, *Hordeum secalinum* [barley], *Secale cereale* [rye], *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida* [oats], *Sorghum bicolor*, *Sorghum halepense*, *Sorghum saccharatum*, *Sorghum vulgare*, *Andropogon drummondii*, *Holcus bicolor*, *Holcus sorghum*, *Sorghum aethiopicum*, *Sorghum arundinaceum*, *Sorghum caffrorum*, *Sorghum cernuum*, *Sorghum dochna*, *Sorghum drummondii*, *Sorghum durra*, *Sorghum guineense*, *Sorghum lanceolatum*, *Sorghum nervosum*, *Sorghum saccharatum*, *Sorghum subglabrescens*, *Sorghum verticilliflorum*, *Sorghum vulgare*, *Holcus halepensis*, *Sorghum miliaceum*, *Panicum militaceum* [millet], *Oryza sativa*, *Oryza latifolia* [rice], *Zea mays* [maize] *Triticum aestivum*, *Triticum durum*, *Triticum turgidum*, *Triticum hybernum*, *Triticum macha*, *Triticum sativum* or *Triticum vulgare* [wheat], Rubiaceae, such as the genus *Coffea*, for example the genera and species *Coffea* spp., *Coffea arabica*, *Coffea canephora* or *Coffea liberica* [coffee], Scrophulariaceae, such as the genus *Verbascum*, for example the genera and species *Verbascum blattaria*, *Verbascum chaixii*, *Verbascum densiflorum*, *Verbascum lagurus*, *Verbascum longifolium*, *Verbascum lychnitis*, *Verbascum nigrum*, *Verbascum olympicum*, *Verbascum phlomoides*, *Verbascum phoenicum*, *Verbascum pulverulentum* or *Verbascum thapsus* [verbascum], Solanaceae, such as the genera *Capsicum*, *Nicotiana*, *Solanum*, *Lycopersicon*, for example the genera and species *Capsicum annuum*, *Capsicum annuum* var. *glabrusculum*, *Capsicum frutescens* [pepper], *Capsicum annuum* [paprika], *Nicotiana tabacum*, *Nicotiana alata*, *Nicotiana attenuata*, *Nicotiana glauca*, *Nicotiana langsdorffii*, *Nicotiana obtusifolia*, *Nicotiana quadrivalvis*, *Nicotiana repanda*, *Nicotiana rustica*, *Nicotiana sylvestris* [tobacco], *Solanum tuberosum* [potato], *Solanum melongena* [eggplant] *Lycopersicon esculentum*, *Lycopersicon lycopersicum*, *Lycopersicon pyriformis*,

*Solanum integrifolium* or *Solanum lycopersicum* [tomato], Sterculiaceae, such as the genus *Theobroma*, for example the genus and species *Theobroma cacao* [cacao] or Theaceae, such as the genus *Camellia*, for example the genus and species *Camellia sinensis* [tea].

Plants which are especially advantageously used in the process according to the invention are plants which belong to the oil-producing plants, that is to say which are used for the production of oil, such as oilseed or oil crop plants which comprise large amounts of lipid compounds, such as peanut, oilseed rape, canola, sunflower, safflower (*Carthamus tinctoria*), poppy, mustard, hemp, castor-oil plant, olive, sesame, *Calendula*, *Punica*, evening primrose, verbascum, thistle, wild roses, hazelnut, almond, *macadamia*, avocado, bay, pumpkin/squash, linseed, soybean, pistachios, borage, trees (oil palm, coconut or walnut) or arable crops such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, *Tagetes*, Solanaceae plants such as potato, tobacco, eggplant and tomato, *Vicia* species, pea, alfalfa or bushy plants (coffee, cacao, tea), *Salix* species, and perennial grasses and fodder crops. Preferred plants according to the invention are oil crop plants such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, *Calendula*, *Punica*, evening primrose, pumpkin/squash, linseed, soybean, borage, trees (oil palm, coconut). Especially preferred are plants which are high in C18:1-, C18:2- and/or C18:3-fatty acids, such as oilseed rape, canola, *Brassica juncea*, *Camelina sativa*, *Orychophragmus*, sunflower, safflower, tobacco, verbascum, sesame, cotton, pumpkin/squash, poppy, evening primrose, walnut, linseed, hemp or thistle. Very especially preferred plants are plants such as rapeseed, canola, safflower, sunflower, poppy, mustard, hemp, evening primrose, walnut, linseed or hemp. Other preferred plants are castor bean, sesame, olive, *calendula*, *punica*, hazel nut, maize, almond, *macadamia*, cotton, avocado, pumpkin, laurel, pistachio, oil palm, peanut, soybean, marigold, coffee, tobacco, cacao and borage

For the production of further  $\omega$ -6- and/or  $\omega$ -3-fatty acids it is advantageously to introduce further nucleic fatty acid sequences, which encode other enzymes of the fatty acids synthesis chain such as preferably  $\Delta$ -5-elongase(s) and/or  $\Delta$ -4-desaturase(s) [for the purposes of the present invention, the plural is understood as comprising the singular and vice versa]. Other Genes of the fatty acid or lipid metabolism, which can be introduced are selected from the group consisting of acyl-CoA dehydrogenase(s), acyl-ACP [=acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyl transferase(s), acyl-CoA:lysophospholipid acyltransferases, fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allenoxide synthases, hydroperoxide lyases or fatty acid elongase(s). Preferred nucleic acid sequences, which can be used in addition in the inventive process, are disclosed in the sequence protocol of WO2005/012316 and in Table 1 of the specification of said application, these sequences are hereby incorporated by reference.

Transgenic plants are to be understood as meaning single plant cells, certain tissues, organs or parts of plants and their cultures on solid media or in liquid culture, parts of plants and entire plants such as plant cell cultures, protoplasts from plants, callus cultures or plant tissues such as leaves, stem, shoots, seeds, flowers, roots, tubers etc. Said transgenic plants can be cultivated for example on solid or liquid culture medium, in soil or in hydroponics. Plants in the sense of the invention also include plant cells and certain tissues, organs

and parts of plants in all their phenotypic forms such as anthers, fibers, root hairs, stalks, embryos, calli, cotyledons, petioles, harvested material, plant tissue, reproductive tissue such as seeds and cell cultures which are derived from the actual transgenic plant and/or can be used for bringing about the transgenic plant. In this context, the seed comprises all parts of the seed such as the seed coats, epidermal cells, seed cells, endosperm or embryonic tissue.

For the purposes of the invention, "transgenic" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette (=gene construct) or a vector comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, gene constructs or vectors as described herein according to the invention, all those constructions brought about by recombinant methods in which either

- a) the nucleic acid sequence according to the invention, or
- b) a genetic control sequence which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or
- c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. The natural genetic environment is understood as meaning the natural genomic or chromosomal locus in the original plant or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, most preferably at least 5000 bp. A naturally occurring expression cassette—for example the naturally occurring combination of the natural promoter of the nucleic acid sequences with the corresponding  $\Delta$ 12-desaturase and  $\Delta$ 15-desaturase-,  $\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- and/or  $\Delta$ 5-desaturase-genes—becomes a transgenic expression cassette when this expression cassette is modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in U.S. Pat. No. 5,565,350 or WO 00/15815.

A transgenic plant for the purposes of the invention is therefore understood as meaning, as above, that the nucleic acids used in the process are not at their natural locus in the genome of a plant, it being possible for the nucleic acids to be expressed homologously or heterologously. However, as mentioned, transgenic also means that, while the nucleic acids according to the invention are at their natural position in the genome of a plant, the sequence has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified. Transgenic is preferably understood as meaning the expression of the nucleic acids according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acids takes place. Preferred transgenic organisms are oilseed crops.

After cultivation transgenic plants which are used in the inventive process can be brought to the market without isolating the  $\omega$ -6- and/or  $\omega$ -3-fatty acids preferably the arachidonic and/or eicosapentaenoic acid. Preferably the  $\omega$ -6- and/or  $\omega$ -3-fatty acids are isolated from the plant in the form of their free fatty acids, their lipids or oils. The purification can be done by conventional methods such as squeezing and extraction of the plants or other methods instead of the extraction such as distillation, crystallization at low temperatures,

chromatography or a combination of said methods. Advantageously the plants are grinded, heated and/or vaporized before the squeezing and extraction procedure. As solvent for the extraction solvents such as hexane or other solvents having a similar extraction behavior are used. The isolated oils are further purified by acidification with for example phosphoric acid. The free fatty acids are produced from said oils or lipids by hydrolysis. Charcoal or diatom earth is used to remove dyes from the fluid. In another preferred embodiment of the inventive process the alkyl ester of the fatty acids are produced from the oils and lipids by transesterification with an enzyme or with conventional chemistry. A preferred method is the production of the alkyl ester in the presence of alcoholates of the corresponding lower alcohols (C1 to C10 alcohols such as methanol, ethanol, propanol, butanol, hexanol etc.) such as methanolate or ethanolate. Therefore as the skilled worker knows the alcohol in the presence of a catalytic amount of a base such as NaOH or KOH is added to the oils or lipids.

In a preferred form of the inventive process the lipids can be obtained in the usual manner after the plants have been grown. To this end, the organisms can first be harvested and then disrupted, or they can be used directly. In the case of plant cells, plant tissue or plant organs, "growing" is understood as meaning, for example, the cultivation on or in a nutrient medium, or of the intact plant on or in a substrate, for example in a hydroponic culture, potting compost or on arable land. It is advantageous to extract the lipids with suitable solvents such as apolar solvents, for example hexane, or polar solvents, for example ethanol, isopropanol, or mixtures such as hexane/isopropanol, phenol/chloroform/isoamyl alcohol, at temperatures between 0° C. and 80° C., preferably between 20° C. and 50° C. As a rule, the biomass is extracted with an excess of solvent, for example with an excess of solvent to biomass of 1:4. The solvent is subsequently removed, for example by distillation. The extraction may also be carried out with supercritical CO<sub>2</sub>. After the extraction, the remainder of the biomass can be removed, for example, by filtration. Standard methods for the extraction of fatty acids from plants and microorganisms are described in Bligh et al. (Can. J. Biochem. Physiol. 37, 1959: 911-917) or Vick et al. (Plant Physiol. 69, 1982: 1103-1108).

The crude oil thus obtained can then be purified further, for example by removing cloudiness by adding polar solvents such as acetone or apolar solvents such as chloroform, followed by filtration or centrifugation. Further purification via columns or other techniques is also possible.

To obtain the free fatty acids from the triglycerides, the latter are hydrolyzed in the customary manner, for example using NaOH or KOH.

In the inventive process oils, lipids and/or free fatty acids or fractions thereof are produced. Said products can be used for the production of feed and food products, cosmetics or pharmaceuticals.

The oils, lipids, LCPUFAs or fatty acid compositions produced according to the inventive process can be used in the manner with which the skilled worker is familiar for mixing with other oils, lipids, fatty acids or fatty acid mixtures of animal origin, such as, for example, fish oils and/or microbial oils such as from *Mortierella* or *Cryptocodium*. These oils, lipids, fatty acids or fatty acid mixtures, which are composed of vegetable, microbial and/or animal constituents, may also be used for the preparation of feedstuffs, foodstuffs, cosmetics or pharmaceuticals.

The term "oil", "lipid" or "fat" is understood as meaning a fatty acid mixture comprising unsaturated, saturated, preferably esterified, fatty acid(s). The oil, lipid, fat, fatty acid and/or fatty acid composition is preferably high in polyunsaturated (PUFA and/or LCPUFA) free and/or, advantageously, esterified fatty acid(s), in particular oleic acid, linoleic acid,  $\alpha$ -linolenic acid, arachidonic acid and/or eicosatetraenoic acid.

Transgenic plants which comprise the LCPUFAs synthesized in the process according to the invention can also advantageously be marketed directly without there being any need for the oils, lipids or fatty acids synthesized to be isolated.

However, the LCPUFAs produced in the process according to the invention can also be isolated from the plants as described above, in the form of their oils, fats, lipids and/or free fatty acids. Polyunsaturated fatty acids produced by this process can be obtained by harvesting the crop in which they grow, or from the field. This can be done via pressing or extraction of the plant parts, preferably the plant seeds. In this context, the oils, fats, lipids and/or free fatty acids can be obtained by what is known as cold-beating or cold-pressing without applying heat. To allow for greater ease of disruption of the plant parts, specifically the seeds, they are previously comminuted, steamed or roasted. The seeds, which have been pretreated in this manner can subsequently be pressed or extracted with solvents such as warm hexane. The solvent is subsequently removed. In the case of microorganisms, the latter are, after harvesting, for example extracted directly without further processing steps or else, after disruption, extracted via various methods with which the skilled worker is familiar. In this manner, more than 96% of the compounds produced in the process can be isolated. Thereafter, the resulting products are processed further, i.e. refined. In this process, substances such as the plant mucilages and suspended matter are first removed. What is known as desliming can be effected enzymatically or, for example, chemico-physically by addition of acid such as phosphoric acid. Thereafter, the free fatty acids are removed by treatment with a base, for example sodium hydroxide solution. The resulting product is washed thoroughly with water to remove the alkali remaining in the product and then dried. To remove the pigment remaining in the product, the products are subjected to bleaching, for example using filler's earth or active charcoal. At the end, the product is deodorized, for example using steam.

The preferred biosynthesis site of the fatty acids, oils, lipids or fats in the plants which are advantageously used is, for example, in general the seed or cell strata of the seed, so that seed-specific expression of the nucleic acids used in the process makes sense. However, it is obvious that the biosynthesis of fatty acids, oils or lipids need not be limited to the seed tissue, but can also take place in a tissue-specific manner in all the other parts of the plant, for example in epidermal cells or in the tubers.

In principle, the LCPUFAs produced by the process according to the invention in the organisms used in the process can be increased in two different ways. Advantageously, the pool of free polyunsaturated fatty acids and/or the content of the esterified polyunsaturated fatty acids produced via the process can be enlarged. Advantageously, the pool of esterified polyunsaturated fatty acids in the transgenic plants is enlarged by the process according to the invention.

In principle all nucleic acids encoding polypeptides with  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase and/or  $\Delta$ -5-desaturase activity can be used in the inventive process. Preferably the nucleic acid sequences can be isolated for example from microorganism or plants such as fungi like *Mortierella*, algae like *Euglena*, *Cryptocodium* or *Isochrysis*, diatoms like

*Phaeodactylum*, protozoa like amoeba such as *Acanthamoeba* or *Perkinsus* or mosses like *Physcomitrella* or *Ceratodon*, but also non-human animals such as *Caenorhabditis* are possible as source for the nucleic acid sequences. Advantageous nucleic acid sequences according to the invention which encode polypeptides having a  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase and/or  $\Delta$ -5-desaturase activity are originate from microorganisms or plants, advantageously *Phaeodactylum tricorutum*, *Ceratodon purpureus*, *Physcomitrella patens*, *Euglena gracilis*, *Acanthamoeba castellanii*, *Perkinsus marinus* or *Isochrysis galbana*. Thus, the co expression of a C18-specific  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase, a C18-specific  $\Delta$ -9 elongase, a C20-specific  $\Delta$ -8-desaturase and a C20-specific  $\Delta$ -5-desaturase leads to the formation of Arachidonic acid (C20:6n-4,  $\Delta$ 5, 8, 11, 14) and/or Eicosapentaenoic acid (C20:3n-5,  $\Delta$ 5, 8, 11, 14, 17). Most preferred are the sequences mentioned in the sequence protocol.

In another embodiment the invention furthermore relates to isolated nucleic acid sequences encoding polypeptides with  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase-,  $\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- and/or  $\Delta$ -5-desaturase-activity.

In one embodiment the invention relates to an isolated nucleic acid sequence which encodes a polypeptide having a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 22 which encode polypeptides having at least 40% homology to the sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity.

This inventive  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase is able to desaturate C16-fatty acids having at least one double bond in the fatty acid chain and/or C18-fatty acids having at least one double bond in the fatty acid chain. Preferably C16- and/or C18-fatty acids having only one double bond in the fatty acid chain are desaturated. This activity leads to an increase in flux from precursor fatty acids such as C18-fatty acids towards C18-fatty acids having more than one double bond in the fatty acid chain such as linoleic and/or linolenic acid. C18-fatty acids are more preferred in the reaction than C16-fatty acids. C18-fatty acids are more than doubled preferred.

In another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a  $\Delta$ -9-elongase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 11;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 12;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 11 which encode polypeptides having at least 70% homology to the sequence as depicted in SEQ ID NO: 12 and which polypeptides having  $\Delta$ -9-elongase activity.

In yet another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a  $\Delta$ -8-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 which encode polypeptides having at least 70% homology to the sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8 and which polypeptides having  $\Delta$ -8-desaturase activity.

Further in another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a  $\Delta$ -5-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17 which encode polypeptides having at least 70% homology to the sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18 and which polypeptides having  $\Delta$ -5-desaturase activity.

By derivative(s) of the sequences according to the invention is meant, for example, functional homologues of the polypeptides or enzymes encoded by SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which exhibit the same said specific enzymatic activity. This specific enzymatic activity allows advantageously the synthesis of LCPUFAs of the  $\omega$ -6- and/or  $\omega$ -3-pathway of the fatty acid synthesis chain such as ARA and/or EPA. The said sequences encode enzymes which exhibit  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase-,  $\Delta$ -9-elongase-,  $\Delta$ -8-desaturase- and/or  $\Delta$ -5-desaturase-activity.

The enzyme according to the invention,  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase and/or  $\Delta$ -5-desaturase, advantageously either elongates fatty acid chains with 18 carbon atoms (see SEQ ID NO: 11) or introduces a double bond into fatty acid residues of glycerolipids, free fatty acids or acyl-CoA fatty acids at position C<sub>8</sub>-C<sub>9</sub> (see SEQ ID NO: 3, 5 or 7) or at position C<sub>5</sub>-C<sub>6</sub> (see SEQ ID NO: 15 or 17) or at position C<sub>12</sub>-C<sub>13</sub> and C<sub>15</sub>-C<sub>16</sub> of the fatty acid chain (see SEQ ID NO: 19, 21 or 23).

The inventive nucleic acid molecules, for example a nucleic acid molecule with a nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or of a part thereof can be isolated using molecular-biological standard techniques and the sequence information provided herein. Also, for example a homologous sequence or homologous, conserved sequence regions can be identified at the DNA or amino acid level with the aid of comparative algorithms. They can be used as hybridization probe and standard hybridization techniques (such as, for example, those described in Sambrook et al., Molecular Cloning: A Laboratory Manual. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) for isolating further nucleic acid sequences which can be used in the process. Moreover, a nucleic acid molecule comprising a complete sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ



ID NO: 21 or SEQ ID NO: 23 or a part thereof can be isolated by polymerase chain reaction, where oligonucleotide primers which are used on the basis of this sequence or parts thereof (for example a nucleic acid molecule comprising the complete sequence or part thereof can be isolated by polymerase chain reaction using oligonucleotide primers which have been generated based on this same sequence). For example, mRNA can be isolated from cells (for example by means of the guanidinium thiocyanate extraction method of Chirgwin et al. (1979) *Biochemistry* 18:5294-5299) and cDNA by means of reverse transcriptase (for example Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, Md., or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, Fla.). Synthetic oligonucleotide primers for the amplification by means of polymerase chain reaction can be generated based on one of the sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or with the aid of the amino acid sequences detailed in SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24. A nucleic acid according to the invention can be amplified by standard PCR amplification techniques using cDNA or, alternatively, genomic DNA as template and suitable oligonucleotide primers. The nucleic acid amplified thus can be cloned into a suitable vector and characterized by means of DNA sequence analysis. Oligonucleotides, which correspond to a desaturase nucleotide sequence can be generated by standard synthetic methods, for example using an automatic DNA synthesizer.

Homologs of the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase nucleic acid sequences with the sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 means, for example, allelic variants with at least approximately 50 or 60%, preferably at least approximately 60 or 70%, more preferably at least approximately 70 or 80%, 90% or 95% and even more preferably at least approximately 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity or homology with a nucleotide sequence shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or its homologs, derivatives or analogs or parts thereof. Furthermore, isolated nucleic acid molecules of a nucleotide sequence which hybridize with one of the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or with a part thereof, for example hybridized under stringent conditions. A part thereof is understood as meaning, in accordance with the invention, that at least 25 base pairs (=bp), 50 bp, 75 bp, 100 bp, 125 bp or 150 bp, preferably at least 175 bp, 200 bp, 225 bp, 250 bp, 275 bp or 300 bp, especially preferably 350 bp, 400 bp, 450 bp, 500 bp or more base pairs are used for the hybridization. It is also possible and advantageous to use the full sequence. Allelic variants comprise in particular functional variants which can be obtained by deletion, insertion or substitution of nucleotides from/into the sequence detailed in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, it being intended, however, that the enzyme activity of the resulting proteins which are synthesized is advantageously retained for the insertion of one or more genes. Proteins which retain the enzymatic activity of

the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase, i.e. whose activity is essentially not reduced, means proteins with at least 10%, preferably 20%, especially preferably 30%, very especially preferably 40% of the original enzyme activity in comparison with the protein encoded by SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23. The homology was calculated over the entire amino acid or nucleic acid sequence region. The skilled worker has available a series of programs which are based on various algorithms for the comparison of various sequences. Here, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981))], which are part of the GCG software packet [Genetics Computer Group, 575 Science Drive, Madison, Wis., USA 53711 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program GAP and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for the sequence alignments.

Homologs of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 means for example also bacterial, fungal and plant homologs, truncated sequences, single-stranded DNA or RNA of the coding and noncoding DNA sequence.

Homologs of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 also means derivatives such as, for example, promoter variants. The promoters upstream of the nucleotide sequences detailed can be modified by one or more nucleotide exchanges, by insertion(s) and/or deletion(s) without the functionality or activity of the promoters being adversely affected, however. It is furthermore possible that the modification of the promoter sequence enhances their activity or that they are replaced entirely by more active promoters, including those from heterologous organisms.

In a further embodiment, derivatives of the nucleic acid molecule according to the invention represented in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 encode proteins with at least 40%, advantageously approximately 50 or 60%, advantageously at least approximately 60 or 70% and more preferably at least approximately 70 or 80%, 80 to 90%, 90 to 95% and most preferably at least approximately 96%, 97%, 98%, 99% or more homology (=identity) with a complete amino acid sequence of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24. The homology was calculated over the entire amino acid or nucleic acid sequence region. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981))], which are part of the GCG software packet [Genetics Computer Group, 575 Science Drive, Madison, Wis., USA 53711 (1991)], were used for the sequence alignment. The sequence homology values



which are indicated above as a percentage were determined over the entire sequence region using the program BestFit and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for the sequence alignments.

Moreover, the invention comprises nucleic acid molecules which differ from one of the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 (and parts thereof) owing to the degeneracy of the genetic code and which thus encode the same  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase as those encoded by the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23.

In addition to the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, the skilled worker will recognize that DNA sequence polymorphisms which lead to changes in the amino acid sequences of the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase may exist within a population. These genetic polymorphisms in the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase gene may exist between individuals within a population owing to natural variation. These natural variants usually bring about a variance of 1 to 5% in the nucleotide sequence of the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase gene. Each and every one of these nucleotide variations and resulting amino acid polymorphisms in the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase or  $\Delta$ -5-desaturase which are the result of natural variation and do not modify the functional activity are to be encompassed by the invention.

The nucleic acid sequence(s) according to the invention (for purposes of the application the singular encompasses the plural and vice versa) or fragments thereof may advantageously be used for isolating other genomic sequences via homology screening.

The said derivatives may be isolated, for example, from other organisms, eukaryotic organisms such as plants, especially mosses, algae, dinoflagellates, protozoa or fungi.

Allele variants include in particular functional variants obtainable by deletion, insertion or substitution of nucleotides in the sequences depicted in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 the enzymatic activity of the derived synthesized proteins being retained.

Starting from the DNA sequence described in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or parts of said sequences such DNA sequences can be isolated using, for example, normal hybridization methods or the PCR technique from other eukaryotes such as those identified above for example. These DNA sequences hybridize under standard conditions with the said sequences. For hybridization use is advantageously made of short oligonucleotides of the conserved regions of an average length of about 15 to 70 bp, preferably of about 17 to 60 bp, more preferably of about 19 to 50 bp, most preferably of about 20 to 40 bp, for example, which can be determined by comparisons with other desaturase or elongase genes in the man-

ner known to those skilled in the art. The histidine box sequences are advantageously employed. However, longer fragments of the nucleic acids according to the invention or the complete sequences may also be used for hybridization. Depending on the nucleic acid employed: oligonucleotide, longer fragment or complete sequence, or depending on which type of nucleic acid, DNA or RNA, is used for hybridization these standard conditions vary. Thus, for example, the melting temperatures of DNA:DNA hybrids are approximately 10° C. lower than those of DNA:RNA hybrids of the same length.

By standard conditions is meant, for example, depending on the nucleic acid in question temperatures between 42° C. and 58° C. in an aqueous buffer solution having a concentration of between 0.1 and 5×SSC (1×SSC=0.15 M NaCl, 15 mM sodium citrate, pH 7.2) or additionally in the presence of 50% formamide, such as by way of example 42° C. in 5×SSC, 50% formamide. Hybridization conditions for DNA:DNA hybrids are advantageously 0.1×SSC and temperatures between approximately 20° C. and 45° C., preferably between approximately 30° C. and 45° C. For DNA:RNA hybrids the hybridization conditions are advantageously 0.1×SSC and temperatures between approximately 30° C. and 55° C., preferably between approximately 45° C. and 55° C. These specified temperatures for hybridization are melting temperature values calculated by way of example for a nucleic acid having a length of approximately 100 nucleotides and a G+C content of 50% in the absence of formamide. The experimental conditions for DNA hybridization are described in relevant genetics textbooks such as by way of example Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989, and may be calculated by formulae known to those skilled in the art, for example as a function of the length of the nucleic acids, the nature of the hybrids or the G+C content. Those skilled in the art may draw on the following textbooks for further information on hybridization: Ausubel et al. (eds), 1985, Current Protocols in Molecular Biology, John Wiley & Sons, New York; Hames and Higgins (eds), 1985, Nucleic Acids Hybridization: A Practical Approach, IRL Press at Oxford University Press, Oxford; Brown (ed), 1991, Essential Molecular Biology: A Practical Approach, IRL Press at Oxford University Press, Oxford.

Furthermore, by derivatives is meant homologues of the sequences SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, for example eukaryotic homologues, truncated sequences, single-stranded DNA of the encoding and nonencoding DNA sequence or RNA of the encoding and nonencoding DNA sequence.

In addition, by homologues of the sequences SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 is meant derivatives such as by way of example promoter variants. These variants may be modified by one or more nucleotide exchanges, by insertion(s) and/or deletion(s) without, however, adversely affecting the functionality or efficiency of the promoters. Furthermore, the promoters can have their efficiency increased by altering their sequence or be completely replaced by more effective promoters even of foreign organisms.

By derivatives is also advantageously meant variants whose nucleotide sequence has been altered in the region from -1 to -2000 ahead of the start codon in such a way that the gene expression and/or the protein expression is modified, preferably increased. Furthermore, by derivatives is also meant variants, which have been modified at the 3' end.

The nucleic acid sequences according to the invention which encode a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase, a  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or a  $\Delta$ -5-desaturase may be produced by synthesis or obtained naturally or contain a mixture of synthetic and natural DNA components as well as consist of various heterologous  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase gene segments from different organisms. In general, synthetic nucleotide sequences are produced with codons, which are preferred by the corresponding host organisms, plants for example. This usually results in optimum expression of the heterologous gene. These codons preferred by plants may be determined from codons having the highest protein frequency, which are expressed in most of the plant species of interest. An example concerning the bacterium *Corynebacterium glutamicum* is provided in Wada et al. (1992) *Nucleic Acids Res.* 20:2111-2118). Such experiments can be carried out using standard methods and are known to the person skilled in the art.

Functionally equivalent sequences which encode the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase gene are those derivatives of the sequence according to the invention which despite differing nucleotide sequence still possess the desired functions, that is to say the enzymatic activity and specific selectivity of the proteins. That means such functionally equivalent sequences have an biological or enzymatic activity, which is at least 10%, preferably at least 20%, 30%, 40% or 50% especially preferably at least 60%, 70%, 80% or 90% and very especially at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or more of the activity of the proteins/enzymes encoded by the inventive sequences. Thus, functional equivalents include naturally occurring variants of the sequences described herein as well as artificial ones, e.g. artificial nucleotide sequences adapted to the codon use of a plant which have been obtained by chemical synthesis.

In addition, artificial DNA sequences are suitable, provided, as described above, they mediate the desired property, for example an increase in the content of  $\Pi$ -12-,  $\Pi$ -15-,  $\Pi$ -8- and/or  $\Delta$ -5-double bonds in fatty acids and an elongation of C18-fatty acids having a  $\Delta$ -9-double bond in fatty acids, oils or lipids in plants that produce mature seeds preferably in crop plants by over expression of the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase gene. Such artificial DNA sequences can exhibit  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase activity, for example by back-translation of proteins constructed by means of molecular modeling, or be determined by in vitro selection. Possible techniques for in vitro evolution of DNA to modify or improve the DNA sequences are described in Patten, P. A. et al., *Current Opinion in Biotechnology* 8, 724-733 (1997) or in Moore, J. C. et al., *Journal of Molecular Biology* 272, 336-347 (1997). Particularly suitable are encoding DNA sequences which are obtained by back-translation of a polypeptide sequence in accordance with the codon use specific to the host plant. Those skilled in the art familiar with the methods of plant genetics can easily determine the specific codon use by computer analyses of other known genes of the plant to be transformed.

Other suitable equivalent nucleic acid sequences, which may be mentioned are sequences that encode fusion proteins, a component of the fusion protein being a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase polypeptide and/or a  $\Delta$ -9 elongase polypeptide or a functionally equivalent part thereof. The second part of the fusion protein can be, for example, another polypeptide having

enzymatic activity or an antigenic polypeptide sequence by means of which it is possible to demonstrate  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase expression (e.g. myc tag or his tag). Preferably, however, this is a regulatory protein sequence, such as by way of example a signal sequence for the endoplasmic reticulum (=ER) which directs the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase protein and/or the  $\Delta$ -9-elongase protein to the desired point of action, or regulatory sequences which influence the expression of the nucleic acid sequence according to the invention, such as promoters or terminators. In another preferred embodiment the second part of the fusion protein is a plastidial targeting sequence as described by Napier J. A. [Targeting of foreign proteins to the chloroplast, *Methods Mol. Biol.*, 49, 1995: 369-376]. A preferred used vector comprising said plastidial targeting sequence is disclosed by Colin Lazarus [Guerineau F., Woolston S., Brooks L., Mullineaux P. "An expression cassette for targeting foreign proteins into chloroplast; *Nucleic. Acids Res.*, December 9, 16 (23), 1988: 11380].

Advantageously, the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Pi$ -8-desaturase and/or  $\Delta$ -5-desaturase genes in the method according to the invention may be combined with other genes for fatty acid biosynthesis as described above. Examples of such genes are the acyl transferases, other desaturases or elongases such as  $\Delta$ -4-desaturases or  $\omega$ -3- and/or  $\omega$ -6-specific desaturases) and/or such as  $\Delta$ -5-elongases to mention only some of them. For in vivo and especially in vitro synthesis combination with e.g. NADH cytochrome B5 reductases, which can take up or release reduction equivalents is advantageous.

By the amino acid sequences according to the invention is meant proteins which contain an amino acid sequence depicted in the sequences SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 or a sequence obtainable there from by substitution, inversion, insertion or deletion of one or more amino acid groups (such sequences are derivatives of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24), whereas the enzymatic activities of the proteins depicted in SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 being retained or not substantially reduced, that is they still possess the same enzymatic specificity. By "not substantially reduced" or "the same enzymatic activity" is meant all enzymes which still exhibit at least 10%, 20%, 30%, 40% or 50%, preferably at least 60%, 70%, 80% or 90% particularly preferably at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, of the enzymatic activity of the initial enzyme obtained from the wild type source organism such as organisms of the genus *Physcomitrella*, *Ceratodon*, *Borago*, *Thraustochytrium*, *Schizochytrium*, *Phytophthora*, *Mortierella*, *Caenorhabditis*, *Aleuritia*, *Muscarioidides*, *Isochrysis*, *Phaeodactylum*, *Cryptocodinium*, *Acanthamoeba* or *Euglena* preferred source organisms are organisms such as the species *Euglena gracilis*, *Isochrysis galbana*, *Phaeodactylum tricoratum*, *Caenorhabditis elegans*, *Thraustochytrium*, *Phytophthora infestans*, *Ceratodon purpureus*, *Isochrysis galbana*, *Aleuritia farinosa*, *Muscarioidides vialii*, *Mortierella alpina*, *Borago officinalis* or *Physcomitrella patens*. For the estimation of an enzymatic activity, which is "not substantially reduced" or which has the "same enzymatic activity" the enzymatic activity of the derived sequences are determined and compared with the wild type enzyme activities. In doing

this, for example, certain amino acids may be replaced by others having similar physicochemical properties (space filling, basicity, hydrophobicity, etc.). For example, arginine residues are exchanged for lysine residues, valine residues for isoleucine residues or aspartic acid residues for glutamic acid residues. However, one or more amino acids may also be swapped in sequence, added or removed, or a plurality of these measures may be combined with one another.

By derivatives is also meant functional equivalents, which in particular also contain natural or artificial mutations of an originally isolated sequence encoding a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase, a  $\Delta$ -9-elongase, a  $\Delta$ -8-desaturase and/or a  $\Delta$ -5-desaturase, which continue to exhibit the desired function, that is the enzymatic activity and substrate selectivity thereof is not substantially reduced. Mutations comprise substitutions, additions, deletions, exchanges or insertions of one or more nucleotide residues. Thus, for example, the present invention also encompasses those nucleotide sequences, which are obtained by modification of the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase nucleotide sequence, the  $\Delta$ -8-desaturase nucleotide sequence, the  $\Delta$ -5-desaturase nucleotide sequence and/or the  $\Delta$ -9-elongase nucleotide sequence used in the inventive processes. The aim of such a modification may be, e.g., to further bind the encoding sequence contained therein or also, e.g., to insert further restriction enzyme interfaces.

Functional equivalents also include those variants whose function by comparison as described above with the initial gene or gene fragment is weakened (=not substantially reduced) or reinforced (=enzyme activity higher than the activity of the initial enzyme, that is activity is higher than 100%, preferably higher than 110%, 120%, 130%, 140% or 150%, particularly preferably higher than 200% or more).

At the same time the nucleic acid sequence may, for example, advantageously be a DNA or cDNA sequence. Suitable encoding sequences for insertion into an expression cassette according to the invention include by way of example those which encode a  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase, a  $\Delta$ -8-desaturase and/or a  $\Delta$ -5-desaturase with the sequences described above and lend the host the ability to overproduce fatty acids, oils or lipids having double bonds in the  $\Delta$ -12-,  $\Delta$ -15-,  $\Delta$ -8-position and  $\Delta$ -5-position, it being advantageous when at the same time fatty acids having at least four double bonds are produced. These sequences may be of homologous or heterologous origin.

By the gene construct (=nucleic acid construct or fragment or expression cassette) according to the invention is meant the sequences specified in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which result from the genetic code and/or derivatives thereof which are functionally linked with one or more regulation signals advantageously to increase the gene expression and which control the expression of the encoding sequence in the host cell. These regulatory sequences should allow the selective expression of the genes and the protein expression. Depending on the host plant this may mean, for example, that the gene is expressed and/or overexpressed only after induction or that it is expressed and/or overexpressed immediately. Examples of these regulatory sequences are sequences to which inducers or repressors bind and in this way regulate the expression of the nucleic acid. In addition to these new regulation sequences or instead of these sequences the natural regulation of these sequences ahead of the actual structural genes may still be present and optionally have been genetically modified so that natural regulation was switched off and the expression of the genes increased. However, the gene construct can also be built up more simply, that is no additional regulation sig-

nals have been inserted ahead of the nucleic acid sequence or derivatives thereof and the natural promoter with its regulation has not been removed. Instead of this the natural regulation sequence was mutated in such a way that no further regulation ensues and/or the gene expression is heightened. These modified promoters in the form of part sequences (=promoter containing parts of the nucleic acid sequences according to the invention) can also be brought on their own ahead of the natural gene to increase the activity. In addition, the gene construct may advantageously also contain one or more so-called enhancer sequences functionally linked to the promoter which allow enhanced expression of the nucleic acid sequence. At the 3' end of the DNA sequences additional advantageous sequences may also be inserted, such as further regulatory elements or terminators. The SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and/or SEQ ID NO: 23 gene may be present in one or more copies in the gene construct (=expression cassette).

As described above, the regulatory sequences or factors can preferably positively influence and so increase the gene expression of the introduced genes. Thus, reinforcement of the regulatory elements advantageously on the transcription level may be effected by using powerful transcription signals such as promoters and/or enhancers. However, in addition reinforcement of translation is also possible, for example by improving the stability of the mRNA.

Suitable promoters in the expression cassette are in principle all promoters which can control the expression of foreign genes in microorganisms like protozoa such as amoeba, ciliates, algae such as green, brown, red or blue algae such as Euglena, bacteria such as gram-positive or gram-negative bacteria, yeasts such as *Saccharomyces*, *Pichia* or *Schizosaccharomyces* or fungi such as *Mortierella*, *Thraustochytrium* or *Schizochytrium* or plants such as *Aleuritia*, advantageously in plants or fungi. Such microorganisms are generally used to clone the inventive genes and possible other genes of the fatty acid biosynthesis chain for the production of fatty acids according to the inventive process. Use is preferably made in particular of plant promoters or promoters derived from a plant virus. Advantageous regulation sequences for the method according to the invention are found for example in promoters such as *cos*, *tac*, *trp*, *tet*, *trp-tet*, *lpp*, *lac*, *lpp-lac*, *lacI<sup>cr</sup>*, *T7*, *T5*, *T3*, *gal*, *trc*, *ara*, *SP6*,  $\lambda$ -*P<sub>R</sub>* or in  $\lambda$ -*P<sub>L</sub>* promoters which are employed advantageously in gram-negative bacteria. Other advantageous regulation sequences are found, for example, in the gram-positive promoters *amy* and *SPO2*, in the yeast or fungal promoters *ADC1*, *MF $\alpha$* , *AC*, *P-60*, *CYC1*, *GAPDH*, *TEF*, *rp28*, *ADH* or in the plant promoters *CaMV/35S* [Franck et al., *Cell* 21 (1980) 285-294], *SSU*, *OCS*, *lib4*, *STLS1*, *B33*, *nos* (=Nopaline Synthase Promoter) or in the ubiquitin or phaseolin promoter. The expression cassette may also contain a chemically inducible promoter by means of which the expression of the exogenous  $\Delta$ -12- and  $\Delta$ -15-,  $\Delta$ -8- and/or  $\Delta$ -5-desaturase gene and/or the  $\Delta$ -9-elongase gene in the microorganism and/or plant can be controlled advantageously in the plants at a particular time. Advantageous plant promoters of this type are by way of example the *PRP1* promoter [Ward et al., *Plant. Mol. Biol.* 22 (1993), 361-366], a promoter inducible by benzenesulfonamide (EP 388 186), a promoter inducible by tetracycline [Gatz et al., (1992) *Plant J.* 2, 397-404], a promoter inducible by salicylic acid (WO 95/19443), a promoter inducible by abscisic acid (EP 335 528) and a promoter inducible by ethanol or cyclohexanone (WO 93/21334). Other examples of plant promoters, which can advantageously be used are the promoter of cytosolic FBPase from potato, the *ST-LSI* promoter from

potato (Stockhaus et al., EMBO J. 8 (1989) 2445-245), the promoter of phosphoribosyl pyrophosphate amidotransferase from *Glycine max* (see also gene bank accession number U87999) or a nodiene-specific promoter as described in EP 249 676. Particularly advantageous are those plant promoters, which ensure expression in tissues or plant parts/organs in which fatty acid biosynthesis or the precursor stages thereof occurs, as in endosperm or in the developing embryo for example. Particularly noteworthy are advantageous promoters, which ensure seed-specific expression such as by way of example the USP promoter or derivatives thereof, the LEB4 promoter, the phaseolin promoter or the napin promoter. The particularly advantageous USP promoter cited according to the invention or its derivatives mediate very early gene expression in seed development [Baeumlein et al., Mol Gen Genet, 1991, 225 (3): 459-67]. Other advantageous seed-specific promoters which may be used for monocotyledonous or dicotyledonous plants are the promoters suitable for dicotyledons such as napin gene promoters, likewise cited by way of example, from oilseed rape (U.S. Pat. No. 5,608,152), the oleosin promoter from *Arabidopsis* (WO 98/45461), the phaseolin promoter from *Phaseolus vulgaris* (U.S. Pat. No. 5,504,200), the Bce4 promoter from *Brassica* (WO 91/13980) or the leguminous B4 promoter (LeB4, Baeumlein et al., Plant J., 2, 2, 1992: 233-239) or promoters suitable for monocotyledons such as the promoters of the lpt2 or lpt1 gene in barley (WO 95/15389 and WO 95/23230) or the promoters of the barley hordeine gene, the rice glutelin gene, the rice oryzin gene, the rice prolamins gene, the wheat gliadin gene, the white glutelin gene, the corn zein gene, the oats glutelin gene, the sorghum kasirin gene or the rye secalin gene which are described in WO99/16890.

Furthermore, particularly preferred are those promoters, which ensure the expression in tissues or plant parts in which, for example, the biosynthesis of fatty acids, oils and lipids or the precursor stages thereof takes place. Particularly noteworthy are promoters, which ensure a seed-specific expression. Noteworthy are the promoter of the napin gene from oilseed rape (U.S. Pat. No. 5,608,152), the USP promoter from *Vicia faba* (USP=unknown seed protein, Baeumlein et al., Mol Gen Genet, 1991, 225 (3): 459-67), the promoter of the oleosin gene from *Arabidopsis* (WO 98/45461), the phaseolin promoter (U.S. Pat. No. 5,504,200) or the promoter of the legumin B4 gene (LeB4; Baeumlein et al., 1992, Plant Journal, 2 (2): 233-9). Other promoters to be mentioned are that of the lpt2 or lpt1 gene from barley (WO 95/15389 and WO 95/23230), which mediate seed-specific expression in monocotyledonous plants. Other advantageous seed specific promoters are promoters such as the promoters from rice, corn or wheat disclosed in WO 99/16890 or Amy32b, Amy6-6 or aleurain (U.S. Pat. No. 5,677,474), Bce4 (rape, U.S. Pat. No. 5,530,149), glycinin (soy bean, EP 571 741), phosphoenol pyruvate carboxylase (soy bean, JP 06/62870), ADR12-2 (soy bean, WO 98/08962), isocitratlyase (rape, U.S. Pat. No. 5,689,040) or  $\beta$ -amylase (barley, EP 781 849).

As described above, the expression construct (=gene construct, nucleic acid construct) may contain yet other genes, which are to be introduced into the microorganism or plant. These genes can be subject to separate regulation or be subject to the same regulation region as the  $\Delta$ -12- and  $\Delta$ -15-desaturase gene and/or the  $\Delta$ -8- and/or  $\Delta$ -5-desaturase gene and/or the  $\Delta$ -9-elongase gene. These genes are by way of example other biosynthesis genes, advantageously for fatty acid biosynthesis, which allow increased synthesis. Examples which may be mentioned are the genes for example of the  $\Delta$ -9-,  $\Delta$ -4-desaturase,  $\Delta$ -5-elongase,  $\alpha$ -ketoacyl reductases,  $\alpha$ -ketoacyl synthases, elongases or the various

hydroxylases and acyl-ACP thioesterases. The desaturase and elongase genes are advantageously used in the nucleic acid construct.

In principle all natural promoters with their regulation sequences can be used like those named above for the expression cassette according to the invention and the method according to the invention. Over and above this, synthetic promoters may also advantageously be used.

In the preparation of an a gene construct various DNA fragments can be manipulated in order to obtain a nucleotide sequence, which usefully reads in the correct direction and is equipped with a correct reading raster. To connect the DNA fragments (=nucleic acids according to the invention) to one another adaptors or linkers may be attached to the fragments.

The promoter and the terminator regions can usefully be provided in the transcription direction with a linker or polylinker containing one or more restriction points for the insertion of this sequence. Generally, the linker has 1 to 10, mostly 1 to 8, preferably 2 to 6, restriction points. In general the size of the linker inside the regulatory region is less than 100 bp, frequently less than 60 bp, but at least 5 bp. The promoter may be native or homologous as well as foreign or heterologous to the host organism, for example to the host plant. In the 5'-3' transcription direction the expression cassette contains the promoter, a DNA sequence which encodes a  $\Delta$ -12- and  $\Delta$ -15-desaturase gene, a  $\Delta$ -8-desaturase gene, a  $\Delta$ -5-desaturase gene and/or a  $\Delta$ -9-elongase gene and a region for transcription termination. Different termination regions can be exchanged for one another in any desired fashion.

Furthermore, manipulations, which provide suitable restriction interfaces or which remove excess DNA or restriction interfaces can be employed. Where insertions, deletions or substitutions, such as transitions and transversions, come into consideration, in vitro mutagenesis, primer repair, restriction or ligation may be used. In suitable manipulations such as restriction, chewing back or filling of overhangs for blunt ends complementary ends of the fragments can be provided for the ligation.

For an advantageous high expression the attachment of the specific ER retention signal SEKDEL inter alia can be of importance (Schouten, A. et al., Plant Mol. Biol. 30 (1996), 781-792). In this way the average expression level is tripled or even quadrupled. Other retention signals, which occur naturally in plant and animal proteins located in the ER may also be employed for the construction of the cassette. In another preferred embodiment a plastidial targeting sequence is used as described by Napier J. A. [Targeting of foreign proteins to the chloroplast, Methods Mol. Biol., 49, 1995: 369-376]. A preferred used vector comprising said plastidial targeting sequence is disclosed by Colin Lazarus [Guerineau F., Woolston S., Brooks L., Mullineaux P. "An expression cassette for targeting foreign proteins into chloroplast; Nucleic. Acids Res., December 9, 16 (23), 1988: 11380].

Preferred polyadenylation signals are plant polyadenylation signals, preferably those which substantially correspond to T-DNA polyadenylation signals from *Agrobacterium tumefaciens*, in particular gene 3 of the T-DNA (octopin synthase) of the Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3 (1984), 835 et seq.) or corresponding functional equivalents.

An expression cassette/gene construct is produced by fusion of a suitable promoter with a suitable  $\Delta$ -12- and  $\Delta$ -15-desaturase DNA sequence, a suitable  $\Delta$ -8- and/or  $\Delta$ -5-desaturase DNA sequence and/or a suitable  $\Delta$ -9-elongase DNA sequence together with a polyadenylation signal by common recombination and cloning techniques as described, for example, in T. Maniatis, E. F. Fritsch and J. Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Har-

bor Laboratory, Cold Spring Harbor, N.Y. (1989) as well as in T. J. Silhavy, M. L. Berman and L. W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and in Ausubel, F. M. et al., Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley-Interscience (1987).

The DNA sequences encoding the nucleic acid sequences used in the inventive processes such as the  $\Delta$ -12- and  $\Delta$ -15-desaturase from *Acanthamoeba castellanii* or *Perkinsus marinus*,  $\Delta$ -8-desaturase from *Euglena gracilis*, *Acanthamoeba castellanii* or *Perkinsus marinus*, the  $\Delta$ -9-elongase from *Isochrysis galbana* or *Acanthamoeba castellanii* and/or the  $\Delta$ -5-desaturase for example from *Thraustochytrium*, *Acanthamoeba castellanii* or *Perkinsus marinus* or other organisms such as *Caenorhabditis elegans*, *Mortierella alpina*, *Borage officinalis* or *Physcomitrella patens* contain all the sequence characteristics needed to achieve correct localization of the site of fatty acid, lipid or oil biosynthesis. Accordingly, no further targeting sequences are needed per se. However, such localization may be desirable and advantageous and hence artificially modified or reinforced so that such fusion constructs are also a preferred advantageous embodiment of the invention.

Particularly preferred are sequences, which ensure targeting in plastids. Under certain circumstances targeting into other compartments (reported in: Kermode, Crit. Rev. Plant Sci. 15, 4 (1996), 285-423) may also be desirable, e.g. into vacuoles, the mitochondrion, the endoplasmic reticulum (ER), peroxisomes, lipid structures or due to lack of corresponding operative sequences retention in the compartment of origin, the cytosol.

Advantageously, the nucleic acid sequences according to the invention or the gene construct together with at least one reporter gene are cloned into a gene construct, which is introduced into the organism via a vector or directly into the genome. This reporter gene should allow easy detection via a growth, fluorescence, chemical, bioluminescence or resistance assay or via a photometric measurement. Examples of reporter genes which may be mentioned are antibiotic- or herbicide-resistance genes, hydrolase genes, fluorescence protein genes, bioluminescence genes, sugar or nucleotide metabolic genes or biosynthesis genes such as the Ura3 gene, the Ilv2 gene, the luciferase gene, the  $\beta$ -galactosidase gene, the gfp gene, the 2-desoxyglucose-6-phosphate phosphatase gene, the  $\beta$ -glucuronidase gene,  $\beta$ -lactamase gene, the neomycin phosphotransferase gene, the hygromycin phosphotransferase gene or the BASTA (=gluphosinate-resistance) gene. These genes permit easy measurement and quantification of the transcription activity and hence of the expression of the genes. In this way genome positions may be identified which exhibit differing productivity.

In a preferred embodiment an gene construct comprises upstream, i.e. at the 5' end of the encoding sequence, a promoter and downstream, i.e. at the 3' end, a polyadenylation signal and optionally other regulatory elements which are operably linked to the intervening encoding sequence for  $\Delta$ -12- and  $\Delta$ -15-desaturase,  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase and/or  $\Delta$ -5-desaturase DNA sequence. By an operable linkage is meant the sequential arrangement of promoter, encoding sequence, terminator and optionally other regulatory elements in such a way that each of the regulatory elements can fulfill its function in the expression of the encoding sequence in due manner. The sequences preferred for operable linkage are targeting sequences for ensuring subcellular localization in plastids. However, targeting sequences for ensuring subcellular localization in the mitochondrion, in the endoplasmic reticulum (=ER), in the nucleus, in oil corpuscles or other

compartments may also be employed as well as translation promoters such as the 5' lead sequence in tobacco mosaic virus (Gallie et al., Nucl. Acids Res. 15 (1987), 8693-8711).

An expression cassette/gene construct may, for example, contain a constitutive promoter or a tissue-specific promoter (preferably the USP or napin promoter) the gene to be expressed and the ER retention signal. For the ER retention signal the KDEL amino acid sequence (lysine, aspartic acid, glutamic acid, leucine) or the KKX amino acid sequence (lysine-lysine-X-stop, wherein X means every other known amino acid) is preferably employed.

For expression in a prokaryotic or eukaryotic host organism, for example a microorganism such as a fungus or a plant such as an oil crop the expression cassette is advantageously inserted into a vector such as by way of example a plasmid, a phage or other DNA which allows optimum expression of the genes in the host organism. Examples of suitable plasmids are: in *E. coli* pLG338, pACYC184, pBR series such as e.g. pBR322, pUC series such as pUC18 or pUC19, M113 mp series, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III<sup>1-13</sup>-B1,  $\lambda$ gt11 or pBdCI; in *Streptomyces* pIJ101, pIJ364, pIJ702 or pIJ361; in *Bacillus* pUB110, pC194 or pBD214; in *Corynebacterium* pSA77 or pAJ667; in fungi pALS1, pIL2 or pBB116; other advantageous fungal vectors are described by Romanos, M. A. et al., [(1992) "Foreign gene expression in yeast: a review", *Yeast* 8: 423-488] and by van den Hondel, C. A. M. J. J. et al. [(1991) "Heterologous gene expression in filamentous fungi" as well as in More Gene Manipulations in Fungi [J. W. Bennet & L. L. Lasure, eds., pp. 396-428: Academic Press: San Diego] and in "Gene transfer systems and vector development for filamentous fungi" [van den Hondel, C. A. M. J. J. & Punt, P. J. (1991) in: Applied Molecular Genetics of Fungi, Peberdy, J. F. et al., eds., pp. 1-28, Cambridge University Press: Cambridge]. Examples of advantageous yeast promoters are 2  $\mu$ M, pAG-1, YEp6, YEp13 or pEMBLyE23. Examples of algal or plant promoters are pLGV23, pGHIac<sup>+</sup>, pBIN19, pAK2004, pVKH or pDH51 (see Schmidt, R. and Willmitzer, L., 1988). The vectors identified above or derivatives of the vectors identified above are a small selection of the possible plasmids. Further plasmids are well known to those skilled in the art and may be found, for example, in the book Cloning Vectors (Eds. Pouwels P. H. et al. Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). Suitable plant vectors are described inter alia in "Methods in Plant Molecular Biology and Biotechnology" (CRC Press), Ch. 6/7, pp. 71-119. Advantageous vectors are known as shuttle vectors or binary vectors which replicate in *E. coli* and *Agrobacterium*.

By vectors is meant with the exception of plasmids all other vectors known to those skilled in the art such as by way of example phages, viruses such as SV40, CMV, baculovirus, adenovirus, transposons, IS elements, phasmids, phagemids, cosmids, linear or circular DNA. These vectors can be replicated autonomously in the host organism or be chromosomally replicated, chromosomal replication being preferred.

In a further embodiment of the vector the gene construct according to the invention may also advantageously be introduced into the organisms in the form of a linear DNA and be integrated into the genome of the host organism by way of heterologous or homologous recombination. This linear DNA may be composed of a linearized plasmid or only of the expression cassette as vector or the nucleic acid sequences according to the invention.

In a further advantageous embodiment the nucleic acid sequence according to the invention can also be introduced into an organism on its own.

If in addition to the nucleic acid sequence according to the invention further genes are to be introduced into the organism, all together with a reporter gene in a single vector or each single gene with a reporter gene in a vector in each case can be introduced into the organism, whereby the different vectors can be introduced simultaneously or successively.

The vector advantageously contains at least one copy of the nucleic acid sequences according to the invention and/or the expression cassette (=gene construct) according to the invention.

By way of example the plant expression cassette can be installed in the pRT transformation vector ((a) Toepfer et al., 1993, *Methods Enzymol.*, 217: 66-78; (b) Toepfer et al. 1987, *Nucl. Acids. Res.* 15: 5890 ff.).

Alternatively, a recombinant vector (=expression vector) can also be transcribed and translated in vitro, e.g. by using the T7 promoter and the T7 RNA polymerase.

Expression vectors employed in prokaryotes frequently make use of inducible systems with and without fusion proteins or fusion oligopeptides, wherein these fusions can ensue in both N-terminal and C-terminal manner or in other useful domains of a protein. Such fusion vectors usually have the following purposes: i.) to increase the RNA expression rate; ii.) to increase the achievable protein synthesis rate; iii.) to increase the solubility of the protein; iv.) or to simplify purification by means of a binding sequence usable for affinity chromatography. Proteolytic cleavage points are also frequently introduced via fusion proteins, which allow cleavage of a portion of the fusion protein and purification. Such recognition sequences for proteases are recognized, e.g. factor Xa, thrombin and enterokinase.

Typical advantageous fusion and expression vectors are pGEX [Pharmacia Biotech Inc; Smith, D. B. and Johnson, K. S. (1988) *Gene* 67: 31-40], pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) which contains glutathione S-transferase (GST), maltose binding protein or protein A.

Other examples of *E. coli* expression vectors are pTrc [Amann et al., (1988) *Gene* 69:301-315] and pET vectors [Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 60-89; Stratagene, Amsterdam, The Netherlands].

Other advantageous vectors for use in yeast are pYepSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES derivatives (Invitrogen Corporation, San Diego, Calif.). Vectors for use in filamentous fungi are described in: van den Hondel, C. A. M. J. J. & Punt, P. J. (1991) "Gene transfer systems and vector development for filamentous fungi", in: *Applied Molecular Genetics of Fungi*, J. F. Peberdy, et al., eds., pp. 1-28, Cambridge University Press: Cambridge.

Alternatively, insect cell expression vectors can also be advantageously utilized, e.g. for expression in Sf 9 cells. These are e.g. the vectors of the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

Furthermore, plant cells or algal cells can advantageously be used for gene expression. Examples of plant expression vectors may be found in Becker, D., et al. (1992) "New plant binary vectors with selectable markers located proximal to the left border", *Plant Mol. Biol.* 20: 1195-1197 or in Bevan, M. W. (1984) "Binary *Agrobacterium* vectors for plant transformation", *Nucl. Acid. Res.* 12: 8711-8721.

The host plant (=transgenic plant) advantageously contains at least one copy of the nucleic acid according to the invention and/or of the gene construct according to the invention.

The introduction of the nucleic acids according to the invention, the gene construct or the vector into organisms, plants for example, can in principle be done by all of the methods known to those skilled in the art. The introduction of the nucleic acid sequences gives rise to recombinant or transgenic plants.

To introduce the nucleic acids used in the process, the latter are advantageously amplified and ligated in the known manner. Preferably, a procedure following the protocol for Pfu DNA polymerase or a Pfu/Taq DNA polymerase mixture is followed. The primers are selected taking into consideration the sequence to be amplified. The primers should advantageously be chosen in such a way that the amplificate comprises the entire codogenic sequence from the start codon to the stop codon. After the amplification, the amplificate is expediently analyzed. For example, a gel-electrophoretic separation can be carried out, which is followed by a quantitative and a qualitative analysis. Thereafter, the amplificate can be purified following a standard protocol (for example Qiagen). An aliquot of the purified amplificate is then available for the subsequent cloning step. Suitable cloning vectors are mentioned above and generally known to the skilled worker. These include, in particular, vectors which are capable of replication in microbial systems, that is to say mainly vectors which ensure efficient cloning in yeasts or fungi and which make possible the stable transformation of plants. Those, which must be mentioned, again herein in particular are various binary and cointegrated vector systems, which are suitable for the T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they comprise at least the vir genes required for the *Agrobacterium*-mediated transformation and the T-DNA-delimiting sequences (T-DNA border). These vector systems advantageously also comprise further cis-regulatory regions such as promoters and terminator sequences and/or selection markers, by means of which suitably transformed organisms can be identified. While in the case of cointegrated vector systems vir genes and T-DNA sequences are arranged on the same vector, binary systems are based on at least two vectors, one of which bears vir genes, but no T-DNA, while a second one bears T-DNA, but no vir gene. Owing to this fact, the last-mentioned vectors are relatively small, easy to manipulate and to replicate both in *E. coli* and in *Agrobacterium*. These binary vectors include vectors from the series pBIB-HYG, pPZP, pBecks, pGreen. In accordance with the invention, Bin19, pBI101, pBinAR, pGPTV and pCAMBIA are used by preference. An overview of the binary vectors and their use is found in Hellens et al, *Trends in Plant Science* (2000) 5, 446-451. In order to prepare the vectors, the vectors can first be linearized with restriction endonuclease(s) and then modified enzymatically in a suitable manner. Thereafter, the vector is purified, and an aliquot is employed for the cloning step. In the cloning step, the enzymatically cleaved and, if appropriate, purified amplificate is cloned with vector fragments, which have been prepared in a similar manner, using ligase. In this context, a particular nucleic acid construct, or vector or plasmid construct, can have one or else more than one codogenic gene segment. The codogenic gene segments in these constructs are preferably linked operably with regulatory sequences. The regulatory sequences include, in particular, plant sequences such as the above-described promoters and terminator sequences. The constructs can advantageously be stably propagated in microorganisms, in particular in *E. coli* and *Agrobacterium tumefaciens*, under selective conditions and make possible the transfer of heterologous DNA into plants or microorganisms.

The nucleic acids used in the process, the inventive nucleic acids and gene constructs, can be introduced into organisms such as microorganisms or advantageously plants, advantageously using cloning vectors, and thus be used in the transformation of plants such as those which are published and cited in: Plant Molecular Biology and Biotechnology (CRC Press, Boca Raton, Fla.), Chapter 6/7, p. 71-119 (1993); F. F. White, Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, 15-38; B. Jenés et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press (1993), 128-143; Potrykus, Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991), 205-225. Thus, the nucleic acids, the inventive nucleic acids and nucleic acid constructs, and/or vectors used in the process can be used for the recombinant modification of a broad spectrum of organisms, advantageously plants, so that the latter become better and/or more efficient PUFA and/or LCPUFA producers.

In the case of microorganisms, those skilled in the art can find appropriate methods for the introduction of the inventive nucleic acid sequences, the gene construct or the vector in the textbooks by Sambrook, J. et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, by F. M. Ausubel et al. (1994) Current protocols in molecular biology, John Wiley and Sons, by D. M. Glover et al., DNA Cloning Vol. 1, (1995), IRL Press (ISBN 019-963476-9), by Kaiser et al. (1994) Methods in Yeast Genetics, Cold Spring Harbor Laboratory Press or Guthrie et al. Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, 1994, Academic Press.

The transfer of foreign genes into the genome of a plant is called transformation. In doing this the methods described for the transformation and regeneration of plants from plant tissues or plant cells are utilized for transient or stable transformation. Suitable methods are protoplast transformation by poly(ethylene glycol)-induced DNA uptake, the "biolistic" method using the gene cannon—referred to as the particle bombardment method, electroporation, the incubation of dry embryos in DNA solution, microinjection and gene transfer mediated by *Agrobacterium*. Said methods are described by way of example in B. Jenés et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S. D. Kung and R. Wu, Academic Press (1993) 128-143 and in Potrykus Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991) 205-225). The nucleic acids or the construct to be expressed is preferably cloned into a vector, which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., Nucl. Acids Res. 12 (1984) 8711). *Agrobacteria* transformed by such a vector can then be used in known manner for the transformation of plants, in particular of crop plants such as by way of example tobacco plants, for example by bathing bruised leaves or chopped leaves in an agrobacterial solution and then culturing them in suitable media. The transformation of plants by means of *Agrobacterium tumefaciens* is described, for example, by Höfgen and Willmitzer in Nucl. Acid Res. (1988) 16, 9877 or is known inter alia from F. F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S. D. Kung and R. Wu, Academic Press, 1993, pp. 15-38.

*Agrobacteria* transformed by an expression vector according to the invention may likewise be used in known manner for the transformation of plants such as test plants like *Arabidopsis* or crop plants such as cereal crops, corn, oats, rye, barley, wheat, soybean, rice, cotton, sugar beet, canola, sunflower, flax, hemp, potatoes, tobacco, tomatoes, carrots,

paprika, oilseed rape, tapioca, cassava, arrowroot, *tagetes*, alfalfa, lettuce and the various tree, nut and vine species, in particular of oil-containing crop plants such as soybean, peanut, castor oil plant, sunflower, corn, cotton, flax (linseed), oilseed rape, poppy, mustard, sesame, almond, *macadamia*, olive, *calendula*, *punica*, hazel nut, avocado, pumpkin, walnut, laurel, pistachio, *Orychophragmus*, marigold, borage, primrose, canola, evening primrose, hemp, coconut, oil palm, safflower (*Carthamus tinctorius*), coffee or cocoa bean, e.g. by bathing bruised leaves or chopped leaves in an agrobacterial solution and then culturing them in suitable media. For the production of LCPUFAs, for example arachidonic acid and/or eicosapentaenoic acid, borage, linseed, sunflower, safflower, *Brassica napus*, *Brassica juncea*, *Camelina sativa* or *Orychophragmus* are advantageously suitable.

The genetically modified plant cells may be regenerated by all of the methods known to those skilled in the art. Appropriate methods can be found in the publications referred to above by S. D. Kung and R. Wu, Potrykus or Höfgen and Willmitzer.

Accordingly, a further aspect of the invention relates to transgenic organisms transformed by at least one nucleic acid sequence, expression cassette or vector according to the invention as well as cells, cell cultures, tissue, parts—such as, for example, leaves, roots, etc. in the case of plant organisms—or reproductive material derived from such organisms. The terms "host organism", "host cell", "recombinant (host) organism" and "transgenic (host) cell" are used here interchangeably. Of course these terms relate not only to the particular host organism or the particular target cell but also to the descendants or potential descendants of these organisms or cells. Since, due to mutation or environmental effects certain modifications may arise in successive generations, these descendants need not necessarily be identical with the parental cell but nevertheless are still encompassed by the term as used here.

Suitable organisms or host organisms for the nucleic acid, gene construct or vector according to the invention are advantageously in principle all plants, which are able to synthesize fatty acids, especially unsaturated fatty acids or are suitable for the expression of recombinant genes as described above. Further examples which may be mentioned are plants such as *Arabidopsis*, Asteraceae such as *Calendula* or crop plants such as soybean, peanut, castor oil plant, sunflower, corn, cotton, flax, oilseed rape, coconut, oil palm, safflower (*Carthamus tinctorius*) or cocoa bean, bacteria such as the genus *Escherichia*, yeasts such as the genus *Saccharomyces*. Preference is given to organisms which can naturally synthesize oils in relatively large quantities such as fungi like *Mortierella alpina*, *Pythium insidiosum* or plants such as soybean, oilseed rape, coconut, oil palm, safflower, flax, castor oil plant, *Calendula*, peanut, cocoa bean or sunflower, or yeasts such as *Saccharomyces cerevisiae* and particular preference is given to the family of the Brassicaceae such as oilseed rape, soybean, flax, sunflower, *Calendula*, *Mortierella* or *Saccharomyces cerevisiae*.

Further useful host cells are identified in: Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990).

Usable expression strains, e.g. those exhibiting a relatively low protease activity, are described in: Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990) 119-128.

A further object of the invention as described relates to the use of an expression cassette containing DNA sequences encoding a  $\Delta$ -12- and  $\Delta$ -15-desaturase, a  $\Delta$ -9-elongase, a  $\Delta$ -8-desaturase and/or a  $\Delta$ -5-desaturase gene or DNA



sequences hybridizing therewith for the transformation of plant cells, tissues or parts of plants. The aim of use is to increase the content of fatty acids, oils or lipids having an increased content of double bonds.

In doing so, depending on the choice of promoter, the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase, the  $\Delta$ -8-desaturase and/or the  $\Delta$ -5-desaturase gene can be expressed specifically in the leaves, in the seeds, the nodules, in roots, in the stem or other parts of the plant, preferably in leaves and/or seeds. Those transgenic plants overproducing fatty acids, oils or lipids according to the invention, the reproductive material thereof, together with the plant cells, tissues or parts thereof are a further object of the present invention.

The expression cassette or the nucleic acid sequences according to the invention containing a  $\Delta$ -12- and  $\Delta$ -15-desaturase, a  $\Delta$ -9-elongase, a  $\Delta$ -8-desaturase and/or a  $\Delta$ -5-desaturase gene sequence can, moreover, also be employed for the transformation of the organisms identified by way of example above such as bacteria, cyanobacteria, yeasts, filamentous fungi, ciliates and algae with the objective of increasing the content of fatty acids, oils or lipids according to the invention.

Within the framework of the present invention is the increase of the content of fatty acids, oils or lipids possessing a higher amount of  $\omega$ -3-fatty acids in comparison to  $\omega$ -6-fatty acids such as eicosapentaenoic acid in comparison to arachidonic acid, due to functional over expression of the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase, the  $\Delta$ -8-desaturase and/or the  $\Delta$ -5-desaturase gene in the plant according to the invention, advantageously in the trans-genic oilseed plants according to the invention, by comparison with the non genetically modified initial plants at least for the duration of at least one plant generation.

The preferred locus of biosynthesis, of fatty acids, oils or lipids for example, is generally the seed or cell layers of the seed so that a seed-specific expression of the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase, the  $\Delta$ -8-desaturase and/or the  $\Delta$ -5-desaturase gene is appropriate. It is, however, obvious that the biosynthesis of fatty acids, oils or lipids need not be limited to the seed tissue but rather can also occur in tissue-specific manner in all other parts of the plant—in epidermis cells or in the nodules for example.

A constitutive expression of the exogenous  $\Delta$ -12- and  $\Delta$ -15-desaturase,  $\Delta$ -9-elongase,  $\Delta$ -8-desaturase and/or  $\Delta$ -5-desaturase gene is, moreover, advantageous. On the other hand, however, an inducible expression may also appear desirable.

The efficiency of the expression of the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase, the  $\Delta$ -8-desaturase and/or the  $\Delta$ -5-desaturase gene can be determined, for example, in vitro by shoot meristem propagation. In addition, an expression of the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase, the  $\Delta$ -8-desaturase and/or the  $\Delta$ -5-desaturase gene modified in nature and level and its effect on fatty acid, oil or lipid biosynthesis performance can be tested on test plants in greenhouse trials.

An additional object of the invention comprises transgenic plants transformed by an expression cassette containing a  $\Delta$ -12- and  $\Delta$ -15-desaturase, a  $\Delta$ -9-elongase, a  $\Delta$ -8-desaturase and/or a  $\Delta$ -5-desaturase gene sequence according to the invention or DNA sequences hybridizing therewith, as well as transgenic cells, tissue, parts and reproduction material of such plants. Particular preference is given in this case to transgenic crop plants such as by way of example barley, wheat, rye, oats, corn, soybean, rice, cotton, sugar beet, the family of the Brassicaceae such as oilseed rape and canola,

sunflower, flax, hemp, thistle, potatoes, tobacco, tomatoes, tapioca, cassava, arrowroot, alfalfa, lettuce and the various tree, nut and vine species.

For the purposes of the invention plants are mono- and dicotyledonous plants that produce mature seeds.

A further refinement according to the invention are transgenic plants as described above which contain the nucleic acid sequences, the gene construct and/or vector of the invention.

The invention is explained in more detail by the following examples.

## EXAMPLES

### Example 1

#### General Cloning Methods

The cloning methods, such as by way of example restriction cleavages, agarose gel electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linkage of DNA fragments, transformation of *Escherichia coli* cells, culture of bacteria and sequence analysis of recombinant DNA, were carried out as described in Sambrook et al. (1989) (Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6).

### Example 2

#### Sequence Analysis of Recombinant DNA

Sequencing of recombinant DNA molecules was done using a laser fluorescence DNA sequencer from the ABI company by the method of Sanger (Sanger et al. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467). Fragments resulting from a polymerase chain reaction were sequenced and checked to prevent polymerase errors in the constructs to be expressed.

### Example 3

#### Cloning of the PUFA Specific Desaturases from *Acanthamoeba castellanii* (=SEQ ID NO: 3, 5, 15, 19 and 21)

*Acanthamoeba castellanii* (Eukaryota; Protista; Sarcostomatophora; Sarcodina; Rhizopodea; Lobosa) is an amoeba species, which is a common species in the soil. *Acanthamoeba castellanii* can grow vegetative over a broad temperature range (10 to 32° C.). *A. castellanii* is able to de novo synthesize linoleic acid and C20 n-6 fatty acids.

*A. castellanii* (ATTC 30010) was grown at 30° C. on a medium containing 0.75% (w/v) peptone, 1.5% (w/v) glucose and 0.75% (w/v) yeast extract according to the reference of Jones et al. [Temperature-induced membrane-lipid adaptation in *Acanthamoeba castellanii*. Biochem J. 1993, 290: 273-278]. The cell cultures were grown under shaking (200 U/min) and harvested with a centrifuge at 250×g, 5 min, 4° C., after they have reached a cell density of 5×10<sup>6</sup>-10<sup>7</sup> (measured in a Fuchs-Rosenthal Haemozytometer).

The total mRNA was isolated from said harvested cells with the aid of the RNeasy plant mini Kit (Qiagen). cDNA was synthesized from the total mRNA with the SMART RACE cDNA amplification kit (Clontech) according to the instructions of the manufacturer.



For the isolation of new desaturase genes the following degenerated primers were used for the amplification:

Deg1 : (SEQ ID NO: 53)  
5' -GGITGG(C/T/A) TIGGICA(T/C) GA(T/C) (GT) (CT) I (GT)  
(GC) ICA-3'

Deg2 : (SEQ ID NO: 54)  
5' -GG(A/G) AA(TCGA) AG(A/G) TG(A/G) TG(T/C) TC(A/G/T)  
AT(T/C) TG-3'

The aforementioned primers were used for the amplification in combination with the 3'-adapter-primer of the SMART RACE cDNA amplification kit.

The following protocol was used for the amplification:

- a) 2 min at 95° C.,
- b) 30 sec at 94° C.  
30 sec at 55-72° C.  
2 min at 72° C.
- c) 10 min at 72° C.

PCR amplicons were cloned and sequenced according to the instructions of the manufacturer (pTOPO, Invitrogen). The sequence information was used for the production of full-length clones. For the cloning of the full-length clones 5'- and 3'-specific primers were synthesized. Said primers were used for the amplification in the SMART RACE cDNA amplification kit (Clontech) and the amplicons were cloned into the pTOPO vector (Invitrogen)

Three sequences were identified, which show low similarities to desaturase genes.

In addition according to [Zank et al. 2002, Plant Journal 31:255-268] sequence 9Ac ( $\Delta$ -9-Elongase from *Acanthamoeba*, SEQ ID NO: 11) could be identified, which shows low similarities to elongase genes.

TABLE 1

<i>Acanthamoeba castellanii</i> desaturase sequences		
Gene	Nucleotide bp	SEQ ID NO:
12Ac ( $\Delta$ -12/ $\Delta$ 15-Desaturase from <i>Acanthamoeba</i> )	1224 bp	19, 21
8Ac ( $\Delta$ -8-Desaturase from <i>Acanthamoeba</i> )	1374 bp	3, 5
5Ac ( $\Delta$ -5-Desaturase from <i>Acanthamoeba</i> )	1353 bp	15

## Example 4

Cloning of the PUFA Specific Desaturases from *Perkinsus marinus* (=SEQ ID NO: 7, 17 and 23)

*Perkinsus marinus*, which belongs to the Protista, is a parasite in seashells. *P. marinus* is able to synthesize LCPUFAs such as arachidonic acid (20:4). The LCPUFAs are produced according to the present work over the  $\Delta$ -8-/ $\Delta$ -5-fatty acid pathway (see FIG. 1).

*P. marinus* was grown at 28° C. as disclosed by La Peyre et al. (J. Eukaryot. Microbiol. 1993, 40: 304-310).

The total mRNA was isolated from said harvested cells with the aid of the RNeasy plant mini Kit (Qiagen). cDNA was synthesized from the total mRNA with the SMART RACE cDNA amplification kit (Clontech) according to the instructions of the manufacturer.

For the isolation of new desaturase genes the following degenerated primers were used for the amplification:

Deg1 : (SEQ ID NO: 53)  
5' -GGITGG(C/T/A) TIGGICA(T/C) GA(T/C) (GT) (CT) I (GT)  
(GC) ICA-3'

Deg2 : (SEQ ID NO: 54)  
5' -GG(A/G) AA(TCGA) AG(A/G) TG(A/G) TG(T/C) TC(A/G/T)  
AT(T/C) TG-3'

The aforementioned primers were used for the amplification in combination with the 3'-adapter-primer of the SMART RACE cDNA amplification kit.

The following protocol was used for the amplification:

- d) 2 min at 95° C.,
- e) 30 sec at 94° C.  
30 sec at 55-72° C.  
2 min at 72° C.
- f) 10 min at 72° C.

Number of cycles: 30

PCR amplicons were cloned and sequenced according to the instructions of the manufacturer (pTOPO, Invitrogen). The sequence information was used for the production of full-length clones. For the cloning of the full-length clones 5'- and 3'-specific primers were synthesized. Said primers were used for the amplification in the SMART RACE cDNA amplification kit (Clontech) and the amplicons were cloned into the pTOPO vector (Invitrogen) Three sequences were identified, which show low similarities to desaturase genes.

TABLE 2

<i>Perkinsus marinus</i> desaturase sequences		
Gene	Nucleotide bp	SEQ ID NO:
12Pm ( $\Delta$ -12-Desaturase from <i>Perkinsus</i> )	1254 bp	23
8Pm ( $\Delta$ -8-Desaturase from <i>Perkinsus</i> )	1236 bp	7
5Pm ( $\Delta$ -5-Desaturase from <i>Perkinsus</i> )	1374 bp	17

50

## Example 5

Cloning of Expression Plasmids for the Heterologous Expression of *A. castellanii* and *P. marinus* Genes in Yeasts

For the heterologous expression in yeasts the respective sequences were PCR amplified and with the restriction enzymes KpnI-SacI the resulting sequences were cloned into the yeast vector pYES2 (Invitrogen). For the amplification specific primers (see table 3 below) were used. Only the open reading frames of the PUFA genes were amplified. In addition restriction cleavage sites were attached to the nucleic acid sequences. At the 5'-end a KpnI side and a so named Kozak sequence (Cell, 1986, 44: 283-292) was added. To the 3'-end a SacI side was attached.

TABLE 3

Primers for the amplification of the nucleic acid sequences of the desaturases			
Gen	bp	primer	SEQ ID NO:
12Ac	1224	Fwd: GGTACCATGGCGATCACGACGACGACAC	25
		Rvs: GAGCTCCTAGTGGCCTTGCCGTGCTTGATCTCC	26
8Ac	1374	Fwd: GGTACCATGGTCTCACAACCCCGGCCCTC	27
		Rvs: GGAGCTCTCAGTCTCAGCACCCATCTTC	28
5Ac	1353	Fwd: GGTACCATGGCCACCGCATCTGCATC	29
		Rvs: GGAGCTTTAGCCGTAGTAGGCCTCCTT	30
9Ac	891	Fwd: GGTACCATGGCGGCTGCGACGGCGAC	31
		Rvs: GGAGCTTTAGTCGTGCTTCTCTTGGG	32
12Pm	1254	Fwd: GGTACCATGACCCAAACTGAGGTCCA	33
		Rvs: GGAGCTCTAACGAGAAGTGCAGGCGT	34
8Pm	1236	Fwd: GGTACCATGTCTTCTCTTACCCTCTA	35
		Rvs: GGAGCTCTATTCCACTATGGCAACAG	36
5Pm	1374	Fwd: GGTACCATGACTACTTCAACCACTAC	37
		Rvs: GGAGCTCTACCTAGCAAGCAATCTCT	38

Composition of the PCR mix (50 µl)

5.00 µL	Template cDNA
5.00 µL	10x Puffer (Advantage-Polymerase) + 25 mM MgCl <sub>2</sub>
5.00 µL	2 mM dNTP
1.25 µL	each primer (10 pmol/µL of the 5'-ATG as well as of the 3'-stopp primer)
0.50 µL	Advantage polymerase

The Advantage polymerase from Clontech was employed. PCR Protocol

Addition temperature: 1 min at 55° C.  
Denaturing temperature: 1 min at 94° C.  
Elongation temperature: 2 min at 72° C.  
Number of cycles: 35

The PCR products and the vector pYES2 were incubated with the restriction enzymes KpnI and SacI for 1 h at 37° C. Afterwards a ligation reaction was done with the Rapid Ligation Kit (Roche) according to the instructions of the manufacturer. The reaction mixture was than used for the transformation of *E. coli* DH5α cells (Invitrogen) again according to the instructions of the manufacturer. Positive clones were identified with PCR (reaction scheme as described above).

The plasmid DNA was isolated (Qiagen Dneasy) and the resulting plasmids were checked by sequencing and transformed with the lithium acetate method into the *Saccharomyces* strain W303-1A. As a control the plasmid pYES2 (vector without insert) was transformed in parallel. The transformed yeasts were selected on complete minimal dropout uracil medium (CMdum) agar plates supplemented with 2% glucose, but without uracil.

To express the genes from *A. castellanii* and *P. marinus*, precultures consisting of in each case 5 ml of CMdum dropout uracil liquid medium supplemented with 2% (w/v) raffinose, but without uracil were initially inoculated with the selected transformants and incubated for 2 days at 30° C. and 200 rpm. Then, 5 ml of CMdum (without uracil) liquid medium supplemented with 2% of raffinose and 300 µM of various fatty acids were inoculated with the precultures to an OD<sub>600</sub> of 0.05. Expression was induced by the addition of 2% (w/v) of galactose. The cultures were incubated for a further 96 hours at 22° C.

Example 6

Cloning of Expression Plasmids for the Expression in Plants

To transform plants, a further transformation vector based on pBIN19-35S (Bevan M. (1984) Binary *Agrobacterium* vectors for plant transformation. Nucl. Acids Res. 18:203) was generated. To this end, BamHI-XbaI cleavage sites were inserted at the 5' and 3' end of the coding sequences, using PCR. The corresponding primer sequences were derived from the 5' and 3' regions of the respective nucleic acid sequence (see table 4).

TABLE 4

Primers for the expression in plants			
Gen	bp	primer	SEQ ID NO:
12Ac	1224	Fwd: GGATCCACCATGGCGATCACGACGACGACAGACAC	39
		Rvs: GGTCTAGACTAGTGGCCTTGCCGTGCTTGATCTCC	40
8Ac	1374	Fwd: GGATCCAGGATGGTCTCACAACCCCGGCCCTC	41
		Rvs: GGTCTAGATCAGTTCTCAGCACCCATCTTC	42
5Ac	1353	Fwd: GGATCCATGGCCACCGCATCTGCATC	43
		Rvs: GGTCTAGATTAGCCGTAGTAGGCCTCCTT	44
9Ac	891	Fwd: GGATCCATGGCGGCTGCGACGGCGAC	45
		Rvs: GGTCTAGATTAGTCGTGCTTCTCTTGGG	46
12Pm	1254	Fwd: GGATCCATGACCCAAACTGAGGTCCA	47
		Rvs: GGTCTAGACTAACGAGAAGTGCAGGCGT	48

TABLE 4-continued

Primers for the expression in plants			
Gen	bp	primer	SEQ ID NO:
8Pm	1236	Fwd: GGATCCATGTCCTTCTTACCCTCTA	49
		Rvs: GGTCTAGACTATTCCACTATGGCAACAG	50
5Pm	1374	Fwd: GGATCCATGACTACTTCAACCACTAC	51
		Rvs: GGTCTAGACTACCTAGCAAGCAATCTCT	52

Composition of the PCR Mix (50 µl):

5.00 µl template cDNA

5.00 µl 10× buffer (Advantage polymerase)+25 mM MgCl<sub>2</sub>

5.00 µl 2 mM dNTP

1.25 µl of each primer (10 pmol/µl)

0.50 µl Advantage polymerase

The Advantage polymerase from Clontech was employed.

PCR Reaction Conditions:

Annealing temperature: 1 min 55° C.

Denaturation temperature: 1 min 94° C.

Elongation temperature: 2 min 72° C.

Number of cycles: 35

The PCR products as well as the vector pBin19-35S were incubated with the restriction enzymes BamHI and XbaI for 16 hours at 37° C. Afterwards a ligation reaction was done with the Rapid Ligation Kit (Roche) according to the instructions of the manufacturer. The reaction mixture was than used for the transformation of *E. coli* DH5α cells (Invitrogen) again according to the instructions of the manufacturer. Positive clones were identified with PCR (reaction scheme as described above) and the plasmid DNA was isolated (Qiagen Dneasy). The resulting plasmids were checked by sequencing and transformed by electroporation into *Agrobacterium tumefaciens* GC3101. Afterwards the transformants were plated on 2% YEB Medium agar plates with kanamycin. Kanamycin tolerant cells were picked and used for the transformation of *Arabidopsis thaliana*.

#### Example 7

##### Expression of *A. castellanii* and *P. marinus* Genes in Yeasts

Yeasts which had been transformed with the plasmids pYES2, pYES-12Ac, pYES-8Ac, pYES2-5Ac, pYES2-9Ac, pYES2-12Pm, pYES2-8Pm and pYES2-5Pm as described in Example 5 were analyzed as follows:

The yeast cells from the main cultures were harvested by centrifugation (100×g, 5 min, 20° C.) and washed with 100 mM NaHCO<sub>3</sub>, pH 8.0 to remove residual medium and fatty acids. Starting with the yeast cell sediments, fatty acid methyl esters (FAMES) were prepared by acid methanolysis. To this end, the cell sediments were incubated for one hour at 80° C. together with 2 ml of 1 N methanolic sulfuric acid and 2% (v/v) of dimethoxypropane. The FAMES were extracted twice with petroleum ether (PE). To remove nonderivatized fatty acids, the organic phases were washed in each case once with 2 ml of 100 mM NaHCO<sub>3</sub>, pH 8.0 and 2 ml of distilled water. Thereafter, the PE phases were dried with Na<sub>2</sub>SO<sub>4</sub>, evaporated under argon and taken up in 100 µl of PE. The samples

were separated on a DB-23 capillary column (30 m, 0.25 mm, 0.25 µm, Agilent) in a Hewlett-Packard 6850 gas chromatograph equipped with flame ionization detector. The conditions for the GLC analysis were as follows: the oven temperature was programmed from 50° C. to 250° C. with a rate of 5° C./min and finally 10 min at 250° C. (holding).

The signals were identified by comparing the retention times with corresponding fatty acid standards (Sigma). The methodology is described for example in Napier and Michaelson, 2001, *Lipids*. 36 (8):761-766; Sayanova et al., 2001, *Journal of Experimental Botany*. 52 (360):1581-1585, Sperling et al., 2001, *Arch. Biochem. Biophys.* 388 (2):293-298 and Michaelson et al., 1998, *FEBS Letters*. 439 (3):215-218.

#### Example 8

##### Functional Characterization of the Genes of *A. castellanii*

The substrate activity and specificity of the genes were determined after expression and after feeding various fatty acids. The substrate specificity of the desaturases after expressions in yeasts can be determined by feeding various different fatty acids. Specific examples for the determination of the specificity and activity are disclosed for example in WO 93/11245, WO 94/11516, WO 93/06712, U.S. Pat. No. 5,614,393, U.S. Pat. No. 5,614,393, WO 96/21022, WO0021557 and WO 99/27111, Qiu et al. 2001, *J. Biol. Chem.* 276, 31561-31566 for Δ4-desaturases, Hong et al. 2002, *Lipids* 37, 863-868 for Δ5-desaturases. WO2005/012316 teaches such a method for example in example 18 in more detail.

##### a) Characterization of the Gene 12Ac:

First the construct pYES-12Ac was tested in yeasts without feeding fatty acids. Astonishingly it was shown in comparison to the control vector pYES2 (vector without insert) that even without feeding fatty acids new fatty acids are detectable in the yeasts (FIGS. 2 A and B).

FIGS. 2 A and B show a comparison of the fatty acid profile between the control (construct pYES2 without insert, FIG. 2A) and the construct pYES2-12Ac (FIG. 2B), which contains the *Acanthamoeba castellanii* gene for the Δ-12-/Δ-15-desaturase. The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3, whereas the unusual fatty acids 16:2n-4 and 16:3n-1 are formed for the C16 fatty acids. For the C18 fatty acids linoleic and linolenic acid (18:2n-6 and 18:2n-3) are formed.

According to the new synthesized fatty acids it is possible to identify the gene product of the nucleic acid sequence as a Δ-12-desaturase. The enzyme is able to desaturate C18:1 and C16:1 as substrate to the corresponding C18:2 and C16:2 fatty acids. The conversion rate of C18:1 (40.0%) is higher than the rate of the C16:1 (15.8%) conversion. That means the conversion rate of C18:1 is more than double than the conversion rate of the C16:1.

The conversion rate of the desaturase was calculated according to the following formula:

$$\frac{\text{Substrate}}{(\text{Substrate} + \text{Product}) \times 100}$$

The result of the formula is given as percentage value.

Furthermore the enzyme shows in addition a clear Δ-15-desaturase-activity. That means also that products of the

$\Delta$ -12-desaturase reaction, which are C16:2 and/or C18:2 are further desaturated to C16:3 and/or C18:3.

b) Characterization of the Gene 8Ac:

According to different sequence alignments (Blast) performed with the sequence SEQ ID NO: 3 (8Ac sequence) with different data bases (NCBI-BLAST: at ncbi.nlm.nih.gov/BLAST/) the encoded protein sequence is most likely a putative  $\Delta$ -5-desaturase.

Sequences with significant similarities	(bits)	Value
gi 16033740 gb AAL13311.1 delta-5 fatty acid desaturase [P...]	176	1e-42
gi 50882495 gb AAT85663.1 polyunsaturated fatty acid delta...	170	6e-41
gi 4150956 dbj BAA37090.1 delta 5 fatty acid desaturase [D...]	156	9e-37
gi 23894018 embl CAD53323.1 delta 5 fatty acid desaturase [L...]	156	1e-36
gi 33466346 gb AAQ19605.1 delta-4 fatty acid desaturase [E...]	150	7e-35
gi 5263169 dbj BAA81814.1 fatty acid desaturase [Dictyoste...]	149	1e-34
gi 25956288 gb AAN75707.1 delta 4-desaturase [Thraustochyt...]	142	1e-32
gi 25956290 gb AAN75708.1 delta 4-desaturase [Thraustochyt...]	139	1e-31
gi 25956294 gb AAN75710.1 delta 4-desaturase [Thraustochyt...]	139	1e-31
gi 25956292 gb AAN75709.1 delta 4-desaturase [Thraustochyt...]	138	2e-31
gi 20069125 gb AAM09688.1 delta-4 fatty acid desaturase [T...]	138	3e-31
gi 39545945 gb AAR28035.1 delta-5 desaturase [Mortierella...]	136	9e-31
gi 3859488 gb AAC72755.1 delta-5 fatty acid desaturase [Mo...]	135	2e-30
gi 41017070 sp O74212 FAD5_MORAP Delta-5 fatty acid desatur...]	130	7e-29
gi 48854274 ref ZP_00308437.1 COG3239: Fatty acid desatur...]	114	4e-24
gi 48854276 ref ZP_00308439.1 COG3239: Fatty acid desatur...]	114	7e-24

According to this putative activity different fatty acids were fed (18:2, 18:3, 20:3n-6, 20:4n-3). None of said fatty acids were desaturated by the enzyme. This result clearly shows that the protein encoded by the 8Ac gene has neither a  $\Delta$ -5-desaturase activity nor a  $\Delta$ -6-desaturase activity.

Unexpectedly after feeding of the fatty acids 20:2n-6 and 20:3n-3 it could be shown, that the 8Ac sequence encodes a  $\Delta$ -8-desaturase (see FIGS. 3 A, 3 B, 4 A and 4 B).

FIGS. 3 A and B shows the fatty acid profile of yeasts transformed with the construct pYES2 as control (FIG. 3 A) and pYES2-8Ac (FIG. 3 B) and fed with the fatty acid C20:2<sup>A11,14</sup>. The respective fatty acids are market.

FIGS. 4 A and B shows the fatty acid profile of yeast transformed with the construct pYES2 (FIG. 4 A) as control and pYES2-8Ac (FIG. 4 B) and fed with the fatty acid C20:3<sup>A11,14,17</sup>. The respective fatty acids are market.

The protein encoded by 8Ac sequence is therefore a  $\Delta$ -8-desaturase. The conversion rates for the fatty acids C20:2 and C20:3 are 15.2% and 17.5% respectively. This is absolutely astonishing as the 8Ac sequence, which has some similarities to "front-end" desaturases, has a different conserved region of the characteristic Cyt b5 motif His-Pro-Gly-Gly (HPGG, SEQ ID NO: 55), which is necessary for building the Heme domain. In general mutations in said domain lead to depletion of the enzymatic activity (Sayanova et al. 1999, Plant Physiol 121 (2):641-646). The amino acid sequence of this new  $\Delta$ -8-desaturase shows unexpected differences to known "front-

end" desaturases. Instead of the HPGG motif this desaturase shows the motif HPAG (see SEQ ID NO: 3), which is due to an alanine in position 44 of the sequence. Sayanova et al. 1999, Plant Physiol 121 (2):641-646 has shown that such a change of the motif from HPPG to HPAG leads to inactive enzymes. Therefore the activity of the new  $\Delta$ -8-desaturase is even more astonishing.

For the further improvement of the activity of the  $\Delta$ -8-desaturase, the sequence of the enzyme was mutagenized. The following primer.

8AcMf  
CAAGTACCACCCGGCGGCAGCAGGGCCA (SEQ ID NO: 56)  
and  
8AcMr  
TGGCCCTGCTGCCGCCGGGTGGTACTTG (SEQ ID NO: 57)

were used together with the site directed mutagenesis Kit (Stratagene) for the mutagenesis according to the instructions of the manufacturer of the  $\Delta$ -8-desaturase. The mutagenesis was afterwards checked by sequencing. Due to the mutagenesis the nucleotide sequences 124-CACCCGGCCGGC was changed to 124-CACCCGGCGGC, which leads to a change from Alanine to Glycine in position 44 of the nucleic acid sequence shown in SEQ ID NO: 3. The resulting sequence is shown in SEQ ID NO: 5. As already described for the sequence of 8Ac the mutated sequence 8AcM was also cloned into the vector pYES2 and transformed into yeast. Yeast transformed either with the vector pYES-8Ac or pYES2-8AcM were grown and fed in parallel with different fatty acids (see table 5). The results of the feeding are shown in table 5. The mutated enzyme 8AcM shows in comparison to the wild type enzyme 8Ac an increased activity towards the fatty acid C20:2. This is a two fold increase of the activity. The mutation has no influence of the activity with the fatty acid C20:3 as substrate. This clearly shows that with the mutation the activity of the  $\Delta$ -8-desaturase can be influenced in a very specific manner.

TABLE 5

Fatty acid conversion rate of yeasts transformed with pYES-8Ac or pYES2-8AcM		
Plasmid	Fatty acid C20:2	Fatty acid C20:3
pYES-8Ac	15.2%	17.5%
pYES2-8AcM	30.0%	17.2%

The mutated  $\Delta$ -8-desaturase 8AcM and its derivatives are especially useful alone or in combination with the  $\Delta$ -12- and  $\Delta$ -15-desaturase, the  $\Delta$ -9-elongase and the  $\Delta$ -5-desaturase for the synthesis of arachidonic acid.

c) Characterization of the Gene 5Pm:

The constructs pYES2 and pYES-5Pm were transformed into yeasts grown in parallel as described. Afterwards 250  $\mu$ M of different fatty acids were fed. During this feeding experiments it can be shown that fatty acids such as C16:0, C16:1, C18:0, C18:1, C18:2n-6, C20:2n-6 or C22:4n-6 are not desaturated by the protein encoded by the 5Pm sequence. Whereas the substrate C20:3n-6 was desaturated by the enzyme (see FIGS. 5 A and 5 B). FIGS. 5 A and 5 B clearly shows that the enzyme produces arachidonic acid during the transformation of the fatty acid substrate C20:3n-6. No new fatty acid is produced by the control (FIG. 5 A). The desaturation of the fatty acid substrate C20:3n-6 to arachidonic acid is due to a  $\Delta$ -5-desaturase activity, which is encoded by the

5Pm sequence (SEQ ID NO: 17). The conversion rate calculated according to the equation mentioned above is 15.4%.

FIGS. 5 A and 5 B shows the comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (FIG. 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (FIG. 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).

d) Characterization of the Genes 5Ac, 9Ac, 12Pm and 8Pm:

According to sequence comparisons it was able to identify the sequences 5Ac, 12Pm and 8Pm as desaturases having a  $\Delta$ -5-desaturase,  $\Delta$ -12-desaturase and  $\Delta$ -8-desaturase activity. For the sequence 9Ac we were able to show a  $\Delta$ -9-elongase activity.

In combination with the 12Ac and 8Ac gene the complete set of enzymes from *A. castellanii*, which is necessary for the synthesis for arachidonic (C20:4n-6) or eicosapentaenoic acid could be identified. In addition further genes for the synthesis of said aforementioned fatty acids are isolated from *P. marinus*. With the aid of said genes the PUFA and/or LCPUFA content can be further improved. For the synthesis of arachidonic acid or eicosapentaenoic acid said genes can be introduced in plants or microorganism (see example 8).

#### Example 8

##### Generation of Transgenic Plants

a) Generation of Transgenic Oilseed Rape Plants (Modified Method of Moloney et al., 1992, Plant Cell Reports, 8:238-242)

Binary vectors in *Agrobacterium tumefaciens* C58C1: pGV2260 or *Escherichia coli* (Deblaere et al, 1984, Nucl. Acids. Res. 13, 4777-4788) can be used for generating transgenic oilseed rape plants. To transform oilseed rape plants (Var. *Drakkar*, NPZ Nordeutsche Pflanzenzucht, Hohenlieth, Germany), a 1:50 dilution of an overnight culture of a positively transformed agrobacterial colony in Murashige-Skoog medium (Murashige and Skoog 1962 Physiol. Plant. 15, 473) supplemented with 3% sucrose (3MS medium) is used. Petioles or hypocotyls of freshly germinated sterile oilseed rape plants (in each case approx. 1 cm<sup>2</sup>) are incubated with a 1:50 agrobacterial dilution for 5-10 minutes in a Petri dish. This is followed by 3 days of coincubation in the dark at 25° C. on 3MS medium supplemented with 0.8% Bacto agar. The cultures are then grown for 3 days at 16 hours light/8 hours dark and the cultivation is continued in a weekly rhythm on MS medium supplemented with 500 mg/l Claforan (cefotaxim sodium), 50 mg/l kanamycin, 20 µM benzylaminopurine (BAP), now supplemented with 1.6 g/l of glucose. Growing shoots are transferred to MS medium supplemented with 2% sucrose, 250 mg/l Claforan and 0.8% Bacto agar. If no roots develop after three weeks, 2-indolebutyric acid was added to the medium as growth hormone for rooting.

Regenerated shoots are obtained on 2MS medium supplemented with kanamycin and Claforan; after rooting, they are transferred to compost and, after growing on for two weeks in a controlled-environment cabinet or in the greenhouse, allowed to flower, and mature seeds are harvested and analyzed by lipid analysis for elongase and/or desaturase expression, such as  $\Delta$ -12- and  $\Delta$ -15-desaturase,  $\Delta$ -8-desaturase,  $\Delta$ -9-elongase or  $\Delta$ -5-desaturase activity. In this manner, lines with elevated contents of PUFAs and/or LCPUFAs can be identified.

b) Generation of Transgenic Linseed Plants

Transgenic linseed plants can be generated for example by the method of Bell et al., 1999, In Vitro Cell. Dev. Biol.-Plant.

35 (6):456-465 by means of particle bombardment. In general, linseed was transformed by an agrobacteria-mediated transformation, for example by the method of Mlynarova et al. (1994), Plant Cell Report 13: 282-285.

c) Generation of Transgenic *Arabidopsis* Plants

Binary plasmids were transferred to *A. tumefaciens* strain GV3101 by electroporation and kanamycin-resistant colonies were selected in all cases. Wildtype Col0 or transgenic line CA1-9, containing the coding region of *I. galbana* elongating activity, IgASE1 [Qi, B., Beaudoin, F., Fraser, T., Stobart, A. K., Napier, J. A. and Lazarus, C. M. (2002) Identification of a cDNA encoding a novel C18-D9 polyunsaturated fatty acid-specific elongating activity from the docosa-hexaenoic acid (DHA)-producing microalga, *Isochrysis galbana*. *FEBS Lett.* 510, 159-65] was used as the host for transformation with *A. castellanii*  $\Delta^8$  desaturase gene. *A. tumefaciens*-mediated transformation was performed as described in Bechthold et al. [(1993) In planta *Agrobacterium*-mediated gene transfer by infiltration of *Arabidopsis thaliana* plants. C.R. Acad. Sci. Ser. III Sci. Vie., 316, 1194-1199.] and seeds from dipped plants were spread on Murashige and Skoog medium containing 50 µg ml<sup>-1</sup> kanamycin.

#### Example 9

##### Lipid Extraction from Leaves

The effect of the genetic modification in plants, fungi, algae, ciliates or on the production of a desired compound (such as a fatty acid) can be determined by growing the modified microorganisms or the modified plant under suitable conditions (such as those described above) and analyzing the medium and/or the cellular components for the elevated production of desired product (i.e. of the lipids or a fatty acid). These analytical techniques are known to the skilled worker and comprise spectroscopy, thin-layer chromatography, various types of staining methods, enzymatic and microbiological methods and analytical chromatography such as high-performance liquid chromatography (see, for example, Ullman, Encyclopedia of Industrial Chemistry, Vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon, A., et al., (1987) "Applications of HPLC in Biochemistry" in: Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 17; Rehm et al. (1993) Biotechnology, Vol. 3, Chapter III: "Product recovery and purification", p. 469-714, VCH: Weinheim; Belter, P. A., et al. (1988) Bioseparations: downstream processing for Biotechnology, John Wiley and Sons; Kennedy, J. F., and Cabral, J. M. S. (1992) Recovery processes for biological Materials, John Wiley and Sons; Shaeiwitz, J. A., and Henry, J. D. (1988) Biochemical Separations, in: Ullmann's Encyclopedia of Industrial Chemistry, Vol. B3; Chapter 11, p. 1-27, VCH: Weinheim; and Dechow, F. J. (1989) Separation and purification techniques in biotechnology, Noyes Publications).

In addition to the abovementioned processes, plant lipids are extracted from plant material as described by Cahoon et al. (1999) Proc. Natl. Acad. Sci. USA 96 (22):12935-12940 and Browse et al. (1986) Analytic Biochemistry 152:141-145. The qualitative and quantitative analysis of lipids or fatty acids is described by Christie, William W., Advances in Lipid Methodology, Ayr/Scotland: Oily Press (Oily Press Lipid Library; 2); Christie, William W., Gas Chromatography and Lipids. A Practical Guide—Ayr, Scotland: Oily Press, 1989, Repr. 1992, IX, 307 pp. (Oily Press Lipid Library; 1); "Progress in Lipid Research, Oxford: Pergamon Press, 1 (1952)-16 (1977) under the title: Progress in the Chemistry of Fats and Other Lipids CODEN.

One example is the analysis of fatty acids (abbreviations: FAME, fatty acid methyl ester; GC-MS, gas liquid chromatography/mass spectrometry; TAG, triacylglycerol; TLC, thin-layer chromatography).

The unambiguous detection for the presence of fatty acid products can be obtained by analyzing recombinant organisms using analytical standard methods: GC, GC-MS or TLC, as described on several occasions by Christie and the references therein (1997, in: *Advances on Lipid Methodology*, Fourth Edition: Christie, Oily Press, Dundee, 119-169; 1998, *Gaschromatographie-Massenspektrometrie-Verfahren* [Gas chromatography/mass spectrometric methods], *Lipide* 33:343-353).

The material to be analyzed can be disrupted by sonication, grinding in a glass mill, liquid nitrogen and grinding or via other applicable methods. After disruption, the material must be centrifuged. The sediment is resuspended in distilled water, heated for 10 minutes at 100° C., cooled on ice and recentrifuged, followed by extraction for one hour at 90° C. in 0.5 M sulfuric acid in methanol with 2% dimethoxypropane, which leads to hydrolyzed oil and lipid compounds, which give transmethylated lipids. These fatty acid methyl esters are extracted in petroleum ether and finally subjected to a GC analysis using a capillary column (Chrompack, WCOT Fused Silica, CP-Wax-52 CB, 25 µm, 0.32 mm) at a temperature gradient of between 170° C. and 240° C. for 20 minutes and 5 minutes at 240° C. The identity of the resulting fatty acid methyl esters must be defined using standards, which are available from commercial sources (i.e. Sigma).

Plant material is initially homogenized mechanically by comminuting in a pestle and mortar to make it more amenable to extraction.

This is followed by heating at 100° C. for 10 minutes and, after cooling on ice, by resedimentation. The cell sediment is hydrolyzed for one hour at 90° C. with 1 M methanolic sulfuric acid and 2% dimethoxypropane, and the lipids are transmethylated. The resulting fatty acid methyl esters (FAMES) are extracted in petroleum ether. The extracted FAMES are analyzed by gas liquid chromatography using a capillary column (Chrompack, WCOT Fused Silica, CP-Wax-52 CB, 25 m, 0.32 mm) and a temperature gradient of from 170° C. to 240° C. in 20 minutes and 5 minutes at 240° C. The identity of the fatty acid methyl esters is confirmed by

comparison with corresponding FAME standards (Sigma). The identity and position of the double bond can be analyzed further by suitable chemical derivatization of the FAME mixtures, for example to give 4,4-dimethoxyoxazoline derivatives (Christie, 1998) by means of GC-MS.

Leaf material from transgenic *Arabidopsis thaliana* Col0 and supertransformants of transgenic line CA1-9 both transformed with the construct pBIN1935S-8Ac were analyzed by gas chromatography of methyl ester derivatives as described above. Identities were confirmed by GC-MS and co-migration with authentic standards. The conversion rates are shown in the following table 6:

TABLE 6

Conversion rate with AcD8 (delta-8-desaturase from <i>Acanthamoeba castellanii</i> ) of different substrates		
fatty acids	% of total fatty acids	% conversion of substrate
20:2 <sup>Δ11, 14</sup>	1.1	—
20:3 <sup>Δ8, 11, 14</sup>	1.9	63
20:2 <sup>Δ11, 14, 17</sup>	1.3	—
20:2 <sup>Δ8, 11, 14, 17</sup>	0.8	40

FIG. 6 shows the result with the line CA1-9. In the double transgenic *Arabidopsis* a clear activity of Ac8 can be shown by the conversion of the present 20:2<sup>Δ11,14</sup> or 20:3<sup>Δ11,14,17</sup> into 20:3<sup>Δ8,11,14</sup> or 20:4<sup>Δ8,11,14,17</sup>, the precursors of arachidonic acid or eicosapentaenoic acid.

Additionally Acyl-CoA profiles were done from the *Arabidopsis* leaves of *Arabidopsis* wild type (FIG. 7 A), *Arabidopsis* Δ9elo (FIG. 7 B) and *Arabidopsis* Δ9eloΔ8des (FIG. 7 C) using the method of Larson et al. [Plant J. 2002 November; 32 (4):519-27]. Results from the measurements are shown in FIG. 7 and demonstrate again the functionality of 8Ac in plants.

## EQUIVALENTS

Many equivalents of the specific embodiments according to the invention described herein can be identified or found by the skilled worker resorting simply to routine experiments. These equivalents are intended to be within the scope of the patent claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 57

<210> SEQ ID NO 1

<211> LENGTH: 1266

<212> TYPE: DNA

<213> ORGANISM: *Euglena gracilis*

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(1266)

<223> OTHER INFORMATION: Delta-8-Desaturase

<400> SEQUENCE: 1

```

atg aag tca aag cgc caa gcg ctt ccc ctt aca att gat gga aca aca      48
Met Lys Ser Lys Arg Gln Ala Leu Pro Leu Thr Ile Asp Gly Thr Thr
1           5           10

tat gat gtg tct gcc tgg gtc aat ttc cac cct ggt ggt gcg gaa att      96
Tyr Asp Val Ser Ala Trp Val Asn Phe His Pro Gly Gly Ala Glu Ile
                20                25                30

ata gag aat tac caa gga agg gat gcc act gat gcc ttc atg gtt atg      144

```

-continued

Ile	Glu	Asn	Tyr	Gln	Gly	Arg	Asp	Ala	Thr	Asp	Ala	Phe	Met	Val	Met	
		35					40					45				
cac	tct	caa	gaa	gcc	ttc	gac	aag	ctc	aag	cgc	atg	ccc	aaa	atc	aat	192
His	Ser	Gln	Glu	Ala	Phe	Asp	Lys	Leu	Lys	Arg	Met	Pro	Lys	Ile	Asn	
	50					55					60					
ccc	agt	tct	gag	ttg	cca	ccc	cag	gct	gca	gtg	aat	gaa	gct	caa	gag	240
Pro	Ser	Ser	Glu	Leu	Pro	Pro	Gln	Ala	Ala	Val	Asn	Glu	Ala	Gln	Glu	
	65				70					75					80	
gat	ttc	cgg	aag	ctc	cga	gaa	gag	ttg	atc	gca	act	ggc	atg	ttt	gat	288
Asp	Phe	Arg	Lys	Leu	Arg	Glu	Glu	Leu	Ile	Ala	Thr	Gly	Met	Phe	Asp	
			85						90					95		
gcc	tcc	ccc	ctc	tgg	tac	tca	tac	aaa	atc	agc	acc	aca	ctg	ggc	ctt	336
Ala	Ser	Pro	Leu	Trp	Tyr	Ser	Tyr	Lys	Ile	Ser	Thr	Thr	Thr	Leu	Gly	Leu
			100					105						110		
gga	gtg	ctg	ggt	tat	ttc	ctg	atg	ggt	cag	tat	cag	atg	tat	ttc	att	384
Gly	Val	Leu	Gly	Tyr	Phe	Leu	Met	Val	Gln	Tyr	Gln	Met	Tyr	Phe	Ile	
		115					120					125				
ggg	gca	gtg	ttg	ctt	ggg	atg	cac	tat	caa	cag	atg	ggc	tgg	ctt	tct	432
Gly	Ala	Val	Leu	Leu	Gly	Met	His	Tyr	Gln	Gln	Met	Gly	Trp	Leu	Ser	
	130					135					140					
cat	gac	att	tgc	cac	cac	cag	act	ttc	aag	aac	cgg	aac	tgg	aac	aac	480
His	Asp	Ile	Cys	His	His	Gln	Thr	Phe	Lys	Asn	Arg	Asn	Trp	Asn	Asn	
	145				150					155					160	
ctc	gtg	gga	ctg	gta	ttt	ggc	aat	ggg	ctg	caa	ggt	ttt	tcc	gtg	aca	528
Leu	Val	Gly	Leu	Val	Phe	Gly	Asn	Gly	Leu	Gln	Gly	Phe	Ser	Val	Thr	
			165					170						175		
tgc	tgg	aag	gac	aga	cac	aat	gca	cat	cat	tgc	gca	acc	aat	ggt	caa	576
Cys	Trp	Lys	Asp	Arg	His	Asn	Ala	His	His	Ser	Ala	Thr	Asn	Val	Gln	
		180						185					190			
ggg	cac	gac	cct	gat	att	gac	aac	ctc	ccc	ctc	tta	gcc	tgg	tct	gag	624
Gly	His	Asp	Pro	Asp	Ile	Asp	Asn	Leu	Pro	Leu	Leu	Ala	Trp	Ser	Glu	
		195				200						205				
gat	gac	gtc	aca	cgg	gcg	tca	cgg	att	tcc	cgc	aag	ctc	att	cag	ttc	672
Asp	Asp	Val	Thr	Arg	Ala	Ser	Pro	Ile	Ser	Arg	Lys	Leu	Ile	Gln	Phe	
	210					215					220					
cag	cag	tat	tat	ttc	ttg	gtc	atc	tgt	atc	ttg	ttg	cgg	ttc	att	tgg	720
Gln	Gln	Tyr	Tyr	Phe	Leu	Val	Ile	Cys	Ile	Leu	Leu	Arg	Phe	Ile	Trp	
	225				230					235					240	
tgt	ttc	cag	agc	gtg	ttg	acc	gtg	cgc	agt	ctg	aag	gac	aga	gat	aac	768
Cys	Phe	Gln	Ser	Val	Leu	Thr	Val	Arg	Ser	Leu	Lys	Asp	Arg	Asp	Asn	
			245					250						255		
caa	ttc	tat	cgc	tct	cag	tat	aag	aag	gag	gcc	att	ggc	ctc	gcc	ctg	816
Gln	Phe	Tyr	Arg	Ser	Gln	Tyr	Lys	Lys	Glu	Ala	Ile	Gly	Leu	Ala	Leu	
			260					265					270			
cat	tgg	aca	ttg	aag	gcc	ctg	ttc	cac	tta	ttc	ttt	atg	ccc	agc	atc	864
His	Trp	Thr	Leu	Lys	Ala	Leu	Phe	His	Leu	Phe	Phe	Met	Pro	Ser	Ile	
		275				280						285				
ctc	aca	tgc	ctg	ttg	gta	ttt	ttc	ggt	tgc	gag	ctg	ggt	ggc	ggc	ttc	912
Leu	Thr	Ser	Leu	Leu	Val	Phe	Phe	Val	Ser	Glu	Leu	Val	Gly	Gly	Phe	
	290					295						300				
ggc	att	gcg	atc	gtg	gtg	ttc	atg	aac	cac	tac	cca	ctg	gag	aag	atc	960
Gly	Ile	Ala	Ile	Val	Val	Phe	Met	Asn	His	Tyr	Pro	Leu	Glu	Lys	Ile	
	305				310					315					320	
ggg	gac	tgc	gtc	tgg	gat	ggc	cat	gga	ttc	tgc	ggt	ggc	cag	atc	cat	1008
Gly	Asp	Ser	Val	Trp	Asp	Gly	His	Gly	Phe	Ser	Val	Gly	Gln	Ile	His	
			325					330						335		
gag	acc	atg	aac	att	cgg	cga	ggg	att	atc	aca	gat	tgg	ttt	ttc	gga	1056
Glu	Thr	Met	Asn	Ile	Arg	Arg	Gly	Ile	Ile	Thr	Asp	Trp	Phe	Phe	Gly	
			340					345					350			
ggc	ttg	aac	tac	cag	atc	gag	cac	cat	ttg	tgg	cgc	acc	ctc	cct	cgc	1104





-continued

Leu Thr Ser Leu Leu Val Phe Phe Val Ser Glu Leu Val Gly Gly Phe  
 290 295 300  
 Gly Ile Ala Ile Val Val Phe Met Asn His Tyr Pro Leu Glu Lys Ile  
 305 310 315 320  
 Gly Asp Ser Val Trp Asp Gly His Gly Phe Ser Val Gly Gln Ile His  
 325 330 335  
 Glu Thr Met Asn Ile Arg Arg Gly Ile Ile Thr Asp Trp Phe Phe Gly  
 340 345 350  
 Gly Leu Asn Tyr Gln Ile Glu His His Leu Trp Pro Thr Leu Pro Arg  
 355 360 365  
 His Asn Leu Thr Ala Val Ser Tyr Gln Val Glu Gln Leu Cys Gln Lys  
 370 375 380  
 His Asn Leu Pro Tyr Arg Asn Pro Leu Pro His Glu Gly Leu Val Ile  
 385 390 395 400  
 Leu Leu Arg Tyr Leu Ala Val Phe Ala Arg Met Ala Glu Lys Gln Pro  
 405 410 415  
 Ala Gly Lys Ala Leu  
 420

<210> SEQ ID NO 3  
 <211> LENGTH: 1374  
 <212> TYPE: DNA  
 <213> ORGANISM: Acanthamoeba castellanii  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1374)  
 <223> OTHER INFORMATION: Delta-8-Desaturase

<400> SEQUENCE: 3

atg gtc ctc aca acc ccg gcc ctc aac ctg aag aag gaa cga acg tcg Met Val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Glu Arg Thr Ser 1 5 10 15	48
ttc acc cag gag gag ctt tcc aag ctc tgg gtc ctt cac ggc cag gtg Phe Thr Gln Glu Glu Leu Ser Lys Leu Trp Val Leu His Gly Gln Val 20 25 30	96
tac gat ttc acc gac ttt gtc aag tac cac ccg gcc ggc agc agg gcc Tyr Asp Phe Thr Asp Phe Val Lys Tyr His Pro Ala Gly Ser Arg Ala 35 40 45	144
atc ctg ctc ggc cgt ggc cgt gat tgt acc gtg ctc ttc gag tcc tac Ile Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Glu Ser Tyr 50 55 60	192
cac aca gtc ctg cct tcc gat gct ctt ctc gag aag tac cgc gtc tct His Thr Val Leu Pro Ser Asp Ala Leu Leu Glu Lys Tyr Arg Val Ser 65 70 75 80	240
gct ccc aac gcc aag ctc gag gag agc cgg tca gcc aag ctg ttc tcg Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser 85 90 95	288
ttc gag gag ggt agc ttc tac cga acc ctc aag cag cga acg cgc gag Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gln Arg Thr Arg Glu 100 105 110	336
tac ttc aag acc aac aac ctg agc acc aag gcc acc acg atg gag gtc Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val 115 120 125	384
atc tac ttc gtg gcc acc atc ctc agc atc tac ttc tgc acg tgg gcc Ile Tyr Phe Val Ala Thr Ile Leu Ser Ile Tyr Phe Cys Thr Trp Ala 130 135 140	432
gcc ttc gtg cag ggt tcc ctc atc gcc gct gtc ctt cac gga gtg ggc Ala Phe Val Gln Gly Ser Leu Ile Ala Ala Val Leu His Gly Val Gly 145 150 155 160	480

-continued

cgt gcg atc tgt atc ata caa ccg act cat gcg act tcg cac tac gcc	528
Arg Ala Ile Cys Ile Ile Gln Pro Thr His Ala Thr Ser His Tyr Ala	
165 170 175	
atg ttc cgc tca gtg tgg ctc aac cag tgg gcc tac agg atc tcc atg	576
Met Phe Arg Ser Val Trp Leu Asn Gln Trp Ala Tyr Arg Ile Ser Met	
180 185 190	
gcc gtc agc gga tcg tcg ccg gcc cag tgg acc acc aag cac gtc atc	624
Ala Val Ser Gly Ser Ser Pro Ala Gln Trp Thr Thr Lys His Val Ile	
195 200 205	
aac cat cac gtc gag acc aac ctg tgc ccc acc gat gac gac acc atg	672
Asn His His Val Glu Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met	
210 215 220	
tac ccc atc aag cgc atc ctg cac gag ttc cct cgt ctg ttc ttc cac	720
Tyr Pro Ile Lys Arg Ile Leu His Glu Phe Pro Arg Leu Phe Phe His	
225 230 235 240	
aag tac cag cac atc tac atc tgg ctg gtg tac ccc tac acc acc atc	768
Lys Tyr Gln His Ile Tyr Ile Trp Leu Val Tyr Pro Tyr Thr Thr Ile	
245 250 255	
ttg tgg cac ttc tcc aac ctg gcc aag ctc gcc ctc ggc gcc gct cgc	816
Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg	
260 265 270	
ggt cag atg tac gag ggt atc gcc aag gtg agc caa gag acc tcg ggt	864
Gly Gln Met Tyr Glu Gly Ile Ala Lys Val Ser Gln Glu Thr Ser Gly	
275 280 285	
gac tgg gtg gag acg gcc atg acg ctg ttc ttc ttc acg ttc tcc cgt	912
Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Phe Thr Phe Ser Arg	
290 295 300	
ctg ctg ctg ccc ttc ctg tgc ctg ccc ttc acc acg gcc gcc gcg gtg	960
Leu Leu Leu Pro Phe Leu Cys Leu Pro Phe Thr Thr Ala Ala Ala Val	
305 310 315 320	
ttc ctg ctc tcc gag tgg acc tgc tcg acc tgg ttc gcg ctg cag ttc	1008
Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gln Phe	
325 330 335	
gcc gtg agc cac gag gtc gac gag tgc gtc gag cac gag aag tcg gtc	1056
Ala Val Ser His Glu Val Asp Glu Cys Val Glu His Glu Lys Ser Val	
340 345 350	
ctc gac acc ctc aag gcc aac gag gcc aag ggc atc gtc aac cag ggc	1104
Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly Ile Val Asn Gln Gly	
355 360 365	
ggc ctc gtc gac tgg ggc gcg cac cag gtt cgg gcc tcg cac aac tac	1152
Gly Leu Val Asp Trp Gly Ala His Gln Val Arg Ala Ser His Asn Tyr	
370 375 380	
tct gcc gac tcc ctg ctg tcg ctc cac ttc agc ggt ggc ctc aac ctt	1200
Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu	
385 390 395 400	
cag atc gag cac cac ctc ttc ccc tcc gtc cac tac act cac tac cct	1248
Gln Ile Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro	
405 410 415	
gcc ccg tcc aag att gtg cag cag acg tgc aag gag ttc aac ttg ccc	1296
Ala Pro Ser Lys Ile Val Gln Gln Thr Cys Lys Glu Phe Asn Leu Pro	
420 425 430	
tgc act ctg tcg ccg tcg atg atg ggt gcc gtg acc aag cac tac cac	1344
Cys Thr Leu Ser Pro Ser Met Met Gly Ala Val Thr Lys His Tyr His	
435 440 445	
cag ctc aag aag atg ggt gct gag aac tga	1374
Gln Leu Lys Lys Met Gly Ala Glu Asn	
450 455	

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 457

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Acanthamoeba castellanii

&lt;400&gt; SEQUENCE: 4

Met Val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Glu Arg Thr Ser  
 1 5 10 15  
 Phe Thr Gln Glu Glu Leu Ser Lys Leu Trp Val Leu His Gly Gln Val  
 20 25 30  
 Tyr Asp Phe Thr Asp Phe Val Lys Tyr His Pro Ala Gly Ser Arg Ala  
 35 40 45  
 Ile Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Glu Ser Tyr  
 50 55 60  
 His Thr Val Leu Pro Ser Asp Ala Leu Leu Glu Lys Tyr Arg Val Ser  
 65 70 75 80  
 Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser  
 85 90 95  
 Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gln Arg Thr Arg Glu  
 100 105 110  
 Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val  
 115 120 125  
 Ile Tyr Phe Val Ala Thr Ile Leu Ser Ile Tyr Phe Cys Thr Trp Ala  
 130 135 140  
 Ala Phe Val Gln Gly Ser Leu Ile Ala Ala Val Leu His Gly Val Gly  
 145 150 155 160  
 Arg Ala Ile Cys Ile Ile Gln Pro Thr His Ala Thr Ser His Tyr Ala  
 165 170 175  
 Met Phe Arg Ser Val Trp Leu Asn Gln Trp Ala Tyr Arg Ile Ser Met  
 180 185 190  
 Ala Val Ser Gly Ser Ser Pro Ala Gln Trp Thr Thr Lys His Val Ile  
 195 200 205  
 Asn His His Val Glu Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met  
 210 215 220  
 Tyr Pro Ile Lys Arg Ile Leu His Glu Phe Pro Arg Leu Phe Phe His  
 225 230 235 240  
 Lys Tyr Gln His Ile Tyr Ile Trp Leu Val Tyr Pro Tyr Thr Thr Ile  
 245 250 255  
 Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg  
 260 265 270  
 Gly Gln Met Tyr Glu Gly Ile Ala Lys Val Ser Gln Glu Thr Ser Gly  
 275 280 285  
 Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Phe Thr Phe Ser Arg  
 290 295 300  
 Leu Leu Leu Pro Phe Leu Cys Leu Pro Phe Thr Thr Ala Ala Ala Val  
 305 310 315 320  
 Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gln Phe  
 325 330 335  
 Ala Val Ser His Glu Val Asp Glu Cys Val Glu His Glu Lys Ser Val  
 340 345 350  
 Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly Ile Val Asn Gln Gly  
 355 360 365  
 Gly Leu Val Asp Trp Gly Ala His Gln Val Arg Ala Ser His Asn Tyr  
 370 375 380  
 Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu  
 385 390 395 400  
 Gln Ile Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro

-continued

405										410					415				
Ala	Pro	Ser	Lys	Ile	Val	Gln	Gln	Thr	Cys	Lys	Glu	Phe	Asn	Leu	Pro				
			420					425					430						
Cys	Thr	Leu	Ser	Pro	Ser	Met	Met	Gly	Ala	Val	Thr	Lys	His	Tyr	His				
		435					440					445							
Gln	Leu	Lys	Lys	Met	Gly	Ala	Glu	Asn											
	450					455													
<210> SEQ ID NO 5																			
<211> LENGTH: 1374																			
<212> TYPE: DNA																			
<213> ORGANISM: Acanthamoeba castellanii																			
<220> FEATURE:																			
<221> NAME/KEY: CDS																			
<222> LOCATION: (1)..(1374)																			
<223> OTHER INFORMATION: Delta-8-Desaturase																			
<400> SEQUENCE: 5																			
atg	gtc	ctc	aca	acc	ccg	gcc	ctc	aac	ctg	aag	aag	gaa	cga	acg	tcg				48
Met	Val	Leu	Thr	Thr	Pro	Ala	Leu	Asn	Leu	Lys	Lys	Glu	Arg	Thr	Ser				
1			5					10						15					
ttc	acc	cag	gag	gag	ctt	tcc	aag	ctc	tgg	gtc	ctt	cac	ggc	cag	gtg				96
Phe	Thr	Gln	Glu	Glu	Leu	Ser	Lys	Leu	Trp	Val	Leu	His	Gly	Gln	Val				
			20					25					30						
tac	gat	ttc	acc	gac	ttt	gtc	aag	tac	cac	ccg	ggc	ggc	agc	agg	gcc				144
Tyr	Asp	Phe	Thr	Asp	Phe	Val	Lys	Tyr	His	Pro	Gly	Gly	Ser	Arg	Ala				
	35					40						45							
atc	ctg	ctc	ggc	cgt	ggc	cgt	gat	tgt	acc	gtg	ctc	ttc	gag	tcc	tac				192
Ile	Leu	Leu	Gly	Arg	Gly	Arg	Asp	Cys	Thr	Val	Leu	Phe	Glu	Ser	Tyr				
	50					55					60								
cac	aca	gtc	ctg	cct	tcc	gat	gct	ctt	ctc	gag	aag	tac	cgc	gtc	tct				240
His	Thr	Val	Leu	Pro	Ser	Asp	Ala	Leu	Leu	Glu	Lys	Tyr	Arg	Val	Ser				
	65				70					75					80				
gct	ccc	aac	gcc	aag	ctc	gag	gag	agc	cgg	tca	gcc	aag	ctg	ttc	tcg				288
Ala	Pro	Asn	Ala	Lys	Leu	Glu	Glu	Ser	Arg	Ser	Ala	Lys	Leu	Phe	Ser				
			85					90					95						
ttc	gag	gag	ggt	agc	ttc	tac	cga	acc	ctc	aag	cag	cga	acg	cgc	gag				336
Phe	Glu	Glu	Gly	Ser	Phe	Tyr	Arg	Thr	Leu	Lys	Gln	Arg	Thr	Arg	Glu				
			100					105					110						
tac	ttc	aag	acc	aac	aac	ctg	agc	acc	aag	gcc	acc	acg	atg	gag	gtc				384
Tyr	Phe	Lys	Thr	Asn	Asn	Leu	Ser	Thr	Lys	Ala	Thr	Thr	Met	Glu	Val				
		115				120						125							
atc	tac	ttc	gtg	gcc	acc	atc	ctc	agc	atc	tac	ttc	tgc	acg	tgg	gcc				432
Ile	Tyr	Phe	Val	Ala	Thr	Ile	Leu	Ser	Ile	Tyr	Phe	Cys	Thr	Trp	Ala				
	130					135						140							
gcc	ttc	gtg	cag	ggt	tcc	ctc	atc	gcc	gct	gtc	ctt	cac	gga	gtg	ggc				480
Ala	Phe	Val	Gln	Gly	Ser	Leu	Ile	Ala	Ala	Val	Leu	His	Gly	Val	Gly				
	145				150					155					160				
cgt	gcg	atc	tgt	atc	ata	caa	ccg	act	cat	gcg	act	tcg	cac	tac	gcc				528
Arg	Ala	Ile	Cys	Ile	Ile	Gln	Pro	Thr	His	Ala	Thr	Ser	His	Tyr	Ala				
			165					170						175					
atg	ttc	cgc	tca	gtg	tgg	ctc	aac	cag	tgg	gcc	tac	agg	atc	tcc	atg				576
Met	Phe	Arg	Ser	Val	Trp	Leu	Asn	Gln	Trp	Ala	Tyr	Arg	Ile	Ser	Met				
			180					185					190						
gcc	gtc	agc	gga	tcg	tcg	ccg	gcc	cag	tgg	acc	acc	aag	cac	gtc	atc				624
Ala	Val	Ser	Gly	Ser	Ser	Pro	Ala	Gln	Trp	Thr	Thr	Lys	His	Val	Ile				
		195				200						205							
aac	cat	cac	gtc	gag	acc	aac	ctg	tgc	ccc	acc	gat	gac	gac	acc	atg				672
Asn	His	His	Val	Glu	Thr	Asn	Leu	Cys	Pro	Thr	Asp	Asp	Asp	Thr	Met				
	210					215					220								
tac	ccc	atc	aag	cgc	atc	ctg	cac	gag	ttc	cct	cgt	ctg	ttc	ttc	cac				720

-continued

Tyr	Pro	Ile	Lys	Arg	Ile	Leu	His	Glu	Phe	Pro	Arg	Leu	Phe	Phe	His	
225					230					235					240	
aag	tac	cag	cac	atc	tac	atc	tgg	ctg	gtg	tac	ccc	tac	acc	acc	atc	768
Lys	Tyr	Gln	His	Ile	Tyr	Ile	Trp	Leu	Val	Tyr	Pro	Tyr	Thr	Thr	Ile	
			245						250					255		
ttg	tgg	cac	ttc	tcc	aac	ctg	gcc	aag	ctc	gcc	ctc	ggc	gcc	gct	cgc	816
Leu	Trp	His	Phe	Ser	Asn	Leu	Ala	Lys	Leu	Ala	Leu	Gly	Ala	Ala	Arg	
			260					265					270			
ggg	cag	atg	tac	gag	ggg	atc	gcc	aag	gtg	agc	caa	gag	acc	tcg	ggg	864
Gly	Gln	Met	Tyr	Glu	Gly	Ile	Ala	Lys	Val	Ser	Gln	Glu	Thr	Ser	Gly	
		275					280					285				
gac	tgg	gtg	gag	acg	gcc	atg	acg	ctg	ttc	ttc	ttc	acg	ttc	tcc	cgt	912
Asp	Trp	Val	Glu	Thr	Ala	Met	Thr	Leu	Phe	Phe	Phe	Thr	Phe	Ser	Arg	
	290					295					300					
ctg	ctg	ctg	ccc	ttc	ctg	tgc	ctg	ccc	ttc	acc	acg	gcc	gcc	gcg	gtg	960
Leu	Leu	Leu	Pro	Phe	Leu	Cys	Leu	Pro	Phe	Thr	Thr	Ala	Ala	Ala	Val	
305					310					315					320	
ttc	ctg	ctc	tcc	gag	tgg	acc	tgc	tcg	acc	tgg	ttc	gcg	ctg	cag	ttc	1008
Phe	Leu	Leu	Ser	Glu	Trp	Thr	Cys	Ser	Thr	Trp	Phe	Ala	Leu	Gln	Phe	
			325						330					335		
gcc	gtg	agc	cac	gag	gtc	gac	gag	tgc	gtc	gag	cac	gag	aag	tcg	gtc	1056
Ala	Val	Ser	His	Glu	Val	Asp	Glu	Cys	Val	Glu	His	Glu	Lys	Ser	Val	
			340					345					350			
ctc	gac	acc	ctc	aag	gcc	aac	gag	gcc	aag	ggc	atc	gtc	aac	cag	ggc	1104
Leu	Asp	Thr	Leu	Lys	Ala	Asn	Glu	Ala	Lys	Gly	Ile	Val	Asn	Gln	Gly	
		355				360						365				
ggc	ctc	gtc	gac	tgg	ggc	gcg	cac	cag	ggt	egg	gcc	tcg	cac	aac	tac	1152
Gly	Leu	Val	Asp	Trp	Gly	Ala	His	Gln	Val	Arg	Ala	Ser	His	Asn	Tyr	
	370				375						380					
tct	gcc	gac	tcc	ctg	ctg	tcg	ctc	cac	ttc	agc	ggg	ggc	ctc	aac	ctt	1200
Ser	Ala	Asp	Ser	Leu	Leu	Ser	Leu	His	Phe	Ser	Gly	Gly	Leu	Asn	Leu	
385					390					395					400	
cag	atc	gag	cac	cac	ctc	ttc	ccc	tcc	gtc	cac	tac	act	cac	tac	cct	1248
Gln	Ile	Glu	His	His	Leu	Phe	Pro	Ser	Val	His	Tyr	Thr	His	Tyr	Pro	
				405					410					415		
gcc	ccg	tcc	aag	att	gtg	cag	cag	acg	tgc	aag	gag	ttc	aac	ttg	ccc	1296
Ala	Pro	Ser	Lys	Ile	Val	Gln	Gln	Thr	Cys	Lys	Glu	Phe	Asn	Leu	Pro	
			420					425					430			
tgc	act	ctg	tcg	ccg	tcg	atg	atg	ggg	gcc	gtg	acc	aag	cac	tac	cac	1344
Cys	Thr	Leu	Ser	Pro	Ser	Met	Met	Gly	Ala	Val	Thr	Lys	His	Tyr	His	
		435					440					445				
cag	ctc	aag	aag	atg	ggg	gct	gag	aac	tga							1374
Gln	Leu	Lys	Lys	Met	Gly	Ala	Glu	Asn								
	450				455											

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 457

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Acanthamoeba castellanii

&lt;400&gt; SEQUENCE: 6

Met	Val	Leu	Thr	Thr	Pro	Ala	Leu	Asn	Leu	Lys	Lys	Glu	Arg	Thr	Ser	
1				5					10					15		
Phe	Thr	Gln	Glu	Glu	Leu	Ser	Lys	Leu	Trp	Val	Leu	His	Gly	Gln	Val	
			20					25					30			
Tyr	Asp	Phe	Thr	Asp	Phe	Val	Lys	Tyr	His	Pro	Gly	Gly	Ser	Arg	Ala	
		35					40					45				
Ile	Leu	Leu	Gly	Arg	Gly	Arg	Asp	Cys	Thr	Val	Leu	Phe	Glu	Ser	Tyr	
	50					55					60					
His	Thr	Val	Leu	Pro	Ser	Asp	Ala	Leu	Leu	Glu	Lys	Tyr	Arg	Val	Ser	

-continued

---

65	70	75	80
Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser	85	90	95
Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gln Arg Thr Arg Glu	100	105	110
Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val	115	120	125
Ile Tyr Phe Val Ala Thr Ile Leu Ser Ile Tyr Phe Cys Thr Trp Ala	130	135	140
Ala Phe Val Gln Gly Ser Leu Ile Ala Ala Val Leu His Gly Val Gly	145	150	155
Arg Ala Ile Cys Ile Ile Gln Pro Thr His Ala Thr Ser His Tyr Ala	165	170	175
Met Phe Arg Ser Val Trp Leu Asn Gln Trp Ala Tyr Arg Ile Ser Met	180	185	190
Ala Val Ser Gly Ser Ser Pro Ala Gln Trp Thr Thr Lys His Val Ile	195	200	205
Asn His His Val Glu Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met	210	215	220
Tyr Pro Ile Lys Arg Ile Leu His Glu Phe Pro Arg Leu Phe Phe His	225	230	235
Lys Tyr Gln His Ile Tyr Ile Trp Leu Val Tyr Pro Tyr Thr Thr Ile	245	250	255
Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg	260	265	270
Gly Gln Met Tyr Glu Gly Ile Ala Lys Val Ser Gln Glu Thr Ser Gly	275	280	285
Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Phe Thr Phe Ser Arg	290	295	300
Leu Leu Leu Pro Phe Leu Cys Leu Pro Phe Thr Thr Ala Ala Ala Val	305	310	315
Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gln Phe	325	330	335
Ala Val Ser His Glu Val Asp Glu Cys Val Glu His Glu Lys Ser Val	340	345	350
Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly Ile Val Asn Gln Gly	355	360	365
Gly Leu Val Asp Trp Gly Ala His Gln Val Arg Ala Ser His Asn Tyr	370	375	380
Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu	385	390	395
Gln Ile Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro	405	410	415
Ala Pro Ser Lys Ile Val Gln Gln Thr Cys Lys Glu Phe Asn Leu Pro	420	425	430
Cys Thr Leu Ser Pro Ser Met Met Gly Ala Val Thr Lys His Tyr His	435	440	445
Gln Leu Lys Lys Met Gly Ala Glu Asn	450	455	

<210> SEQ ID NO 7  
 <211> LENGTH: 1236  
 <212> TYPE: DNA  
 <213> ORGANISM: Perkinsus marinus  
 <220> FEATURE:

-continued

&lt;221&gt; NAME/KEY: CDS

&lt;222&gt; LOCATION: (1)..(1236)

&lt;223&gt; OTHER INFORMATION: Delta-8-Desaturase

&lt;400&gt; SEQUENCE: 7

```

atg tct tct ctt acc ctc tac aga ggc ccc ttt tcc cga atg gtg ctc      48
Met Ser Ser Leu Thr Leu Tyr Arg Gly Pro Phe Ser Arg Met Val Leu
1           5           10          15

cct cgt cag gaa atc tgc atc gat ggt cgc ata tac gat gtc act gag      96
Pro Arg Gln Glu Ile Cys Ile Asp Gly Arg Ile Tyr Asp Val Thr Glu
          20          25          30

ttc atc aat cgt cat cca ggt ggt aag att atc ctc ttc caa gtt ggt     144
Phe Ile Asn Arg His Pro Gly Gly Lys Ile Ile Leu Phe Gln Val Gly
          35          40          45

gct gat gcc act gat gct ttt cgt gag ttt cat gct ggc agt gag aag     192
Ala Asp Ala Thr Asp Ala Phe Arg Glu Phe His Ala Gly Ser Glu Lys
          50          55          60

gca gag aag atc ctc aaa acc cta cca tcc cgt gat gat gac ggt act     240
Ala Glu Lys Ile Leu Lys Thr Leu Pro Ser Arg Asp Asp Asp Gly Thr
          65          70          75          80

ttc ctt cct tca acc caa cgc tcc atc atg gat gat ttc aaa cgc cta     288
Phe Leu Pro Ser Thr Gln Arg Ser Ile Met Asp Asp Phe Lys Arg Leu
          85          90          95

aga gat gac ctc gtc agc aga ggt gtc ttc aag cca agc gtc atg cat     336
Arg Asp Asp Leu Val Ser Arg Gly Val Phe Lys Pro Ser Val Met His
          100         105         110

ggt gta tac cgc tgc ttg gaa gtc gtt gct ctc tat ctc att ggc ttc     384
Val Val Tyr Arg Cys Leu Glu Val Val Ala Leu Tyr Leu Ile Gly Phe
          115         120         125

tat ttg gct ctg tgc acc agt aat gtg tac gtt ggg tgt gct gta ctt     432
Tyr Leu Ala Leu Cys Thr Ser Asn Val Tyr Val Gly Cys Ala Val Leu
          130         135         140

ggt gta gct caa ggt cgt gct ggt tgg ttg atg cat gaa gga ggt cat     480
Gly Val Ala Gln Gly Arg Ala Gly Trp Leu Met His Glu Gly Gly His
          145         150         155         160

cac tct ctg act ggt aac tgg aaa gtt gac cag ttc ctc caa gaa cta     528
His Ser Leu Thr Gly Asn Trp Lys Val Asp Gln Phe Leu Gln Glu Leu
          165         170         175

ttt ttc ggc att ggt tgt ggt atg tca gct gcg tgg tgg cgc aat gca     576
Phe Phe Gly Ile Gly Cys Gly Met Ser Ala Ala Trp Trp Arg Asn Ala
          180         185         190

cac aac aag cat cac gct gct cct cag cat tta ggg aaa gat gtt gat     624
His Asn Lys His His Ala Ala Pro Gln His Leu Gly Lys Asp Val Asp
          195         200         205

ctc gag aca ttg cct ctg gtc gcc ttc aat aag gcc gta ctt cga ggc     672
Leu Glu Thr Leu Pro Leu Val Ala Phe Asn Lys Ala Val Leu Arg Gly
          210         215         220

cgt cta ccg tct gtc tgg atc aga tca caa gct gtg tgc ttt gca ccg     720
Arg Leu Pro Ser Val Trp Ile Arg Ser Gln Ala Val Cys Phe Ala Pro
          225         230         235         240

ata tca aca cta ctg gta tcg ttc ttt tgg caa ttc tac cta cac ccg     768
Ile Ser Thr Leu Leu Val Ser Phe Phe Trp Gln Phe Tyr Leu His Pro
          245         250         255

agg cat att att agg aca ggt cga cga atg gag tct ttc tgg cta ctc     816
Arg His Ile Ile Arg Thr Gly Arg Arg Met Glu Ser Phe Trp Leu Leu
          260         265         270

gta cgc tac tta gtt att gtg tac ctc ggg ttc agc tat gga ttg gta     864
Val Arg Tyr Leu Val Ile Val Tyr Leu Gly Phe Ser Tyr Gly Leu Val
          275         280         285

tcg gtc ttg tta tgt tac atc gca agt gtg cat gtt ggt ggt atg tac     912
Ser Val Leu Leu Cys Tyr Ile Ala Ser Val His Val Gly Gly Met Tyr

```

-continued

290	295	300	
atc ttt gta cac ttc gct cta tca cat aca cat tta cct gtc att aac			960
Ile Phe Val His Phe Ala Leu Ser His Thr His Leu Pro Val Ile Asn			
305	310	315	320
cag cat ggt aga gct aac tgg ttg gaa tac gca tct aag cac aca gtt			1008
Gln His Gly Arg Ala Asn Trp Leu Glu Tyr Ala Ser Lys His Thr Val			
	325	330	335
aat gtg tca act aac aat tat ttc gtc aca tgg ctc atg agt tat ttg			1056
Asn Val Ser Thr Asn Asn Tyr Phe Val Thr Trp Leu Met Ser Tyr Leu			
	340	345	350
aat tat caa ata gag cat cat ctc ttc ccg tca tgt ccc cag ttt aga			1104
Asn Tyr Gln Ile Glu His His Leu Phe Pro Ser Cys Pro Gln Phe Arg			
	355	360	365
ttc cct ggt tac gtc agt atg agg gtt cga gaa ttt ttt cat aag cat			1152
Phe Pro Gly Tyr Val Ser Met Arg Val Arg Glu Phe Phe His Lys His			
	370	375	380
gga ttg aag tat aac gag gtc ggc tat cta cat gca ctc aat ctc aca			1200
Gly Leu Lys Tyr Asn Glu Val Gly Tyr Leu His Ala Leu Asn Leu Thr			
	385	390	400
ttt tca aat ctg gct gct gtt gcc ata gtg gaa tag			1236
Phe Ser Asn Leu Ala Val Ala Ile Val Glu			
	405	410	

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 411

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Perkinsus marinus

&lt;400&gt; SEQUENCE: 8

Met Ser Ser Leu Thr Leu Tyr Arg Gly Pro Phe Ser Arg Met Val Leu			
1	5	10	15
Pro Arg Gln Glu Ile Cys Ile Asp Gly Arg Ile Tyr Asp Val Thr Glu			
	20	25	30
Phe Ile Asn Arg His Pro Gly Gly Lys Ile Ile Leu Phe Gln Val Gly			
	35	40	45
Ala Asp Ala Thr Asp Ala Phe Arg Glu Phe His Ala Gly Ser Glu Lys			
	50	55	60
Ala Glu Lys Ile Leu Lys Thr Leu Pro Ser Arg Asp Asp Asp Gly Thr			
	65	70	75
Phe Leu Pro Ser Thr Gln Arg Ser Ile Met Asp Asp Phe Lys Arg Leu			
	85	90	95
Arg Asp Asp Leu Val Ser Arg Gly Val Phe Lys Pro Ser Val Met His			
	100	105	110
Val Val Tyr Arg Cys Leu Glu Val Val Ala Leu Tyr Leu Ile Gly Phe			
	115	120	125
Tyr Leu Ala Leu Cys Thr Ser Asn Val Tyr Val Gly Cys Ala Val Leu			
	130	135	140
Gly Val Ala Gln Gly Arg Ala Gly Trp Leu Met His Glu Gly Gly His			
	145	150	155
His Ser Leu Thr Gly Asn Trp Lys Val Asp Gln Phe Leu Gln Glu Leu			
	165	170	175
Phe Phe Gly Ile Gly Cys Gly Met Ser Ala Ala Trp Trp Arg Asn Ala			
	180	185	190
His Asn Lys His His Ala Ala Pro Gln His Leu Gly Lys Asp Val Asp			
	195	200	205
Leu Glu Thr Leu Pro Leu Val Ala Phe Asn Lys Ala Val Leu Arg Gly			
	210	215	220



-continued

---

Arg Leu Pro Ser Val Trp Ile Arg Ser Gln Ala Val Cys Phe Ala Pro  
 225 230 235 240

Ile Ser Thr Leu Leu Val Ser Phe Phe Trp Gln Phe Tyr Leu His Pro  
 245 250 255

Arg His Ile Ile Arg Thr Gly Arg Arg Met Glu Ser Phe Trp Leu Leu  
 260 265 270

Val Arg Tyr Leu Val Ile Val Tyr Leu Gly Phe Ser Tyr Gly Leu Val  
 275 280 285

Ser Val Leu Leu Cys Tyr Ile Ala Ser Val His Val Gly Gly Met Tyr  
 290 295 300

Ile Phe Val His Phe Ala Leu Ser His Thr His Leu Pro Val Ile Asn  
 305 310 315 320

Gln His Gly Arg Ala Asn Trp Leu Glu Tyr Ala Ser Lys His Thr Val  
 325 330 335

Asn Val Ser Thr Asn Asn Tyr Phe Val Thr Trp Leu Met Ser Tyr Leu  
 340 345 350

Asn Tyr Gln Ile Glu His His Leu Phe Pro Ser Cys Pro Gln Phe Arg  
 355 360 365

Phe Pro Gly Tyr Val Ser Met Arg Val Arg Glu Phe Phe His Lys His  
 370 375 380

Gly Leu Lys Tyr Asn Glu Val Gly Tyr Leu His Ala Leu Asn Leu Thr  
 385 390 395 400

Phe Ser Asn Leu Ala Ala Val Ala Ile Val Glu  
 405 410

<210> SEQ ID NO 9  
 <211> LENGTH: 777  
 <212> TYPE: DNA  
 <213> ORGANISM: Isochrysis galbana  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(777)  
 <223> OTHER INFORMATION: Delta-9-Elongase

<400> SEQUENCE: 9

atg gcc ctc gca aac gac gcg gga gag cgc atc tgg gcg gct gtg acc	48
Met Ala Leu Ala Asn Asp Ala Gly Glu Arg Ile Trp Ala Ala Val Thr	
1 5 10 15	
gac ccg gaa atc ctc att ggc acc ttc tcg tac ttg cta ctc aaa ccg	96
Asp Pro Glu Ile Leu Ile Gly Thr Phe Ser Tyr Leu Leu Leu Lys Pro	
20 25 30	
ctg ctc cgc aat tcc ggg ctg gtg gat gag aag aag ggc gca tac agg	144
Leu Leu Arg Asn Ser Gly Leu Val Asp Glu Lys Lys Gly Ala Tyr Arg	
35 40 45	
acg tcc atg atc tgg tac aac gtt ctg ctg gcg ctc ttc tct gcg ctg	192
Thr Ser Met Ile Trp Tyr Asn Val Leu Leu Ala Leu Phe Ser Ala Leu	
50 55 60	
agc ttc tac gtg acg gcg acc gcc ctc ggc tgg gac tat ggt acg ggc	240
Ser Phe Tyr Val Thr Ala Thr Ala Leu Gly Trp Asp Tyr Gly Thr Gly	
65 70 75 80	
gcg tgg ctg cgc agg caa acc ggc gac aca ccg cag ccg ctc ttc cag	288
Ala Trp Leu Arg Arg Gln Thr Gly Asp Thr Pro Gln Pro Leu Phe Gln	
85 90 95	
tgc ccg tcc ccg gtt tgg gac tcg aag ctc ttc aca tgg acc gcc aag	336
Cys Pro Ser Pro Val Trp Asp Ser Lys Leu Phe Thr Trp Thr Ala Lys	
100 105 110	
gca ttc tat tac tcc aag tac gtg gag tac ctc gac acg gcc tgg ctg	384
Ala Phe Tyr Tyr Ser Lys Tyr Val Glu Tyr Leu Asp Thr Ala Trp Leu	
115 120 125	

-continued

---

```

agg gtc tcc ttt ctc cag gcc ttc cac cac ttt ggc gcg ccg tgg gat 432
Arg Val Ser Phe Leu Gln Ala Phe His His Phe Gly Ala Pro Trp Asp
130 135 140

gtg tac ctc ggc att cgg ctg cac aac gag ggc gta tgg atc ttc atg 480
Val Tyr Leu Gly Ile Arg Leu His Asn Glu Gly Val Trp Ile Phe Met
145 150 155 160

ttt ttc aac tcg ttc att cac acc atc atg tac acc tac tac ggc ctc 528
Phe Phe Asn Ser Phe Ile His Thr Ile Met Tyr Thr Tyr Tyr Gly Leu
165 170 175

acc gcc gcc ggg tat aag ttc aag gcc aag ccg ctc atc acc gcg atg 576
Thr Ala Ala Gly Tyr Lys Phe Lys Ala Lys Pro Leu Ile Thr Ala Met
180 185 190

cag atc tgc cag ttc gtg ggc gcc ttc ctg ttg gtc tgg gac tac atc 624
Gln Ile Cys Gln Phe Val Gly Gly Phe Leu Leu Val Trp Asp Tyr Ile
195 200 205

aac gtc ccc tgc ttc aac tcg gac aaa ggg aag ttg ttc agc tgg gct 672
Asn Val Pro Cys Phe Asn Ser Asp Lys Gly Lys Leu Phe Ser Trp Ala
210 215 220

ttc aac tat gca tac gtc ggc tcg gtc ttc ttg ctc ttc tgc cac ttt 720
Phe Asn Tyr Ala Tyr Val Gly Ser Val Phe Leu Leu Phe Cys His Phe
225 230 235 240

ttc tac cag gac aac ttg gca acg aag aaa tcg gcc aag gcg ggc aag 768
Phe Tyr Gln Asp Asn Leu Ala Thr Lys Lys Ser Ala Lys Ala Gly Lys
245 250 255

cag ctc tag 777
Gln Leu

```

```

<210> SEQ ID NO 10
<211> LENGTH: 258
<212> TYPE: PRT
<213> ORGANISM: Isochrysis galbana

<400> SEQUENCE: 10

```

```

Met Ala Leu Ala Asn Asp Ala Gly Glu Arg Ile Trp Ala Ala Val Thr
1 5 10 15

Asp Pro Glu Ile Leu Ile Gly Thr Phe Ser Tyr Leu Leu Leu Lys Pro
20 25 30

Leu Leu Arg Asn Ser Gly Leu Val Asp Glu Lys Lys Gly Ala Tyr Arg
35 40 45

Thr Ser Met Ile Trp Tyr Asn Val Leu Leu Ala Leu Phe Ser Ala Leu
50 55 60

Ser Phe Tyr Val Thr Ala Thr Ala Leu Gly Trp Asp Tyr Gly Thr Gly
65 70 75 80

Ala Trp Leu Arg Arg Gln Thr Gly Asp Thr Pro Gln Pro Leu Phe Gln
85 90 95

Cys Pro Ser Pro Val Trp Asp Ser Lys Leu Phe Thr Trp Thr Ala Lys
100 105 110

Ala Phe Tyr Tyr Ser Lys Tyr Val Glu Tyr Leu Asp Thr Ala Trp Leu
115 120 125

Arg Val Ser Phe Leu Gln Ala Phe His His Phe Gly Ala Pro Trp Asp
130 135 140

Val Tyr Leu Gly Ile Arg Leu His Asn Glu Gly Val Trp Ile Phe Met
145 150 155 160

Phe Phe Asn Ser Phe Ile His Thr Ile Met Tyr Thr Tyr Tyr Gly Leu
165 170 175

Thr Ala Ala Gly Tyr Lys Phe Lys Ala Lys Pro Leu Ile Thr Ala Met
180 185 190

Gln Ile Cys Gln Phe Val Gly Gly Phe Leu Leu Val Trp Asp Tyr Ile

```



-continued

```

cag att ttg caa atg gtg ctc ggc gtg gcc atc atc acc atc tcc gcc      720
Gln Ile Leu Gln Met Val Leu Gly Val Ala Ile Ile Thr Ile Ser Ala
225                               230                               235                               240

ggc tgc gag aag gtg gac ccc atc gga acg acc ttc ggc tac ttt att      768
Gly Cys Glu Lys Val Asp Pro Ile Gly Thr Thr Phe Gly Tyr Phe Ile
                               245                               250                               255

tat ttc tcg ttc ttc gtc ctc ttc tgc aag ttc ttc tac tac cgc tac      816
Tyr Phe Ser Phe Phe Val Leu Phe Cys Lys Phe Phe Tyr Tyr Arg Tyr
                               260                               265                               270

atc gcc acg ccc gcc aag aag ccc gag gcc gcc gcc aag tgc cca gcc      864
Ile Ala Thr Pro Ala Lys Lys Pro Glu Ala Ala Ala Lys Ser Pro Ala
                               275                               280                               285

acc aag ccc aag agg aag cac gac taa                                  891
Thr Lys Pro Lys Arg Lys His Asp
                               290                               295

```

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 296

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Acanthamoeba castellanii

&lt;400&gt; SEQUENCE: 12

```

Met Ala Ala Ala Thr Ala Thr Thr Ala Thr Thr Ala Val Met Glu Gln
1           5           10           15

Val Pro Ile Thr Glu Ala Ile Phe Arg Pro Asp Leu Trp Val Gly Arg
20           25           30

Asp Gln Trp Glu Ala Asn Ala Val Ser Phe Val Trp Arg Tyr Trp Trp
35           40           45

Phe Phe Leu Val Met Gly Val Ala Tyr Leu Pro Ile Ile Phe Gly Leu
50           55           60

Lys Tyr Trp Met Lys Asp Arg Pro Ala Phe Asn Leu Arg Arg Pro Leu
65           70           75           80

Ile Leu Trp Asn Ile Phe Met Ala Thr Phe Ser Thr Ala Gly Phe Leu
85           90           95

Ser Ile Val Tyr Pro Leu Ile Glu Asn Trp Val Tyr Pro Gly Gly Gly
100          105          110

Leu Thr Pro His Glu Phe Ile Cys Ser Ala Ser Tyr Ser Tyr Lys Phe
115          120          125

Gly Asp Cys Ala Ile Trp Val Phe Leu Phe Asn Met Ser Lys Ile Leu
130          135          140

Glu Phe Val Asp Thr Ile Phe Ile Val Pro Arg Lys Thr His Leu Gly
145          150          155          160

Phe Leu His Tyr Tyr His His Ile Ile Thr Tyr Ser Phe Cys Leu Tyr
165          170          175

Ala Gly Gln Tyr Met His His Tyr Asn Cys Gly Gly Tyr Phe Phe Cys
180          185          190

Leu Met Asn Phe Phe Val His Gly Ile Met Tyr Phe Tyr Tyr Ala Leu
195          200          205

Arg Ser Met Gly Phe Arg Pro Ser Phe Asp Ile Gly Ile Thr Phe Leu
210          215          220

Gln Ile Leu Gln Met Val Leu Gly Val Ala Ile Ile Thr Ile Ser Ala
225          230          235          240

Gly Cys Glu Lys Val Asp Pro Ile Gly Thr Thr Phe Gly Tyr Phe Ile
245          250          255

Tyr Phe Ser Phe Phe Val Leu Phe Cys Lys Phe Phe Tyr Tyr Arg Tyr
260          265          270

```

-continued

Ile Ala Thr Pro Ala Lys Lys Pro Glu Ala Ala Ala Lys Ser Pro Ala  
 275 280 285

Thr Lys Pro Lys Arg Lys His Asp  
 290 295

<210> SEQ ID NO 13  
 <211> LENGTH: 1320  
 <212> TYPE: DNA  
 <213> ORGANISM: Thraustrochytrium sp.  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1320)  
 <223> OTHER INFORMATION: Delta-5-Desaturase

<400> SEQUENCE: 13

atg ggc aag ggc agc gag ggc cgc agc gcg gcg cgc gag atg acg gcc Met Gly Lys Gly Ser Glu Gly Arg Ser Ala Ala Arg Glu Met Thr Ala 1 5 10 15	48
gag gcg aac ggc gac aag cgg aaa acg att ctg atc gag ggc gtc ctg Glu Ala Asn Gly Asp Lys Arg Lys Thr Ile Leu Ile Glu Gly Val Leu 20 25 30	96
tac gac gcg acg aac ttt aag cac ccg ggc ggt tcg atc atc aac ttc Tyr Asp Ala Thr Asn Phe Lys His Pro Gly Gly Ser Ile Ile Asn Phe 35 40 45	144
ttg acc gag ggc gag gcc ggc gtg gac gcg acg cag gcg tac cgc gag Leu Thr Glu Gly Glu Ala Gly Val Asp Ala Thr Gln Ala Tyr Arg Glu 50 55 60	192
ttt cat cag cgg tcc ggc aag gcc gac aag tac ctc aag tcg ctg ccg Phe His Gln Arg Ser Gly Lys Ala Asp Lys Tyr Leu Lys Ser Leu Pro 65 70 75 80	240
aag ctg gat gcg tcc aag gtg gag tcg cgg ttc tcg gcc aaa gag cag Lys Leu Asp Ala Ser Lys Val Glu Ser Arg Phe Ser Ala Lys Glu Gln 85 90 95	288
gcg cgg cgc gac gcc atg acg cgc gac tac gcg gcc ttt cgc gag gag Ala Arg Arg Asp Ala Met Thr Arg Asp Tyr Ala Ala Phe Arg Glu Glu 100 105 110	336
ctc gtc gcc gag ggg tac ttt gac ccg tcg atc ccg cac atg att tac Leu Val Ala Glu Gly Tyr Phe Asp Pro Ser Ile Pro His Met Ile Tyr 115 120 125	384
cgc gtc gtg gag atc gtg gcg ctc ttc gcg ctc tcg ttc tgg ctc atg Arg Val Val Glu Ile Val Ala Leu Phe Ala Leu Ser Phe Trp Leu Met 130 135 140	432
tcc aag gcc tcg ccc acc tcg ctc gtg ctg gcc gtg gtg atg aac ggc Ser Lys Ala Ser Pro Thr Ser Leu Val Leu Gly Val Val Met Asn Gly 145 150 155 160	480
att gcg cag ggc cgc tgc ggc tgg gtc atg cac gag atg ggc cac ggg Ile Ala Gln Gly Arg Cys Gly Trp Val Met His Glu Met Gly His Gly 165 170 175	528
tcg ttc acg ggc gtc atc tgg ctc gac gac cgg atg tgc gag ttc ttc Ser Phe Thr Gly Val Ile Trp Leu Asp Asp Arg Met Cys Glu Phe Phe 180 185 190	576
tac gcc gtc gcc tgc gcc atg agc ggg cac tac tgg aag aac cag cac Tyr Gly Val Gly Cys Gly Met Ser Gly His Tyr Trp Lys Asn Gln His 195 200 205	624
agc aag cac cac gcc gcg ccc aac cgc ctc gag cac gat gtc gat ctc Ser Lys His His Ala Ala Pro Asn Arg Leu Glu His Asp Val Asp Leu 210 215 220	672
aac acg ctg ccc ctg gtc gcc ttt aac gag cgc gtc gtg cgc aag gtc Asn Thr Leu Pro Leu Val Ala Phe Asn Glu Arg Val Val Arg Lys Val 225 230 235 240	720
aag ccg gga tcg ctg ctg gcg ctc tgg ctg cgc gtg cag gcg tac ctc Lys Pro Gly Ser Leu Leu Ala Leu Trp Leu Arg Val Gln Ala Tyr Leu	768

-continued

245	250	255	
ttt gcg ccc gtc tcg tgc ctg ctc atc ggc ctt ggc tgg acg ctc tac Phe Ala Pro Val Ser Cys Leu Leu Ile Gly Leu Gly Trp Thr Leu Tyr			816
260	265	270	
ctg cac ccg cgc tac atg ctg cgc acc aag cgg cac atg gag ttc gtc Leu His Pro Arg Tyr Met Leu Arg Thr Lys Arg His Met Glu Phe Val			864
275	280	285	
tgg atc ttc gcg cgc tac att ggc tgg ttc tcg ctc atg ggc gct ctc Trp Ile Phe Ala Arg Tyr Ile Gly Trp Phe Ser Leu Met Gly Ala Leu			912
290	295	300	
ggc tac tcg ccg ggc acc tcg gtc ggg atg tac ctg tgc tcg ttc ggc Gly Tyr Ser Pro Gly Thr Ser Val Gly Met Tyr Leu Cys Ser Phe Gly			960
305	310	315	320
ctc ggc tgc att tac att ttc ctg cag ttc gcc gtc agc cac acg cac Leu Gly Cys Ile Tyr Ile Phe Leu Gln Phe Ala Val Ser His Thr His			1008
325	330	335	
ctg ccg gtg acc aac ccg gag gac cag ctg cac tgg ctc gag tac gcg Leu Pro Val Thr Asn Pro Glu Asp Gln Leu His Trp Leu Glu Tyr Ala			1056
340	345	350	
gcc gac cac acg gtg aac att agc acc aag tcc tgg ctc gtc acg tgg Ala Asp His Thr Val Asn Ile Ser Thr Lys Ser Trp Leu Val Thr Trp			1104
355	360	365	
tgg atg tcg aac ctg aac ttt cag atc gag cac cac ctc ttc ccc acg Trp Met Ser Asn Leu Asn Phe Gln Ile Glu His His Leu Phe Pro Thr			1152
370	375	380	
gcg ccg cag ttc cgc ttc aag gaa atc agt cct cgc gtc gag gcc ctc Ala Pro Gln Phe Arg Phe Lys Glu Ile Ser Pro Arg Val Glu Ala Leu			1200
385	390	395	400
ttc aag cgc cac aac ctc ccg tac tac gac ctg ccc tac acg agc gcg Phe Lys Arg His Asn Leu Pro Tyr Tyr Asp Leu Pro Tyr Thr Ser Ala			1248
405	410	415	
gtc tcg acc acc ttt gcc aat ctt tat tcc gtc ggc cac tcg gtc ggc Val Ser Thr Thr Phe Ala Asn Leu Tyr Ser Val Gly His Ser Val Gly			1296
420	425	430	
gcc gac acc aag aag cag gac tga Ala Asp Thr Lys Lys Gln Asp			1320
435			

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 439

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Thraustrochytrium sp.

&lt;400&gt; SEQUENCE: 14

Met Gly Lys Gly Ser Glu Gly Arg Ser Ala Ala Arg Glu Met Thr Ala  
1 5 10 15Glu Ala Asn Gly Asp Lys Arg Lys Thr Ile Leu Ile Glu Gly Val Leu  
20 25 30Tyr Asp Ala Thr Asn Phe Lys His Pro Gly Gly Ser Ile Ile Asn Phe  
35 40 45Leu Thr Glu Gly Glu Ala Gly Val Asp Ala Thr Gln Ala Tyr Arg Glu  
50 55 60Phe His Gln Arg Ser Gly Lys Ala Asp Lys Tyr Leu Lys Ser Leu Pro  
65 70 75 80Lys Leu Asp Ala Ser Lys Val Glu Ser Arg Phe Ser Ala Lys Glu Gln  
85 90 95Ala Arg Arg Asp Ala Met Thr Arg Asp Tyr Ala Ala Phe Arg Glu Glu  
100 105 110

Leu Val Ala Glu Gly Tyr Phe Asp Pro Ser Ile Pro His Met Ile Tyr

-continued

---

115	120	125			
Arg Val Val Glu Ile Val Ala Leu Phe Ala Leu Ser Phe Trp Leu Met					
130	135	140			
Ser Lys Ala Ser Pro Thr Ser Leu Val Leu Gly Val Val Met Asn Gly					
145	150	155			160
Ile Ala Gln Gly Arg Cys Gly Trp Val Met His Glu Met Gly His Gly					
	165	170			175
Ser Phe Thr Gly Val Ile Trp Leu Asp Asp Arg Met Cys Glu Phe Phe					
	180	185			190
Tyr Gly Val Gly Cys Gly Met Ser Gly His Tyr Trp Lys Asn Gln His					
	195	200			205
Ser Lys His His Ala Ala Pro Asn Arg Leu Glu His Asp Val Asp Leu					
	210	215			220
Asn Thr Leu Pro Leu Val Ala Phe Asn Glu Arg Val Val Arg Lys Val					
	225	230			240
Lys Pro Gly Ser Leu Leu Ala Leu Trp Leu Arg Val Gln Ala Tyr Leu					
	245	250			255
Phe Ala Pro Val Ser Cys Leu Leu Ile Gly Leu Gly Trp Thr Leu Tyr					
	260	265			270
Leu His Pro Arg Tyr Met Leu Arg Thr Lys Arg His Met Glu Phe Val					
	275	280			285
Trp Ile Phe Ala Arg Tyr Ile Gly Trp Phe Ser Leu Met Gly Ala Leu					
	290	295			300
Gly Tyr Ser Pro Gly Thr Ser Val Gly Met Tyr Leu Cys Ser Phe Gly					
	305	310			320
Leu Gly Cys Ile Tyr Ile Phe Leu Gln Phe Ala Val Ser His Thr His					
	325	330			335
Leu Pro Val Thr Asn Pro Glu Asp Gln Leu His Trp Leu Glu Tyr Ala					
	340	345			350
Ala Asp His Thr Val Asn Ile Ser Thr Lys Ser Trp Leu Val Thr Trp					
	355	360			365
Trp Met Ser Asn Leu Asn Phe Gln Ile Glu His His Leu Phe Pro Thr					
	370	375			380
Ala Pro Gln Phe Arg Phe Lys Glu Ile Ser Pro Arg Val Glu Ala Leu					
	385	390			395
Phe Lys Arg His Asn Leu Pro Tyr Tyr Asp Leu Pro Tyr Thr Ser Ala					
	405	410			415
Val Ser Thr Thr Phe Ala Asn Leu Tyr Ser Val Gly His Ser Val Gly					
	420	425			430
Ala Asp Thr Lys Lys Gln Asp					
	435				

<210> SEQ ID NO 15  
 <211> LENGTH: 1353  
 <212> TYPE: DNA  
 <213> ORGANISM: Acanthamoeba castellanii  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1353)  
 <223> OTHER INFORMATION: Delta-5-Desaturase

<400> SEQUENCE: 15

atg gcc acc gca tct gca tcc aac gtt ctc cgc ctg ccc gga gag gga	48
Met Ala Thr Ala Ser Ala Ser Asn Val Leu Arg Leu Pro Gly Glu Gly	
1	5 10 15

ctc gcg act ggc ctc gag cag ctc gag tgg gcc gaa gtg cag aag cac	96
Leu Ala Thr Gly Leu Glu Gln Leu Glu Trp Ala Glu Val Gln Lys His	

-continued

20				25				30								
aac	acg	cgc	gag	agc	tcg	tgg	ctg	gtg	att	aac	gac	cag	gtg	tac	gac	144
Asn	Thr	Arg	Glu	Ser	Ser	Trp	Leu	Val	Ile	Asn	Asp	Gln	Val	Tyr	Asp	
		35				40					45					
atc	acc	aac	ttc	ggc	cgg	cgc	cat	ccc	ggt	ggc	aag	gta	atc	tac	cac	192
Ile	Thr	Asn	Phe	Gly	Arg	Arg	His	Pro	Gly	Gly	Lys	Val	Ile	Tyr	His	
		50				55					60					
tac	gcg	ggt	caa	gat	gcc	acg	gac	tcg	ttt	cgg	gct	ctt	cac	ccc	gat	240
Tyr	Ala	Gly	Gln	Asp	Ala	Thr	Asp	Ser	Phe	Arg	Ala	Leu	His	Pro	Asp	
		65				70					75				80	
tcc	gcc	ctg	gtg	atg	aag	tat	ctc	aag	ccc	ctc	ctc	atc	ggt	caa	gtg	288
Ser	Ala	Leu	Val	Met	Lys	Tyr	Leu	Lys	Pro	Leu	Leu	Ile	Gly	Gln	Val	
			85								90				95	
gca	ccc	ggc	tca	tcc	acc	gca	gca	tcg	att	gtt	gat	ggc	gcc	cgc	ccg	336
Ala	Pro	Gly	Ser	Ser	Thr	Ala	Ala	Ser	Ile	Val	Asp	Gly	Ala	Arg	Pro	
			100								105				110	
gcg	ccc	tcg	gca	ttc	gta	gag	gaa	ttc	aga	cag	gtg	cgc	aaa	gaa	ttc	384
Ala	Pro	Ser	Ala	Phe	Val	Glu	Glu	Phe	Arg	Gln	Val	Arg	Lys	Glu	Phe	
			115				120								125	
gag	gag	cag	ggc	ctg	ttc	gag	gcc	agc	tgg	tcc	ttc	ttc	ttc	ggg	atg	432
Glu	Glu	Gln	Gly	Leu	Phe	Glu	Ala	Ser	Trp	Ser	Phe	Phe	Phe	Gly	Met	
			130				135								140	
ctg	gcc	cac	atc	ttc	ctg	ctc	gag	gct	gcc	gcc	tac	tac	agc	atc	aag	480
Leu	Ala	His	Ile	Phe	Leu	Leu	Glu	Ala	Ala	Ala	Tyr	Tyr	Ser	Ile	Lys	
			145			150					155				160	
ctg	ctg	ggc	aac	agt	tgg	ccc	gtc	tac	ctc	ctc	gcc	gtc	ggc	ctc	ctc	528
Leu	Leu	Gly	Asn	Ser	Trp	Pro	Val	Tyr	Leu	Leu	Ala	Val	Gly	Leu	Leu	
			165								170				175	
gcc	act	gcc	cag	gca	cag	gcc	ggc	tgg	ctc	cag	cac	gat	tgt	ggg	cac	576
Ala	Thr	Ala	Gln	Ala	Gln	Ala	Gly	Trp	Leu	Gln	His	Asp	Cys	Gly	His	
			180												190	
ttg	tcc	gtg	ttc	aag	aag	tcg	aag	tgg	aac	cat	tgg	atg	cac	tac	atc	624
Leu	Ser	Val	Phe	Lys	Lys	Ser	Lys	Trp	Asn	His	Trp	Met	His	Tyr	Ile	
			195				200								205	
gtc	atc	tgc	cac	atc	aag	ggc	gcc	tcg	cga	gcc	tgg	tgg	aac	tgg	cgt	672
Val	Ile	Cys	His	Ile	Lys	Gly	Ala	Ser	Arg	Ala	Trp	Trp	Asn	Trp	Arg	
			210				215								220	
cac	ttt	gag	cac	cac	gca	aag	ccc	aac	gtg	gtg	cgc	aag	gac	ccc	gac	720
His	Phe	Glu	His	His	Ala	Lys	Pro	Asn	Val	Val	Arg	Lys	Asp	Pro	Asp	
			225			230					235				240	
atc	acc	ttc	ccc	aac	ctc	ttc	ctt	ctc	ggc	gac	cac	ctg	acg	cgc	aag	768
Ile	Thr	Phe	Pro	Asn	Leu	Phe	Leu	Leu	Gly	Asp	His	Leu	Thr	Arg	Lys	
			245								250				255	
tgg	gcc	aag	gcc	aag	aag	gga	gtg	atg	ccc	tac	aac	aag	cag	cac	ctc	816
Trp	Ala	Lys	Ala	Lys	Lys	Gly	Val	Met	Pro	Tyr	Asn	Lys	Gln	His	Leu	
			260								265				270	
tac	tgg	tgg	gct	ttc	ccc	ccg	ctc	ctg	ctg	ccc	gtc	tac	ttc	cac	tac	864
Tyr	Trp	Trp	Ala	Phe	Pro	Pro	Leu	Leu	Leu	Pro	Val	Tyr	Phe	His	Tyr	
			275				280								285	
gac	aac	att	cga	tac	gtc	ttc	cag	cac	aag	cac	tgg	tgg	gac	ctc	ttc	912
Asp	Asn	Ile	Arg	Tyr	Val	Phe	Gln	His	Lys	His	Trp	Trp	Asp	Leu	Phe	
			290				295								300	
tgg	atc	gcc	acg	ttc	ttc	gcg	aag	cac	ttc	acg	ctc	tac	ggc	ccg	ctg	960
Trp	Ile	Ala	Thr	Phe	Phe	Ala	Lys	His	Phe	Thr	Leu	Tyr	Gly	Pro	Leu	
			305			310					315				320	
atg	ggc	ggc	tgg	ggc	gcg	ttc	tgg	ttc	tac	atg	ctg	gtg	cgc	acg	gtc	1008
Met	Gly	Gly	Trp	Gly	Ala	Phe	Trp	Phe	Tyr	Met	Leu	Val	Arg	Thr	Val	
			325								330				335	
gag	agc	cac	tgg	ttc	aca	tgg	gtg	acc	cag	atg	aac	cac	atc	ccc	atg	1056
Glu	Ser	His	Trp	Phe	Thr	Trp	Val	Thr	Gln	Met	Asn	His	Ile	Pro	Met	



-continued

340	345	350	
cac gtc gac aac gac cgc gag ctg gac tgg ccc acc ctg cag ggt ctc His Val Asp Asn Asp Arg Glu Leu Asp Trp Pro Thr Leu Gln Gly Leu 355 360 365			1104
gcc acg tgc aac gtc gag ggc agc ctc ttc aac gac tgg ttc acg ggc Ala Thr Cys Asn Val Glu Gly Ser Leu Phe Asn Asp Trp Phe Thr Gly 370 375 380			1152
cac ctc aac tac cag atc gag cac cac ctc ttc ccc acc atg ccc cgc His Leu Asn Tyr Gln Ile Glu His His Leu Phe Pro Thr Met Pro Arg 385 390 395 400			1200
cac aac tac gcg gtg gcc aac aag aag gtc cag gcc ctc tac aag aag His Asn Tyr Ala Val Ala Asn Lys Lys Val Gln Ala Leu Tyr Lys Lys 405 410 415			1248
cac ggc gtg ccg atg cag acc aag ggc ctc atc gaa gcc ttc gcc gac His Gly Val Pro Met Gln Thr Lys Gly Leu Ile Glu Ala Phe Ala Asp 420 425 430			1296
atc gtc aag tcg ctc gag cac tat ggt gag gtg tgg aag gag gcc tac Ile Val Lys Ser Leu Glu His Tyr Gly Glu Val Trp Lys Glu Ala Tyr 435 440 445			1344
tac ggc taa Tyr Gly 450			1353
 <210> SEQ ID NO 16 <211> LENGTH: 450 <212> TYPE: PRT <213> ORGANISM: Acanthamoeba castellanii			
 <400> SEQUENCE: 16			
Met Ala Thr Ala Ser Ala Ser Asn Val Leu Arg Leu Pro Gly Glu Gly 1 5 10 15			
Leu Ala Thr Gly Leu Glu Gln Leu Glu Trp Ala Glu Val Gln Lys His 20 25 30			
Asn Thr Arg Glu Ser Ser Trp Leu Val Ile Asn Asp Gln Val Tyr Asp 35 40 45			
Ile Thr Asn Phe Gly Arg Arg His Pro Gly Gly Lys Val Ile Tyr His 50 55 60			
Tyr Ala Gly Gln Asp Ala Thr Asp Ser Phe Arg Ala Leu His Pro Asp 65 70 75 80			
Ser Ala Leu Val Met Lys Tyr Leu Lys Pro Leu Leu Ile Gly Gln Val 85 90 95			
Ala Pro Gly Ser Ser Thr Ala Ala Ser Ile Val Asp Gly Ala Arg Pro 100 105 110			
Ala Pro Ser Ala Phe Val Glu Glu Phe Arg Gln Val Arg Lys Glu Phe 115 120 125			
Glu Glu Gln Gly Leu Phe Glu Ala Ser Trp Ser Phe Phe Phe Gly Met 130 135 140			
Leu Ala His Ile Phe Leu Leu Glu Ala Ala Ala Tyr Tyr Ser Ile Lys 145 150 155 160			
Leu Leu Gly Asn Ser Trp Pro Val Tyr Leu Leu Ala Val Gly Leu Leu 165 170 175			
Ala Thr Ala Gln Ala Gln Ala Gly Trp Leu Gln His Asp Cys Gly His 180 185 190			
Leu Ser Val Phe Lys Lys Ser Lys Trp Asn His Trp Met His Tyr Ile 195 200 205			
Val Ile Cys His Ile Lys Gly Ala Ser Arg Ala Trp Trp Asn Trp Arg 210 215 220			

-continued

His Phe Glu His His Ala Lys Pro Asn Val Val Arg Lys Asp Pro Asp  
 225 230 235 240  
 Ile Thr Phe Pro Asn Leu Phe Leu Leu Gly Asp His Leu Thr Arg Lys  
 245 250 255  
 Trp Ala Lys Ala Lys Lys Gly Val Met Pro Tyr Asn Lys Gln His Leu  
 260 265 270  
 Tyr Trp Trp Ala Phe Pro Pro Leu Leu Leu Pro Val Tyr Phe His Tyr  
 275 280 285  
 Asp Asn Ile Arg Tyr Val Phe Gln His Lys His Trp Trp Asp Leu Phe  
 290 295 300  
 Trp Ile Ala Thr Phe Phe Ala Lys His Phe Thr Leu Tyr Gly Pro Leu  
 305 310 315 320  
 Met Gly Gly Trp Gly Ala Phe Trp Phe Tyr Met Leu Val Arg Thr Val  
 325 330 335  
 Glu Ser His Trp Phe Thr Trp Val Thr Gln Met Asn His Ile Pro Met  
 340 345 350  
 His Val Asp Asn Asp Arg Glu Leu Asp Trp Pro Thr Leu Gln Gly Leu  
 355 360 365  
 Ala Thr Cys Asn Val Glu Gly Ser Leu Phe Asn Asp Trp Phe Thr Gly  
 370 375 380  
 His Leu Asn Tyr Gln Ile Glu His His Leu Phe Pro Thr Met Pro Arg  
 385 390 395 400  
 His Asn Tyr Ala Val Ala Asn Lys Lys Val Gln Ala Leu Tyr Lys Lys  
 405 410 415  
 His Gly Val Pro Met Gln Thr Lys Gly Leu Ile Glu Ala Phe Ala Asp  
 420 425 430  
 Ile Val Lys Ser Leu Glu His Tyr Gly Glu Val Trp Lys Glu Ala Tyr  
 435 440 445  
 Tyr Gly  
 450

<210> SEQ ID NO 17  
 <211> LENGTH: 1374  
 <212> TYPE: DNA  
 <213> ORGANISM: Perkinsus marinus  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1374)  
 <223> OTHER INFORMATION: Delta-5-Desaturase

<400> SEQUENCE: 17

atg act act tca acc act act gtg caa cta caa gaa gac ctg tca agt	48
Met Thr Thr Ser Thr Thr Thr Val Gln Leu Gln Glu Asp Leu Ser Ser	
1 5 10 15	
ggg gac cag aac gcc cac ccc agt cca agc cga gct act cct agt gtt	96
Gly Asp Gln Asn Ala His Pro Ser Pro Ser Arg Ala Thr Pro Ser Val	
20 25 30	
ggg gat act aag gag gat gcg agg gtt gtt atc aaa cta ttt ggt aca	144
Gly Asp Thr Lys Glu Asp Ala Arg Val Val Ile Lys Leu Phe Gly Thr	
35 40 45	
tgg gtt gat gtt aca gct tgg ttg aat gac cat cct ggt ggt tct aaa	192
Trp Val Asp Val Thr Ala Trp Leu Asn Asp His Pro Gly Gly Ser Lys	
50 55 60	
gtg ctc aga gca ttc aac aag aag gac gcg act gat gct gtt atg gcc	240
Val Leu Arg Ala Phe Asn Lys Lys Asp Ala Thr Asp Ala Val Met Ala	
65 70 75 80	
atg cac act gat gaa gct atc aag cgc atc atc aga ttt tca aat gtg	288
Met His Thr Asp Glu Ala Ile Lys Arg Ile Ile Arg Phe Ser Asn Val	
85 90 95	

-continued

gtc tcc tcg gcc ccc atc aac gcc tct att ggt gat gtc cag gtt att Val Ser Ser Ala Pro Ile Asn Ala Ser Ile Gly Asp Val Gln Val Ile 100 105 110	336
gag aaa tct cta tcg aga gaa cag ttg atg tat tac aag ctc cgc act Glu Lys Ser Leu Ser Arg Glu Gln Leu Met Tyr Tyr Lys Leu Arg Thr 115 120 125	384
ctt gct aga aac cag gcc tgg ttt caa agc aat cta tta tac gaa gga Leu Ala Arg Asn Gln Gly Trp Phe Gln Ser Asn Leu Leu Tyr Glu Gly 130 135 140	432
gtg aaa gca atg ata gcc ttc ggt ttg ctc atc atc ggg ttt gct act Val Lys Ala Met Ile Ala Phe Gly Leu Leu Ile Ile Gly Phe Ala Thr 145 150 155 160	480
ctc tac ttc gac tat ggt att tgg tca acc gca ctg ata ggt ttc gct Leu Tyr Phe Asp Tyr Gly Ile Trp Ser Thr Ala Leu Ile Gly Phe Ala 165 170 175	528
tgg ttt cag ctg ggg tgg ttg gga cat gac tgg tct cat cat aca gct Trp Phe Gln Leu Gly Trp Leu Gly His Asp Trp Ser His His Thr Ala 180 185 190	576
cta cca aag tct act act aac tgt gcg aac tac aat gac tat ctt ggc Leu Pro Lys Ser Thr Thr Asn Cys Ala Asn Tyr Asn Asp Tyr Leu Gly 195 200 205	624
tgg ctt act ggt ttg gct aga ggg aat aca ctt ctg tgg tgg aaa cta Trp Leu Thr Gly Leu Ala Arg Gly Asn Thr Leu Leu Trp Trp Lys Leu 210 215 220	672
agg cat aat act cat cac gtg ctg acc aat cag tac gag aat gat cct Arg His Asn Thr His His Val Leu Thr Asn Gln Tyr Glu Asn Asp Pro 225 230 235 240	720
gat ata cta act caa cca ccg ttg cat ttt ttc gag gac ttc gat gtt Asp Ile Leu Thr Gln Pro Pro Leu His Phe Phe Glu Asp Phe Asp Val 245 250 255	768
ggt aat gtg aac aga tat caa gct gtc tac tat cta cca atg cta act Gly Asn Val Asn Arg Tyr Gln Ala Val Tyr Tyr Leu Pro Met Leu Thr 260 265 270	816
cta ctg cat cta ttt tgg ttg tac gag tcg gta ttg gtt tgc ttg aga Leu Leu His Leu Phe Trp Leu Tyr Glu Ser Val Leu Val Cys Leu Arg 275 280 285	864
caa agt aag tct att aat aga tac aac cgt atg cat gcc cgg agg gat Gln Ser Lys Ser Ile Asn Arg Tyr Asn Arg Met His Ala Arg Arg Asp 290 295 300	912
acc gta gct ttg gta ctt cac ata ctc att gtt ggc atc ata tcg tac Thr Val Ala Leu Val Leu His Ile Leu Ile Val Gly Ile Ile Ser Tyr 305 310 315 320	960
acc agt ggt aag tat ttg ctc atc ctt ctg gcc tac atg ctt agt ggc Thr Ser Gly Lys Tyr Leu Leu Ile Leu Leu Ala Tyr Met Leu Ser Gly 325 330 335	1008
ttt cta act gct gtt gtt gta ttt gcc agc cac tac aac gag cct agg Phe Leu Thr Ala Val Val Val Phe Ala Ser His Tyr Asn Glu Pro Arg 340 345 350	1056
gta gct tct ggt gaa tcc tta tca ctc gtt cgt cag aca ttg tta acc Val Ala Ser Gly Glu Ser Leu Ser Leu Val Arg Gln Thr Leu Leu Thr 355 360 365	1104
act atc aat ata ggc tca ttc agt gat act cat tgg gag aag aag ttg Thr Ile Asn Ile Gly Ser Phe Ser Asp Thr His Trp Glu Lys Lys Leu 370 375 380	1152
tgg ttc tat cta act ggt ggt ctt aat atg caa atc gag cat cat ctc Trp Phe Tyr Leu Thr Gly Gly Leu Asn Met Gln Ile Glu His His Leu 385 390 395 400	1200
ttc cca aca atg ccc cgc cat aat ctt ccg aag aca act ttt ctg gtc Phe Pro Thr Met Pro Arg His Asn Leu Pro Lys Thr Thr Phe Leu Val 405 410 415	1248

-continued

```

aag tca cta gcc cag gag cta gga ctg cca tac aag gaa acc aac att 1296
Lys Ser Leu Ala Gln Glu Leu Gly Leu Pro Tyr Lys Glu Thr Asn Ile
      420                      425                      430

gtc agt tta acc aag gcg gcc gtt act act ttg cat cat aat gct ctg 1344
Val Ser Leu Thr Lys Ala Ala Val Thr Thr Leu His His Asn Ala Leu
      435                      440                      445

cgt aac atc gag aga ttg ctt gct agg tag 1374
Arg Asn Ile Glu Arg Leu Leu Ala Arg
      450                      455

```

```

<210> SEQ ID NO 18
<211> LENGTH: 457
<212> TYPE: PRT
<213> ORGANISM: Perkinsus marinus

```

```

<400> SEQUENCE: 18

```

```

Met Thr Thr Ser Thr Thr Thr Val Gln Leu Gln Glu Asp Leu Ser Ser
1      5      10      15

Gly Asp Gln Asn Ala His Pro Ser Pro Ser Arg Ala Thr Pro Ser Val
      20      25      30

Gly Asp Thr Lys Glu Asp Ala Arg Val Val Ile Lys Leu Phe Gly Thr
      35      40      45

Trp Val Asp Val Thr Ala Trp Leu Asn Asp His Pro Gly Gly Ser Lys
      50      55      60

Val Leu Arg Ala Phe Asn Lys Lys Asp Ala Thr Asp Ala Val Met Ala
      65      70      75      80

Met His Thr Asp Glu Ala Ile Lys Arg Ile Ile Arg Phe Ser Asn Val
      85      90      95

Val Ser Ser Ala Pro Ile Asn Ala Ser Ile Gly Asp Val Gln Val Ile
      100     105     110

Glu Lys Ser Leu Ser Arg Glu Gln Leu Met Tyr Tyr Lys Leu Arg Thr
      115     120     125

Leu Ala Arg Asn Gln Gly Trp Phe Gln Ser Asn Leu Leu Tyr Glu Gly
      130     135     140

Val Lys Ala Met Ile Ala Phe Gly Leu Leu Ile Ile Gly Phe Ala Thr
      145     150     155     160

Leu Tyr Phe Asp Tyr Gly Ile Trp Ser Thr Ala Leu Ile Gly Phe Ala
      165     170     175

Trp Phe Gln Leu Gly Trp Leu Gly His Asp Trp Ser His His Thr Ala
      180     185     190

Leu Pro Lys Ser Thr Thr Asn Cys Ala Asn Tyr Asn Asp Tyr Leu Gly
      195     200     205

Trp Leu Thr Gly Leu Ala Arg Gly Asn Thr Leu Leu Trp Trp Lys Leu
      210     215     220

Arg His Asn Thr His His Val Leu Thr Asn Gln Tyr Glu Asn Asp Pro
      225     230     235     240

Asp Ile Leu Thr Gln Pro Pro Leu His Phe Phe Glu Asp Phe Asp Val
      245     250     255

Gly Asn Val Asn Arg Tyr Gln Ala Val Tyr Tyr Leu Pro Met Leu Thr
      260     265     270

Leu Leu His Leu Phe Trp Leu Tyr Glu Ser Val Leu Val Cys Leu Arg
      275     280     285

Gln Ser Lys Ser Ile Asn Arg Tyr Asn Arg Met His Ala Arg Arg Asp
      290     295     300

Thr Val Ala Leu Val Leu His Ile Leu Ile Val Gly Ile Ile Ser Tyr
      305     310     315     320

```

-continued

Thr Ser Gly Lys Tyr Leu Leu Ile Leu Leu Ala Tyr Met Leu Ser Gly  
 325 330 335  
 Phe Leu Thr Ala Val Val Val Phe Ala Ser His Tyr Asn Glu Pro Arg  
 340 345 350  
 Val Ala Ser Gly Glu Ser Leu Ser Leu Val Arg Gln Thr Leu Leu Thr  
 355 360 365  
 Thr Ile Asn Ile Gly Ser Phe Ser Asp Thr His Trp Glu Lys Lys Leu  
 370 375 380  
 Trp Phe Tyr Leu Thr Gly Gly Leu Asn Met Gln Ile Glu His His Leu  
 385 390 395 400  
 Phe Pro Thr Met Pro Arg His Asn Leu Pro Lys Thr Thr Phe Leu Val  
 405 410 415  
 Lys Ser Leu Ala Gln Glu Leu Gly Leu Pro Tyr Lys Glu Thr Asn Ile  
 420 425 430  
 Val Ser Leu Thr Lys Ala Ala Val Thr Thr Leu His His Asn Ala Leu  
 435 440 445  
 Arg Asn Ile Glu Arg Leu Leu Ala Arg  
 450 455

<210> SEQ ID NO 19  
 <211> LENGTH: 1224  
 <212> TYPE: DNA  
 <213> ORGANISM: Acanthamoeba castellanii  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1224)  
 <223> OTHER INFORMATION: Delta-12/Delta-15-Desaturase

<400> SEQUENCE: 19

atg act att act act acc cag acc ttg aac cag aag gct gct aag aag	48
Met Thr Ile Thr Thr Gln Thr Leu Asn Gln Lys Ala Ala Lys Lys	
1 5 10 15	
gga gga aag gag agg gct cca att att cca aag gag aac gct cca ttc	96
Gly Gly Lys Glu Arg Ala Pro Ile Ile Pro Lys Glu Asn Ala Pro Phe	
20 25 30	
act ttg gga cag atc aag gga gct atc cca cct cat ctc ttc aag cac	144
Thr Leu Gly Gln Ile Lys Gly Ala Ile Pro Pro His Leu Phe Lys His	
35 40 45	
tcc atg ttg aag tct ttc tcc tac ttg gga gtg gat ttg ttg gag tct	192
Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser	
50 55 60	
acc atc tgg ttg ttc ctc atc ttg tac ttg gat gga ctc act aag gag	240
Thr Ile Trp Leu Phe Leu Ile Leu Tyr Leu Asp Gly Leu Thr Lys Glu	
65 70 75 80	
aac acc ttg ttg aac tgg act tgc tgg gtt gca tac tgg ttg tac caa	288
Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln	
85 90 95	
gga ttg act tgg act gga att tgg gtg ttg gct cat gag tgt gga cat	336
Gly Leu Thr Trp Thr Gly Ile Trp Val Leu Ala His Glu Cys Gly His	
100 105 110	
gga gga ttc gtt gct caa gag tgg ttg aac gat acc gtg ggt ttc att	384
Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe Ile	
115 120 125	
ttc cat acc gtg ctc tac gtt cca tac ttc tcc tgg aag ttc tct cat	432
Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His	
130 135 140	
gct aag cac cat cac tac acc aac cac atg act aag gat gag cca ttc	480
Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe	
145 150 155 160	

-continued

gtg cca cat aca atc act cca gag caa agg gct aaa gtg gat caa gga	528
Val Pro His Thr Ile Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly	
165 170 175	
gag ttg cca cat cca aac aag cca tcc ctc ttc gct ttc tac gag aga	576
Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg	
180 185 190	
tgg gtg atc cca ttc gtg atg ttg ttc ttg gga tgg cca ctc tac ttg	624
Trp Val Ile Pro Phe Val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu	
195 200 205	
tct atc aac gct tct gga cca cca aag aag gag ttg gtt tcc cac tac	672
Ser Ile Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr	
210 215 220	
gat cca aag gct tcc atc ttc aac aag aaa gat tgg tgg aag atc ttg	720
Asp Pro Lys Ala Ser Ile Phe Asn Lys Lys Asp Trp Trp Lys Ile Leu	
225 230 235 240	
ctc tct gat ttg gga ttg gtt gct tgg act ttg gct ttg tgg aag ttg	768
Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu	
245 250 255	
gga gag act ttc gga ttc gga ttg gtg gct gct ctt tac att cca cca	816
Gly Glu Thr Phe Gly Phe Gly Leu Val Ala Ala Leu Tyr Ile Pro Pro	
260 265 270	
gtg ctc gtt acc aac tct tac ttg gtg gct atc acc ttc ttg caa cac	864
Val Leu Val Thr Asn Ser Tyr Leu Val Ala Ile Thr Phe Leu Gln His	
275 280 285	
acc gat gat atc ctc cca cat tac gat gct act gag tgg act tgg ttg	912
Thr Asp Asp Ile Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu	
290 295 300	
aga gga gct ttg tgc act gtg gat aga tct ttg gga tgg ttc gga gat	960
Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp	
305 310 315 320	
tac aag acc cat cac atc gtt gat act cat gtg acc cac cac atc ttc	1008
Tyr Lys Thr His His Ile Val Asp Thr His Val Thr His His Ile Phe	
325 330 335	
tct tac ctc cca ttc tat aac gct gag gag gct act aag gct att aag	1056
Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala Ile Lys	
340 345 350	
cca gtg ttg aag gag tat cac tgc gag gat aag aga gga ttc ttc cac	1104
Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His	
355 360 365	
ttc tgg tac ttg ttc ttc aag acc gct gct gag aac tct gtt gtg gat	1152
Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp	
370 375 380	
aac gag acc aac aag tcc cca gga atc ttc tac ttc ttc agg gag gag	1200
Asn Glu Thr Asn Lys Ser Pro Gly Ile Phe Tyr Phe Phe Arg Glu Glu	
385 390 395 400	
att aag cac gga aag gct cat tga	1224
Ile Lys His Gly Lys Ala His	
405	
<210> SEQ ID NO 20	
<211> LENGTH: 407	
<212> TYPE: PRT	
<213> ORGANISM: Acanthamoeba castellanii	
<400> SEQUENCE: 20	
Met Thr Ile Thr Thr Thr Gln Thr Leu Asn Gln Lys Ala Ala Lys Lys	
1 5 10 15	
Gly Gly Lys Glu Arg Ala Pro Ile Ile Pro Lys Glu Asn Ala Pro Phe	
20 25 30	
Thr Leu Gly Gln Ile Lys Gly Ala Ile Pro Pro His Leu Phe Lys His	
35 40 45	

-continued

---

Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser  
 50 55 60  
 Thr Ile Trp Leu Phe Leu Ile Leu Tyr Leu Asp Gly Leu Thr Lys Glu  
 65 70 75 80  
 Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln  
 85 90 95  
 Gly Leu Thr Trp Thr Gly Ile Trp Val Leu Ala His Glu Cys Gly His  
 100 105 110  
 Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe Ile  
 115 120 125  
 Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His  
 130 135 140  
 Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe  
 145 150 155 160  
 Val Pro His Thr Ile Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly  
 165 170 175  
 Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg  
 180 185 190  
 Trp Val Ile Pro Phe Val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu  
 195 200 205  
 Ser Ile Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr  
 210 215 220  
 Asp Pro Lys Ala Ser Ile Phe Asn Lys Lys Asp Trp Trp Lys Ile Leu  
 225 230 235 240  
 Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu  
 245 250 255  
 Gly Glu Thr Phe Gly Phe Gly Leu Val Ala Ala Leu Tyr Ile Pro Pro  
 260 265 270  
 Val Leu Val Thr Asn Ser Tyr Leu Val Ala Ile Thr Phe Leu Gln His  
 275 280 285  
 Thr Asp Asp Ile Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu  
 290 295 300  
 Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp  
 305 310 315 320  
 Tyr Lys Thr His His Ile Val Asp Thr His Val Thr His His Ile Phe  
 325 330 335  
 Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala Ile Lys  
 340 345 350  
 Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His  
 355 360 365  
 Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp  
 370 375 380  
 Asn Glu Thr Asn Lys Ser Pro Gly Ile Phe Tyr Phe Phe Arg Glu Glu  
 385 390 395 400  
 Ile Lys His Gly Lys Ala His  
 405

<210> SEQ ID NO 21  
 <211> LENGTH: 1224  
 <212> TYPE: DNA  
 <213> ORGANISM: Acanthamoeba castellanii  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1224)  
 <223> OTHER INFORMATION: Delta-12/Delta-15-Desaturase

-continued

&lt;400&gt; SEQUENCE: 21

atg acg atc acg acg acg cag aca ctg aat cag aag gca gcc aag aag	48
Met Thr Ile Thr Thr Thr Gln Thr Leu Asn Gln Lys Ala Ala Lys Lys	
1 5 10 15	
ggc gga aag gag cgc gct ccg atc att ccc aag gag aac gcc ccc ttc	96
Gly Gly Lys Glu Arg Ala Pro Ile Ile Pro Lys Glu Asn Ala Pro Phe	
20 25 30	
act ctg ggc cag atc aag ggc gcc att cct ccg cat ctc ttc aag cac	144
Thr Leu Gly Gln Ile Lys Gly Ala Ile Pro Pro His Leu Phe Lys His	
35 40 45	
agc atg ctc aaa tcc ttc agc tat ctg ggc gtg gat ctg ctg gag agc	192
Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser	
50 55 60	
acc atc tgg ctc ttc ctc atc ctc tac ctc gac gcc ctc acc aag gag	240
Thr Ile Trp Leu Phe Leu Ile Leu Tyr Leu Asp Gly Leu Thr Lys Glu	
65 70 75 80	
aac acg ctc ctc aac tgg act tgc tgg gtt gcg tac tgg ctc tac cag	288
Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln	
85 90 95	
ggt ctg acc tgg act ggc att tgg gtg ctg gcc cac gag tgt ggc cat	336
Gly Leu Thr Trp Thr Gly Ile Trp Val Leu Ala His Glu Cys Gly His	
100 105 110	
ggc ggc ttc gtg gcg cag gag tgg ctc aac gac acg gtc gcc ttc atc	384
Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe Ile	
115 120 125	
ttc cac acc gtc ctc tac gtg ccc tac ttc tcg tgg aag ttc tcc cac	432
Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His	
130 135 140	
gcc aag cac cac cac tac acc aac cac atg aca aag gac gag ccc ttc	480
Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe	
145 150 155 160	
gtg ccc cac acc atc acc cct gag cag agg gcc aag gtc gac cag ggc	528
Val Pro His Thr Ile Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly	
165 170 175	
gag ctg ccc cac ccc aac aag ccc tcc ctc ttc gcc ttc tac gaa agg	576
Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg	
180 185 190	
tgg gtc atc ccc ttc gtc atg ctc ttc ctc gcc tgg ccg ctc tac ctg	624
Trp Val Ile Pro Phe Val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu	
195 200 205	
tcc atc aac gcc tct ggc cct ccc aag aag gag ctt gtg tcc cac tac	672
Ser Ile Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr	
210 215 220	
gac ccc aaa gcc agc atc ttc aac aag aag gac tgg tgg aag atc ctt	720
Asp Pro Lys Ala Ser Ile Phe Asn Lys Lys Asp Trp Trp Lys Ile Leu	
225 230 235 240	
ctc tct gac ctc ggc ctt gtg gcg tgg acc ctg gcc ctc tgg aag ctg	768
Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu	
245 250 255	
ggc gag acc ttc ggc ttc ggt ctc gtg gcc gcc ctc tac att ccg ccc	816
Gly Glu Thr Phe Gly Phe Gly Leu Val Ala Ala Leu Tyr Ile Pro Pro	
260 265 270	
gtg ctg gtg acc aac tcc tac ctg gtg gcc atc acc ttc ctc cag cac	864
Val Leu Val Thr Asn Ser Tyr Leu Val Ala Ile Thr Phe Leu Gln His	
275 280 285	
acc gac gac att ctg ccc cac tac gac gcc acc gag tgg acc tgg ctc	912
Thr Asp Asp Ile Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu	
290 295 300	
agg ggt gct ctc tgc act gtt gat cgt tcg ctg gcc tgg ttc gcc gac	960
Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp	



-continued

305	310	315	320	
tac aag acg cac cac atc gtc gac acc cac gtg acg cac cac atc ttc				1008
Tyr Lys Thr His His Ile Val Asp Thr His Val Thr His His Ile Phe				
	325	330	335	
tcg tac ctg ccg ttc tac aac gcc gag gag gcc acc aag gcc atc aag				1056
Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala Ile Lys				
	340	345	350	
ccc gtg ctc aag gag tac cac tgc gag gac aag cgt ggc ttc ttc cac				1104
Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His				
	355	360	365	
ttc tgg tat ctg ttc ttc aag acc gcc gcc gag aac agc gtt gtc gac				1152
Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp				
	370	375	380	
aac gag acc aac aag agc ccc gcc atc ttc tac ttc ttc cgg gag gag				1200
Asn Glu Thr Asn Lys Ser Pro Gly Ile Phe Tyr Phe Phe Arg Glu Glu				
	385	390	395	400
atc aag cac ggc aag gcc cac tag				1224
Ile Lys His Gly Lys Ala His				
	405			
<210> SEQ ID NO 22				
<211> LENGTH: 407				
<212> TYPE: PRT				
<213> ORGANISM: Acanthamoeba castellanii				
<400> SEQUENCE: 22				
Met Thr Ile Thr Thr Thr Gln Thr Leu Asn Gln Lys Ala Ala Lys Lys				
1	5	10	15	
Gly Gly Lys Glu Arg Ala Pro Ile Ile Pro Lys Glu Asn Ala Pro Phe				
	20	25	30	
Thr Leu Gly Gln Ile Lys Gly Ala Ile Pro Pro His Leu Phe Lys His				
	35	40	45	
Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser				
	50	55	60	
Thr Ile Trp Leu Phe Leu Ile Leu Tyr Leu Asp Gly Leu Thr Lys Glu				
	65	70	75	80
Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln				
	85	90	95	
Gly Leu Thr Trp Thr Gly Ile Trp Val Leu Ala His Glu Cys Gly His				
	100	105	110	
Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe Ile				
	115	120	125	
Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His				
	130	135	140	
Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe				
	145	150	155	160
Val Pro His Thr Ile Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly				
	165	170	175	
Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg				
	180	185	190	
Trp Val Ile Pro Phe Val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu				
	195	200	205	
Ser Ile Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr				
	210	215	220	
Asp Pro Lys Ala Ser Ile Phe Asn Lys Lys Asp Trp Trp Lys Ile Leu				
	225	230	235	240
Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu				



-continued

tgg cag tac tcc cat gcg agg cac cat aag ttc acc aac cac gct act	480
Trp Gln Tyr Ser His Ala Arg His His Lys Phe Thr Asn His Ala Thr	
145 150 155 160	
aag ggt gag agc cat gtc ccc agc ctg gaa agt gag atg ggc gta ttc	528
Lys Gly Glu Ser His Val Pro Ser Leu Glu Ser Glu Met Gly Val Phe	
165 170 175	
agt cgt ata cag aag gcc ctg gag ggt tat ggt ctc gat gat gtc ttc	576
Ser Arg Ile Gln Lys Ala Leu Glu Gly Tyr Gly Leu Asp Asp Val Phe	
180 185 190	
cca gtc ttc cct ata gtg atg ctc ctg gtt ggg tat cct gtg tat ctc	624
Pro Val Phe Pro Ile Val Met Leu Leu Val Gly Tyr Pro Val Tyr Leu	
195 200 205	
ttc tgg aat gca tca ggt ggg cgt gtg ggc tac gat cgc cgt ccg tac	672
Phe Trp Asn Ala Ser Gly Gly Arg Val Gly Tyr Asp Arg Arg Pro Tyr	
210 215 220	
agc gac act aag cca tct cat ttc aat ccc aac ggt ggc ctt ttc cct	720
Ser Asp Thr Lys Pro Ser His Phe Asn Pro Asn Gly Gly Leu Phe Pro	
225 230 235 240	
cct tat atg aga gag aaa gtc ctc ctt agt gga gtt ggc tgt agc ata	768
Pro Tyr Met Arg Glu Lys Val Leu Leu Ser Gly Val Gly Cys Ser Ile	
245 250 255	
acc ctc ctt att ttg gcc tat tgt gct ggg agg gta ggc ctt agc agt	816
Thr Leu Leu Ile Leu Ala Tyr Cys Ala Gly Arg Val Gly Leu Ser Ser	
260 265 270	
gta ttg ttg tgg tat ggt tgt ccc tac ctt atg acc aac gcc tgg cta	864
Val Leu Leu Trp Tyr Gly Cys Pro Tyr Leu Met Thr Asn Ala Trp Leu	
275 280 285	
acg ctg tat acc tcc cta cag cac acg cat gaa gga gtc ccc cat tat	912
Thr Leu Tyr Thr Ser Leu Gln His Thr His Glu Gly Val Pro His Tyr	
290 295 300	
ggc gat gag gct ttc acc ttc atc aga ggt gcc tta gct tct atc gat	960
Gly Asp Glu Ala Phe Thr Phe Ile Arg Gly Ala Leu Ala Ser Ile Asp	
305 310 315 320	
cgt cca ccg tat ggc att ttc tct acg cat ttt cac cac gaa att ggc	1008
Arg Pro Pro Tyr Gly Ile Phe Ser Thr His Phe His His Glu Ile Gly	
325 330 335	
acc act cat gtt ctg cac cac att gat tct agg atc ccc tgt tac cat	1056
Thr Thr His Val Leu His His Ile Asp Ser Arg Ile Pro Cys Tyr His	
340 345 350	
gct aga gaa gcc act gat gct atc aag cct att ctg ggg gat tac tat	1104
Ala Arg Glu Ala Thr Asp Ala Ile Lys Pro Ile Leu Gly Asp Tyr Tyr	
355 360 365	
agg gag gat ggt act cct ata gta aag gca ttt ttg aag gtc cac aga	1152
Arg Glu Asp Gly Thr Pro Ile Val Lys Ala Phe Leu Lys Val His Arg	
370 375 380	
gag tgc aag ttc atc gga ggc ctc aac ggc gtc cag ttt tac cgt cct	1200
Glu Cys Lys Phe Ile Gly Gly Leu Asn Gly Val Gln Phe Tyr Arg Pro	
385 390 395 400	
ggg cag cgg ccg cag cag cag ccc tgc ggc agc aac gct cgc act tct	1248
Gly Gln Arg Pro Gln Gln Gln Pro Cys Gly Ser Asn Ala Arg Thr Ser	
405 410 415	
cgt tag	1254
Arg	

<210> SEQ ID NO 24

<211> LENGTH: 417

<212> TYPE: PRT

<213> ORGANISM: Perkinsus marinus

<400> SEQUENCE: 24

Met Thr Gln Thr Glu Val Gln Ala Gly Pro Cys Arg Asp Gly Arg Asn

-continued

---

1                    5                    10                    15  
 Leu Lys Ser Glu Ala Asp Val Lys Gly Phe Thr Ala Glu Glu Phe Thr  
                   20                    25                    30  
 Lys Val Gly Pro Ser Val Cys Ala Ile Gln Ser Ala Ile Pro Met His  
                   35                    40                    45  
 Cys Arg Asp Arg Ser Leu Ser Arg Ser Val Leu Cys Val Ile Arg Asp  
                   50                    55                    60  
 Leu Leu Tyr Ile Thr Ala Cys Ala Ala Val Gln Tyr Ser Leu Leu Ala  
                   65                    70                    75                    80  
 Leu Val Pro Pro Asp Ser Thr Leu Leu Arg Ala Val Leu Trp Gly Val  
                   85                    90                    95  
 Tyr Ile Phe Trp Gln Gly Val Phe Phe Thr Gly Ile Trp Val Met Gly  
                   100                    105                    110  
 His Glu Cys Gly His Gly Ala Phe Ser Pro Tyr Ser Met Leu Asn Asp  
                   115                    120                    125  
 Ser Ile Gly Phe Val Leu His Ser Ala Leu Leu Val Pro Tyr Phe Ser  
                   130                    135                    140  
 Trp Gln Tyr Ser His Ala Arg His His Lys Phe Thr Asn His Ala Thr  
                   145                    150                    155                    160  
 Lys Gly Glu Ser His Val Pro Ser Leu Glu Ser Glu Met Gly Val Phe  
                   165                    170                    175  
 Ser Arg Ile Gln Lys Ala Leu Glu Gly Tyr Gly Leu Asp Asp Val Phe  
                   180                    185                    190  
 Pro Val Phe Pro Ile Val Met Leu Leu Val Gly Tyr Pro Val Tyr Leu  
                   195                    200                    205  
 Phe Trp Asn Ala Ser Gly Gly Arg Val Gly Tyr Asp Arg Arg Pro Tyr  
                   210                    215                    220  
 Ser Asp Thr Lys Pro Ser His Phe Asn Pro Asn Gly Gly Leu Phe Pro  
                   225                    230                    235                    240  
 Pro Tyr Met Arg Glu Lys Val Leu Leu Ser Gly Val Gly Cys Ser Ile  
                   245                    250                    255  
 Thr Leu Leu Ile Leu Ala Tyr Cys Ala Gly Arg Val Gly Leu Ser Ser  
                   260                    265                    270  
 Val Leu Leu Trp Tyr Gly Cys Pro Tyr Leu Met Thr Asn Ala Trp Leu  
                   275                    280                    285  
 Thr Leu Tyr Thr Ser Leu Gln His Thr His Glu Gly Val Pro His Tyr  
                   290                    295                    300  
 Gly Asp Glu Ala Phe Thr Phe Ile Arg Gly Ala Leu Ala Ser Ile Asp  
                   305                    310                    315                    320  
 Arg Pro Pro Tyr Gly Ile Phe Ser Thr His Phe His His Glu Ile Gly  
                   325                    330                    335  
 Thr Thr His Val Leu His His Ile Asp Ser Arg Ile Pro Cys Tyr His  
                   340                    345                    350  
 Ala Arg Glu Ala Thr Asp Ala Ile Lys Pro Ile Leu Gly Asp Tyr Tyr  
                   355                    360                    365  
 Arg Glu Asp Gly Thr Pro Ile Val Lys Ala Phe Leu Lys Val His Arg  
                   370                    375                    380  
 Glu Cys Lys Phe Ile Gly Gly Leu Asn Gly Val Gln Phe Tyr Arg Pro  
                   385                    390                    395                    400  
 Gly Gln Arg Pro Gln Gln Gln Pro Cys Gly Ser Asn Ala Arg Thr Ser  
                   405                    410                    415  
 Arg

-continued

---

<210> SEQ ID NO 25  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 25  
 ggtaccatgg cgatcacgac gacgcagaca c 31

<210> SEQ ID NO 26  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 26  
 gagctcctag tgggccttgc cgtgcttgat ctcc 34

<210> SEQ ID NO 27  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 27  
 ggtaccatgg tcctcacaac cccggccctc 30

<210> SEQ ID NO 28  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 28  
 ggagctctca gttctcagca cccatcttc 29

<210> SEQ ID NO 29  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 29  
 ggtaccatgg ccaccgcacg tgcacg 26

<210> SEQ ID NO 30  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 30  
 ggagctttag ccgtagtagg cctcctt 27

<210> SEQ ID NO 31  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 31

-continued

---

ggtaccatgg cggctgcgac ggcgac 26

<210> SEQ ID NO 32  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 32

ggagctttag tcgtgcttcc tcttggg 27

<210> SEQ ID NO 33  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 33

ggtaccatga cccaaactga ggtcca 26

<210> SEQ ID NO 34  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 34

ggagctctaa cgagaagtgc gagcgt 26

<210> SEQ ID NO 35  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 35

ggtaccatgt cttctcttac cctcta 26

<210> SEQ ID NO 36  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 36

ggagctctat tccactatgg caacag 26

<210> SEQ ID NO 37  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
<400> SEQUENCE: 37

ggtaccatga ctacttcaac cactac 26

<210> SEQ ID NO 38  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 38

ggagctctac cttagcaagca atctct 26

<210> SEQ ID NO 39  
 <211> LENGTH: 34  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 39

ggatccacca tggcgatcac gacgacgcag acac 34

<210> SEQ ID NO 40  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 40

ggtctagact agtgggcctt gccgtgcttg atctcc 36

<210> SEQ ID NO 41  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 41

ggatccagga tggctctcac aaccccgcc ctc 33

<210> SEQ ID NO 42  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 42

ggtctagatc agttctcagc acccatcttc 30

<210> SEQ ID NO 43  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 43

ggatccatgg ccaccgcac tgcac 26

<210> SEQ ID NO 44  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 44

ggtctagatt agccgtagta ggcctcctt 29

-continued

---

<210> SEQ ID NO 45  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 45  
  
ggatccatgg cggctgcgac ggcgac 26

<210> SEQ ID NO 46  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 46  
  
ggtctagatt agtcgtgctt cctcttggg 29

<210> SEQ ID NO 47  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 47  
  
ggatccatga cccaaactga ggtcca 26

<210> SEQ ID NO 48  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 48  
  
ggtctagact aacgagaagt gcgagcgt 28

<210> SEQ ID NO 49  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 49  
  
ggatccatgt cttctcttac cctcta 26

<210> SEQ ID NO 50  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 50  
  
ggtctagact attccactat ggcaacag 28

<210> SEQ ID NO 51  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: primer  
  
<400> SEQUENCE: 51



-continued

---

ggatccatga ctacttcaac cactac 26

<210> SEQ ID NO 52  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <400> SEQUENCE: 52

ggtctagact acctagcaag caatctct 28

<210> SEQ ID NO 53  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (3)..(3)  
 <223> OTHER INFORMATION: n is inosine  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (9)..(9)  
 <223> OTHER INFORMATION: n is inosine  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (12)..(12)  
 <223> OTHER INFORMATION: n is inosine  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (21)..(21)  
 <223> OTHER INFORMATION: n is inosine  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (24)..(24)  
 <223> OTHER INFORMATION: n is inosine  
 <400> SEQUENCE: 53

ggntgghtng gncaygayky nksnca 26

<210> SEQ ID NO 54  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: primer  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (6)..(6)  
 <223> OTHER INFORMATION: n is a, c, g, or t  
 <400> SEQUENCE: 54

ggraanagrt grtgytcdat ytg 23

<210> SEQ ID NO 55  
 <211> LENGTH: 4  
 <212> TYPE: PRT  
 <213> ORGANISM: Unknown  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Cyt b5 motif  
 <400> SEQUENCE: 55

His Pro Gly Gly  
 1

<210> SEQ ID NO 56  
 <211> LENGTH: 29  
 <212> TYPE: DNA

-continued

---

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 56

caagtaccac ccgggcgcca gcaggcca

29

<210> SEQ ID NO 57

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: primer

<400> SEQUENCE: 57

tgccctgct gccgccggg tggacttg

29

---

We claim:

1. A process for the production of arachidonic acid or eicosapentaenoic acid or arachidonic acid and eicosapentaenoic acid in transgenic plants that produce mature seeds with a content of at least 1% by weight of said compounds referred to the total lipid content of said organism which comprises:

a) introducing at least one nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -12-desaturase- and  $\Delta$ -15-desaturase-activity, and

b) introducing at least one second nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -9-elongase-activity, and

c) introducing at least one third nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a  $\Delta$ -8-desaturase-activity, and

d) introducing at least one fourth nucleic acid sequence, which encodes a polypeptide having a  $\Delta$ -5-desaturase-activity, and

e) cultivating and harvesting of said transgenic plant, wherein the nucleic acid sequence which encodes a polypeptide having  $\Delta$ -8-desaturase activity comprises a nucleic acid sequence selected from the group consisting of

a) the nucleic acid sequence of SEQ ID NO: 3 or 5;

b) a nucleic acid sequence encoding the polypeptide sequence of SEQ ID NO: 4 or 6; and

c) a nucleic acid sequence encoding a polypeptide having  $\Delta$ -8-desaturase activity and having at least 90% homology to the sequence of SEQ ID NO: 4 or 6.

2. The process of claim 1, wherein the nucleic acid sequence which encodes a polypeptide having  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity,  $\Delta$ -9-elongase, or  $\Delta$ -5-desaturase activity comprises a nucleic acid sequence selected from the group consisting of

a) a nucleic acid sequence depicted in SEQ ID NO: 1, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23, encoding a polypeptide sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24, and

b) a nucleic acid sequence encoding a polypeptide having at least 50% homology to the sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO:

20 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptide has  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase activity,  $\Delta$ -9-elongase, or  $\Delta$ -5-desaturase activity.

25 3. The process of claim 1, wherein the transgenic plant is an oilseed plant.

30 4. The process of claim 1, wherein the transgenic plant that produces mature seeds is selected from the group consisting of the plant families of Anacardiaceae, Asteraceae, Apiaceae, Boraginaceae, Brassicaceae, Cannabaceae, Elaeagnaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Leguminosae, Linaceae, Lythraeae, Malvaceae, Onagraceae, Palmae, Poaceae, Rubiaceae, Scrophulariaceae, Solanaceae, Sterculiaceae, and Theaceae.

35 5. The process of claim 1, wherein the transgenic plant that produces mature seeds is selected from the group consisting of the plant genera of *Pistacia*, *Mangifera*, *Anacardium*, *Calendula*, *Carthamus*, *Centaurea*, *Cichorium*, *Cynara*, *Helianthus*, *Lactuca*, *Locusta*, *Tagetes*, *Valeriana*, *Borago*, *Daucus*, *Brassica*, *Camelina*, *Melanosinapis*, *Sinapis*, *Arabidopsis*, *Orychophragmus*, *Cannabis*, *Elaeagnus*, *Manihot*, *Janipha*, *Jatropha*, *Ricinus*, *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicago*, *Glycine*, *Dolichos*, *Phaseolus*, *Pelargonium*, *Cocos*, *Oleum*, *Juglans*, *Wallia*, *Arachis*, *Linum*, *Punica*, *Gossypium*, *Camissonia*, *Oenothera*, *Elaeis*, *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea*, *Triticum*, *Coffea*, *Verbascum*, *Capsicum*, *Nicotiana*, *Solanum*, *Lycopersicon*, *Theobroma*, and *Camellia*.

45 6. The process of claim 1, wherein the transgenic plant is selected from the group consisting of rapeseed, poppy, mustard, hemp, castor bean, sesame, olive, *calendula*, *punica*, hazel nut, maize, almond, *macadamia*, cotton, avocado, pumpkin, walnut, laurel, pistachio, primrose, canola, evening primrose, oil palm, peanut, linseed, soybean, safflower, marigold, coffee, tobacco, cacao, sunflower, and borage.

50 7. The process of claim 1, wherein the arachidonic acid or eicosapentaenoic acid or arachidonic acid and eicosapentaenoic acid is isolated in the form of their oils, lipids, or free fatty acids.

55 8. The process of claim 1, wherein arachidonic acid and eicosapentaenoic acid is produced in at least a 1:2 ratio.

60 9. The process of claim 1, wherein the arachidonic acid and eicosapentaenoic acid are produced in a content of at least 5% by weight referred to the total lipid content.

65 10. The process of claim 1, wherein the  $\Delta$ -12-desaturase and  $\Delta$ -15-desaturase used in the process desaturates C16 or

## 119

C18-fatty acids having one double bond in the fatty acid chain or C16 and C18-fatty acids having one double bond in the fatty acid chain.

11. An isolated nucleic acid sequence comprising a nucleotide sequence which encodes a  $\Delta$ -8-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 3 or 5;
- b) a nucleic acid sequence encoding a polypeptide sequence as depicted in SEQ ID NO: 4 or 6; and
- c) a nucleic acid sequence encoding a polypeptide having at least 90% homology to the sequence as depicted in SEQ ID NO: 4 or 6 and which polypeptide has  $\Delta$ -8-desaturase activity.

12. A gene construct comprising the isolated nucleic acid of claim 11, where the nucleic acid is functionally linked to one or more regulatory signals.

13. The gene construct of claim 12, whose gene expression is increased by the regulatory signals.

14. A vector comprising the gene construct of claim 13.

15. A vector comprising the nucleic acid of claim 11 or a gene construct comprising said nucleic acid wherein the nucleic acid is functionally linked to one or more regulatory sequence.

16. A transgenic plant comprising

- a) the nucleic acid of claim 11,
- b) a gene construct comprising said nucleic acid wherein the nucleic acid is functionally linked to one or more regulatory sequence, or

## 120

c) a vector comprising said nucleic acid or said gene construct.

17. The transgenic plant of claim 16, wherein the plant is an oilseed plant.

18. A transgenic plant comprising the gene construct of claim 13 or a vector comprising the gene construct.

19. The process of claim 1, wherein the nucleic acid which encodes a polypeptide having  $\Delta$ -8-desaturase activity comprises a nucleic acid sequence encoding a polypeptide having at least 95% homology to the sequence of SEQ ID NO: 4 or 6.

20. The process of claim 1, wherein the nucleic acid which encodes a polypeptide having  $\Delta$ -8-desaturase activity comprises the nucleic acid sequence of SEQ ID NO: 3 or 5 or a nucleic acid sequence encoding the polypeptide sequence of SEQ ID NO: 4 or 6.

21. The isolated nucleic acid sequence of claim 11, wherein the nucleotide sequence comprises a nucleic acid sequence encoding a polypeptide having at least 95% homology to the sequence of SEQ ID NO: 4 or 6.

22. The isolated nucleic acid sequence of claim 11, wherein the nucleotide sequence comprises the nucleic acid sequence of SEQ ID NO: 3 or 5 or a nucleic acid sequence encoding the polypeptide sequence of SEQ ID NO: 4 or 6.

\* \* \* \* \*