

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2007 (19.04.2007)

PCT

(10) International Publication Number
WO 2007/042510 A2

(51) **International Patent Classification:**
C12N 9/02 (2006.01) C12N 15/82 (2006.01)
C12N 9/10 (2006.01)

(21) **International Application Number:**
PCT/EP2006/067223

(22) **International Filing Date:** 10 October 2006 (10.10.2006)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
0520843.4 13 October 2005 (13.10.2005) GB

(71) **Applicant (for all designated States except US):** Rothamsted Research [-/GB]; A15 2jq, Harpenden (GB).

(72) **Inventors; and**

(75) **Inventors/Applicants (for US only):** NAPIER, Johnathan A. [GB/GB]; The Wilderness, Butchers Lane, Preston, Hertfordshire SG4 7tr (GB). SAYANOVA, Olga [GB/GB]; Rothamsted Research, Harpenden, Herts A15 2jq (GB).

(74) **Agent:** POPP, Andreas; BASF Aktiengesellschaft, 67056 Ludwigshafen (DE).

(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) **Title:** PROCESS FOR THE PRODUCTION OF ARACHIDONIC ACID AND/OR EICOSAPENTAENOIC ACID

(57) **Abstract:** The present invention relates to a new process for the production of arachidonic acid and/or eicosapentaenoic acid in plants through the co-expression of a Δ -12/ Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase and a Δ -5-desaturase and a process for the production of lipids or oils having an increased content of unsaturated fatty acids, in particular ω -3 and ω -6 fatty acids having at least two double bonds and a 18 or 20 carbon atom chain length. Preferably the arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio. The invention furthermore relates to the production of a transgenic plants, preferably a transgenic crop plant, having an increased content of arachidonic acid and/or eicosapentaenoic acid, oils or lipids containing C β - or C20-fatty acids with a double bond in position Δ 5, 8, 9, 11, 12, 14, 15 or of the fatty acid produced, respectively due to the expression of the Δ -12/ Δ -15-desaturase, of the Δ -9-elongase, of the Δ -8-desaturase and of the Δ -5-desaturase in the plant. The expression of the inventive Δ -12/ Δ -15-desaturase leads preferably to linoleic acid and linolenic acid as products having a double bond in the position Δ 9, 12 and 15 of the fatty acid. The invention additionally relates to specific nucleic acid sequences encoding for proteins with Δ -12/ Δ -15-desaturase-, Δ -9-elongase-, Δ -8-desaturase- or Δ -5-desaturase- activity, nucleic acid constructs, vectors and transgenic plants containing said nucleic acid sequences.

WO 2007/042510 A2

Process for the production of arachidonic acid and/or eicosapentaenoic acid

Description

The present invention relates to a new process for the production of arachidonic acid and/or eicosapentaenoic acid in plants through the co-expression of a Δ -12-/ Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase and a Δ -5-desaturase and a process for
5 the production of lipids or oils having an increased content of unsaturated fatty acids, in particular ω -3 and ω -6 fatty acids having at least two double bonds and a 18 or 20 carbon atom chain length. Preferably the arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio.

10 The invention furthermore relates to the production of a transgenic plants, preferably a transgenic crop plant, having an increased content of arachidonic acid and/or eicosapentaenoic acid, oils or lipids containing C_{18} - or C_{20} - fatty acids with a double bond in position Δ . 5, 8, 9, 11, 12, 14, 15 or 17 of the fatty acid produced, respectively
15 due to the expression of the Δ -12-/ Δ -15-desaturase, of the Δ -9-elongase, of the Δ -8-desaturase and of the Δ -5-desaturase in the plant. The expression of the inventive Δ -12-/ Δ -15-desaturase leads preferably to linoleic acid and α -linolenic acid as products having a double bond in the position Δ . 9, 12 and 15 of the fatty acid.

The invention additionally relates to specific nucleic acid sequences encoding for proteins with Δ -12-/ Δ -15-desaturase-, Δ -9-elongase-, Δ -8-desaturase- or Δ -5-desaturase-
20 activity, nucleic acid constructs, vectors and transgenic plants containing said nucleic acid sequences.

Plants and especially oil crops have been used for centuries as sources for edible and non-edible products. There are written records and archaeological excavations that oil crops such as linseed, olive and sesame were widespread use at least six thousand
25 years ago.

Non-edible products of oilseed crops such as rapeseed were used and included in lubricants, oil lamps, and cosmetics such as soaps. Oil crops differ in their cultural, economic and utilization characteristics, for example rapeseed and linseed are adapted to relatively cool climates, whereas oil palm and coconut are adapted to
30 warm and damp climates. Some plants are a real oilseed plant that means the main product of such plants is the oil, whereas in case of others such as cotton or soybean

the oil is more or less a side product. The oils of different plants are basically characterized by their individual fatty acid pattern.

Fatty acids and triglycerides have numerous applications in the food industry, animal nutrition, cosmetics and in the drug sector. Depending on whether they are free saturated or unsaturated fatty acids or triglycerides with an increased content of saturated or unsaturated fatty acids, they are suitable for the most varied applications; thus, for example, long chain polyunsaturated fatty acids (= LCPUFAs) are added to infant formula to increase its nutritional value. The various fatty acids and triglycerides are mainly obtained from microorganisms such as *Mortierella* or from oil-producing plants such as soybean, oilseed rape, sunflower and others, where they are usually obtained in the form of their triacylglycerides. Alternatively, they are obtained advantageously from animals, such as fish. The free fatty acids are prepared advantageously by hydrolysis.

Whether oils with unsaturated or with saturated fatty acids are preferred depends on the intended purpose; thus, for example, lipids with unsaturated fatty acids, specifically polyunsaturated fatty acids, are preferred in human nutrition since they have a positive effect on the cholesterol level in the blood and thus on the possibility of heart disease. They are used in a variety of dietetic foodstuffs or medicaments. In addition PUFAs are commonly used in food, feed and in the cosmetic industry. Poly unsaturated ω -3- and/or ω -6-fatty acids are an important part of animal feed and human food. Because of the common composition of human food poly unsaturated ω -3-fatty acids, which are an essential component of fish oil, should be added to the food to increase the nutritional value of the food; thus, for example, poly unsaturated fatty acids such as Docosahexaenoic acid (= DHA, C₂₂^{6⁴ 7¹⁰ 13¹⁶ 19}) or Eicosapentaenoic acid (= EPA, C₂₀^{5⁸ 11¹⁴ 17}) are added as mentioned above to infant formula to increase its nutritional value. Whereas DHA has a positive effect on the brain development of babies. The addition of poly unsaturated ω -3-fatty acids is preferred as the addition of poly unsaturated ω -6-fatty acids like Arachidonic acid (= ARA, C₂₀^{4⁵ 8¹¹ 14}) to common food have an undesired effect for example on rheumatic diseases such as rheumatoid arthritis. Poly unsaturated ω -3- and ω -6-fatty acids are precursor of a family of paracrine hormones called eicosanoids such as prostaglandins which are products of the metabolism of Dihomo- γ -linoleic acid, ARA or EPA. Eicosanoids are in-

volved in the regulation of lipolysis, the initiation of inflammatory responses, the regulation of blood circulation and pressure and other central functions of the body. Eicosanoids comprise prostaglandins, leukotrienes, thromboxanes, and prostacyclins. ω -3-fatty acids seem to prevent arteriosclerosis and cardiovascular diseases primarily
5 by regulating the levels of different eicosanoids. Other Eicosanoids are the thromboxanes and leukotrienes, which are products of the metabolism of ARA or EPA.

Principally microorganisms such as *Mortierella* or oil producing plants such as soybean, rapeseed or sunflower or algae such as *Cryptocodinium* or *Phaeodactylum* are a common source for oils containing PUFAs, where they are usually obtained in the
10 form of their triacyl glycerides. Alternatively, they are obtained advantageously from animals, such as fish. The free fatty acids are prepared advantageously by hydrolysis with a strong base such as potassium or sodium hydroxide.

Plant oils are in general rich in fatty acids such as monounsaturated fatty acids like oleic acid or poly unsaturated fatty acids (= PUFA) like linoleic or linolenic acid. LCPU-
15 FAs like arachidonic acid or eicosapentaenoic acid are rarely found in plants exceptions are some *Nephelium* and *Salvia* species in which arachidonic acid is found and some *Santalum* species in which eicosapentaenoic acid is found. The LCPUFA Docosahexaenoic acid is not found in plants. LCPUFAs such as DHA, EPA, ARA, Dihomo- γ -linoleic acid ($C_{20,3}^{\Delta 8,11,14}$) or Docosapentaenoic acid (= DPA, $C_{22,5}^{\Delta 7,10,13,16,19}$) are not
20 produced by oil producing plants such as soybean, rapeseed, safflower or sunflower. A natural sources for said fatty acids are fish for example herring, salmon, sardine, redfish, eel, carp, trout, halibut, mackerel, pike-perch, tuna or algae.

Approximately 80% of the oils and fats are used in the food industry. Nearly about 84 % of all world wide used vegetable oils are stemming from only six crops/oil crops,
25 which are soybean, oil palm, rapeseed, sunflower, cottonseed, and groundnut.

On account of their positive properties there has been no shortage of attempts in the past to make available genes which participate in the synthesis of fatty acids or triglycerides for the production of oils in various organisms having a modified content of unsaturated fatty acids. Thus, in WO 91/13972 and its US equivalent a Δ -9-
30 desaturase is described. In WO 93/1 1245 a Δ -15-desaturase and in WO 94/1 1516 a Δ -12-desaturase is claimed. WO 00/34439 discloses a Δ -5- and a Δ -8-desaturase.

Other desaturases are described, for example, in EP-A-O 550 162, WO 94/18337, WO 97/30582, WO 97/21340, WO 95/18222, EP-A-O 794 250, Stukey et al., J. Biol. Chem., 265, 1990: 20144-20149, Wada et al., Nature 347, 1990: 200-203 or Huang et al., Lipids 34, 1999: 649-659. To date, however, the various desaturases have
5 been only inadequately characterized biochemically since the enzymes in the form of membrane-bound proteins are isolable and characterizable only with very great difficulty (McKeon et al., Methods in Enzymol. 71, 1981 : 275-277, Wang et al., Plant Physiol. Biochem., 26, 1988: 777-792). Generally, membrane-bound desaturases are characterized by introduction into a suitable organism, which is then investigated for
10 enzyme activity by means of analysis of starting materials and products. Δ -6-Desaturases are described in WO 93/06712, US 5,614,393, US 5614393, WO 96/21022, WO0021557 and WO 99/271 11 and their application to production in transgenic organisms is also described, e.g. in WO 9846763, WO 9846764 and WO 9846765. At the same time the expression of various fatty acid biosynthesis
15 genes, as in WO 9964616 or WO 9846776, and the formation of poly-unsaturated fatty acids is also described and claimed. With regard to the effectiveness of the expression of desaturases and their effect on the formation of polyunsaturated fatty acids it may be noted that through expression of a desaturases and elongases as described to date only low contents of poly-unsaturated fatty acids/lipids, such as by way
20 of example eicosapentaenoic or arachidonic acid, have been achieved. Therefore, an alternative and more effective pathway with higher product yield is desirable.

Accordingly, there is still a great demand for new and more suitable genes, which encode enzymes, which participate in the biosynthesis of unsaturated fatty acids and make it possible to produce certain fatty acids specifically on an industrial scale without unwanted byproducts forming. In the selection of genes for biosynthesis two characteristics above all are particularly important. On the one hand, there is as ever a
25 need for improved processes for obtaining the highest possible contents of polyunsaturated fatty acids. Advantageously genes should be as selective as possible and should if possible have more than one activity in the fatty acid biosynthesis chain.

30 Accordingly, it is an object of the present invention to provide further genes of desaturase and elongase enzymes for the synthesis of polyunsaturated fatty acids in plants preferably in oilseed plants and to use them in a commercial process for the produc-

tion of PUFAs especially LCPUFAs. Said process should increase LCPUFA content in plants as much as possible preferably in seeds of an oil producing plant.

We have found that a process for the production of arachidonic acid or eicosapentaenoic acid achieves this object or arachidonic acid and eicosapentaenoic acid in transgenic plants that produces mature seeds with a content of at least 1 % by weight of said compounds referred to the total lipid content of said organism, which comprises the following steps:

- 10 a) introduction of at least one nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -12-desaturase and Δ -15-desaturase activity, and
- b) introduction of at least one second nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -9-elongase activity, and
- c) introduction of at least one third nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -8-desaturase activity, and
- 15 d) introduction of at least a one fourth nucleic acid sequence, which encodes a polypeptide having a Δ -5-desaturase activity, and
- e) cultivating and harvesting of said transgenic plant.

According to the invention the used nucleic acid sequences are isolated nucleic sequences coding for polypeptides having a Δ -12-desaturase- and Δ -15-desaturase-, Δ 9-elongase-, Δ -8 desaturase- or Δ 5-desaturase-activity.

Advantageously nucleic acid sequences are used in the abovementioned process of the invention, which encode polypeptides having Δ -12-desaturase and Δ -15-desaturase activity, Δ -8-desaturase, Δ -9-elongase or Δ -5-desaturase activity and which are selected from the group consisting of

- 25 a) a nucleic acid sequence depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO:13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23, and

- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 according to the degeneracy of the genetic code,
- c) derivatives of the nucleic acid sequences depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, , SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which encode polypeptides having at least 50 % homology to the sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having Δ -12-desaturase and Δ -15-desaturase activity, Δ -8-desaturase, Δ -9-elongase or Δ -5-desaturase activity.
- 15 In the inventive process the nucleic acid sequence encoding the bifunctional Δ -12-desaturase- and Δ -15-desaturase-enzyme leads to an increased flux from oleic acid (C18:1 Δ 9) to linolenic acid (C18:3 Δ^9 Δ^{12} Δ^{15}) and thereby to an increase of ω -3-fatty acids in comparison to the ω -6-fatty acids. Furthermore this bifunctional enzyme acts on C16-fatty acids having one double bond in the fatty acid molecule as well as on C18-
- 20 fatty acids having one double bond in the fatty acid molecule. This leads to a further increase in flux from precursor fatty acids such as C 18 fatty acids such as oleic acid towards C18 fatty acids such as linoleic and linolenic acid. This is especially of advantage in plants such as oilseed plants having a high content of oleic acid like such as those from the family of the Brassicaceae, such as the genus Brassica, for example
- 25 oilseed rape or canola; the family of the Elaeagnaceae, such as the genus Elaeagnus, for example the genus and species *Olea europaea*, or the family Fabaceae, such as the genus Glycine, for example the genus and species *Glycine max*, which are high in oleic acid. But also in other plants such oilseed plants like Brassica juncea, Camelina sativa, sunflower or safflower and all other plants mentioned herein this
- 30 leads to a higher amount of ω -3-fatty acids. By using said inventive nucleic acid sequence and the activity of its gene product ω -3-fatty acids to the ω -6-fatty acids are produced in at least a 1:2 ratio, preferably in at least a 1:3 or 1:4 ratio, more prefera-

bly in at least a 1:5 or 1:6 ratio. That means especially arachidonic acid and eicosapentaenoic acid are produced in at least a 1:2 ratio, preferably in at least a 1:3 or 1:4 ratio, more preferably in at least a 1:5 or 1:6 ratio.

In particular ω -3-fatty acids or ω -6-fatty acids molecules are produced in the inventive process, arachidonic acid and eicosapentaenoic acid are most preferred produced. We have found that this object is advantageously achieved by the combined expression of four isolated nucleic acid sequences according to the invention which encode for polypeptides having the following activities: a polypeptide with Δ -12-desaturase- and Δ -15-desaturase-activity, a polypeptide with a C18- Δ -9-elongase-activity, a polypeptide with C20- Δ -8-desaturase-activity and a C20- Δ -5-desaturase-activity. This objective was achieved in particular by the co-expression of the isolated nucleic acid sequences according to the invention. C 18 fatty acids with a single double bond in Δ -9-position are desaturated a first time to linoleic acid by the Δ -12-desaturase and Δ -15-desaturase and thereafter a second time to linolenic acid by the same enzyme advantageously used in the inventive process. The produced C 18 fatty acids linoleic and linolenic acid both having a double bond in Δ -9-position are then elongated by the Δ -9-elongase, which is advantageously used in the inventive process. By the Δ -8-desaturase used in the process a double bond in Δ -8-position is introduced into C20 fatty acids. In addition a double bond is introduced into the produced fatty acid molecules in Δ -5-position by the Δ -5-desaturase. The end products of the whole enzymatic reaction are arachidonic acid and eicosapentaenoic acid.

The ω -3-fatty acids or ω -6-fatty acids, preferably ω -3-fatty acids produced in the process are advantageously bound in membrane lipids and/or triacylglycerides or mixtures of different glycerides, but may also occur in the plants as free fatty acids or else bound in the form of other fatty acid esters.

The fatty acid esters with ω -3-fatty acids or ω -6-fatty acids especially arachidonic acid and eicosapentaenoic acid molecules can be isolated in the form of an oil or lipid, for example in the form of compounds such as sphingolipids, phosphoglycerides, lipids, glycolipids such as glycosphingolipids, phospholipids such as phosphatidylethanolamine, phosphatidylcholine, phosphatidylserine, phosphatidylglycerol, phosphatidylinositol or diphosphatidylglycerol, monoacylglycerides, diacylglycerides, triacylglycerides or other fatty acid esters such as the acetyl-coenzyme A esters from the plants

which have been used for the preparation of the fatty acid esters; preferably, they are isolated in the form of their diacylglycerides, triacylglycerides and/or in the form of phosphatidylcholine, especially preferably in the form of the triacylglycerides. In addition to these esters, the LCPUFAs are also present in the plants, advantageously in the oilseed plants as free fatty acids or bound in other compounds. As a rule, the various abovementioned compounds (fatty acid esters and free fatty acids) are present in the plants with an approximate distribution of 80 to 90% by weight of triglycerides, 2 to 5% by weight of diglycerides, 5 to 10% by weight of monoglycerides, 1 to 5% by weight of free fatty acids, 2 to 8% by weight of phospholipids, the total of the various compounds amounting to 100% by weight.

In the inventive process(es) [the singular shall include the plural and vice versa] the LCPUFAs are produced in a content of at least 1 % by weight, preferably at least 2, 3, 4 or 5 % by weight, more preferably at least 6, 7, 8, or 9 % by weight, most preferably 10, 20 or 30 % by weight referred to the total lipid content of the plant used in the process. That means Arachidonic acid and eicosapentaenoic acid are produced in a content of at least 1 % by weight, preferably at least 2, 3, 4 or 5 % by weight, more preferably at least 6, 7, 8, or 9 % by weight, most preferably 10, 20 or 30 % by weight referred to the total lipid content. Preferred starting material for the inventive process is oleic acid (C18:1), which is transformed to the preferred end products ARA or EPA. As for the inventive process plants are used the product of the process is not a product of one pure substance per se. It is a mixture of different substances where one or more compounds are the major product and others are only contained as side products. Advantageously the side products shall not exceed 20 % by weight referred to the total lipid content of the plant, preferably the side products shall not exceed 15 % by weight, more preferably they shall not exceed 10 % by weight, most preferably they shall not exceed 5 % by weight. In the event that a mixture of different fatty acids such as ARA and EPA are the product of the inventive process said fatty acids can be further purified by method known by a person skilled in the art such as distillation, extraction, crystallization at low temperatures, chromatography or a combination of said methods. These chemically pure fatty acids or fatty acid compositions are advantageous for applications in the food industry sector, the cosmetic sector and especially the pharmacological industry sector.

Fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise 6 to 15% of palmitic acid, 1 to 6% of stearic acid, 7 to 85% of oleic acid, 0.5 to 8% of vaccenic acid, 0.1 to 1% of arachic acid, 7 to 25% of saturated fatty acids, 8 to 85% of monounsaturated fatty acids and 60 to 85% of polyunsaturated fatty acids including LCPUFAs, in each case based on 100% and on the total fatty acid content of the organisms. Advantageous LCPUFAs, which are present in the fatty acid esters or fatty acid mixtures are preferably at least 1%, 2%, 3%, 4% or 5% by weight of arachidonic acid and/or preferably at least 5%, 6%, 7%, 8%, 9% or 10% by weight of eicosapentaenoic acid, based on the total fatty acid content.

Moreover, the fatty acid esters or fatty acid mixtures which have been produced by the process of the invention advantageously comprise fatty acids selected from the group of the fatty acids erucic acid (13-docosaenoic acid), sterculic acid (9,10-methyleneoctadec-9-enoic acid), malvalic acid (8,9-methyleneheptadec-8-enoic acid), chaulmoogric acid (cyclopentenedodecanoic acid), furan fatty acid (9,12-epoxyoctadeca-9,11-dienoic acid), vernolic acid (9,10-epoxyoctadec-12-enoic acid), tariric acid (6-octadecynoic acid), 6-nonadecynoic acid, santalbic acid (α -11-octadecen-9-ynoic acid), 6,9-octadecenynoic acid, pyruvic acid (α -10-heptadecen-8-ynoic acid), crepenynic acid (9-octadecen-12-ynoic acid), 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diynoic acid, petroselenic acid (cis-6-octadecenoic acid), 9c,12t-octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid), catalpic acid (9t11t13c-octadecatrienoic acid), eleostearic acid (9c11t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), punitic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid), pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or coriolic acid (13-hydroxy-9c,11t-octadecadienoic acid). The abovementioned fatty acids are, as a rule, advantageously only found in traces in the fatty acid esters or fatty acid mixtures produced by the process according to the invention, that is to say that, based on the total fatty acids, they occur to less than 30%, preferably to less than 25%, 24%, 23%, 22% or 21%, especially preferably to less than 20%, 15%, 10%, 9%, 8%, 7%, 6% or 5%, very especially preferably to less than 4%, 3%, 2% or 1%. In a further preferred form of the invention, these abovementioned fatty acids occur to less than 0.9%, 0.8%, 0.7%, 0.6% or 0.5%, especially preferably to less than 0.4%, 0.3%, 0.2%, 0.1%, based on the total fatty acids. The fatty

acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise less than 0.1 %, based on the total fatty acids, and/or no butyric acid, no cholesterol, no clupanodonic acid (= docosapentaenoic acid, C22:5^{Δ4,8,12,15,21}) and no nisinic acid (tetracosahexaenoic acid, C23:6^{Δ3,8,12,15,18,21}).

- 5 The isolated nucleic acid sequences used in the process according to the invention encode proteins or parts of these, where the proteins or the individual protein or parts thereof comprise(s) an amino acid sequence with sufficient homology to an amino acid sequence which is shown in the sequences SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 and SEQ ID NO: 24 so that
- 10 the proteins or parts thereof retain a Δ -12-desaturase and Δ -15-desaturase-, Δ -9-elongase-, Δ -8-desaturase- and/or Δ -5-desaturase activity. The proteins or parts thereof which is/are encoded by the nucleic acid molecule(s) preferably retains their essential enzymatic activity and the ability of participating in the metabolism of
- 15 compounds required for the synthesis of cell membranes or lipid bodies in organisms, advantageously in plants, or in the transport of molecules across these membranes. Advantageously, the proteins encoded by the nucleic acid molecules have at least approximately 50%, preferably at least approximately 60% and more preferably at least approximately 70%, 80% or 90% and most preferably at least approximately
- 20 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity with the amino acid sequences shown in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 and SEQ ID NO: 24. For the purposes of the invention, homology or homologous is understood as
- 25 meaning identity or identical, respectively.

The homology was calculated over the entire amino acid or nucleic acid sequence region. The skilled worker has available a series of programs which are based on various algorithms for the comparison of various sequences. Here, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. The

30 program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489

(1981)], which are part of the GCG software packet [Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program GAP and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000
5 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for the sequence alignments.

Moreover, in the process of the invention advantageously nucleic acid sequences are used which differ from one of the nucleotide sequences shown in SEQ ID NO: 1, SEQ
10 ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23 (and parts thereof) owing to the degeneracy of the genetic code and which thus encode the same Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase as those encoded by the nucleotide sequences
15 shown in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23.

Suitable plants for the production in the process according to the invention are, in principle all plants that produces mature seeds especially crop plants such as oilseed
20 plants.

Plants which are suitable are, in principle, all those plants which are capable of synthesizing fatty acids and that produce mature seeds, such as all dicotyledonous or monocotyledonous plants. Advantageous plants are selected from the group consisting of the plant families Anacardiaceae, Asteraceae, Apiaceae, Boraginaceae, Brassicaceae, Cannabaceae, Elaeagnaceae, Euphorbiaceae, Fabaceae, Geraniaceae,
25 Gramineae, Juglandaceae, Leguminosae, Linaceae, Lythraeae, Malvaceae, Onagraceae, Palmae, Poaceae, Rubiaceae, Scrophulariaceae, Solanaceae, Sterculiaceae and Theaceae or vegetable plants or ornamentals. More preferred plants are selected from the group consisting of the plant genera of Pistacia, Mangifera, Anacardium, Calendula, Carthamus, Centaurea, Cichorium, Cynara, Helianthus, Lactuca,
30 Locusta, Tagetes, Valeriana, Borago, Daucus, Brassica, Camelina, Melanosinapis, Sinapis, Arabidopsis, Orychophragmus, Cannabis, Elaeagnus, Manihot, Janipha,

Jatropha, Ricinus, Pisum, Albizia, Cathormion, Feuillea, Inga, Pithecolobium, Acacia, Mimosa, Medicago, Glycine, Dolichos, Phaseolus, Pelargonium, Cocos, Oleum, Juglans, Wallia, Arachis, *Linum*, Punica, Gossypium, Camissonia, Oenothera, Elaeis, Hordeum, Secale, Avena, Sorghum, Andropogon, Holcus, Panicum, Oryza, Zea, Triticum, Coffea, Verbascum, Capsicum, Nicotiana, Solanum, Lycopersicon, Theobroma and Camellia.

Examples which may be mentioned are the following plants selected from the group consisting of Anacardiaceae such as the genera Pistacia, Mangifera, Anacardium, for example the genus and species Pistacia vera [pistachio], Mangifera indica [mango] or Anacardium occidentale [cashew], Asteraceae, such as the genera Calendula, Carthamus, Centaurea, Cichorium, Cynara, Helianthus, Lactuca, Locusta, Tagetes, Valeriana, for example the genus and species Calendula officinalis [common marigold], Carthamus tinctorius [safflower], Centaurea cyanus [cornflower], Cichorium intybus [chicory], Cynara scolymus [artichoke], Helianthus annuus [sunflower], Lactuca sativa, *Lactuca crispera*, *Lactuca esculenta*, *Lactuca scariola L. ssp. sativa*, *Lactuca scariola L. var. integrata*, *Lactuca scariola L. var. integrifolia*, *Lactuca sativa subsp. romana*, *Locusta communis*, *Valeriana locusta* [salad vegetables], Tagetes lucida, Tagetes erecta or Tagetes tenuifolia [african or french marigold], Apiaceae, such as the genus Daucus, for example the genus and species Daucus carota [carrot], Boraginaceae, such as the genus Borago, for example the genus and species Borago officinalis [borage], Brassicaceae, such as the genera Brassica, Camelina, *Melanosinapis*, Sinapis, Arabidopsis, for example the genera and species Brassica napus, Brassica rapa ssp. [oilseed rape], Sinapis arvensis Brassica juncea, Brassica juncea var. juncea, Brassica juncea var. crispifolia, Brassica juncea var. foliosa, Brassica nigra, *Brassica sinapioides*, Camelina sativa, *Melanosinapis communis* [mustard], Brassica oleracea [fodder beet] or Arabidopsis thaliana, Cannabaceae, such as the genus Cannabis, such as the genus and species Cannabis sativa [hemp], Elaeagnaceae, such as the genus Elaeagnus, for example the genus and species Olea europaea [olive], Euphorbiaceae, such as the genera Manihot, *Janipha*, *Jatropha*, Ricinus, for example the genera and species Manihot utilissima, *Janipha manihot*, *Jatropha manihot*, *Manihot aipil*, *Manihot dulcis*, *Manihot manihot*, *Manihot melanobasis*, Manihot esculenta [cassava] or Ricinus communis [castor-oil plant], Fabaceae, such as the genera Pisum, Albizia, *Cathormion*, *Feuillea*, Inga,

Pithecolobium, Acacia, Mimosa, Medicago, Glycine, Dolichos, Phaseolus, soybean, for example the genera and species *Pisum sativum*, *Pisum arvense*, *Pisum humile* [pea], *Albizia berteriana*, *Albizia julibrissin*, *Albizia lebeck*, *Acacia berteriana*, *Acacia littoralis*, *Albizia berteriana*, *Albizia berteriana*, *Cathormion berteriana*, *Feuillea berteriana*, *Inga fragrans*, *Pithecellobium berterianum*, *Pithecellobium fragrans*, *Pithecolobium berterianum*, *Pseudalbizia berteriana*, *Acacia julibrissin*, *Acacia nemu*, *Albizia nemu*, *Feuillea julibrissin*, *Mimosa julibrissin*, *Mimosa speciosa*, *Sericanrda julibrissin*, *Acacia lebeck*, *Acacia macrophylla*, *Albizia lebeck*, *Feuillea lebeck*, *Mimosa lebeck*, *Mimosa speciosa*, *Medicago sativa*, *Medicago falcata*, *Medicago varia* [alfalfa] *Glycine max* *Dolichos soja*, *Glycine gracilis*, *Glycine hispida*, *Phaseolus max*, *Soja hispida* or *Soja max* [soybean], Geraniaceae, such as the genera *Pelargonium*, *Cocos*, *Oleum*, for example the genera and species *Cocos nucifera*, *Pelargonium grossularioides* or *Oleum cocois* [coconut], Gramineae, such as the genus *Saccharum*, for example the genus and species *Saccharum officinarum*, *Juglandaceae*, such as the genera *Juglans*, *Wallia*, for example the genera and species *Juglans regia*, *Juglans ailanthifolia*, *Juglans sieboldiana*, *Juglans cinerea*, *Wallia cinerea*, *Juglans bixbyi*, *Juglans californica*, *Juglans hindsii*, *Juglans intermedia*, *Juglans jamaicensis*, *Juglans major*, *Juglans microcarpa*, *Juglans nigra* or *Wallia nigra* [walnut], Leguminosae, such as the genus *Arachis*, for example the genus and species *Arachis hypogaea* [peanut], Linaceae, such as the genera *Adenolinum*, for example the genera and species *Linum usitatissimum*, *Linum humile*, *Linum austriacum*, *Linum bienne*, *Linum angustifolium*, *Linum catharticum*, *Linum flavum*, *Linum grandiflorum*, *Adenolinum grandiflorum*, *Linum lewisii*, *Linum narbonense*, *Linum perenne*, *Linum perenne var. lewisii*, *Linum pratense* or *Linum trigynum* [linseed], Lythraeae, such as the genus *Punica*, for example the genus and species *Punica granatum* [pomegranate], Malvaceae, such as the genus *Gossypium*, for example the genera and species *Gossypium hirsutum*, *Gossypium arboreum*, *Gossypium barbadense*, *Gossypium herbaceum* or *Gossypium thurberi* [cotton], Onagraceae, such as the genera *Camissonia*, *Oenothera*, for example the genera and species *Oenothera biennis* or *Camissonia brevipes* [evening primrose], Palmae, such as the genus *Elaeis*, for example the genus and species *Elaeis guineensis* [oil palm], Poaceae, such as the genera *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea* (maize), *Triticum*, for example the genera

and species *Hordeum vulgare*, *Hordeum jubatum*, *Hordeum murinum*, *Hordeum secalinum*, *Hordeum distichon* *Hordeum aegiceras*, *Hordeum hexastichon*, *Hordeum hexastichum*, *Hordeum irregulare*, *Hordeum sativum*, *Hordeum secalinum* [barley], *Secale cereale* [rye], *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida* [oats], *Sorghum bicolor*, *Sorghum halepense*, *Sorghum saccharatum*, *Sorghum vulgare*, *Andropogon drummondii*, *Holcus bicolor*, *Holcus sorghum*, *Sorghum aethiopicum*, *Sorghum arundinaceum*, *Sorghum caffrorum*, *Sorghum cernuum*, *Sorghum dochna*, *Sorghum drummondii*, *Sorghum durra*, *Sorghum guineense*, *Sorghum lanceolatum*, *Sorghum nervosum*, *Sorghum saccharatum*, *Sorghum subglabrescens*, *Sorghum verticilliflorum*, *Sorghum vulgare*, *Holcus halepensis*, *Sorghum miliaceum*, *Panicum militaceum* [millet], *Oryza sativa*, *Oryza latifolia* [rice], *Zea mays* [maize] *Triticum aestivum*, *Triticum durum*, *Triticum turgidum*, *Triticum hybernum*, *Triticum macha*, *Triticum sativum* or *Triticum vulgare* [wheat], Rubiaceae, such as the genus *Coffea*, for example the genera and species *Coffea* spp., *Coffea arabica*, *Coffea canephora* or *Coffea liberica* [coffee], Scrophulariaceae, such as the genus *Verbascum*, for example the genera and species *Verbascum blattaria*, *Verbascum chaixii*, *Verbascum densiflorum*, *Verbascum lagurus*, *Verbascum longifolium*, *Verbascum lychnitis*, *Verbascum nigrum*, *Verbascum olympicum*, *Verbascum phlomoides*, *Verbascum phoenicum*, *Verbascum pulverulentum* or *Verbascum thapsus* [verbascum], Solanaceae, such as the genera *Capsicum*, *Nicotiana*, *Solanum*, *Lycopersicon*, for example the genera and species *Capsicum annuum*, *Capsicum annuum* var. *glabriusculum*, *Capsicum frutescens* [pepper], *Capsicum annuum* [paprika], *Nicotiana tabacum*, *Nicotiana alata*, *Nicotiana attenuata*, *Nicotiana glauca*, *Nicotiana langsdorffii*, *Nicotiana obtusifolia*, *Nicotiana quadrivalvis*, *Nicotiana repanda*, *Nicotiana rustica*, *Nicotiana sylvestris* [tobacco], *Solanum tuberosum* [potato], *Solanum melongena* [eggplant] *Lycopersicon esculentum*, *Lycopersicon lycopersicum*, *Lycopersicon pyriforme*, *Solanum integrifolium* or *Solanum lycopersicum* [tomato], Sterculiaceae, such as the genus *Theobroma*, for example the genus and species *Theobroma cacao* [cacao] or Theaceae, such as the genus *Camellia*, for example the genus and species *Camellia sinensis* [tea].

Plants which are especially advantageously used in the process according to the invention are plants which belong to the oil-producing plants, that is to say which are

used for the production of oil, such as oilseed or oil crop plants which comprise large amounts of lipid compounds, such as peanut, oilseed rape, canola, sunflower, safflower (*Carthamus tinctoria*), poppy, mustard, hemp, castor-oil plant, olive, sesame, Calendula, Punica, evening primrose, verbascum, thistle, wild roses, hazelnut, almond, macadamia, avocado, bay, pumpkin/squash, linseed, soybean, pistachios, borage, trees (oil palm, coconut or walnut) or arable crops such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, Tagetes, Solanaceae plants such as potato, tobacco, eggplant and tomato, Vicia species, pea, alfalfa or bushy plants (coffee, cacao, tea), SaNx species, and perennial grasses and fodder crops. Preferred plants according to the invention are oil crop plants such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, Calendula, Punica, evening primrose, pumpkin/squash, linseed, soybean, borage, trees (oil palm, coconut). Especially preferred are plants which are high in C18:1-, C18:2- and/or C18:3-fatty acids, such as oilseed rape, canola, Brassica juncea, Camelina sativa, Orychophragmus, sunflower, safflower, tobacco, verbascum, sesame, cotton, pumpkin/squash, poppy, evening primrose, walnut, linseed, hemp or thistle. Very especially preferred plants are plants such as rapeseed, canola, safflower, sunflower, poppy, mustard, hemp, evening primrose, walnut, linseed or hemp. Other preferred plants are castor bean, sesame, olive, calendula, punica, hazel nut, maize, almond, macadamia, cotton, avocado, pumpkin, laurel, pistachio, oil palm, peanut, soybean, marigold, coffee, tobacco, cacao and borage

For the production of further ω -6- and/or ω -3-fatty acids it is advantageously to introduce further nucleic fatty acid sequences, which encode other enzymes of the fatty acids synthesis chain such as preferably Δ -5-elongase(s) and/or Δ -4-desaturase(s) [for the purposes of the present invention, the plural is understood as comprising the singular and vice versa]. Other Genes of the fatty acid or lipid metabolism, which can be introduced are selected from the group consisting of acyl-CoA dehydrogenase(s), acyl-ACP [= acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyl transferase(s), acyl-CoA:lysophospholipid acyltransferases, fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allenoxide synthases, hydroperoxide lyases or fatty acid elongase(s). Preferred nucleic acid sequences, which can be used in addition in the inventive

process, are disclosed in the sequence protocol of WO2005/012316 and in Table 1 of the specification of said application, these sequences are hereby incorporated by reference.

Transgenic plants are to be understood as meaning single plant cells, certain tissues,
5 organs or parts of plants and their cultures on solid media or in liquid culture, parts of plants and entire plants such as plant cell cultures, protoplasts from plants, callus cultures or plant tissues such as leaves, stem, shoots, seeds, flowers, roots, tubers etc. Said transgenic plants can be cultivated for example on solid or liquid culture medium, in soil or in hydroponics. Plants in the sense of the invention also include plant cells
10 and certain tissues, organs and parts of plants in all their phenotypic forms such as anthers, fibers, root hairs, stalks, embryos, calli, cotyledons, petioles, harvested material, plant tissue, reproductive tissue such as seeds and cell cultures which are derived from the actual transgenic plant and/or can be used for bringing about the transgenic plant. In this context, the seed comprises all parts of the seed such as the seed coats,
15 epidermal cells, seed cells, endosperm or embryonic tissue.

For the purposes of the invention, "transgenic" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette (= gene construct) or a vector comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, gene constructs or vectors as described herein according to
20 the invention, all those constructions brought about by recombinant methods in which either

- a) the nucleic acid sequence according to the invention, or
- b) a genetic control sequence which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or
- 25 c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. The natural genetic environment is understood as meaning the
30 natural genomic or chromosomal locus in the original plant or the presence in a

genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, most preferably at least 5000 bp. A naturally occurring expression cassette - for example the naturally occurring combination of the natural promoter of the nucleic acid sequences with the corresponding Δ 12-desaturase and Δ 15-desaturase-, Δ 9-elongase-, Δ 8-desaturase- and/or Δ 5-desaturase-genes - becomes a transgenic expression cassette when this expression cassette is modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in US 5,565,350 or WO 00/15815.

A transgenic plant for the purposes of the invention is therefore understood as meaning, as above, that the nucleic acids used in the process are not at their natural locus in the genome of a plant, it being possible for the nucleic acids to be expressed homologously or heterologously. However, as mentioned, transgenic also means that, while the nucleic acids according to the invention are at their natural position in the genome of a plant, the sequence has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified. Transgenic is preferably understood as meaning the expression of the nucleic acids according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acids takes place. Preferred transgenic organisms are oilseed crops.

After cultivation transgenic plants which are used in the inventive process can be brought to the market without isolating the ω -6- and/or ω -3-fatty acids preferably the arachidonic and/or eicosapentaenoic acid. Preferably the ω -6- and/or ω -3-fatty acids are isolated from the plant in the form of their free fatty acids, their lipids or oils. The purification can be done by conventional methods such as squeezing and extraction of the plants or other methods instead of the extraction such as distillation, crystallization at low temperatures, chromatography or a combination of said methods. Advantageously the plants are grinded, heated and/or vaporized before the squeezing and extraction procedure. As solvent for the extraction solvents such as hexane or other solvents having a similar extraction behavior are used. The isolated oils are further purified by acidification with for example phosphoric acid. The free fatty acids are produced from said oils or lipids by hydrolysis. Charcoal or diatom earth is used to remove dyes from the fluid. In another preferred embodiment of the inventive process the alkyl ester of the fatty acids are produced from the oils and lipids by transesterification with an enzyme or with conventional chemistry. A preferred method is the production of the alkyl ester in the presence of alcoholates of the corresponding lower alcohols (C1 to C10 alcohols such as methanol, ethanol, propanol, butanol, hexanol etc.) such as methanolate or ethanolate. Therefore as the skilled worker knows the alcohol in the presence of a catalytic amount of a base such as NaOH or KOH is added to the oils or lipids.

In a preferred form of the inventive process the lipids can be obtained in the usual manner after the plants have been grown. To this end, the organisms can first be harvested and then disrupted, or they can be used directly. In the case of plant cells, plant tissue or plant organs, "growing" is understood as meaning, for example, the cultivation on or in a nutrient medium, or of the intact plant on or in a substrate, for example in a hydroponic culture, potting compost or on arable land. It is advantageous to extract the lipids with suitable solvents such as apolar solvents, for example hexane, or polar solvents, for example ethanol, isopropanol, or mixtures such as hexane/isopropanol, phenol/chloroform/isoamyl alcohol, at temperatures between 0°C and 80°C, preferably between 20°C and 50°C. As a rule, the biomass is extracted with an excess of solvent, for example with an excess of solvent to biomass of 1:4. The solvent is subsequently removed, for example by distillation. The extraction may also be carried out with supercritical CO₂. After the extraction, the remainder of the biomass can be removed, for example, by filtration. Standard methods for the extraction of fatty acids from plants and microorganisms are described in Bligh et al. (Can. J. Biochem. Physiol. 37, 1959: 911-917) or Vick et al. (Plant Physiol. 69, 1982: 1103-1108).

The crude oil thus obtained can then be purified further, for example by removing cloudiness by adding polar solvents such as acetone or apolar solvents such as chloroform, followed by filtration or centrifugation. Further purification via columns or other techniques is also possible.

To obtain the free fatty acids from the triglycerides, the latter are hydrolyzed in the customary manner, for example using NaOH or KOH.

In the inventive process oils, lipids and/or free fatty acids or fractions thereof are produced. Said products can be used for the production of feed and food products, cosmetics or pharmaceuticals.

The oils, lipids, LCPUFAs or fatty acid compositions produced according to the inventive process can be used in the manner with which the skilled worker is familiar for mixing with other oils, lipids, fatty acids or fatty acid mixtures of animal origin, such as, for example, fish oils and/or microbial oils such as from *Mortierella* or *Cryptocodium*. These oils, lipids, fatty acids or fatty acid mixtures, which are

composed of vegetable, microbial and/or animal constituents, may also be used for the preparation of feedstuffs, foodstuffs, cosmetics or pharmaceuticals.

The term "oil", "lipid" or "fat" is understood as meaning a fatty acid mixture comprising unsaturated, saturated, preferably esterified, fatty acid(s). The oil, lipid, fat, fatty acid
5 and/or fatty acid composition is preferably high in polyunsaturated (PUFA and/or LCPUFA) free and/or, advantageously, esterified fatty acid(s), in particular oleic acid, linoleic acid, α -linolenic acid, arachidonic acid and/or eicosatetraenoic acid.

Transgenic plants which comprise the LCPUFAs synthesized in the process according to the invention can also advantageously be marketed directly without there being any
10 need for the oils, lipids or fatty acids synthesized to be isolated.

However, the LCPUFAs produced in the process according to the invention can also be isolated from the plants as described above, in the form of their oils, fats, lipids and/or free fatty acids. Polyunsaturated fatty acids produced by this process can be obtained by harvesting the crop in which they grow, or from the field. This can be
15 done via pressing or extraction of the plant parts, preferably the plant seeds. In this context, the oils, fats, lipids and/or free fatty acids can be obtained by what is known as cold-beating or cold-pressing without applying heat. To allow for greater ease of disruption of the plant parts, specifically the seeds, they are previously comminuted, steamed or roasted. The seeds, which have been pretreated in this manner can
20 subsequently be pressed or extracted with solvents such as warm hexane. The solvent is subsequently removed. In the case of microorganisms, the latter are, after harvesting, for example extracted directly without further processing steps or else, after disruption, extracted via various methods with which the skilled worker is familiar. In this manner, more than 96% of the compounds produced in the process can be
25 isolated. Thereafter, the resulting products are processed further, i.e. refined. In this process, substances such as the plant mucilages and suspended matter are first removed. What is known as desliming can be effected enzymatically or, for example, chemico-physically by addition of acid such as phosphoric acid. Thereafter, the free fatty acids are removed by treatment with a base, for example sodium hydroxide
30 solution. The resulting product is washed thoroughly with water to remove the alkali remaining in the product and then dried. To remove the pigment remaining in the product, the products are subjected to bleaching, for example using filler's earth or

active charcoal. At the end, the product is deodorized, for example using steam.

The preferred biosynthesis site of the fatty acids, oils, lipids or fats in the plants which are advantageously used is, for example, in general the seed or cell strata of the seed, so that seed-specific expression of the nucleic acids used in the process makes
5 sense. However, it is obvious that the biosynthesis of fatty acids, oils or lipids need not be limited to the seed tissue, but can also take place in a tissue-specific manner in all the other parts of the plant, for example in epidermal cells or in the tubers.

In principle, the LCPUFAs produced by the process according to the invention in the organisms used in the process can be increased in two different ways.

10 Advantageously, the pool of free polyunsaturated fatty acids and/or the content of the esterified polyunsaturated fatty acids produced via the process can be enlarged. Advantageously, the pool of esterified polyunsaturated fatty acids in the transgenic plants is enlarged by the process according to the invention.

In principle all nucleic acids encoding polypeptides with Δ -8-desaturase, Δ -9-elongase
15 and/or Δ -5-desaturase activity can be used in the inventive process. Preferably the nucleic acid sequences can be isolated for example from microorganism or plants such as fungi like *Mortierella*, algae like *Euglena*, *Cryptocodium* or *Isochrysis*, diatoms like *Phaeodactylum*, protozoa like amoeba such as *Acanthamoeba* or *Perkinsus* or mosses like *Physcomitrella* or *Ceratodon*, but also non-human animals such as
20 *Caenorhabditis* are possible as source for the nucleic acid sequences. Advantageous nucleic acid sequences according to the invention which encode polypeptides having a Δ -8-desaturase, Δ -9-elongase and/or Δ -5-desaturase activity are originate from microorganisms or plants, advantageously *Phaeodactylum tricornutum*, *Ceratodon purpureus*, *Physcomitrella patens*, *Euglena gracilis*, *Acanthamoeba castellanii*, *Perkinsus*
25 *marinus* or *Isochrysis galbana*. Thus, the co expression of a C18-specific Δ -12-desaturase and Δ -15-desaturase, a C18-specific Δ -9 elongase, a C20-specific Δ -8-desaturase and a C20-specific Δ -5-desaturase leads to the formation of Arachidonic acid (C20:6n-4, Δ 5, 8, 11, 14) and/or Eicosapentaenoic acid (C20:3n-5, Δ 5, 8, 11, 14, 17). Most preferred are the sequences mentioned in the sequence protocol.

30 In another embodiment the invention furthermore relates to isolated nucleic acid sequences encoding polypeptides with Δ -12-desaturase and Δ -15-desaturase-, Δ -9-

elongase-, Δ -8-desaturase- and/or Δ -5-desaturase-activity.

In one embodiment the invention relates to an isolated nucleic acid sequence which encodes a polypeptide having a Δ -12-desaturase and Δ -15-desaturase activity selected from the group consisting of

- 5 a) a nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23;
 - b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24;
 - 10 c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 22 which encode polypeptides having at least 40 % homology to the sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having Δ -12-desaturase and Δ -15-desaturase activity.
- 15 This inventive Δ -12-desaturase and Δ -15-desaturase is able to desaturate C16-fatty acids having at least one double bond in the fatty acid chain and/or C18-fatty acids having at least one double bond in the fatty acid chain. Preferably C16- and/or C18-fatty acids having only one double bond in the fatty acid chain are desaturated. This activity leads to an increase in flux from precursor fatty acids such as C18-fatty acids
- 20 towards C18-fatty acids having more than one double bond in the fatty acid chain such as linoleic and/or linolenic acid. C18-fatty acids are more preferred in the reaction than C16-fatty acids. C18-fatty acids are more than doubled preferred.

In another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -9-elongase selected from the

25 group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 11;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 12;

- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 11 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 12 and which polypeptides having Δ -9-elongase activity.

In yet another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -8-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8 and which polypeptides having Δ -8-desaturase activity.

Further in another embodiment the invention relates to an isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -5-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18;
- c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18 and which polypeptides having Δ -5-desaturase activity.

By derivative(s) of the sequences according to the invention is meant, for example, functional homologues of the polypeptides or enzymes encoded by SEQ ID NO: 3,

SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which exhibit the same said specific enzymatic activity. This specific enzymatic activity allows advantageously the synthesis of LCPUFAs of the ω -6- and/or ω -3-pathway of the fatty acid synthesis chain such as ARA and/or EPA. The said sequences encode enzymes which exhibit Δ -12-desaturase and Δ -15-desaturase-, Δ -9-elongase-, Δ -8-desaturase- and/or Δ -5-desaturase-activity.

The enzyme according to the invention, Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase and/or Δ -5-desaturase, advantageously either elongates fatty acid chains with 18 carbon atoms (see SEQ ID NO: 11) or introduces a double bond into fatty acid residues of glycerolipids, free fatty acids or acyl-CoA fatty acids at position C₈-C₉ (see SEQ ID NO: 3, 5 or 7) or at position C₅-C₆ (see SEQ ID NO: 15 or 17) or at position C_{i2}-C_{i3} and C_{i5}-C_{i6} of the fatty acid chain (see SEQ ID NO: 19, 21 or 23).

The inventive nucleic acid molecules, for example a nucleic acid molecule with a nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or of a part thereof can be isolated using molecular-biological standard techniques and the sequence information provided herein. Also, for example a homologous sequence or homologous, conserved sequence regions can be identified at the DNA or amino acid level with the aid of comparative algorithms. They can be used as hybridization probe and standard hybridization techniques (such as, for example, those described in Sambrook et al., Molecular Cloning: A Laboratory Manual. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) for isolating further nucleic acid sequences which can be used in the process. Moreover, a nucleic acid molecule comprising a complete sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or a part thereof can be isolated by polymerase chain reaction, where oligonucleotide primers which are used on the basis of this sequence or parts thereof (for example a nucleic acid molecule comprising the complete sequence or part thereof can be isolated by polymerase chain reaction using oligonucleotide primers which have been generated based on this

same sequence). For example, mRNA can be isolated from cells (for example by means of the guanidinium thiocyanate extraction method of Chirgwin et al. (1979) Biochemistry 18:5294-5299) and cDNA by means of reverse transcriptase (for example Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide primers for the amplification by means of polymerase chain reaction can be generated based on one of the sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or with the aid of the amino acid sequences detailed in SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24. A nucleic acid according to the invention can be amplified by standard PCR amplification techniques using cDNA or, alternatively, genomic DNA as template and suitable oligonucleotide primers. The nucleic acid amplified thus can be cloned into a suitable vector and characterized by means of DNA sequence analysis. Oligonucleotides, which correspond to a desaturase nucleotide sequence can be generated by standard synthetic methods, for example using an automatic DNA synthesizer.

Homologs of the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase nucleic acid sequences with the sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 means, for example, allelic variants with at least approximately 50 or 60%, preferably at least approximately 60 or 70%, more preferably at least approximately 70 or 80%, 90% or 95% and even more preferably at least approximately 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95 %, 96%, 97%, 98%, 99% or more identity or homology with a nucleotide sequence shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or its homologs, derivatives or analogs or parts thereof. Furthermore, isolated nucleic acid molecules of a nucleotide sequence which hybridize with one of the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or with a part thereof, for example hybridized under stringent conditions. A part thereof is understood as meaning, in accordance with the invention, that at least 25

base pairs (= bp), 50 bp, 75 bp, 100 bp, 125 bp or 150 bp, preferably at least 175 bp, 200 bp, 225 bp, 250 bp, 275 bp or 300 bp, especially preferably 350 bp, 400 bp, 450 bp, 500 bp or more base pairs are used for the hybridization. It is also possible and advantageous to use the full sequence. Allelic variants comprise in particular

5 functional variants which can be obtained by deletion, insertion or substitution of nucleotides from/into the sequence detailed in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, it being intended, however, that the enzyme activity of the resulting proteins which are synthesized is advantageously retained for the insertion

10 of one or more genes. Proteins which retain the enzymatic activity of the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase, i.e. whose activity is essentially not reduced, means proteins with at least 10%, preferably 20%, especially preferably 30%, very especially preferably 40% of the original enzyme activity in comparison with the protein encoded by SEQ ID NO: 3,

15 SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23. The homology was calculated over the entire amino acid or nucleic acid sequence region. The skilled worker has available a series of programs which are based on various algorithms for the comparison of various sequences. Here, the algorithms of Needleman and Wunsch or Smith and

20 Waterman give particularly reliable results. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981)), which are part of the GCG software

25 packet [Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 5371 1 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program GAP and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless

30 otherwise specified, these settings were always used as standard settings for the sequence alignments.

Homologs of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 means for example also bacterial, fungal and plant homologs, truncated sequences, single-

stranded DNA or RNA of the coding and noncoding DNA sequence.

Homologs of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 also means derivatives such as, for example, promoter variants. The promoters upstream
5 of the nucleotide sequences detailed can be modified by one or more nucleotide exchanges, by insertion(s) and/or deletion(s) without the functionality or activity of the promoters being adversely affected, however. It is furthermore possible that the modification of the promoter sequence enhances their activity or that they are replaced entirely by more active promoters, including those from heterologous
10 organisms.

In a further embodiment, derivatives of the nucleic acid molecule according to the invention represented in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 encode proteins with at least 40%, advantageously approximately 50 or 60%,
15 advantageously at least approximately 60 or 70% and more preferably at least approximately 70 or 80%, 80 to 90%, 90 to 95% and most preferably at least approximately 96%, 97%, 98%, 99% or more homology (= identity) with a complete amino acid sequence of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO:
20 24. The homology was calculated over the entire amino acid or nucleic acid sequence region. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981)), which are part of the GCG software packet [Genetics Computer
25 Group, 575 Science Drive, Madison, Wisconsin, USA 5371 1 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program BestFit and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings
30 were always used as standard settings for the sequence alignments.

Moreover, the invention comprises nucleic acid molecules which differ from one of the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID

NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 (and parts thereof) owing to the degeneracy of the genetic code and which thus encode the same Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase as those encoded by the nucleotide sequences shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23.

In addition to the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase shown in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, the skilled worker will recognize that DNA sequence polymorphisms which lead to changes in the amino acid sequences of the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase may exist within a population. These genetic polymorphisms in the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase gene may exist between individuals within a population owing to natural variation. These natural variants usually bring about a variance of 1 to 5% in the nucleotide sequence of the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase gene. Each and every one of these nucleotide variations and resulting amino acid polymorphisms in the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase or Δ -5-desaturase which are the result of natural variation and do not modify the functional activity are to be encompassed by the invention.

The nucleic acid sequence(s) according to the invention (for purposes of the application the singular encompasses the plural and vice versa) or fragments thereof may advantageously be used for isolating other genomic sequences via homology screening.

The said derivatives may be isolated, for example, from other organisms, eukaryotic organisms such as plants, especially mosses, algae, dinoflagellates, protozoa or fungi.

Allele variants include in particular functional variants obtainable by deletion, insertion or substitution of nucleotides in the sequences depicted in SEQ ID NO: 3, SEQ ID

NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 the enzymatic activity of the derived synthesized proteins being retained.

Starting from the DNA sequence described in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 or parts of said sequences such DNA sequences can be isolated using, for example, normal hybridization methods or the PCR technique from other eukaryotes such as those identified above for example. These DNA sequences hybridize under standard conditions with the said sequences. For hybridization use is advantageously made of short oligonucleotides of the conserved regions of an average length of about 15 to 70 bp, preferably of about 17 to 60 bp, more preferably of about 19 to 50 bp, most preferably of about 20 to 40 bp, for example, which can be determined by comparisons with other desaturase or elongase genes in the manner known to those skilled in the art. The histidine box sequences are advantageously employed. However, longer fragments of the nucleic acids according to the invention or the complete sequences may also be used for hybridization. Depending on the nucleic acid employed: oligonucleotide, longer fragment or complete sequence, or depending on which type of nucleic acid, DNA or RNA, is used for hybridization these standard conditions vary. Thus, for example, the melting temperatures of DNA:DNA hybrids are approximately 10°C lower than those of DNA:RNA hybrids of the same length.

By standard conditions is meant, for example, depending on the nucleic acid in question temperatures between 42°C and 58°C in an aqueous buffer solution having a concentration of between 0.1 and 5 x SSC (1 X SSC = 0.15 M NaCl, 15 mM sodium citrate, pH 7.2) or additionally in the presence of 50 % formamide, such as by way of example 42°C in 5 x SSC, 50 % formamide. Hybridization conditions for DNA:DNA hybrids are advantageously 0.1 x SSC and temperatures between approximately 20°C and 45°C, preferably between approximately 30°C and 45°C. For DNA:RNA hybrids the hybridization conditions are advantageously 0.1 x SSC and temperatures between approximately 30°C and 55°C, preferably between approximately 45°C and 55°C. These specified temperatures for hybridization are melting temperature values calculated by way of example for a nucleic acid having a length of approximately 100 nu-

cleotides and a G + C content of 50 % in the absence of formamide. The experimental conditions for DNA hybridization are described in relevant genetics textbooks such as by way of example Sambrook et al., ..Molecular Cloning", Cold Spring Harbor Laboratory, 1989, and may be calculated by formulae known to those skilled in the art, for example as a function of the length of the nucleic acids, the nature of the hybrids or the G + C content. Those skilled in the art may draw on the following textbooks for further information on hybridization: Ausubel et al. (eds), 1985, Current Protocols in Molecular Biology, John Wiley & Sons, New York; Hames and Higgins (eds), 1985, Nucleic Acids Hybridization: A Practical Approach, IRL Press at Oxford University Press, Oxford; Brown (ed), 1991, Essential Molecular Biology: A Practical Approach, IRL Press at Oxford University Press, Oxford.

Furthermore, by derivatives is meant homologues of the sequences SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23, for example eukaryotic homologues, truncated sequences, single-stranded DNA of the encoding and nonencoding DNA sequence or RNA of the encoding and nonencoding DNA sequence.

In addition, by homologues of the sequences SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 is meant derivatives such as by way of example promoter variants. These variants may be modified by one or more nucleotide exchanges, by insertion[^] and/or deletion(s) without, however, adversely affecting the functionality or efficiency of the promoters. Furthermore, the promoters can have their efficiency increased by altering there sequence or be completely replaced by more effective promoters even of foreign organisms.

By derivatives is also advantageously meant variants whose nucleotide sequence has been altered in the region from - 1 to -2000 ahead of the start codon in such a way that the gene expression and/or the protein expression is modified, preferably increased. Furthermore, by derivatives is also meant variants, which have been modified at the 3' end.

The nucleic acid sequences according to the invention which encode a Δ -12-desaturase and Δ -15-desaturase, a Δ -9-elongase, a Δ -Sesaturase and/or a Δ -5-

desaturase may be produced by synthesis or obtained naturally or contain a mixture of synthetic and natural DNA components as well as consist of various heterologous Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - δ -Sesaturase and/or Δ -5-desaturase gene segments from different organisms. In general, synthetic nucleotide sequences are produced with codons, which are preferred by the corresponding host organisms, plants for example. This usually results in optimum expression of the heterologous gene. These codons preferred by plants may be determined from codons having the highest protein frequency, which are expressed in most of the plant species of interest. An example concerning the bacterium *Corynebacterium glutamicum* is provided in Wada et al. (1992) *Nucleic Acids Res.* 20:21 11-21 18). Such experiments can be carried out using standard methods and are known to the person skilled in the art.

Functionally equivalent sequences which encode the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - δ -Sesaturase and/or Δ -5-desaturase gene are those derivatives of the sequence according to the invention which despite differing nucleotide sequence still possess the desired functions, that is to say the enzymatic activity and specific selectivity of the proteins. That means such functionally equivalent sequences have an biological or enzymatic activity, which is at least 10%, preferably at least 20%, 30%, 40% or 50% especially preferably at least 60%, 70%, 80% or 90% and very especially at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% or more of the activity of the proteins/enzymes encoded by the inventive sequences. Thus, functional equivalents include naturally occurring variants of the sequences described herein as well as artificial ones, e.g. artificial nucleotide sequences adapted to the codon use of a plant which have been obtained by chemical synthesis.

In addition, artificial DNA sequences are suitable, provided, as described above, they mediate the desired property, for example an increase in the content of Δ -12 Δ -15 Δ - δ and/or Δ -5-double bonds in fatty acids and an elongation of C18-fatty acids having a Δ -9-double bond in fatty acids, oils or lipids in plants that produce mature seeds preferably in crop plants by over expression of the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - δ -Sesaturase and/or Δ -5-desaturase gene. Such artificial DNA sequences can exhibit Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - δ -Sesaturase and/or Δ -5-desaturase activity, for example by back-translation of pro-

- teins constructed by means of molecular modeling, or be determined by in vitro selection. Possible techniques for in vitro evolution of DNA to modify or improve the DNA sequences are described in Patten, P.A. et al., *Current Opinion in Biotechnology* 8, 724-733(1997) or in Moore, J.C. et al., *Journal of Molecular Biology* 272, 336-347 (1997). Particularly suitable are encoding DNA sequences which are obtained by back-translation of a polypeptide sequence in accordance with the codon use specific to the host plant. Those skilled in the art familiar with the methods of plant genetics can easily determine the specific codon use by computer analyses of other known genes of the plant to be transformed.
- 10 Other suitable equivalent nucleic acid sequences, which may be mentioned are sequences that encode fusion proteins, a component of the fusion protein being a Δ -12-desaturase and Δ -15-desaturase, Δ - Δ -Saturase and/or Δ -5-desaturase polypeptide and/or a Δ -9 elongase polypeptide or a functionally equivalent part thereof. The second part of the fusion protein can be, for example, another polypeptide having enzymatic activity or an antigenic polypeptide sequence by means of which it is possible to demonstrate Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - Δ -Saturase and/or Δ -5-desaturase expression (e.g. myc tag or his tag). Preferably, however, this is a regulatory protein sequence, such as by way of example a signal sequence for the endoplasmic reticulum (= ER) which directs the Δ -12-desaturase and Δ -15-
- 15 desaturase, Δ - Δ -Saturase and/or Δ -5-desaturase protein and/or the Δ -9-elongase protein to the desired point of action, or regulatory sequences which influence the expression of the nucleic acid sequence according to the invention, such as promoters or terminators. In another preferred embodiment the second part of the fusion protein is a plastidial targeting sequence as described by Napier J.A. [Targeting of foreign proteins to the chloroplast, *Methods Mol. Biol.*, 49, 1995: 369 - 376]. A preferred used vector comprising said plastidial targeting sequence is disclosed by Colin Lazarus [Guerineau F., Woolston S., Brooks L., Mullineaux P. "An expression cassette for targeting foreign proteins into chloroplast; *Nucleic. Acids Res.*, Dec 9, 16 (23), 1988: 11380].
- 20
- 25
- 30 Advantageously, the Δ -12-desaturase and Δ -15-desaturase, Δ -9-elongase, Δ - Δ -desaturase and/or Δ -5-desaturase genes in the method according to the invention may be combined with other genes for fatty acid biosynthesis as described above.

Examples of such genes are the acyl transferases, other desaturases or elongases such as Δ -4-desaturases or ω -3- and/or ω -6-specific desaturases) and/or such as Δ -5-elongases to mention only some of them. For in vivo and especially in vitro synthesis combination with e.g. NADH cytochrome B5 reductases, which can take up or release reduction equivalents is advantageous.

By the amino acid sequences according to the invention is meant proteins which contain an amino acid sequence depicted in the sequences SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 or a sequence obtainable there from by substitution, inversion, insertion or deletion of one or more amino acid groups (such sequences are derivatives of SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24), whereas the enzymatic activities of the proteins depicted in SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 12, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 being retained or not substantially reduced, that is they still possess the same enzymatic specificity. By „not substantially reduced" or „the same enzymatic activity" is meant all enzymes which still exhibit at least 10%, 20%, 30%, 40% or 50%, preferably at least 60%, 70%, 80% or 90% particularly preferably at least 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more, of the enzymatic activity of the initial enzyme obtained from the wild type source organism such as organisms of the genus *Physcomitrella*, *Ceratodon*, *Borago*, *Thraustochytrium*, *Schizochytrium*, *Phytophthora*, *Mortierella*, *Caenorhabditis*, *Aleuritia*, *Muscarioides*, *Isochrysis*, *Phaeodactylum*, *Crypthecodinium*, *Acanthamoeba* or *Euglena* preferred source organisms are organisms such as the species *Euglena gracilis*, *Isochrysis galbana*, *Phaeodactylum tricorutum*, *Caenorhabditis elegans*, *Thraustochytrium*, *Phytophthora infestans*, *Ceratodon purpureus*, *Isochrysis galbana*, *Aleuritia farinosa*, *Muscarioides vialii*, *Mortierella alpina*, *Borago officinalis* or *Physcomitrella patens*. For the estimation of an enzymatic activity, which is "not substantially reduced" or which has the "same enzymatic activity" the enzymatic activity of the derived sequences are determined and compared with the wild type enzyme activities. In doing this, for example, certain amino acids may be replaced by others having similar physicochemical properties (space filling, basicity, hydrophobicity, etc.). For example, arginine residues are exchanged for lysine residues, valine residues for isoleucine

residues or aspartic acid residues for glutamic acid residues. However, one or more amino acids may also be swapped in sequence, added or removed, or a plurality of these measures may be combined with one another.

By derivatives is also meant functional equivalents, which in particular also contain
5 natural or artificial mutations of an originally isolated sequence encoding a Δ -12-
desaturase and Δ -15-desaturase, a Δ -9-elongase, a Δ - δ -Saturase and/or a Δ -5-
desaturase, which continue to exhibit the desired function, that is the enzymatic activ-
ity and substrate selectivity thereof is not substantially reduced. Mutations comprise
10 substitutions, additions, deletions, exchanges or insertions of one or more nucleotide
residues. Thus, for example, the present invention also encompasses those nucleotide
sequences, which are obtained by modification of the Δ -12-desaturase and Δ -15-
desaturase nucleotide sequence, the Δ -8-desaturase nucleotide sequence, the Δ -5-
desaturase nucleotide sequence and/or the Δ -9-elongase nucleotide sequence used
in the inventive processes. The aim of such a modification may be, e.g., to further
15 bind the encoding sequence contained therein or also, e.g., to insert further restriction
enzyme interfaces.

Functional equivalents also include those variants whose function by comparison as
described above with the initial gene or gene fragment is weakened (= not substan-
tially reduced) or reinforced (= enzyme activity higher than the activity of the initial
20 enzyme, that is activity is higher than 100%, preferably higher than 110%, 120%,
130%, 140% or 150%, particularly preferably higher than 200% or more).

At the same time the nucleic acid sequence may, for example, advantageously be a
DNA or cDNA sequence. Suitable encoding sequences for insertion into an expres-
sion cassette according to the invention include by way of example those which en-
25 code a Δ -12-desaturase and Δ -15-desaturase, a Δ - δ -Saturase and/or a Δ -5-
desaturase with the sequences described above and lend the host the ability to over-
produce fatty acids, oils or lipids having double bonds in the Δ -12-, Δ -15-, Δ -8-position
and Δ -5-position, it being advantageous when at the same time fatty acids having at
least four double bonds are produced. These sequences may be of homologous or
30 heterologous origin.

By the gene construct (= nucleic acid construct or fragment or expression cassette) according to the invention is meant the sequences specified in SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which result from the genetic code and/or derivatives thereof which are functionally linked with one or more regulation signals advantageously to increase the gene expression and which control the expression of the encoding sequence in the host cell. These regulatory sequences should allow the selective expression of the genes and the protein expression. Depending on the host plant this may mean, for example, that the gene is expressed and/or overexpressed only after induction or that it is expressed and/or overexpressed immediately. Examples of these regulatory sequences are sequences to which inducers or repressors bind and in this way regulate the expression of the nucleic acid. In addition to these new regulation sequences or instead of these sequences the natural regulation of these sequences ahead of the actual structural genes may still be present and optionally have been genetically modified so that natural regulation was switched off and the expression of the genes increased. However, the gene construct can also be built up more simply, that is no additional regulation signals have been inserted ahead of the nucleic acid sequence or derivatives thereof and the natural promoter with its regulation has not been removed. Instead of this the natural regulation sequence was mutated in such a way that no further regulation ensues and/or the gene expression is heightened. These modified promoters in the form of part sequences (= promoter containing parts of the nucleic acid sequences according to the invention) can also be brought on their own ahead of the natural gene to increase the activity. In addition, the gene construct may advantageously also contain one or more so-called enhancer sequences functionally linked to the promoter which allow enhanced expression of the nucleic acid sequence. At the 3' end of the DNA sequences additional advantageous sequences may also be inserted, such as further regulatory elements or terminators. The SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and/or SEQ ID NO: 23 gene may be present in one or more copies in the gene construct (=expression cassette).

As described above, the regulatory sequences or factors can preferably positively influence and so increase the gene expression of the introduced genes. Thus, reinforcement of the regulatory elements advantageously on the transcription level may

be effected by using powerful transcription signals such as promoters and/or enhancers. However, in addition reinforcement of translation is also possible, for example by improving the stability of the mRNA.

Suitable promoters in the expression cassette are in principle all promoters which can control the expression of foreign genes in microorganisms like protozoa such as amoeba, ciliates, algae such as green, brown, red or blue algae such as Euglena, bacteria such as gram-positive or gram-negative bacteria, yeasts such as Saccharomyces, Pichia or Schizosaccharomyces or fungi such as Mortierella, Thraustochytrium or Schizochytrium or plants such as Aleuritia, advantageously in plants or fungi. Such microorganisms are generally used to clone the inventive genes and possible other genes of the fatty acid biosynthesis chain for the production of fatty acids according to the inventive process. Use is preferably made in particular of plant promoters or promoters derived from a plant virus. Advantageous regulation sequences for the method according to the invention are found for example in promoters such as cos, tac, trp, tet, trp-tet, lpp, lac, lpp-lac, $lacI^q$ T7, T5, T3, gal, trc, ara, SP6, λ -P_R or in λ -P_L promoters which are employed advantageously in gram-negative bacteria. Other advantageous regulation sequences are found, for example, in the gram-positive promoters amy and SPO2, in the yeast or fungal promoters ADC1, MFa, AC, P-60, CYC1, GAPDH, TEF, rp28, ADH or in the plant promoters CaMV/35S [Franck et al., Cell 21(1980) 285-294], SSU, OCS, lib4, STLS1, B33, nos (= Nopaline Synthase Promoter) or in the ubiquitin or phaseolin promoter. The expression cassette may also contain a chemically inducible promoter by means of which the expression of the exogenous Δ -12- and Δ -15-, Δ -8- and/or Δ -5-desaturase gene and/or the Δ -9-elongase gene in the microorganism and/or plant can be controlled advantageously in the plants at a particular time. Advantageous plant promoters of this type are by way of example the PRP1 promoter [Ward et al., Plant.Mol. Biol. 22 (1993), 361-366], a promoter inducible by benzenesulfonamide (EP 388 186), a promoter inducible by tetracycline [Gatz et al., (1992) Plant J. 2, 397 - 404], a promoter inducible by salicylic acid (WO 95/19443), a promoter inducible by abscisic acid (EP 335 528) and a promoter inducible by ethanol or cyclohexanone (WO 93/21334). Other examples of plant promoters, which can advantageously be used are the promoter of cytosolic FBPase from potato, the ST-LSI promoter from potato (Stockhaus et al., EMBO J. 8 (1989) 2445-245), the promoter of

phosphoribosyl pyrophosphate amidotransferase from *Glycine max* (see also gene bank accession number U87999) or a nodiene-specific promoter as described in EP 249 676. Particularly advantageous are those plant promoters, which ensure expression in tissues or plant parts/organs in which fatty acid biosynthesis or the precursor stages thereof occurs, as in endosperm or in the developing embryo for example. Particularly noteworthy are advantageous promoters, which ensure seed-specific expression such as by way of example the USP promoter or derivatives thereof, the LEB4 promoter, the phaseolin promoter or the napin promoter. The particularly advantageous USP promoter cited according to the invention or its derivatives mediate very early gene expression in seed development [Baeumlein et al., *Mol Gen Genet*, 1991, 225 (3): 459-67]. Other advantageous seed-specific promoters which may be used for monocotyledonous or dicotyledonous plants are the promoters suitable for dicotyledons such as napin gene promoters, likewise cited by way of example, from oilseed rape (US 5,608,152), the oleosin promoter from *Arabidopsis* (WO 98/45461), the phaseolin promoter from *Phaseolus vulgaris* (US 5,504,200), the Bce4 promoter from *Brassica* (WO 91/13980) or the leguminous B4 promoter (LeB4, Baeumlein et al., *Plant J.*, 2, 2, 1992: 233 - 239) or promoters suitable for monocotyledons such as the promoters of the *lpt2* or *lpt1* gene in barley (WO 95/15389 and WO 95/23230) or the promoters of the barley hordeine gene, the rice glutelin gene, the rice oryzin gene, the rice prolamin gene, the wheat gliadin gene, the white glutelin gene, the corn zein gene, the oats glutelin gene, the sorghum kasirin gene or the rye secalin gene which are described in WO99/16890.

Furthermore, particularly preferred are those promoters, which ensure the expression in tissues or plant parts in which, for example, the biosynthesis of fatty acids, oils and lipids or the precursor stages thereof takes place. Particularly noteworthy are promoters, which ensure a seed-specific expression. Noteworthy are the promoter of the napin gene from oilseed rape (US 5,608,152), the USP promoter from *Vicia faba* (USP = unknown seed protein, Baeumlein et al., *Mol Gen Genet*, 1991, 225 (3): 459 - 67), the promoter of the oleosin gene from *Arabidopsis* (WO 98/45461), the phaseolin promoter (US 5,504,200) or the promoter of the legumin B4 gene (LeB4; Baeumlein et al., 1992, *Plant Journal*, 2 (2): 233-9). Other promoters to be mentioned are that of the *lpt2* or *lpt1* gene from barley (WO 95/15389 and WO 95/23230), which mediate seed-specific expression in monocotyledonous plants. Other advantageous seed specific

promoters are promoters such as the promoters from rice, corn or wheat disclosed in WO 99/16890 or Amy32b, Amy6-6 or aleurain (US 5,677,474), Bce4 (rape, US 5,530,149), glycinin (soy bean, EP 571 741), phosphoenol pyruvat carboxylase (soy bean, JP 06/62870), ADR12-2 (soy bean, WO 98/08962), isocitratlyase (rape, US 5,689,040) or β -amylase (barley, EP 781 849).

As described above, the expression construct (= gene construct, nucleic acid construct) may contain yet other genes, which are to be introduced into the microorganism or plant. These genes can be subject to separate regulation or be subject to the same regulation region as the Δ -12- and Δ -15-desaturase gene and/or the Δ -8- and/or Δ -5-desaturase gene and/or the Δ -9-elongase gene. These genes are by way of example other biosynthesis genes, advantageously for fatty acid biosynthesis, which allow increased synthesis. Examples which may be mentioned are the genes for example of the Δ -9-, Δ -4-desaturase, Δ -5-elongase, α -ketoacyl reductases, α -ketoacyl synthases, elongases or the various hydroxylases and acyl-ACP thioesterases. The desaturase and elongase genes are advantageously used in the nucleic acid construct.

In principle all natural promoters with their regulation sequences can be used like those named above for the expression cassette according to the invention and the method according to the invention. Over and above this, synthetic promoters may also advantageously be used.

In the preparation of an a gene construct various DNA fragments can be manipulated in order to obtain a nucleotide sequence, which usefully reads in the correct direction and is equipped with a correct reading raster. To connect the DNA fragments (= nucleic acids according to the invention) to one another adaptors or linkers may be attached to the fragments.

The promoter and the terminator regions can usefully be provided in the transcription direction with a linker or polylinker containing one or more restriction points for the insertion of this sequence. Generally, the linker has 1 to 10, mostly 1 to 8, preferably 2 to 6, restriction points. In general the size of the linker inside the regulatory region is less than 100 bp, frequently less than 60 bp, but at least 5 bp. The promoter may be native or homologous as well as foreign or heterologous to the host organism, for ex-

ample to the host plant. In the 5'-3' transcription direction the expression cassette contains the promoter, a DNA sequence which encodes a Δ -12- and Δ -15-desaturase gene, a Δ -8-desaturase gene, a Δ -5-desaturase gene and/or a Δ -9-elongase gene and a region for transcription termination. Different termination regions can be exchanged for one another in any desired fashion.

Furthermore, manipulations, which provide suitable restriction interfaces or which remove excess DNA or restriction interfaces can be employed. Where insertions, deletions or substitutions, such as transitions and transversions, come into consideration, *in vitro* mutagenesis, primer repair, restriction or ligation may be used. In suitable manipulations such as restriction, chewing back or filling of overhangs for blunt ends complementary ends of the fragments can be provided for the ligation.

For an advantageous high expression the attachment of the specific ER retention signal SEKDEL inter alia can be of importance (Schouten, A. et al., Plant Mol. Biol. 30 (1996), 781-792). In this way the average expression level is tripled or even quadrupled. Other retention signals, which occur naturally in plant and animal proteins located in the ER may also be employed for the construction of the cassette. In another preferred embodiment a plastidial targeting sequence is used as described by Napier J.A. [Targeting of foreign proteins to the chloroplast, Methods Mol. Biol., 49, 1995: 369 - 376]. A preferred used vector comprising said plastidial targeting sequence is disclosed by Colin Lazarus [Guerineau F., Woolston S., Brooks L., Mullineaux P. "An expression cassette for targeting foreign proteins into chloroplast; Nucleic. Acids Res., Dec 9, 16 (23), 1988: 11380].

Preferred polyadenylation signals are plant polyadenylation signals, preferably those which substantially correspond to T-DNA polyadenylation signals from *Agrobacterium tumefaciens*, in particular gene 3 of the T-DNA (octopin synthase) of the Ti plasmid pTiACH5 (Gielen et al., EMBO J.3 (1984), 835 et seq.) or corresponding functional equivalents.

An expression cassette/gene construct is produced by fusion of a suitable promoter with a suitable Δ -12- and Δ -15-desaturase DNA sequence, a suitable Δ -8- and/or Δ -5-desaturase DNA sequence and/or a suitable Δ -9-elongase DNA sequence together with a polyadenylation signal by common recombination and cloning techniques as

described, for example, in T. Maniatis, E.F. Fritsch and J. Sambrook, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989) as well as in T.J. Silhavy, M.L. Berman and L.W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and in
5 Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience (1987).

The DNA sequences encoding the nucleic acid sequences used in the inventive processes such as the Δ -12- and Δ -15-desaturase from *Acanthamoeba castellanii* or *Perkinsus marinus*, Δ -8-desaturase from *Euglena gracilis*, *Acanthamoeba castellanii*
10 or *Perkinsus marinus*, the Δ -9-elongase from *Isochrysis galbana* or *Acanthamoeba castellanii* and/or the Δ -5-desaturase for example from *Thraustochytrium*, *Acanthamoeba castellanii* or *Perkinsus marinus* or other organisms such as *Caenorhabditis elegans*, *Mortierella alpina*, *Borage officinalis* or *Physcomitrella patens* contain all the sequence characteristics needed to achieve correct localization of the site of fatty
15 acid, lipid or oil biosynthesis. Accordingly, no further targeting sequences are needed per se. However, such localization may be desirable and advantageous and hence artificially modified or reinforced so that such fusion constructs are also a preferred advantageous embodiment of the invention.

Particularly preferred are sequences, which ensure targeting in plastids. Under certain
20 circumstances targeting into other compartments (reported in: Kermode, *Crit. Rev. Plant Sci.* 15, 4 (1996), 285-423) may also be desirable, e.g. into vacuoles, the mitochondrion, the endoplasmic reticulum (ER), peroxisomes, lipid structures or due to lack of corresponding operative sequences retention in the compartment of origin, the cytosol.

25 Advantageously, the nucleic acid sequences according to the invention or the gene construct together with at least one reporter gene are cloned into a gene construct, which is introduced into the organism via a vector or directly into the genome. This reporter gene should allow easy detection via a growth, fluorescence, chemical, bioluminescence or resistance assay or via a photometric measurement. Examples of
30 reporter genes which may be mentioned are antibiotic- or herbicide-resistance genes, hydrolase genes, fluorescence protein genes, bioluminescence genes, sugar or nucleotide metabolic genes or biosynthesis genes such as the *Ura3* gene, the *Ilv2* gene,

the luciferase gene, the β -galactosidase gene, the gfp gene, the 2-deoxyglucose-6-phosphate phosphatase gene, the β -glucuronidase gene, β -lactamase gene, the neomycin phosphotransferase gene, the hygromycin phosphotransferase gene or the BASTA (= glufosinate-resistance) gene. These genes permit easy measurement and quantification of the transcription activity and hence of the expression of the genes. In this way genome positions may be identified which exhibit differing productivity.

In a preferred embodiment an gene construct comprises upstream, i.e. at the 5' end of the encoding sequence, a promoter and downstream, i.e. at the 3' end, a polyadenylation signal and optionally other regulatory elements which are operably linked to the intervening encoding sequence for Δ -12- and Δ -15-desaturase, Δ -8-desaturase, Δ -9-elongase and/or Δ -5-desaturase DNA sequence. By an operable linkage is meant the sequential arrangement of promoter, encoding sequence, terminator and optionally other regulatory elements in such a way that each of the regulatory elements can fulfill its function in the expression of the encoding sequence in due manner. The sequences preferred for operable linkage are targeting sequences for ensuring subcellular localization in plastids. However, targeting sequences for ensuring subcellular localization in the mitochondrion, in the endoplasmic reticulum (= ER), in the nucleus, in oil corpuscles or other compartments may also be employed as well as translation promoters such as the 5' lead sequence in tobacco mosaic virus (GalMe et al., Nucl. Acids Res. 15 (1987), 8693 -871 1).

An expression cassette/gene construct may, for example, contain a constitutive promoter or a tissue-specific promoter (preferably the USP or napin promoter) the gene to be expressed and the ER retention signal. For the ER retention signal the KDEL amino acid sequence (lysine, aspartic acid, glutamic acid, leucine) or the KKX amino acid sequence (lysine-lysine-X-stop, wherein X means every other known amino acid) is preferably employed.

For expression in a prokaryotic or eukaryotic host organism, for example a microorganism such as a fungus or a plant such as an oil crop the expression cassette is advantageously inserted into a vector such as by way of example a plasmid, a phage or other DNA which allows optimum expression of the genes in the host organism. Examples of suitable plasmids are: in E. coli pLG338, pACYC184, pBR series such as

e.g. pBR322, pUC series such as pUC18 or pUC19, M113mp series, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-IN¹¹³-B1, λ gt1 1 or pBdCl; in Streptomyces pIJ101, pIJ364, pIJ702 or pIJ361; in Bacillus pUB1 10, pC194 or pBD214; in Corynebacterium pSA77 or pAJ667; in fungi pALS1, pIL2 or pBB1 16;

5 other advantageous fungal vectors are described by Romanos, M.A. et al., [(1992) ..Foreign gene expression in yeast: a review", *Yeast*8: 423-488] and by van den Hondel, C.A.M.J.J. et al. [(1991) ..Heterologous gene expression in filamentous fungi" as well as in More Gene Manipulations in Fungi [J.W. Bennet & L.L. Lasure, eds., pp. 396-428: Academic Press: San Diego] and in „Gene transfer systems and vector

10 development for filamentous fungi" [van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) in: Applied Molecular Genetics of Fungi, Peberdy, J.F. et al., eds., pp. 1-28, Cambridge University Press: Cambridge]. Examples of advantageous yeast promoters are 2 μ M, pAG-1, YEp6, YEpl 3 or pEMBLye23. Examples of algal or plant promoters are pLGV23, pGHlac⁺, pBIN19, pAK2004, pVKH or pDH51 (see Schmidt, R. and

15 Willmitzer, L., 1988). The vectors identified above or derivatives of the vectors identified above are a small selection of the possible plasmids. Further plasmids are well known to those skilled in the art and may be found, for example, in the book Cloning Vectors (Eds. Pouwels P.H. et al. Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). Suitable plant vectors are described inter alia in ..Methods in

20 Plant Molecular Biology and Biotechnology" (CRC Press), Ch. 6/7, pp. 71-1 19. Advantageous vectors are known as shuttle vectors or binary vectors which replicate in *E. coli* and *Agrobacterium*.

By vectors is meant with the exception of plasmids all other vectors known to those skilled in the art such as by way of example phages, viruses such as SV40, CMV,

25 baculovirus, adenovirus, transposons, IS elements, phasmids, phagemids, cosmids, linear or circular DNA. These vectors can be replicated autonomously in the host organism or be chromosomally replicated, chromosomal replication being preferred.

In a further embodiment of the vector the gene construct according to the invention may also advantageously be introduced into the organisms in the form of a linear

30 DNA and be integrated into the genome of the host organism by way of heterologous or homologous recombination. This linear DNA may be composed of a linearized

plasmid or only of the expression cassette as vector or the nucleic acid sequences according to the invention.

In a further advantageous embodiment the nucleic acid sequence according to the invention can also be introduced into an organism on its own.

5 If in addition to the nucleic acid sequence according to the invention further genes are to be introduced into the organism, all together with a reporter gene in a single vector or each single gene with a reporter gene in a vector in each case can be introduced into the organism, whereby the different vectors can be introduced simultaneously or successively.

10 The vector advantageously contains at least one copy of the nucleic acid sequences according to the invention and/or the expression cassette (= gene construct) according to the invention.

By way of example the plant expression cassette can be installed in the pRT transformation vector ((a) Toepfer et al., 1993, *Methods Enzymol.*, 217: 66-78; (b) Toepfer
15 et al. 1987, *Nucl. Acids. Res.* 15: 5890 ff.).

Alternatively, a recombinant vector (= expression vector) can also be transcribed and translated in vitro, e.g. by using the T7 promoter and the T7 RNA polymerase.

Expression vectors employed in prokaryotes frequently make use of inducible systems with and without fusion proteins or fusion oligopeptides, wherein these fusions
20 can ensue in both N-terminal and C-terminal manner or in other useful domains of a protein. Such fusion vectors usually have the following purposes: i.) to increase the RNA expression rate; ii.) to increase the achievable protein synthesis rate; iii.) to increase the solubility of the protein; iv.) or to simplify purification by means of a binding sequence usable for affinity chromatography. Proteolytic cleavage points are also
25 frequently introduced via fusion proteins, which allow cleavage of a portion of the fusion protein and purification. Such recognition sequences for proteases are recognized, e.g. factor Xa, thrombin and enterokinase.

Typical advantageous fusion and expression vectors are pGEX [Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67: 31-40], pMAL (New England Bio-

labs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which contains glutathione S-transferase (GST), maltose binding protein or protein A.

Other examples of E. coli expression vectors are pTrc [Amann et al., (1988) *Gene* 69:301-315] and pET vectors [Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89; Stratagene, Amsterdam, The Netherlands].

Other advantageous vectors for use in yeast are pYepSed (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Ce//* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:1 13-123), and pYES derivatives (Invitrogen Corporation, San Diego, CA). Vectors for use in filamentous fungi are described in: van den Hondel, C.A.M.J.J. & Punt, P.J. (1991) „Gene transfer systems and vector development for filamentous fungi“, in: *Applied Molecular Genetics of Fungi*, J.F. Peberdy, et al., eds., pp. 1-28, Cambridge University Press: Cambridge.

Alternatively, insect cell expression vectors can also be advantageously utilized, e.g. for expression in Sf 9 cells. These are e.g. the vectors of the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

Furthermore, plant cells or algal cells can advantageously be used for gene expression. Examples of plant expression vectors may be found in Becker, D., et al. (1992) „New plant binary vectors with selectable markers located proximal to the left border“, *Plant Mol. Biol.* 20: 1195-1 197 or in Bevan, M.W. (1984) „Binary *Agrobacterium* vectors for plant transformation“, *Nucl. Acid. Res.* 12: 871 1-8721 .

The host plant (= transgenic plant) advantageously contains at least one copy of the nucleic acid according to the invention and/or of the gene construct according to the invention.

The introduction of the nucleic acids according to the invention, the gene construct or the vector into organisms, plants for example, can in principle be done by all of the methods known to those skilled in the art. The introduction of the nucleic acid sequences gives rise to recombinant or transgenic plants.

To introduce the nucleic acids used in the process, the latter are advantageously amplified and ligated in the known manner. Preferably, a procedure following the protocol for Pfu DNA polymerase or a Pfu/Taq DNA polymerase mixture is followed. The primers are selected taking into consideration the sequence to be amplified. The primers should advantageously be chosen in such a way that the amplificate comprises the entire codogenic sequence from the start codon to the stop codon. After the amplification, the amplificate is expediently analyzed. For example, a gel-electrophoretic separation can be carried out, which is followed by a quantitative and a qualitative analysis. Thereafter, the amplificate can be purified following a standard protocol (for example Qiagen). An aliquot of the purified amplificate is then available for the subsequent cloning step. Suitable cloning vectors are mentioned above and generally known to the skilled worker. These include, in particular, vectors which are capable of replication in microbial systems, that is to say mainly vectors which ensure efficient cloning in yeasts or fungi and which make possible the stable transformation of plants. Those, which must be mentioned, again herein in particular are various binary and cointegrated vector systems, which are suitable for the T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they comprise at least the vir genes required for the Agrobacterium-mediated transformation and the T-DNA-delimiting sequences (T-DNA border). These vector systems advantageously also comprise further cis-regulatory regions such as promoters and terminator sequences and/or selection markers, by means of which suitably transformed organisms can be identified. While in the case of cointegrated vector systems vir genes and T-DNA sequences are arranged on the same vector, binary systems are based on at least two vectors, one of which bears vir genes, but no T-DNA, while a second one bears T-DNA, but no vir gene. Owing to this fact, the last-mentioned vectors are relatively small, easy to manipulate and to replicate both in *E. coli* and in Agrobacterium. These binary vectors include vectors from the series pBIB-HYG, pPZP, pBecks, pGreen. In accordance with the invention, Bin19, pBI101, pBinAR, pGPTV and pCAMBIA are used by preference. An overview of the binary vectors and their use is found in Hellens et al, Trends in Plant Science (2000) 5, 446-451. In order to prepare the vectors, the vectors can first be linearized with restriction endonuclease(s) and then modified enzymatically in a suitable manner. Thereafter, the vector is purified, and an aliquot is employed for the cloning step. In the cloning step, the enzymatically cleaved and, if appropriate, purified amplificate is cloned with

- vector fragments, which have been prepared in a similar manner, using ligase. In this context, a particular nucleic acid construct, or vector or plasmid construct, can have one or else more than one codogenic gene segment. The codogenic gene segments in these constructs are preferably linked operably with regulatory sequences. The regulatory sequences include, in particular, plant sequences such as the above-described promoters and terminator sequences. The constructs can advantageously be stably propagated in microorganisms, in particular in *E. coli* and *Agrobacterium tumefaciens*, under selective conditions and make possible the transfer of heterologous DNA into plants or microorganisms.
- 10 The nucleic acids used in the process, the inventive nucleic acids and gene constructs, can be introduced into organisms such as microorganisms or advantageously plants, advantageously using cloning vectors, and thus be used in the transformation of plants such as those which are published and cited in: *Plant Molecular Biology and Biotechnology* (CRC Press, Boca Raton, Florida), Chapter 6/7, p. 71-1 19 (1993); F.F. White, *Vectors for Gene Transfer in Higher Plants*; in: *Transgenic Plants, Vol. 1, Engineering and Utilization*, Ed.: Kung and R. Wu, Academic Press, 1993, 15-38; B. Jenes et al., *Techniques for Gene Transfer*, in: *Transgenic Plants, Vol. 1, Engineering and Utilization*, Ed.: Kung and R. Wu, Academic Press (1993), 128-143; Potrykus, *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42 (1991), 205-225. Thus, the nucleic acids, the inventive nucleic acids and nucleic acid constructs, and/or vectors used in the process can be used for the recombinant modification of a broad spectrum of organisms, advantageously plants, so that the latter become better and/or more efficient PUFA and/or LCPUFA producers.
- 25 In the case of microorganisms, those skilled in the art can find appropriate methods for the introduction of the inventive nucleic acid sequences, the gene construct or the vector in the textbooks by Sambrook, J. et al. (1989) *Molecular cloning: A laboratory manual*, Cold Spring Harbor Laboratory Press, by F.M. Ausubel et al. (1994) *Current protocols in molecular biology*, John Wiley and Sons, by D.M. Glover et al., *DNA Cloning Vol. 1*, (1995), IRL Press (ISBN 019-963476-9), by Kaiser et al. (1994) *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory Press or Guthrie et al. *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, 1994, Academic Press.
- 30

The transfer of foreign genes into the genome of a plant is called transformation. In doing this the methods described for the transformation and regeneration of plants from plant tissues or plant cells are utilized for transient or stable transformation. Suitable methods are protoplast transformation by poly(ethylene glycol)-induced DNA uptake, the „biolistic" method using the gene cannon - referred to as the particle bombardment method, electroporation, the incubation of dry embryos in DNA solution, microinjection and gene transfer mediated by *Agrobacterium*. Said methods are described by way of example in B. Jenes et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press (1993) 128-143 and in Potrykus Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991) 205-225). The nucleic acids or the construct to be expressed is preferably cloned into a vector, which is suitable for transforming *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al., Nucl. Acids Res. 12 (1984) 871-1). *Agrobacteria* transformed by such a vector can then be used in known manner for the transformation of plants, in particular of crop plants such as by way of example tobacco plants, for example by bathing bruised leaves or chopped leaves in an agrobacterial solution and then culturing them in suitable media. The transformation of plants by means of *Agrobacterium tumefaciens* is described, for example, by Hofgen and Willmitzer in Nucl. Acid Res. (1988) 16, 9877 or is known inter alia from F.F. White, Vectors for Gene Transfer in Higher Plants; in Transgenic Plants, Vol. 1, Engineering and Utilization, eds. S.D. Kung and R. Wu, Academic Press, 1993, pp. 15-38.

Agrobacteria transformed by an expression vector according to the invention may likewise be used in known manner for the transformation of plants such as test plants like *Arabidopsis* or crop plants such as cereal crops, corn, oats, rye, barley, wheat, soybean, rice, cotton, sugar beet, canola, sunflower, flax, hemp, potatoes, tobacco, tomatoes, carrots, paprika, oilseed rape, tapioca, cassava, arrowroot, tagetes, alfalfa, lettuce and the various tree, nut and vine species, in particular of oil-containing crop plants such as soybean, peanut, castor oil plant, sunflower, corn, cotton, flax (linseed), oilseed rape, poppy, mustard, sesame, almond, macadamia, olive, calendula, punica, hazel nut, avocado, pumpkin, walnut, laurel, pistachio, *Orychopragmus*, marigold, borage, primrose, canola, evening primrose, hemp, coconut, oil palm, safflower (*Carthamus tinctorius*), coffee or cocoa bean, e.g. by bathing bruised leaves or chopped leaves in an agrobacterial solution and then culturing them in suitable media.

For the production of LCPUFAs, for example arachidonic acid and/or eicosapentaenoic acid, borage, linseed, sunflower, safflower, Brassica napus, Brassica juncea, Camelina sativa or Orychophragmus are advantageously suitable.

5 The genetically modified plant cells may be regenerated by all of the methods known to those skilled in the art. Appropriate methods can be found in the publications referred to above by S.D. Kung and R. Wu, Potrykus or Hofgen and Willmitzer.

Accordingly, a further aspect of the invention relates to transgenic organisms transformed by at least one nucleic acid sequence, expression cassette or vector according to the invention as well as cells, cell cultures, tissue, parts - such as, for example, 10 leaves, roots, etc. in the case of plant organisms - or reproductive material derived from such organisms. The terms „host organism“, „host cell“, „recombinant (host) organism“ and „transgenic (host) cell“ are used here interchangeably. Of course these terms relate not only to the particular host organism or the particular target cell but also to the descendants or potential descendants of these organisms or cells. Since, 15 due to mutation or environmental effects certain modifications may arise in successive generations, these descendants need not necessarily be identical with the parental cell but nevertheless are still encompassed by the term as used here.

Suitable organisms or host organisms for the nucleic acid, gene construct or vector according to the invention are advantageously in principle all plants, which are able to 20 synthesize fatty acids, especially unsaturated fatty acids or are suitable for the expression of recombinant genes as described above. Further examples which may be mentioned are plants such as Arabidopsis, Asteraceae such as Calendula or crop plants such as soybean, peanut, castor oil plant, sunflower, corn, cotton, flax, oilseed rape, coconut, oil palm, safflower (Carthamus tinctorius) or cocoa bean, bacteria such 25 as the genus Escherichia, yeasts such as the genus Saccharomyces. Preference is given to organisms which can naturally synthesize oils in relatively large quantities such as fungi like Mortierella alpina, Pythium insidiosum or plants such as soybean, oilseed rape, coconut, oil palm, safflower, flax, castor oil plant, Calendula, peanut, cocoa bean or sunflower, or yeasts such as Saccharomyces cerevisiae and particular 30 preference is given to the family of the Brassicaceae such as oilseed rape, soybean, flax, sunflower, Calendula, Mortierella or Saccharomyces cerevisiae.

Further useful host cells are identified in: Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990).

Usable expression strains, e.g. those exhibiting a relatively low protease activity, are described in: Gottesman, S., *Gene Expression Technology: Methods in Enzymology*
5 185, Academic Press, San Diego, California (1990) 119-128.

A further object of the invention as described relates to the use of an expression cassette containing DNA sequences encoding a Δ -12- and Δ -15-desaturase, a Δ -9-elongase, a Δ -8-desaturase and/or a Δ -5-desaturase gene or DNA sequences hybridizing therewith for the transformation of plant cells, tissues or parts of plants. The aim
10 of use is to increase the content of fatty acids, oils or lipids having an increased content of double bonds.

In doing so, depending on the choice of promoter, the Δ -12- and Δ -15-desaturase, the Δ -9-elongase, the Δ -8-desaturase and/or the Δ -5-desaturase gene can be expressed specifically in the leaves, in the seeds, the nodules, in roots, in the stem or other parts
15 of the plant, preferably in leaves and/or seeds. Those transgenic plants overproducing fatty acids, oils or lipids according to the invention, the reproductive material thereof, together with the plant cells, tissues or parts thereof are a further object of the present invention.

The expression cassette or the nucleic acid sequences according to the invention
20 containing a Δ -12- and Δ -15-desaturase, a Δ -9-elongase, a Δ -8-desaturase and/or a Δ -5-desaturase gene sequence can, moreover, also be employed for the transformation of the organisms identified by way of example above such as bacteria, cyanobacteria, yeasts, filamentous fungi, ciliates and algae with the objective of increasing the content of fatty acids, oils or lipids according to the invention.

25 Within the framework of the present invention is the increase of the content of fatty acids, oils or lipids possessing a higher amount of ω -3-fatty acids in comparison to ω -6-fatty acids such as eicosapentaenoic acid in comparison to arachidonic acid, due to functional over expression of the Δ -12- and Δ -15-desaturase, the Δ -9-elongase, the Δ -8-desaturase and/or the Δ -5-desaturase gene in the plant according to the inven-
30 tion, advantageously in the transgenic oilseed plants according to the invention, by

comparison with the non genetically modified initial plants at least for the duration of at least one plant generation.

The preferred locus of biosynthesis, of fatty acids, oils or lipids for example, is generally the seed or cell layers of the seed so that a seed-specific expression of the Δ -12- and Δ -15-desaturase, the Δ -9-elongase, the Δ -8-desaturase and/or the Δ -5-
5 desaturase gene is appropriate. It is, however, obvious that the biosynthesis of fatty acids, oils or lipids need not be limited to the seed tissue but rather can also occur in tissue-specific manner in all other parts of the plant - in epidermis cells or in the nodules for example.

10 A constitutive expression of the exogenous Δ -12- and Δ -15-desaturase, Δ -9-elongase, Δ -8-desaturase and/or Δ -5-desaturase gene is, moreover, advantageous. On the other hand, however, an inducible expression may also appear desirable.

The efficiency of the expression of the Δ -12- and Δ -15-desaturase, the Δ -9-elongase, the Δ -8-desaturase and/or the Δ -5-desaturase gene can be determined, for example,
15 *in vitro* by shoot meristem propagation. In addition, an expression of the Δ -12- and Δ -15-desaturase, the Δ -9-elongase, the Δ -8-desaturase and/or the Δ -5-desaturase gene modified in nature and level and its effect on fatty acid, oil or lipid biosynthesis performance can be tested on test plants in greenhouse trials.

An additional object of the invention comprises transgenic plants transformed by an
20 expression cassette containing a Δ -12- and Δ -15-desaturase, a Δ -9-elongase, a Δ -8-desaturase and/or a Δ -5-desaturase gene sequence according to the invention or DNA sequences hybridizing therewith, as well as transgenic cells, tissue, parts and reproduction material of such plants. Particular preference is given in this case to transgenic crop plants such as by way of example barley, wheat, rye, oats, corn, soy-
25 bean, rice, cotton, sugar beet, the family of the Brassicaceae such as oilseed rape and canola, sunflower, flax, hemp, thistle, potatoes, tobacco, tomatoes, tapioca, cassava, arrowroot, alfalfa, lettuce and the various tree, nut and vine species.

For the purposes of the invention plants are mono- and dicotyledonous plants that produce mature seeds.

A further refinement according to the invention are transgenic plants as described above which contain the nucleic acid sequences, the gene construct and/or vector of the invention.

The invention is explained in more detail by the following examples.

5 Examples

Example 1: General cloning methods

The cloning methods, such as by way of example restriction cleavages, agarose gel electrophoresis, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linkage of DNA fragments, transformation of *Escherichia coli* cells, culture of bacteria and sequence analysis of recombinant DNA, were carried
10 out as described in Sambrook et al. (1989) (Cold Spring Harbor Laboratory Press: ISBN 0-87969-309-6).

Example 2: Sequence analysis of recombinant DNA

Sequencing of recombinant DNA molecules was done using a laser fluorescence
15 DNA sequencer from the ABI company by the method of Sanger (Sanger et al. (1977) Proc. Natl. Acad. Sci. USA74, 5463-5467). Fragments resulting from a polymerase chain reaction were sequenced and checked to prevent polymerase errors in the constructs to be expressed.

Example 3: Cloning of the PUFA specific desaturases from *Acanthamoeba castellanii* (= SEQ ID NO: 3, 5, 15, 19 and 21)
20

Acanthamoeba castellanii (Eukaryota; Protista; Sarcomastigophora; Sarcodina; Rhizopodea; Lobosa) is an amoeba species, which is a common species in the soil. *Acanthamoeba castellanii* can grow vegetative over a broad temperature range (10 to 32°C). *A. castellanii* is able to de novo synthesize linoleic acid and C20 n-6 fatty acids.
25 *A. castellanii* (ATTC 30010) was grown at 30°C on a medium containing 0,75% (w/v) peptone, 1,5% (w/v) glucose and 0,75% (w/v) yeast extract according to the reference of Jones et al. [Temperature-induced membrane-lipid adaptation in *Acanthamoeba castellanii*. Biochem J. 1993, 290:273-278]. The cell cultures were grown under shak-

ing (200 U/min) and harvested with a centrifuge at 250 x g, 5 min, 4°C, after they have reached a cell density of $5 \times 10^6 - 10^7$ (measured in a Fuchs-Rosenthal Haemozytometer).

The total mRNA was isolated from said harvested cells with the aid of the RNeasy plant mini Kit (Qiagen). cDNA was synthesized from the total mRNA with the SMART RACE cDNA amplification kit (Clontech) according to the instructions of the manufacturer.

For the isolation of new desaturase genes the following degenerated primers were used for the amplification:

10 Deg1 :

5'- GGITGG(CZTZA)TIGGICA(TZC) GA(TZC)(GT) (CT)I(GT) (GC)ICA-3'

Deg2:

5'- GG(AZG)AA(TCGA)AG(AZG)TG(AZG)TG(TZC)TC(AZGZT)AT(TZC)TG-S'

The aforementioned primers were used for the amplification in combination with the 3'-adapter-primer of the SMART RACE cDNA amplification kit.

The following protocol was used for the amplification:

- 20 a) 2 min at 95°C,
 b) 30 sec at 94 °C
 30 sec at 55-72°C
 2 min at 72 °C
 Number of cycles: 30
 c) 10 min at 72 °C

25 PCR amplicons were cloned and sequenced according to the instructions of the manufacturer (pTOPO, Invitrogen). The sequence information was used for the production of full-length clones. For the cloning of the full-length clones 5'- and 3'-specific primers were synthesized. Said primers were used for the amplification in the SMART RACE cDNA amplification kit (Clontech) and the amplicons were cloned into the pTOPO vector (Invitrogen)

Three sequences were identified, which show low similarities to desaturase genes.

In addition according to [Zank et al. 2002, Plant Journal 31:255-268] sequence 9Ac (Δ -9-Elongase from *Acanthamoeba*, SEQ ID NO: 11) could be identified, which shows low similarities to elongase genes.

Table 1: *Acanthamoeba castellanii* desaturase sequences

Gene	Nucleotide bp	SEQ ID NO:
12Ac (Δ -12/ Δ 15-Desaturase from <i>Acanthamoeba</i>)	1224 bp	19, 21
8Ac (Δ -8-Desaturase from <i>Acanthamoeba</i>)	1374 bp	3, 5
5Ac (Δ -5-Desaturase from <i>Acanthamoeba</i>)	1353 bp	15

Example 4: Cloning of the PUFA specific desaturases from *Perkinsus marinus*
(= SEQ ID NO: 7, 17 and 23)

5 *Perkinsus marinus*, which belongs to the Protista, is a parasite in seashells. *P. marinus* is able to synthesize LCPUFAs such as arachidonic acid (20:4). The LCPUFAs are produced according to the present work over the Δ -8/ Δ -5-fatty acid pathway (see figure 1).

10 *P. marinus* was grown at 28°C as disclosed by La Peyre et al. (J: *Eukaryot. Microbiol.* 1993, 40: 304 - 310).

The total imRNA was isolated from said harvested cells with the aid of the RNeasy plant mini Kit (Qiagen). cDNA was synthesized from the total mRNA with the SMART RACE cDNA amplification kit (Clontech) according to the instructions of the manufacturer.

15 For the isolation of new desaturase genes the following degenerated primers were used for the amplification:

Deg1 :

5'- GGITGG(CZTZA)TIGGICA(TZC) GA(TZC)(GT) (CT)I(GT) (GC)ICA-3'

Deg2:

5'- GG(AZG)AA(TCGA)AG(AZG)TG(AZG)TG(TZC)TC(AZGZT)AT(TZC)TG-S'

The aforementioned primers were used for the amplification in combination with the 3'-adapter-primer of the SMART RACE cDNA amplification kit.

5 The following protocol was used for the amplification:

d) 2 min at 95°C,

e) 30 sec at 94 °C
30 sec at 55-72°C
2 min at 72 °C

10 Number of cycles: 30

f) 10 min at 72 °C

PCR amplicons were cloned and sequenced according to the instructions of the manufacturer (pTOPO, Invitrogen). The sequence information was used for the production of full-length clones. For the cloning of the full-length clones 5'- and 3'-specific primers were synthesized. Said primers were used for the amplification in the SMART RACE cDNA amplification kit (Clontech) and the amplicons were cloned into the pTOPO vector (Invitrogen). Three sequences were identified, which show low similarities to desaturase genes.

Table 2: Perkinsus marinus desaturase sequences

Gene	Nucleotide bp	SEQ ID NO:
12Pm (Δ -12 -Desaturase from Perkinsus)	1254 bp	23
8Pm (Δ -8-Desaturase from Perkinsus)	1236 bp	7
5Pm (Δ -5-Desaturase from Perkinsus)	1374bp	17

Example 5: Cloning of expression plasmids for the heterologous expression of *A. castellanii* and *P. marinus* genes in yeasts

For the heterologous expression in yeasts the respective sequences were PCR amplified and with the restriction enzymes KpnI-SacI the resulting sequences were cloned into the yeast vector pYES2 (Invitrogen). For the amplification specific primers (see table 3 below) were used. Only the open reading frames of the PUFA genes were amplified. In addition restriction cleavage sites were attached to the nucleic acid sequences. At the 5'-end a KpnI site and a so named Kozak sequence (Cell, 1986, 44: 283 - 292) was added. To the 3'-end a SacI site was attached.

10 Table 3: Primers for the amplification of the nucleic acid sequences of the desaturases

Gen	bp	primer	SEQ ID NO:
12Ac	1224	Fwd : GGTACCATGGCGATCACGACGACGCAGACAC	25
		Rvs : GAGCTCCTAGTGGGCCTTGCCGTGCTTGATCTCC	26
8Ac	1374	Fwd : GGTACCATGGTCCTCACAACCCCGGCCCTC	27
		Rvs : GGAGCTCTCAGTTCTCAGCACCCATCTTC	28
5Ac	1353	Fwd: GGTACCATGGCCACCGCATCTGCATC	29
		Rvs: GGAGCTTTAGCCGTAGTAGGCCTCCTT	30
9Ac	891	Fwd : GGTACCATGGCGGCTGCGACGGCGAC	31
		Rvs: GGAGCTTTAGTCGTGCTTCCTCTTGGG	32
12Pm	1254	Fwd : GGTACCATGACCCAAACTGAGGTCCA	33
		Rvs: GGAGCTCTAACGAGAAGTGCGAGCGT	34
8Pm	1236	Fwd : GGTACCATGTCTTCTCTTACCCTCTA	35
		Rvs: GGAGCTCTATTCCACTATGGCAACAG	36
5Pm	1374	Fwd : GGTACCATGACTACTTCAACCACTAC	37
		Rvs: GGAGCTCTACCTAGCAAGCAATCTCT	38

Composition of the PCR mix (50 μ l)

- 5,00 μ L Template cDNA
- 5,00 μ L 10x Puffer (Advantage-Polymerase)+ 25mM MgCl₂
- 5,00 μ L 2mM dNTP
- 5 1,25 μ L each primer (10 pmol/ μ L of the 5'-ATG as well as of the 3'-stopp primer)
- 0,50 μ L Advantage polymerase

The Advantage polymerase from Clontech was employed.

PCR protocol

- 10 Addition temperature: 1 min at 55 °C
- Denaturing temperature: 1 min at 94 °C
- Elongation temperature: 2 min at 72 °C
- Number of cycles: 35

- The PCR products and the vector pYES2 were incubated with the restriction enzymes
- 15 KpnI and SacI for 1 h at 37°C. Afterwards a ligation reaction was done with the Rapid Ligation Kit (Roche) according to the instructions of the manufacturer. The reaction mixture was then used for the transformation of *E. coli* DH5 α cells (Invitrogen) again according to the instructions of the manufacturer. Positive clones were identified with PCR (reaction scheme as described above). The plasmid DNA was isolated (Qiagen
 - 20 Dneasy) and the resulting plasmids were checked by sequencing and transformed with the lithium acetate method into the *Saccharomyces* strain W303-1A. As a control the plasmid pYES2 (vector without insert) was transformed in parallel. The transformed yeasts were selected on complete minimal dropout uracil medium (CMdum) agar plates supplemented with 2% glucose, but without uracil.
 - 25 To express the genes from *A. castellanii* and *P. marinus*, precultures consisting of in each case 5 ml of CMdum dropout uracil liquid medium supplemented with 2% (w/v) raffinose, but without uracil were initially inoculated with the selected transformants and incubated for 2 days at 30°C and 200 rpm. Then, 5 ml of CMdum (without uracil)
 - 30 liquid medium supplemented with 2% of raffinose and 300 μ M of various fatty acids were inoculated with the precultures to an OD₆₀₀ of 0.05. Expression was induced by the addition of 2% (w/v) of galactose. The cultures were incubated for a further 96 hours at 22°C.

Example 6: Cloning of expression plasmids for the expression in plants

To transform plants, a further transformation vector based on pBIN19-35S (Bevan M. (1984) Binary Agrobacterium vectors for plant transformation. Nucl. Acids Res. 18:203) was generated. To this end, BamHI-XbaI cleavage sites were inserted at the 5' and 3' end of the coding sequences, using PCR. The corresponding primer sequences were derived from the 5' and 3' regions of the respective nucleic acid sequence (see table 4).

Table 4: Primers for the expression in plants

Gen	bp	primer	SEQ ID NO:
12Ac	1224	Fwd : GGATCC ACCATGGCGATCACGACGACGCAGACA C	39
		Rvs : GGTCTAG ACTAGTGGGCCTTGCCGTGCTTGATCT CC	40
8Ac	1374	Fwd : GGATCC AGGATGGTCCTCACAACCCCGGCCCTC	41
		Rvs : GGTCTAG ATCAGTTCTCAGCACCCATCTTC	42
5Ac	1353	Fwd: GGATCC ATGGCCACCGCATCTGCATC	43
		Rvs: GGTCTAG ATTAGCCGTAGTAGGCCTCCTT	44
9Ac	891	Fwd : GGATCC ATGGCGGCTGCGACGGCGAC	45
		Rvs: GGTCTAG ATTAGTCGTGCTTCCTCTTGGG	46
12Pm	1254	Fwd : GGATCC ATGACCCAACTGAGGTCCA	47
		Rvs: GGTCTAG ACTAACGAGAAGTGCGAGCGT	48
8Pm	1236	Fwd : GGATCC ATGTCTTCTCTTACCCTCTA	49
		Rvs: GGTCTAG ACTATTCCAATATGGCAACAG	50
5Pm	1374	Fwd : GGATCC ATGACTACTTCAACCACTAC	51
		Rvs: GGTCTAG ACTACCTAGCAAGCAATCTCT	52

Composition of the PCR mix (50 µl):

- 5.00 µl template cDNA
- 5.00 µl 10x buffer (Advantage polymerase)+ 25mM MgCl₂
- 5.00 µl 2mM dNTP
- 5 1.25 µl of each primer (10 pmol/µl)
- 0.50 µl Advantage polymerase

The Advantage polymerase from Clontech was employed.

PCR reaction conditions:

- Annealing temperature: 1 min 55°C
- 10 Denaturation temperature: 1 min 94°C
- Elongation temperature: 2 min 72°C
- Number of cycles: 35

- The PCR products as well as the vector pBin19-35S were incubated with the restriction enzymes BamHI and XbaI for 16 hours at 37°C. Afterwards a ligation
- 15 reaction was done with the Rapid Ligation Kit (Roche) according to the instructions of the manufacturer. The reaction mixture was then used for the transformation of *E. coli* DH5α cells (Invitrogen) again according to the instructions of the manufacturer.

- Positive clones were identified with PCR (reaction scheme as described above) and the plasmid DNA was isolated (Qiagen Dneasy). The resulting plasmids were checked
- 20 by sequencing and transformed by electroporation into *Agrobacterium tumefaciens* GC3101. Afterwards the transformants were plated on 2% YEB Medium agar plates with kanamycin. Kanamycin tolerant cells were picked and used for the transformation of *Arabidopsis thaliana*.

Example 7: Expression of *A. castellanii* and *P. marinus* genes in yeasts

- 25 Yeasts which had been transformed with the plasmids pYES2, pYES-12Ac, pYES-8Ac, pYES2-5Ac, pYES2-9Ac, pYES2-12Pm, pYES2-8Pm and pYES2-5Pm as described in Example 5 were analyzed as follows:

- The yeast cells from the main cultures were harvested by centrifugation (100 x g, 5 min, 20°C) and washed with 100 mM NaHCO₃, pH 8.0 to remove residual medium
- 30 and fatty acids. Starting with the yeast cell sediments, fatty acid methyl esters (FAMES) were prepared by acid methanolysis. To this end, the cell sediments were incubated for one hour at 80°C together with 2 ml of 1 N methanolic sulfuric acid and

2% (v/v) of dimethoxypropane. The FAMES were extracted twice with petroleum ether (PE). To remove nonderivatized fatty acids, the organic phases were washed in each case once with 2 ml of 100 mM NaHCO₃, pH 8.0 and 2 ml of distilled water.

Thereafter, the PE phases were dried with Na₂SO₄, evaporated under argon and
5 taken up in 100 µl of PE. The samples were separated on a DB-23 capillary column (30 m, 0.25 mm, 0.25 µm, Agilent) in a Hewlett-Packard 6850 gas chromatograph equipped with flame ionization detector. The conditions for the GLC analysis were as follows: the oven temperature was programmed from 50°C to 250°C with a rate of 5°C/min and finally 10 min at 250°C (holding).

10 The signals were identified by comparing the retention times with corresponding fatty acid standards (Sigma). The methodology is described for example in Napier and Michaelson, 2001, *Lipids*. 36 (8):761-766; Sayanova et al., 2001, *Journal of Experimental Botany*. 52 (360):1581-1585, Sperling et al., 2001, *Arch. Biochem. Biophys.* 388 (2):293-298 and Michaelson et al., 1998, *FEBS Letters*. 439 (3):215-
15 218.

Example 8: Functional characterization of the genes of *A. castellanii*

The substrate activity and specificity of the genes were determined after expression and after feeding various fatty acids. The substrate specificity of the desaturases after expressions in yeasts can be determined by feeding various different fatty acids.

20 Specific examples for the determination of the specificity and activity are disclosed for example in WO 93/1 1245, WO 94/1 1516, WO 93/06712, US 5,614,393, US5614393, WO 96/21 022, WO0021 557 und WO 99/271 11, Qiu et al. 2001, *J. Biol. Chem.* 276, 31561-31566 for Δ^4 -desaturases, Hong et al. 2002, *Lipids* 37,863-868 for Δ^5 -desaturases. WO2005/012316 teaches such a method for example in example 18 in
25 more detail.

a) Characterization of the gene 12Ac:

First the construct pYES-12Ac was tested in yeasts without feeding fatty acids. Astonishingly it was shown in comparison to the control vector pYES2 (vector without insert) that even without feeding fatty acids new fatty acids are detectable in the yeasts (Fig-
30 ure 2 A and B).

Figure 2 A and B show a comparison of the fatty acid profile between the control (construct pYES2 without insert, Figure 2A) and the construct pYES2-12Ac (Figure 2B), which contains the *Acanthamoeba castellanii* gene for the Δ -12-/ Δ -15-desaturase. The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3, whereas the unusual fatty acids 16:2n-4 and 16:3n-1 are formed for the C 16 fatty acids. For the C 18 fatty acids linoleic and linolenic acid (18:2n-6 and 18:2n-3) are formed.

According to the new synthesized fatty acids it is possible to identify the gene product of the nucleic acid sequence as a Δ -12-desaturase. The enzyme is able to desaturate C18:1 and C16:1 as substrate to the corresponding C18:2 and C16:2 fatty acids. The conversion rate of C18:1 (40,0%) is higher than the rate of the C16:1 (15,8%) conversion. That means the conversion rate of C18:1 is more than double than the conversion rate of the C16:1 .

The conversion rate of the desaturase was calculated according to the following formula:

$$\frac{\text{Substrate}}{(\text{Substrate} + \text{Product})} \times 100$$

The result of the formula is given as percentage value.

Furthermore the enzyme shows in addition a clear Δ -15-desaturase-activity. That means also that products of the Δ -12-desaturase reaction, which are C16:2 and/or C18:2 are further desaturated to C16:3 and/or C 18:3.

b) Characterization of the gene 8Ac:

According to different sequence alignments (Blast) performed with the sequence SEQ ID NO: 3 (8Ac sequence) with different data bases (NCBI-BLAST: <http://www.ncbi.nlm.nih.gov/BLAST/>) the encoded protein sequence is most likely a putative Δ -5-desaturase.

	Sequences with significant similarities	(bits)	Value
	gi 16033740 gb AAL1_331.1.1.1 delta-5 fatty acid desaturase [P...	176	1e-42
	gi 50882495 gb AAT85663.1 polyunsaturated fatty acid delta...	170	6e-41
5	gi 4150956 dbj BAA37090.1 1 delta 5 fatty acid desaturase [D...	156	9e-37
	gi 23894018 emb CAD53323.1 delta 5 fatty acid desaturase [...	156	1e-36
	gi 33466346 gb AAQ1_9605.1.1 delta-4 fatty acid desaturase [E...	150	7e-35
	gi 5263169 dbj BAA81814.1 fatty acid desaturase [Dictyoste...	149	1e-34
	gi 25956288 gb AAN75707.1 delta 4-desaturase [Thraustochyt...	142	1e-32
10	gi 25956290 gb AAN75708.1 1 delta 4-desaturase [Thraustochyt...	139	1e-31
	gi 25956294 gb AAN75710.1 delta 4-desaturase [Thraustochyt...	139	1e-31
	gi 25956292 gb AAN75709.1 delta 4-desaturase [Thraustochyt...	138	2e-31
	gi 20069125 gb AAM09688.1 delta-4 fatty acid desaturase [T...	138	3e-31
	gi 39545945 gb AAR28035.1 delta-5 desaturase [Mortierella ...	136	9e-31
15	gi 3859488 gb AAC72755.1 1 delta-5 fatty acid desaturase [Mo...	135	2e-30
	gi 41017070 sp O74212 FAD5_MORAP Delta-5 fatty acid desatur...	130	7e-29
	gi 48854274 ref ZP_00308437.1 COG3239: Fatty acid desatura...	114	4e-24
	gi 48854276 ref ZP_00308439.1 COG3239: Fatty acid desatura...	114	7e-24

According to this putative activity different fatty acids were fed (18:2, 18:3, 20:3n-6, 20:4n-3). None of said fatty acids were desaturated by the enzyme. This result clearly shows that the protein encoded by the 8Ac gene has neither a Δ -5-desaturase activity nor a Δ -6-desaturase activity.

Unexpectedly after feeding of the fatty acids 20:2n-6 und 20:3n-3 it could be shown, that the 8Ac sequence encodes a Δ -8-desaturase (see figures 3 A, 3 B, 4 A and 4 B).

Figure 3 A and B shows the fatty acid profile of yeasts transformed with the construct pYES2 as control (Figure 3 A) and pYES2-8Ac (Figure 3 B) and fed with the fatty acid C20:2^{A11 14}. The respective fatty acids are market.

Figure 4 A and B shows the fatty acid profile of yeast transformed with the construct pYES2 (Figure 4 A) as control and pYES2-8Ac (Figure 4 B) and fed with the fatty acid C20:3^{A11 14 17}. The respective fatty acids are market.

The protein encoded by 8Ac sequence is therefore a Δ -8-desaturase. The conversion rates for the fatty acids C20:2 and C20:3 are 15,2% and 17,5% respectively. This is absolutely astonishing as the 8Ac sequence, which has some similarities to "front-end" desaturases, has a different conserved region of the characteristic Cyt b5 motif His-Pro-Gly-Gly (HPGG), which is necessary for building the Heme domain. In general mutations in said domain lead to depletion of the enzymatic activity (Sayanova et al. 1999, Plant Physiol 121 (2):641-646). The amino acid sequence of this new Δ -8-desaturase shows unexpected differences to known "front-end" desaturases. Instead of the HPGG motive this desaturase shows the motive HPAG, which is due to an alanine in position 44 of the sequence. Sayanova et al. 1999, Plant Physiol 121(2):641-646 has shown that such a change of the motive from HPPG to HPAG leads to inactive enzymes. Therefore the activity of the new Δ -8-desaturase is even more astonishing.

For the further improvement of the activity of the Δ -8-desaturase, the sequence of the enzyme was mutagenized. The following primer.

8AcMf CAAGTACCACCCGGGCGGCAGCAGGGCCA and
8AcMr TGGCCCTGCTGCCGCCCGGGTGGTACTTG

were used together with the site directed mutagenesis Kit (Stratagene) for the mutagenesis according to the instructions of the manufacturer of the Δ -8-desaturase. The mutagenesis was afterwards checked by sequencing. Due to the mutagenesis the nucleotide sequences 124-CACCCGGGCCGGC was changed to 124-CACCCGGGGCGGC, which leads to a change from Alanine to Glycine in position 44 of the nucleic acid sequence shown in SEQ ID NO: 3. The resulting sequence is shown in SEQ ID NO: 5. As already described for the sequence of 8Ac the mutated sequence 8AcM was also cloned into the vector pYES2 and transformed into yeast. Yeast transformed either with the vector pYES-8Ac or pYES2-8AcM were grown and fed in parallel with different fatty acids (see table 5). The results of the feeding are shown in table 5. The mutated enzyme 8AcM shows in comparison to the wild type enzyme 8Ac an increased activity towards the fatty acid C20:2. This is a two fold increase of the activity. The mutation has no influence of the activity with the fatty acid

C20:3 as substrate. This clearly shows that with the mutation the activity of the Δ -8-desaturase can be influenced in a very specific manner.

Table 5. Fatty acid conversion rate of yeasts transformed with pYES-8Ac or pYES2-8AcM

Plasmid	Fatty acid C20:2	Fatty acid C20:3
pYES-8Ac	15,2 %	17,5 %
pYES2-8AcM	30,0 %	17,2 %

5

The mutated Δ -8-desaturase 8AcM and its derivatives are especially useful alone or in combination with the Δ -12- and Δ -15-desaturase, the Δ -9-elongase and the Δ -5-desaturase for the synthesis of arachidonic acid.

c) Characterization of the gene 5Pm:

10 The constructs pYES2 and pYES-5Pm were transformed into yeasts grown in parallel as described. Afterwards 250 μ M of different fatty acids were fed. During this feeding experiments it can be shown that fatty acids such as C16:0, C16:1, C18:0, C18:1, C18:2n-6, C20:2n-6 or C22:4n-6 are not desaturated by the protein encoded by the 5Pm sequence. Whereas the substrate C20:3n-6 was desaturated by the enzyme
 15 (see figures 5 A and 5 B). Figures 5 A and 5 B clearly shows that the enzyme produces arachidonic acid during the transformation of the fatty acid substrate C20:3n-6. No new fatty acid is produced by the control (Figure 5 A). The desaturation of the fatty acid substrate C20:3n-6 to arachidonic acid is due to a Δ -5-desaturase activity, which is encoded by the 5Pm sequence (SEQ ID NO: 17). The conversion rate calculated
 20 according to the equation mentioned above is 15,4%.

Figure 5 A and 5 B shows the comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (Figure 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (Figure 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).
 25

d) Characterization of the genes 5Ac, 9Ac, 12Pm und 8Pm:

According to sequence comparisons it was able to identify the sequences 5Ac, 12Pm and 8Pm as desaturases having a Δ -5-desaturase, Δ -12-desaturase and Δ -8-desaturase activity. For the sequence 9Ac we were able to show a Δ -9-elongase activity.

In combination with the 12Ac and 8Ac gene the complete set of enzymes from *A. castellanii*, which is necessary for the synthesis for arachidonic (C20:4n-6) or eicosapentaenoic acid could be identified. In addition further genes for the synthesis of said aforementioned fatty acids are isolated from *P. marinus*. With the aid of said genes the PUFA and/or LCPUFA content can be further improved. For the synthesis of arachidonic acid or eicosapentaenoic acid said genes can be introduced in plants or microorganism (see example 8).

Example 8: Generation of transgenic plants

a) Generation of transgenic oilseed rape plants (modified method of Moloney et al., 1992, Plant Cell Reports, 8:238-242)

Binary vectors in *Agrobacterium tumefaciens* C58C1 :pGV2260 or *Escherichia coli* (Deblaere et al, 1984, Nucl. Acids. Res. 13, 4777-4788) can be used for generating transgenic oilseed rape plants. To transform oilseed rape plants (Var. Drakkar, NPZ Nordeutsche Pflanzenzucht, Hohenlieth, Germany), a 1:50 dilution of an overnight culture of a positively transformed agrobacterial colony in Murashige-Skoog medium (Murashige and Skoog 1962 Physiol. Plant. 15, 473) supplemented with 3% sucrose (3MS medium) is used. Petioles or hypocotyls of freshly germinated sterile oilseed rape plants (in each case approx. 1 cm²) are incubated with a 1:50 agrobacterial dilution for 5-10 minutes in a Petri dish. This is followed by 3 days of coincubation in the dark at 25°C on 3MS medium supplemented with 0.8% Bacto agar. The cultures are then grown for 3 days at 16 hours light/8 hours dark and the cultivation is continued in a weekly rhythm on MS medium supplemented with 500 mg/l Claforan (cefotaxim sodium), 50 mg/l kanamycin, 20 μ M benzylaminopurine (BAP), now supplemented with 1.6 g/l of glucose. Growing shoots are transferred to MS medium supplemented with 2% sucrose, 250 mg/l Claforan and 0.8% Bacto agar. If no roots develop after three weeks, 2-indolebutyric acid was added to the medium as growth hormone for

rooting.

Regenerated shoots are obtained on 2MS medium supplemented with kanamycin and Claforan; after rooting, they are transferred to compost and, after growing on for two weeks in a controlled-environment cabinet or in the greenhouse, allowed to flower, and mature seeds are harvested and analyzed by lipid analysis for elongase and/or desaturase expression, such as Δ -12- and Δ -15-desaturase, Δ -8-desaturase, Δ -9-elongase or Δ -5-desaturase activity. In this manner, lines with elevated contents of PUFAs and/or LCPUFAs can be identified.

b) Generation of transgenic linseed plants

10 Transgenic linseed plants can be generated for example by the method of Bell et al., 1999, *In Vitro Cell. Dev. Biol.-Plant.* 35(6):456-465 by means of particle bombardment. In general, linseed was transformed by an agrobacteria-mediated transformation, for example by the method of Mlynarova et al. (1994), *Plant Cell Report* 13: 282-285.

15 c) Generation of transgenic Arabidopsis plants

Binary plasmids were transferred to *A. tumefaciens* strain GV3101 by electroporation and kanamycin-resistant colonies were selected in all cases. Wildtype Col0 or transgenic line CA1-9, containing the coding region of *l. galbana* elongating activity, *lgASEI* [Qi, B., Beaudoin, F., Fraser, T., Stobart, A.K., Napier, J.A. and Lazarus, CM. (2002) Identification of a cDNA encoding a novel C18-D9 polyunsaturated fatty acid-specific elongating activity from the docosahexaenoic acid (DHA)-producing microalga, *Isochrysis galbana*. *FEBS Lett.* 510, 159-65] was used as the host for transformation with *A. castellanii* Δ^8 -desaturase gene. *A. tumefaciens*-mediated transformation was performed as described in Bechthold et al. [(1993) *In planta Agrobacterium-mediated gene transfer by infiltration of Arabidopsis thaliana* plants. *CR. Acad. Sci. Ser. III Sci.Vie.*, 316, 1194-1199.] and seeds from dipped plants were spread on Murashige and Skoog medium containing $50 \mu\text{g ml}^{-1}$ kanamycin.

Example 9: Lipid extraction from leaves

The effect of the genetic modification in plants, fungi, algae, ciliates or on the

production of a desired compound (such as a fatty acid) can be determined by growing the modified microorganisms or the modified plant under suitable conditions (such as those described above) and analyzing the medium and/or the cellular components for the elevated production of desired product (i.e. of the lipids or a fatty acid). These analytical techniques are known to the skilled worker and comprise spectroscopy, thin-layer chromatography, various types of staining methods, enzymatic and microbiological methods and analytical chromatography such as high-performance liquid chromatography (see, for example, Ullman, Encyclopedia of Industrial Chemistry, Vol. A2, p. 89-90 and p. 443-613, VCH: Weinheim (1985); Fallon, A., et al., (1987) "Applications of HPLC in Biochemistry" in: Laboratory Techniques in Biochemistry and Molecular Biology, Vol. 17; Rehm et al. (1993) Biotechnology, Vol. 3, Chapter III: "Product recovery and purification", p. 469-714, VCH: Weinheim; Belter, P.A., et al. (1988) Bioseparations: downstream processing for Biotechnology, John Wiley and Sons; Kennedy, J.F., and Cabral, J.M.S. (1992) Recovery processes for biological Materials, John Wiley and Sons; Shaeiwitz, J.A., and Henry, J.D. (1988) Biochemical Separations, in: Ullmann's Encyclopedia of Industrial Chemistry, Vol. B3; Chapter 11, p. 1-27, VCH: Weinheim; and Dechow, F.J. (1989) Separation and purification techniques in biotechnology, Noyes Publications).

In addition to the abovementioned processes, plant lipids are extracted from plant material as described by Cahoon et al. (1999) Proc. Natl. Acad. Sci. USA 96 (22): 12935-12940 and Browse et al. (1986) Analytic Biochemistry 152:141-145. The qualitative and quantitative analysis of lipids or fatty acids is described by Christie, William W., Advances in Lipid Methodology, Ayr/Scotland: Oily Press (Oily Press Lipid Library; 2); Christie, William W., Gas Chromatography and Lipids. A Practical Guide - Ayr, Scotland: Oily Press, 1989, Repr. 1992, IX, 307 pp. (Oily Press Lipid Library; 1); "Progress in Lipid Research, Oxford: Pergamon Press, 1 (1952) - 16 (1977) under the title: Progress in the Chemistry of Fats and Other Lipids CODEN.

One example is the analysis of fatty acids (abbreviations: FAME, fatty acid methyl ester; GC-MS, gas liquid chromatography/mass spectrometry; TAG, triacylglycerol; TLC, thin-layer chromatography).

The unambiguous detection for the presence of fatty acid products can be obtained by analyzing recombinant organisms using analytical standard methods: GC, GC-MS

or TLC, as described on several occasions by Christie and the references therein (1997, in: Advances on Lipid Methodology, Fourth Edition: Christie, Oily Press, Dundee, 119-169; 1998, Gaschromatographie-Massenspektrometrie-Verfahren [Gas chromatography/mass spectrometric methods], Lipide 33:343-353).

- 5 The material to be analyzed can be disrupted by sonication, grinding in a glass mill, liquid nitrogen and grinding or via other applicable methods. After disruption, the material must be centrifuged. The sediment is resuspended in distilled water, heated for 10 minutes at 100°C, cooled on ice and recentrifuged, followed by extraction for one hour at 90°C in 0.5 M sulfuric acid in methanol with 2% dimethoxypropane, which
10 leads to hydrolyzed oil and lipid compounds, which give transmethylated lipids. These fatty acid methyl esters are extracted in petroleum ether and finally subjected to a GC analysis using a capillary column (Chrompack, WCOT Fused Silica, CP-Wax-52 CB, 25 µm, 0.32 mm) at a temperature gradient of between 170°C and 240°C for 20 minutes and 5 minutes at 240°C. The identity of the resulting fatty acid methyl
15 esters must be defined using standards, which are available from commercial sources (i.e. Sigma).

Plant material is initially homogenized mechanically by comminuting in a pestle and mortar to make it more amenable to extraction.

- This is followed by heating at 100°C for 10 minutes and, after cooling on ice, by
20 resedimentation. The cell sediment is hydrolyzed for one hour at 90°C with 1 M methanolic sulfuric acid and 2% dimethoxypropane, and the lipids are transmethylated. The resulting fatty acid methyl esters (FAMES) are extracted in petroleum ether. The extracted FAMES are analyzed by gas liquid chromatography using a capillary column (Chrompack, WCOT Fused Silica, CP-Wax-52 CB, 25 m,
25 0.32 mm) and a temperature gradient of from 170°C to 240°C in 20 minutes and 5 minutes at 240°C. The identity of the fatty acid methyl esters is confirmed by comparison with corresponding FAME standards (Sigma). The identity and position of the double bond can be analyzed further by suitable chemical derivatization of the FAME mixtures, for example to give 4,4-dimethoxyoxazoline derivatives (Christie,
30 1998) by means of GC-MS.

Leaf material from transgenic *Arabidopsis thaliana* Col0 and super-transformants of

transgenic line CA1-9 both transformed with the construct pBIN1935S-8Ac were analyzed by gas chromatography of methyl ester derivatives as described above. Identities were confirmed by GC-MS and co-migration with authentic standards. The conversion rates are shown in the following table 6:

Table 6: Conversion rate with AcD8 (delta-8-desaturase from *Acanthamoeba castellanii*) of different substrates

fatty acids	% of total fatty acids	% conversion of substrate
20:2 ^{Δ11 14}	1.1	-
20:3 ^{Δ8 11 14}	1.9	63
20:2 ^{Δ11 14 17}	1.3	-
20:2 ^{Δ8 11 14 17}	0.8	40

Figure 6 shows the result with the line CA1-9. In the double transgenic *Arabidopsis* a clear activity of Ac8 can be shown by the conversion of the present 20:2^{Δ11 14} or 20:3^{Δ11 14 17} into 20:3^{Δ8 11 14} or 20:4^{Δ8 11 14 17}, the precursors of arachidonic acid or eicosapentaenoic acid.

Additionally Acyl-CoA profiles were done from the *Arabidopsis* leaves of *Arabidopsis* wild type (Figure 7 A), *Arabidopsis* Δ9elo (Figure 7 B) and *Arabidopsis* Δ9eloΔ8des (Figure 7 C) using the method of Larson et al. [Plant J. 2002 Nov;32(4):519-27]. Results from the measurements are shown in Figure 7 and demonstrate again the functionality of 8Ac in plants.

Equivalents:

Many equivalents of the specific embodiments according to the invention described herein can be identified or found by the skilled worker resorting simply to routine experiments. These equivalents are intended to be within the scope of the patent claims.

What is claimed is:

1. A process for the production of arachidonic acid or eicosapentaenoic acid or arachidonic acid and eicosapentaenoic acid in transgenic plants that produce mature seeds with a content of at least 1 % by weight of said compounds referred to the total lipid content of said organism which comprises the following steps:
 - a) introduction of at least one nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -12-desaturase- and Δ -15-desaturase-activity, and
 - 10 b) introduction of at least one second nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -9-elongase-activity, and
 - c) introduction of at least one third nucleic acid sequence in said transgenic plant, which encodes a polypeptide having a Δ -8-desaturase-activity, and
 - 15 d) introduction of at least a one fourth nucleic acid sequence, which encodes a polypeptide having a Δ -5-desaturase-activity, and
 - e) cultivating and harvesting of said transgenic plant.
2. The process as claimed in claim 1, wherein the nucleic acid sequences which encode polypeptides having Δ -12-desaturase and Δ -15-desaturase activity, Δ -8-desaturase, Δ -9-elongase or Δ -5-desaturase activity are selected from the group consisting of
 - a) a nucleic acid sequence depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 and SEQ ID NO: 23, and
 - 25 b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18,

SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 according to the degeneracy of the genetic code,

- 5 c) derivatives of the nucleic acid sequences depicted in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, , SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 which encode polypeptides having at least 50 % homology to the sequence as depicted in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having Δ -12-desaturase and Δ -15-desaturase activity, Δ -8-desaturase, Δ -9-elongase or Δ -5-desaturase activity.
- 10 3. The process as claimed in claim 1 or 2, wherein the transgenic plant that produces mature seeds is an oilseed plant.
- 15 4. The process as claimed in claim 1 or 2, wherein the transgenic plant that produces mature seeds is selected from the group consisting of the plant families of Anacardiaceae, Asteraceae, Apiaceae, Boraginaceae, Brassicaceae, Cannabaceae, Elaeagnaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Leguminosae, Linaceae, Lythraeae, Malvaceae, Onagraceae, Palmae, Poaceae, Rubiaceae, Scrophulariaceae, Solanaceae, Sterculiaceae and Theaceae.
- 20 5. The process as claimed in claim 1 or 2, wherein the transgenic plant that produces mature seeds is selected from the group consisting of the plant genera of Pistacia, Mangifera, Anacardium, Calendula, Carthamus, Centaurea, Cichorium, Cynara, Helianthus, Lactuca, Locusta, Tagetes, Valeriana, Borago, Daucus, Brassica, Camelina, Melanosinapis, Sinapis, Arabadopsis, Orychophragmus, Cannabis, Elaeagnus, Manihot, Janipha, Jatropha, Ricinus, Pisum, Albizia, Cathormion, Feuillea, Inga, Pithecolobium, Acacia, Mimosa, Medicago, Glycine, Dolichos, Phaseolus, Pelargonium, Cocos, Oleum, Juglans, Wallia, Arachis, 25 *Linum*, Punica, Gossypium, Camissonia, Oenothera, Elaeis, Hordeum, Secale, Avena, Sorghum, Andropogon, Holcus, Panicum, Oryza, Zea, Triticum, Coffea,
- 30

Verbascum, Capsicum, Nicotiana, Solanum, Lycopersicon, Theobroma and Camellia.

6. The process as claimed in any of the claims 1 to 5, wherein the transgenic plant is selected from the group consisting of rapeseed, poppy, mustard, hemp, castor bean, sesame, olive, calendula, punica, hazel nut, maize, almond, macadamia, cotton, avocado, pumpkin, walnut, laurel, pistachio, primrose, canola, evening primrose, oil palm, peanut, linseed, soybean, safflower, marigold, coffee, tobacco, cacao, sunflower and borage.
7. The process as claimed in any of the claims 1 to 6, wherein the arachidonic acid or eicosapentaenoic acid or arachidonic acid and eicosapentaenoic acid is isolated in the form of their oils, lipids or free fatty acids.
8. The process as claimed in any of the claims 1 to 7, wherein arachidonic acid and eicosapentaenoic acid is produced in at least a 1:2 ratio.
9. The process as claimed in any of the claims 1 to 8, wherein the arachidonic acid and eicosapentaenoic acid are produced in a content of at least 5 % by weight referred to the total lipid content.
10. The process as claimed in any of the claims 1 to 9, wherein the Δ -12-desaturase- and Δ -15-desaturase used in the process desaturates C16 or C18-fatty acids having one double bond in the fatty acid chain or C16 and C18-fatty acids having one double bond in the fatty acid chain.
11. An isolated nucleic acid sequence which encodes a polypeptide having a Δ -12-desaturase and Δ -15-desaturase activity selected from the group consisting of
 - a) a nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23;
 - b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24;

- 5 c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 22 which encode polypeptides having at least 40 % homology to the sequence as depicted in SEQ ID NO: 20, SEQ ID NO: 22 or SEQ ID NO: 24 and which polypeptides having Δ -12-desaturase and Δ -15-desaturase activity.
12. An isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -9-elongase selected from the group consisting of
- 10 a) a nucleic acid sequence depicted in SEQ ID NO: 11;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 12;
- 15 c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 11 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 12 and which polypeptides having Δ -9-elongase activity.
13. An isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -8-desaturase selected from the group consisting of
- 20 a) a nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8;
- 25 c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 3, SEQ ID NO: 5 or SEQ ID NO: 7 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 4, SEQ ID NO: 6 or SEQ ID NO: 8 and which polypeptides having Δ -8-desaturase activity.
14. An isolated nucleic acid sequence comprising a nucleotide sequence which encodes a Δ -5-desaturase selected from the group consisting of

- a) a nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17;
- b) a nucleic acid sequence, which, as a result of the degeneracy of the genetic code, can be derived from a polypeptide sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18;
- 5 c) derivatives of the nucleic acid sequence depicted in SEQ ID NO: 15 or SEQ ID NO: 17 which encode polypeptides having at least 70 % homology to the sequence as depicted in SEQ ID NO: 16 or SEQ ID NO: 18 and which polypeptides having Δ -5-desaturase activity.
15. A polypeptide encoded by an isolated nucleic acid sequence as claimed in
10 claims 11 to 14.
16. A gene construct comprising an isolated nucleic acid having the sequence SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 15, SEQ ID NO: 17, SEQ ID NO: 19, SEQ ID NO: 21 or SEQ ID NO: 23 as claimed in
15 claims 11 to 14, where the nucleic acid is functionally linked to one or more regulatory signals.
17. A gene construct as claimed in claim 16, whose gene expression is increased by the regulatory signals.
18. A vector comprising a nucleic acid as claimed in claims 11 to 14 or a gene construct as claimed in claim 17.
- 20 19. A transgenic plant comprising at least one nucleic acid as claimed in claims 11 to 14, a gene construct as claimed in claim 17 or a vector as claimed in claim 18.
20. The transgenic plant as claimed in claim 19, wherein the plant is an oilseed plant.

Figure 1: Biosynthesis pathway to ARA and/or EPA




Figure 2 A: Comparison of the fatty acid profile of yeast transformed with the constructs pYES2 (2A) as control and construct pYES2-12Ac (2B). The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3.




Figure 2B: Comparison of the fatty acid profile of yeast transformed with the constructs pYES2 (2A) as control and construct pYES2-12Ac (2B). The fatty acids are marked. The new fatty acids synthesized are in case of construct pYES2-12Ac (2B) the fatty acids C16:2, C16:3, C18:2 and C18:3.




Figure 3 A: Fatty acid profile of yeasts transformed with the construct pYES2 as control (Figure 3 A) and pYES2-8Ac (Figure 3 B) and fed with the fatty acid C20:2 $\Delta^{11,14}$. The respective fatty acids are marked.




Figure 3 B: Fatty acid profile of yeasts transformed with the construct pYES2 as control (Figure 3 A) and pYES2-8Ac (Figure 3 B) and fed with the fatty acid C20:2 $\Delta^{11,14}$. The respective fatty acids are marked.




Figure 4 A: Fatty acid profile of yeast transformed with the construct pYES2 (Figure 4 A) as control and pYES2-8Ac (Figure 4 B) and fed with the fatty acid C20:3 $\Delta^{11,14,17}$. The respective fatty acids are marked.




Figure 4 B: Fatty acid profile of yeast transformed with the construct pYES2 (Figure 4 A) as control and pYES2-8Ac (Figure 4 B) and fed with the fatty acid C20:3 $\Delta^{11,14,17}$. The respective fatty acids are marked.




Figure 5 A: Comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (Figure 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (Figure 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).




Figure 5 B: Comparison of the fatty acid profile of yeasts transformed with the construct pYES2 as control and fed with the fatty acid C20:3n-6 (Figure 5 A) and with the construct pYES2-5Pm fed with the fatty acid C20:3n-6 (Figure 5 B). The fatty acids are marked. The new synthesized fatty acid is C20:4n-6 (arachidonic acid).





Figure 6: Expression of AcD8 in double transgenic Arabidopsis



7 A) Arabidopsis Wild





Figure 7 A: Expression of the Δ -9-elongase or Δ -9-elongase and Δ -8-desaturase in transgenic Arabidopsis


Figure 7 B: Expression of the Δ -9-elongase or Δ -9-elongase and Δ -8-desaturase in transgenic Arabidopsis

7 B) Arabidopsis Δ 9elo



7 C) Arabidopsis $\Delta 9elo\Delta 8des$

Figure 7 C: Expression of the $\Delta 9$ -elongase or $\Delta 9$ -elongase and $\Delta 8$ -desaturase in transgenic Arabidopsis



PhoenixTemp18528 .tmp .txt
 SEQUENCE LISTING

<110> Rothamsted Research

<120> Process for the production of arachidonic acid and/or eicosapentaenoic acid

<130> PF57175

<140> 20050119

<141> 2005-12-10

<160> 52

<170> Patentln version 3.1

<210> 1

<211> 1266

<212> DNA

<213> Euglena gracilis

<220>

<221> CDS

<222> (1)..(1266)

<223> Delta-8-Desaturase

<400> 1

atg aag tea aag cgc caa gcg ctt ccc ctt aca att gat gga aca aca	48
Met Lys Ser Lys Arg Gin Ala Leu Pro Leu Thr lie Asp Gly Thr Thr	
1 5 10 15	

tat gat gtg tct gcc tgg gtc aat ttc cac cct ggt ggt gcg gaa att	96
Tyr Asp val ser Ala Trp val Asn Phe His Pro Gly Gly Ala Glu lie	
20 25 30	

ata gag aat tac caa gga agg gat gcc act gat gcc ttc atg gtt atg	144
lie Glu Asn Tyr Gin Gly Arg Asp Ala Thr Asp Ala Phe Met Val Met	
35 40 45	

cac tct caa gaa gcc ttc gac aag etc aag cgc atg ccc aaa ate aat	192
His Ser Gin Glu Ala Phe Asp Lys Leu Lys Arg Met Pro Lys lie Asn	
50 55 60	

ccc agt tct gag ttg cca ccc cag get gca gtg aat gaa get caa gag	240
---	-----

PhoenixTemp18528 .tmp .txt

Pro 65	Ser	Ser	Glu	Leu	Pro 70	Pro	Gin	Ala	Ala	Val 75	Asn	Glu	Ala	Gin	Glu 80	
gat	ttc	egg	aag	etc	cga	gaa	gag	ttg	ate	gca	act	ggc	atg	ttt	gat	288
Asp	Phe	Arg	Lys	Leu 85	Arg	Glu	Glu	Leu	lie 90	Ala	Thr	Gly	Met	Phe 95	Asp	
gcc	tec	ccc	etc	tgg	tac	tea	tac	aaa	ate	age	ace	aca	ctg	ggc	ctt	336
Ala	Ser	Pro	Leu 100	Trp	Tyr	Ser	Tyr	Lys 105	lie	Ser	Thr	Thr	Leu 110	Gly	Leu	
gga	gtg	ctg	ggg	tat	ttc	ctg	atg	ggt	cag	tat	cag	atg	tat	ttc	att	384
Gly	Val	Leu 115	Gly	Tyr	Phe	Leu	Met	Val 120	Gin	Tyr	Gin	Met	Tyr 125	Phe	lie	
ggg	gca	gtg	ttg	ctt	ggg	atg	cac	tat	caa	cag	atg	ggc	tgg	ctt	tct	432
Gly	Ala 130	val	Leu	Leu	Gly	Met 135	His	Tyr	Gin	Gin	Met 140	Gly	Trp	Leu	ser	
cat	gac	att	tgc	cac	cac	cag	act	ttc	aag	aac	egg	aac	tgg	aac	aac	480
His	Asp	lie	Cys	His 150	His	Gin	Thr	Phe	Lys 155	Asn	Arg	Asn	Trp	Asn 160	Asn	
etc	gtg	gga	ctg	gta	ttt	ggc	aat	ggg	ctg	caa	ggg	ttt	tec	gtg	aca	528
Leu	val	Gly	Leu 165	val	Phe	Gly	Asn	Gly 170	Leu	Gin	Gly	Phe	ser	val 175	Thr	
tgc	tgg	aag	gac	aga	cac	aat	gca	cat	cat	teg	gca	ace	aat	ggt	caa	576
Cys	Trp	Lys 180	Asp	Arg	His	Asn	Ala	His 185	His	Ser	Ala	Thr	Asn 190	Val	Gin	
ggg	cac	gac	cct	gat	att	gac	aac	etc	ccc	etc	tta	gcc	tgg	tct	gag	624
Gly	His	Asp 195	Pro	Asp	lie	Asp	Asn 200	Leu	Pro	Leu	Leu	Ala 205	Trp	ser	Glu	
gat	gac	gtc	aca	egg	gcg	tea	ccg	att	tec	cgc	aag	etc	att	cag	ttc	672
Asp	Asp 210	Val	Thr	Arg	Ala 215	Ser	Pro	lie	Ser	Arg	Lys 220	Leu	lie	Gin	Phe	
cag	cag	tat	tat	ttc	ttg	gtc	ate	tgt	ate	ttg	ttg	egg	ttc	att	tgg	720
Gin	Gin	Tyr	Tyr	Phe 230	Leu	Val	lie	Cys	lie	Leu	Leu	Arg 235	Phe	lie	Trp 240	
tgt	ttc	cag	age	gtg	ttg	ace	gtg	cgc	agt	ctg	aag	gac	aga	gat	aac	768
Cys	Phe	Gin	Ser 245	Val	Leu	Thr	Val	Arg 250	Ser	Leu	Lys	Asp	Arg	Asp 255	Asn	
caa	ttc	tat	cgc	tct	cag	tat	aag	aag	gag	gcc	att	ggc	etc	gcc	ctg	816
Gin	Phe	Tyr 260	Arg	Ser	Gin	Tyr	Lys 265	Lys	Glu	Ala	lie	Gly	Leu 270	Ala	Leu	
cat	tgg	aca	ttg	aag	gcc	ctg	ttc	cac	tta	ttc	ttt	atg	ccc	age	ate	864
His	Trp	Thr 275	Leu	Lys	Ala	Leu	Phe 280	His	Leu	Phe	Phe	Met 285	Pro	Ser	lie	
etc	aca	teg	ctg	ttg	gta	ttt	ttc	ggt	teg	gag	ctg	ggt	ggc	ggc	ttc	912
Leu	Thr 290	Ser	Leu	Leu	Val	Phe 295	Phe	Val	Ser	Glu	Leu 300	Val	Gly	Gly	Phe	
ggc	att	gcg	ate	gtg	gtg	ttc	atg	aac	cac	tac	cca	ctg	gag	aag	ate	960
Gly	lie 305	Ala	lie	Val	Val 310	Phe	Met	Asn	His	Tyr 315	Pro	Leu	Glu	Lys	lie 320	
ggg	gac	teg	gtc	tgg	gat	ggc	cat	gga	ttc	teg	ggt	ggc	cag	ate	cat	1008
Gly	Asp	Ser 325	Val	Trp	Asp	Gly	His	Gly 330	Phe	Ser	Val	Gly	Gin	lie 335	His	
gag	ace	atg	aac	att	egg	cga	ggg	att	ate	aca	gat	tgg	ttt	ttc	gga	1056
Glu	Thr	Met	Asn	lie	Arg	Arg	Gly	lie	lie	Thr	Asp	Trp	Phe	Phe	Gly	

PhoenixTemp18528 .tmp .txt

```

340                               345                               350
ggc ttg aac tac cag ate gag cac cat ttg tgg ccg ace etc cct cgc      1104
Gly Leu Asn Tyr Gin lie Glu His His Leu Trp Pro Thr Leu Pro Arg
      355                               360                               365

cac aac ctg aca gcg gtt age tac cag gtg gaa cag ctg tgc cag aag      1152
His Asn Leu Thr Ala val ser Tyr Gin val Glu Gin Leu cys Gin Lys
      370                               375                               380

cac aac ctg ccg tat egg aac ccg ctg ccc cat gaa ggg ttg gtc ate      1200
His Asn Leu Pro Tyr Arg Asn Pro Leu Pro His Glu Gly Leu Val lie
      385                               390                               395

ctg ctg cgc tat ctg gcg gtg ttc gcc egg atg gcg gag aag caa ccc      1248
Leu Leu Arg Tyr Leu Ala Val Phe Ala Arg Met Ala Glu Lys Gin Pro
      405                               410                               415

gcg ggg aag get eta taa      1266
Ala Gly Lys Ala Leu
      420

```

<210> 2

<211> 421

<212> PRT

<213> Euglena gracilis

<400> 2

```

Met Lys Ser Lys Arg Gi n Ala Leu Pro Leu Thr lie Asp Gly Thr Thr
1      5      10      15

Tyr Asp Val Ser Ala Trp Val Asn Phe Hi s Pro Gly Gly Al a Gl u lie
      20      25      30

l i e Gl u Asn Tyr Gi n Gly Arg Asp Ala Thr Asp Ala Phe Met Val Met
      35      40      45

Hi s Ser Gi n Gl u Ala Phe Asp Lys Leu Lys Arg Met Pro Lys lie Asn
      50      55      60

Pro Ser Ser Gl u Leu Pro Pro Gi n Ala Ala Val Asn Gl u Al a Gi n Gl u
      65      70      75      80

Asp Phe Arg Lys Leu Arg Gl u Gl u Leu lie 90 Al a Thr Gly Met Phe Asp
      85      90      95

Al a Ser Pro Leu Trp Tyr Ser Tyr Lys lie Ser Thr Thr Leu Gly Leu
      100      105      110

Gly Val Leu Gly Tyr Phe Leu Met Val Gi n Tyr Gi n Met Tyr Phe lie
      115      120      125

Gly Ala Val Leu Leu Gly Met Hi s Tyr Gi n Gi n Met Gly Trp Leu Ser
      130      135      140

```

PhoenixTemp18528 .tmp .txt

His Asp lie Cys His His Gin Thr Phe Lys Asn Arg Asn Trp Asn Asn
 145 150 155 160

Leu Val Gly Leu Val Phe Gly Asn Gly Leu Gin Gly Phe Ser Val Thr
 165 170 175

Cys Trp Lys Asp Arg His Asn Ala His His Ser Ala Thr Asn Val Gin
 180 185 190

Gly His Asp Pro Asp lie Asp Asn Leu Pro Leu Leu Ala Trp Ser Glu
 195 200 205

Asp Asp Val Thr Arg Ala Ser Pro lie Ser Arg Lys Leu lie Gin Phe
 210 215 220

Gin Gin Tyr Tyr Phe Leu Val lie Cys lie Leu Leu Arg Phe lie Trp
 225 230 235 240

Cys Phe Gin Ser Val Leu Thr Val Arg Ser Leu Lys Asp Arg Asp Asn
 245 250 255

Gin Phe Tyr Arg Ser Gin Tyr Lys Lys Glu Ala lie Gly Leu Ala Leu
 260 265 270

His Trp Thr Leu Lys Ala Leu Phe His Leu Phe Phe Met Pro Ser lie
 275 280 285

Leu Thr Ser Leu Leu Val Phe Phe Val Ser Glu Leu Val Gly Gly Phe
 290 295 300

Gly He Ala He Val Val Phe Met Asn His Tyr Pro Leu Glu Lys He
 305 310 315 320

Gly Asp Ser Val Trp Asp Gly His Gly Phe Ser Val Gly Gin lie His
 325 330 335

Glu Thr Met Asn lie Arg Arg Gly lie lie Thr Asp Trp Phe Phe Gly
 340 345 350

Gly Leu Asn Tyr Gin lie Glu His His Leu Trp Pro Thr Leu Pro Arg
 355 360 365

His Asn Leu Thr Ala Val Ser Tyr Gin Val Glu Gin Leu Cys Gin Lys
 370 375 380

His Asn Leu Pro Tyr Arg Asn Pro Leu Pro His Glu Gly Leu Val lie
 385 390 395 400

Leu Leu Arg Tyr Leu Ala val Phe Ala Arg Met Ala Glu Lys Gin Pro
 405 410 415

PhoenixTtempl8528 .tmp .txt

Ala Gly Lys Ala Leu
420

<210> 3

<211> 1374

<212> DNA

<213> Acanthamoeba castellanii

<220>

<221> CDS

<222> (1)..(1374)

<223> Delta-8-Desaturase

<400> 3

atg gtc etc aca ace ccg gcc etc aac ctg aag aag gaa cga acg teg	48
Met Val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Glu Arg Thr Ser	
1 5 10 15	
ttc ace cag gag gag ctt tec aag etc tgg gtc ctt cac ggc cag gtg	96
Phe Thr Gln Glu Glu Leu Ser Lys Leu Trp Val Leu His Gly Gln Val	
20 25 30	
tac gat ttc ace gac ttt gtc aag tac cac ccg gcc ggc age agg gcc	144
Tyr Asp Phe Thr Asp Phe Val Lys Tyr His Pro Ala Gly Ser Arg Ala	
35 40 45	
ate ctg etc ggc cgt ggc cgt gat tgt ace gtg etc ttc gag tec tac	192
lie Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Glu Ser Tyr	
50 55 60	
cac aca gtc ctg cct tec gat get ctt etc gag aag tac cgc gtc tct	240
His Thr Val Leu Pro Ser Asp Ala Leu Leu Glu Lys Tyr Arg Val Ser	
65 70 75 80	
get ccc aac gcc aag etc gag gag age egg tea gcc aag ctg ttc teg	288
Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser	
85 90 95	
ttc gag gag ggt age ttc tac cga ace etc aag cag cga acg cgc gag	336
Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gln Arg Thr Arg Glu	
100 105 110	
tac ttc aag ace aac aac ctg age ace aag gcc ace acg atg gag gtc	384
Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val	
115 120 125	
ate tac ttc gtg gcc ace ate etc age ate tac ttc tgc acg tgg gcc	432
lie Tyr Phe Val Ala Thr lie Leu Ser lie Tyr Phe Cys Thr Trp Ala	
130 135 140	
gcc ttc gtg cag ggt tec etc ate gcc get gtc ctt cac gga gtg ggc	480
Ala Phe Val Gln Gly Ser Leu lie Ala Ala Val Leu His Gly Val Gly	
145 150 155 160	
cgt gcg ate tgt ate ata caa ccg act cat gcg act teg cac tac gcc	528
Arg Ala lie Cys lie lie Gln Pro Thr His Ala Thr Ser His Tyr Ala	
165 170 175	

PhoenixTemp18528 .tmp .txt

atg ttc cgc tea gtg tgg etc aac cag tgg gcc tac agg ate tec atg	576
Met Phe Arg Ser Val Trp Leu Asn Gin Trp Ala Tyr Arg lie Ser Met	
180 185 190	
gcc gtc age gga teg teg ccg gcc cag tgg ace ace aag cac gtc ate	624
Ala Val Ser Gly Ser Ser Pro Ala Gin Trp Thr Thr Lys His Val lie	
195 200 205	
aac cat cac gtc gag ace aac ctg tgc ccc ace gat gac gac ace atg	672
Asn His His Val Glu Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met	
210 215 220	
tac ccc ate aag cgc ate ctg cac gag ttc cct cgt ctg ttc ttc cac	720
Tyr Pro lie Lys Arg lie Leu His Glu Phe Pro Arg Leu Phe Phe His	
225 230 235 240	
aag tac cag cac ate tac ate tgg ctg gtg tac ccc tac ace ace ate	768
Lys Tyr Gin His lie Tyr lie Trp Leu Val Tyr Pro Tyr Thr Thr lie	
245 250 255	
ttg tgg cac ttc tec aac ctg gcc aag etc gcc etc ggc gcc get cgc	816
Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg	
260 265 270	
ggt cag atg tac gag ggt ate gcc aag gtg age caa gag ace teg ggt	864
Gly Gin Met Tyr Glu Gly lie Ala Lys Val Ser Gin Glu Thr Ser Gly	
275 280 285	
gac tgg gtg gag acg gcc atg acg ctg ttc ttc ttc acg ttc tec cgt	912
Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Phe Thr Phe Ser Arg	
290 295 300	
ctg ctg ctg ccc ttc ctg tgc ctg ccc ttc ace acg gcc gcc gcg gtg	960
Leu Leu Leu Pro Phe Leu Cys Leu Pro Phe Thr Thr Ala Ala Ala Val	
305 310 315 320	
ttc ctg etc tec gag tgg ace tgc teg ace tgg ttc gcg ctg cag ttc	1008
Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gin Phe	
325 330 335	
gcc gtg age cac gag gtc gac gag tgc gtc gag cac gag aag teg gtc	1056
Ala Val Ser His Glu Val Asp Glu Cys Val Glu His Glu Lys Ser Val	
340 345 350	
etc gac ace etc aag gcc aac gag gcc aag ggc ate gtc aac cag ggc	1104
Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly lie Val Asn Gin Gly	
355 360 365	
ggc etc gtc gac tgg ggc gcg cac cag gtt egg gcc teg cac aac tac	1152
Gly Leu Val Asp Trp Gly Ala His Gin Val Arg Ala Ser His Asn Tyr	
370 375 380	
tct gcc gac tec ctg ctg teg etc cac ttc age ggt ggc etc aac ctt	1200
Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu	
385 390 395 400	
cag ate gag cac cac etc ttc ccc tec gtc cac tac act cac tac cct	1248
Gin lie Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro	
405 410 415	
gcc ccg tec aag att gtg cag cag acg tgc aag gag ttc aac ttg ccc	1296
Ala Pro Ser Lys lie Val Gin Gin Thr Cys Lys Glu Phe Asn Leu Pro	
420 425 430	
tgc act ctg teg ccg teg atg atg ggt gcc gtg ace aag cac tac cac	1344
Cys Thr Leu ser Pro ser Met Met Gly Ala val Thr Lys His Tyr His	
435 440 445	

Phoeni xtempl8528 .tmp .txt
 cag etc aag aag atg ggt get gag aac tga 1374
 Gin Leu Lys Lys Met gly Ala Glu Asn
 450 455

<210> 4

<211> 457

<212> PRT

<213> Acanthamoeba castellanii

<400> 4

Met Val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Glu Arg Thr Ser
 1 5 10 15

Phe Thr Gin Glu Glu Leu Ser Lys Leu Trp Val Leu His Gly Gin Val
 20 25 30

Tyr Asp Phe Thr Asp Phe Val Lys Tyr His Pro Ala Gly Ser Arg Ala
 35 40 45

Iie Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Glu Ser Tyr
 50 55 60

His Thr Val Leu Pro Ser Asp Ala Leu Leu Glu Lys Tyr Arg Val Ser
 65 70 75 80

Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser
 85 90 95

Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gin Arg Thr Arg Glu
 100 105 110

Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val
 115 120 125

Iie Tyr Phe val Ala Thr lie Leu ser lie Tyr Phe cys Thr Trp Ala
 130 135 140

Ala Phe Val Gin Gly Ser Leu lie Ala Ala Val Leu His Gly Val Gly
 145 150 155 160

Arg Ala lie cys lie Iie Gin Pro Thr His Ala Thr ser His Tyr Ala
 165 170 175

Met Phe Arg Ser Val Trp Leu Asn Gin Trp Ala Tyr Arg Iie Ser Met
 180 185 190

Ala Val Ser Gly Ser Ser Pro Ala Gin Trp Thr Thr Lys His Val lie
 195 200 205

Phoenix\templ8528.tmp.txt

Asn His His Val Glu Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met
 210 215 220

Tyr Pro lie Lys Arg lie Leu His Glu Phe Pro Arg Leu Phe Phe His
 225 230 235 240

Lys Tyr Gin His lie Tyr lie Trp Leu Val Tyr Pro Tyr Thr Thr lie
 245 250 255

Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg
 260 265 270

Gly Gin Met Tyr Glu Gly lie Ala Lys val ser Gin Glu Thr ser Gly
 275 280 285

Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Phe Thr Phe Ser Arg
 290 295 300

Leu Leu Leu Pro Phe Leu cys Leu Pro Phe Thr Thr Ala Ala Ala val
 305 310 315 320

Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gin Phe
 325 330 335

Ala val ser His Glu val Asp Glu cys val Glu His Glu Lys Ser val
 340 345 350

Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly lie Val Asn Gin Gly
 355 360 365

Gly Leu Val Asp Trp Gly Ala His Gin Val Arg Ala Ser His Asn Tyr
 370 375 380

Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu
 385 390 395 400

Gin lie Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro
 405 410 415

Ala Pro Ser Lys lie Val Gin Gin Thr Cys Lys Glu Phe Asn Leu Pro
 420 425 430

Cys Thr Leu Ser Pro Ser Met Met Gly Ala Val Thr Lys His Tyr His
 435 440 445

Gin Leu Lys Lys Met Gly Ala Glu Asn
 450 455

<210> 5

<211> 1374

<212> DNA

PhoenixTemp18528 .tmp .txt

<213> Acanthamoeba castellanii

<220>

<221> CDS

<222> (1)..(1374)

<223> Delta-8-Desaturase

<400> 5

atg gtc etc aca ace ccg gcc etc aac ctg aag aag gaa cga acg teg	48
Met Val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Glu Arg Thr Ser	
1 5 10 15	
ttc ace cag gag gag ctt tec aag etc tgg gtc ctt cac ggc cag gtg	96
Phe Thr Gin Glu Glu Leu Ser Lys Leu Trp Val Leu His Gly Gin Val	
20 25 30	
tac gat ttc ace gac ttt gtc aag tac cac ccg ggc ggc age agg gcc	144
Tyr Asp Phe Thr Asp Phe Val Lys Tyr His Pro Gly Gly Ser Arg Ala	
35 40 45	
ate ctg etc ggc cgt ggc cgt gat tgt ace gtg etc ttc gag tec tac	192
lie Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Glu Ser Tyr	
50 55 60	
cac aca gtc ctg cct tec gat get ctt etc gag aag tac cgc gtc tct	240
His Thr Val Leu Pro Ser Asp Ala Leu Leu Glu Lys Tyr Arg Val Ser	
65 70 75 80	
get ccc aac gcc aag etc gag gag age egg tea gcc aag ctg ttc teg	288
Ala Pro Asn Ala Lys Leu Glu Glu Ser Arg Ser Ala Lys Leu Phe Ser	
85 90 95	
ttc gag gag ggt age ttc tac cga ace etc aag cag cga acg cgc gag	336
Phe Glu Glu Gly Ser Phe Tyr Arg Thr Leu Lys Gin Arg Thr Arg Glu	
100 105 110	
tac ttc aag ace aac aac ctg age ace aag gcc ace acg atg gag gtc	384
Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Ala Thr Thr Met Glu Val	
115 120 125	
ate tac ttc gtg gcc ace ate etc age ate tac ttc tgc acg tgg gcc	432
lie Tyr Phe val Ala Thr lie Leu ser lie Tyr Phe cys Thr Trp Ala	
130 135 140	
gcc ttc gtg cag ggt tec etc ate gcc get gtc ctt cac gga gtg ggc	480
Ala Phe Val Gin Gly Ser Leu lie Ala Ala Val Leu His Gly Val Gly	
145 150 155 160	
cgt gcg ate tgt ate ata caa ccg act cat gcg act teg cac tac gcc	528
Arg Ala lie cys lie lie Gin Pro Thr His Ala Thr ser His Tyr Ala	
165 170 175	
atg ttc cgc tea gtg tgg etc aac cag tgg gcc tac agg ate tec atg	576
Met Phe Arg Ser Val Trp Leu Asn Gin Trp Ala Tyr Arg lie Ser Met	
180 185 190	
gcc gtc age gga teg teg ccg gcc cag tgg ace ace aag cac gtc ate	624
Ala Val Ser Gly Ser Ser Pro Ala Gin Trp Thr Thr Lys His Val lie	
195 200 205	
aac cat cac gtc gag ace aac ctg tgc ccc ace gat gac gac ace atg	672

PhoenixTemp18528.tmp.txt

Asn	His	His	Val	Glu	Thr	Asn	Leu	Cys	Pro	Thr	Asp	Asp	Asp	Thr	Met	
	210					215					220					
tac	ccc	ate	aag	cgc	ate	ctg	cac	gag	ttc	cct	cgt	ctg	ttc	ttc	cac	720
Tyr	Pro	lie	Lys	Arg	lie	Leu	His	Glu	Phe	Pro	Arg	Leu	Phe	Phe	His	
225					230					235					240	
aag	tac	cag	cac	ate	tac	ate	tgg	ctg	gtg	tac	ccc	tac	ace	ace	ate	768
Lys	Tyr	Gin	His	lie	Tyr	lie	Trp	Leu	Val	Tyr	Pro	Tyr	Thr	Thr	lie	
				245					250						255	
ttg	tgg	cac	ttc	tec	aac	ctg	gee	aag	etc	gee	etc	ggc	gee	get	cgc	816
Leu	Trp	His	Phe	Ser	Asn	Leu	Ala	Lys	Leu	Ala	Leu	Gly	Ala	Ala	Arg	
			260					265						270		
ggc	cag	atg	tac	gag	ggc	ate	gee	aag	gtg	age	caa	gag	ace	teg	ggc	864
Gly	Gin	Met	Tyr	Glu	Gly	lie	Ala	Lys	val	ser	Gin	Glu	Thr	ser	Gly	
		275					280						285			
gac	tgg	gtg	gag	acg	gee	atg	acg	ctg	ttc	ttc	ttc	acg	ttc	tec	cgt	912
Asp	Trp	Val	Glu	Thr	Ala	Met	Thr	Leu	Phe	Phe		Thr	Phe	Ser	Arg	
	290					295						300				
ctg	ctg	ctg	ccc	ttc	ctg	tgc	ctg	ccc	ttc	ace	acg	gee	gcc	gcg	gtg	960
Leu	Leu	Leu	Pro	Phe	Leu	cys	Leu	Pro	Phe	Thr	Thr	Ala	Ala	Ala	val	
305					310					315					320	
ttc	ctg	etc	tec	gag	tgg	ace	tgc	teg	ace	tgg	ttc	gcg	ctg	cag	ttc	1008
Phe	Leu	Leu	Ser	Glu	Trp	Thr	Cys	Ser	Thr	Trp	Phe	Ala	Leu	Gin	Phe	
				325					330						335	
gcc	gtg	age	cac	gag	gtc	gac	gag	tgc	gtc	gag	cac	gag	aag	teg	gtc	1056
Ala	val	ser	His	Glu	val	Asp	Glu	cys	val	Glu	His	Glu	Lys	ser	val	
			340					345						350		
etc	gac	ace	etc	aag	gcc	aac	gag	gcc	aag	ggc	ate	gtc	aac	cag	ggc	1104
Leu	Asp	Thr	Leu	Lys	Ala	Asn	Glu	Ala	Lys	Gly	lie	Val	Asn	Gin	Gly	
		355					360						365			
ggc	etc	gtc	gac	tgg	ggc	gcg	cac	cag	ggt	egg	gcc	teg	cac	aac	tac	1152
Gly	Leu	Val	Asp	Trp	Gly	Ala	His	Gin	Val	Arg	Ala	Ser	His	Asn	Tyr	
	370					375					380					
tct	gcc	gac	tec	ctg	ctg	teg	etc	cac	ttc	age	ggc	ggc	etc	aac	ctt	1200
Ser	Ala	Asp	Ser	Leu	Leu	Ser	Leu	His	Phe	Ser	Gly	Gly	Leu	Asn	Leu	
385					390					395					400	
cag	ate	gag	cac	cac	etc	ttc	ccc	tec	gtc	cac	tac	act	cac	tac	cct	1248
Gin	lie	Glu	His	His	Leu	Phe	Pro	Ser	Val	His	Tyr	Thr	His	Tyr	Pro	
				405					410						415	
gcc	ccg	tec	aag	att	gtg	cag	cag	acg	tgc	aag	gag	ttc	aac	ttg	ccc	1296
Ala	Pro	Ser	Lys	lie	Val	Gin	Gin	Thr	Cys	Lys	Glu	Phe	Asn	Leu	Pro	
			420					425						430		
tgc	act	ctg	teg	ccg	teg	atg	atg	ggc	gcc	gtg	ace	aag	cac	tac	cac	1344
Cys	Thr	Leu	Ser	Pro	Ser	Met	Met	Gly	Ala	Val	Thr	Lys	His	Tyr	His	
		435					440					445				
cag	etc	aag	aag	atg	ggc	get	gag	aac	tga							1374
Gin	Leu	Lys	Lys	Met	Gly	Ala	Glu	Asn								
	450					455										

<210> 6

<211> 457

<212> PRT

PhoenixTempl8528 .tmp .txt

<213> Acanthamoeba castellanii

<400> 6

Met val Leu Thr Thr Pro Ala Leu Asn Leu Lys Lys Gl u Arg Thr ser
 1 5 10 15
 Phe Thr Gl n Gl u Gl u Leu Ser Lys Leu Trp Val Leu Hi s Gl y Gl n Val
 20 25 30
 Tyr Asp Phe Thr Asp Phe Val Lys Tyr Hi s Pro Gly Gly Ser Arg Ala
 35 40 45
 I i e Leu Leu Gly Arg Gly Arg Asp Cys Thr Val Leu Phe Gl u Ser Tyr
 50 55 60
 Hi s Thr Val Leu Pro Ser Asp Ala Leu Leu Gl u Lys Tyr Arg Val Ser
 65 70 75 80
 Al a Pro Asn Ala Lys Leu Gl u Gl u Ser Arg Ser Ala Lys Leu Phe Ser
 85 90 95
 Phe Gl u Gl u Gly Ser Phe Tyr Arg Thr Leu Lys Gl n Arg Thr Arg Gl u
 100 105 110
 Tyr Phe Lys Thr Asn Asn Leu Ser Thr Lys Al a Thr Thr Met Gl u Val
 115 120 125
 I i e Tyr Phe Val Ala Thr lie Leu Ser lie Tyr Phe Cys Thr Trp Ala
 130 135 140
 Al a Phe Val Gl n Gly Ser Leu lie Ala Ala Val Leu Hi s Gly Val Gly
 145 150 155 160
 Arg Ala lie Cys lie I i e Gl n Pro Thr Hi s Al a Thr Ser Hi s Tyr Ala
 165 170 175
 Met Phe Arg Ser Val Trp Leu Asn Gl n Trp Al a Tyr Arg I i e Ser Met
 180 185 190
 Al a Val Ser Gly Ser Ser Pro Ala Gl n Trp Thr Thr Lys Hi s Val lie
 195 200 205
 Asn Hi s Hi s Val Gl u Thr Asn Leu Cys Pro Thr Asp Asp Asp Thr Met
 210 215 220
 Tyr Pro lie Lys Arg I i e Leu Hi s Gl u Phe Pro Arg Leu Phe Phe Hi s
 225 230 235 240
 Lys Tyr Gl n Hi s lie Tyr lie Trp Leu Val Tyr Pro Tyr Thr Thr lie
 245 250 255

PhoenixTtempl8528 .tmp .txt

Leu Trp His Phe Ser Asn Leu Ala Lys Leu Ala Leu Gly Ala Ala Arg
 260 265 270

Gly Gin Met Tyr Glu Gly lie Ala Lys Val Ser Gin Glu Thr Ser Gly
 275 280 285

Asp Trp Val Glu Thr Ala Met Thr Leu Phe Phe Thr Phe Ser Arg
 290 295 300

Leu Leu Leu Pro Phe Leu Cys Leu Pro Phe Thr Thr Ala Ala Ala Val
 305 310 315 320

Phe Leu Leu Ser Glu Trp Thr Cys Ser Thr Trp Phe Ala Leu Gin Phe
 325 330 335

Ala Val Ser His Glu Val Asp Glu Cys Val Glu His Glu Lys Ser Val
 340 345 350

Leu Asp Thr Leu Lys Ala Asn Glu Ala Lys Gly lie Val Asn Gin Gly
 355 360 365

Gly Leu Val Asp Trp Gly Ala His Gin Val Arg Ala Ser His Asn Tyr
 370 375 380

Ser Ala Asp Ser Leu Leu Ser Leu His Phe Ser Gly Gly Leu Asn Leu
 385 390 395 400

Gin lie Glu His His Leu Phe Pro Ser Val His Tyr Thr His Tyr Pro
 405 410 415

Ala Pro Ser Lys lie Val Gin Gin Thr Cys Lys Glu Phe Asn Leu Pro
 420 425 430

Cys Thr Leu Ser Pro Ser Met Met Gly Ala Val Thr Lys His Tyr His
 435 440 445

Gin Leu Lys Lys Met Gly Ala Glu Asn
 450 455

<210> 7

<211> 1236

<212> DNA

<213> Perkinsus marinus

<220>

<221> CDS

<222> (D..C1236)

PhoenixTemp18528 .tmp .txt

<223> Delta-8-Desaturase

<400> 7

atg	tct	tct	ctt	ace	etc	tac	aga	ggc	ccc	ttt	tec	cga	atg	gtg	etc	48
Met	ser	ser	Leu	Thr	Leu	Tyr	Arg	Gly	Pro	Phe	ser	Arg	Met	val	Leu	
1			5						10					15		
cct	cg	cag	gaa	ate	tgc	ate	gat	gg	cg	ata	tac	gat	gtc	act	gag	96
Pro	Arg	Gln	Glu	lie	Cys	lie	Asp	Gly	Arg	lie	Tyr	Asp	Val	Thr	Glu	
			20					25					30			
ttc	ate	aat	cg	cat	cca	gg	gg	aag	att	ate	etc	ttc	caa	g	g	144
Phe	lie	Asn	Arg	His	Pro	Gly	Gly	Lys	lie	lie	Leu	Phe	Gln	Val	Gly	
		35				40						45				
get	gat	gee	act	gat	get	ttt	cg	gag	ttt	cat	get	ggc	agt	gag	aag	192
Ala	Asp	Ala	Thr	Asp	Ala	Phe	Arg	Glu	Phe	His	Ala	Gly	Ser	Glu	Lys	
	50					55					60					
gca	gag	aag	ate	etc	aaa	ace	eta	cca	tec	cg	gat	gat	gac	ggt	act	240
Ala	Glu	Lys	lie	Leu	Lys	Thr	Leu	Pro	Ser	Arg	Asp	Asp	Asp	Gly	Thr	
65					70					75					80	
ttc	ctt	cct	tea	ace	caa	cg	tec	ate	atg	gat	gat	ttc	aaa	cg	eta	288
Phe	Leu	Pro	Ser	Thr	Gln	Arg	Ser	lie	Met	Asp	Asp	Phe	Lys	Arg	Leu	
				85					90					95		
aga	gat	gac	etc	gtc	age	aga	gg	gtc	ttc	aag	cca	age	gtc	atg	cat	336
Arg	Asp	Asp	Leu	Val	Ser	Arg	Gly	Val	Phe	Lys	Pro	Ser	Val	Met	His	
			100					105					110			
g	g	tac	cg	tgc	ttg	gaa	gtc	g	get	etc	tat	etc	att	gg	ttc	384
Val	Val	Tyr	Arg	Cys	Leu	Glu	Val	Val	Ala	Leu	Tyr	Leu	lie	Gly	Phe	
		115					120					125				
tat	ttg	get	ctg	tgc	ace	agt	aat	gtg	tac	gtt	ggg	tgt	get	gta	ctt	432
Tyr	Leu	Ala	Leu	Cys	Thr	Ser	Asn	Val	Tyr	Val	Gly	Cys	Ala	Val	Leu	
	130					135					140					
gg	g	get	caa	gg	cg	get	gg	tgg	ttg	atg	cat	gaa	gga	ggt	cat	480
Gly	Val	Ala	Gln	Gly	Arg	Ala	Gly	Trp	Leu	Met	His	Glu	Gly	Gly	His	
145					150				155						160	
cac	tct	ctg	act	gg	aac	tgg	aaa	g	gac	cag	ttc	etc	caa	gaa	eta	528
His	Ser	Leu	Thr	Gly	Asn	Trp	Lys	Val	Asp	Gln	Phe	Leu	Gln	Glu	Leu	
				165					170					175		
ttt	ttc	ggc	att	gg	tgt	gg	atg	tea	get	gcg	tgg	tgg	cg	aat	gca	576
Phe	Phe	Gly	lie	Gly	Cys	Gly	Met	Ser	Ala	Ala	Trp	Trp	Arg	Asn	Ala	
			180				185						190			
cac	aac	aag	cat	cac	get	get	cct	cag	cat	tta	ggg	aaa	gat	g	gat	624
His	Asn	Lys	His	His	Ala	Ala	Pro	Gln	His	Leu	Gly	Lys	Asp	Val	Asp	
		195					200					205				
etc	gag	aca	ttg	cct	ctg	gtc	gee	ttc	aat	aag	gee	gta	ctt	cga	ggc	672
Leu	Glu	Thr	Leu	Pro	Leu	Val	Ala	Phe	Asn	Lys	Ala	Val	Leu	Arg	Gly	
	210					215					220					
cg	eta	ccg	tct	gtc	tgg	ate	aga	tea	caa	get	gtg	tgc	ttt	gca	ccg	720
Arg	Leu	Pro	Ser	Val	Trp	lie	Arg	Ser	Gln	Ala	Val	Cys	Phe	Ala	Pro	
225					230					235					240	
ata	tea	aca	eta	ctg	gta	teg	ttc	ttt	tgg	caa	ttc	tac	eta	cac	ccg	768
lie	Ser	Thr	Leu	Leu	Val	Ser	Phe	Phe	Trp	Gln	Phe	Tyr	Leu	His	Pro	
				245					250					255		

PhoenixTempl8528 .tmp .txt

Ala Glu Lys lie Leu Lys Thr Leu Pro Ser Arg Asp Asp Asp Gly Thr
 65 70 75 80

Phe Leu Pro Ser Thr Gin Arg Ser lie Met Asp Asp Phe Lys Arg Leu
 85 90 95

Arg Asp Asp Leu Val Ser Arg Gly Val Phe Lys Pro Ser Val Met His
 100 105 110

Val Val Tyr Arg Cys Leu Glu Val Val Ala Leu Tyr Leu lie Gly Phe
 115 120 125

Tyr Leu Ala Leu Cys Thr Ser Asn Val Tyr Val Gly Cys Ala Val Leu
 130 135 140

Gly Val Ala Gin Gly Arg Ala Gly Trp Leu Met His Glu Gly Gly His
 145 150 155 160

His Ser Leu Thr Gly Asn Trp Lys Val Asp Gin Phe Leu Gin Glu Leu
 165 170 175

Phe Phe Gly lie Gly Cys Gly Met Ser Ala Ala Trp Trp Arg Asn Ala
 180 185 190

His Asn Lys His His Ala Ala Pro Gin His Leu Gly Lys Asp Val Asp
 195 200 205

Leu Glu Thr Leu Pro Leu Val Ala Phe Asn Lys Ala Val Leu Arg Gly
 210 215 220

Arg Leu Pro ser val Trp lie Arg ser Gin Ala val cys Phe Ala Pro
 225 230 235 240

lie Ser Thr Leu Leu Val Ser Phe Phe Trp Gin Phe Tyr Leu His Pro
 245 250 255

Arg His lie lie Arg Thr Gly Arg Arg Met Glu ser Phe Trp Leu Leu
 260 265 270

Val Arg Tyr Leu Val lie Val Tyr Leu Gly Phe Ser Tyr Gly Leu Val
 275 280 285

Ser val Leu Leu cys Tyr lie Ala ser val His val Gly Gly Met Tyr
 290 295 300

lie Phe Val His Phe Ala Leu Ser His Thr His Leu Pro Val lie Asn
 305 310 315 320

Gin His Gly Arg Ala Asn Trp Leu Glu Tyr Ala Ser Lys His Thr Val
 325 330 335

PhoenixTempl8528.tmp.txt

Asn Val Ser Thr Asn Asn Tyr Phe Val Thr Trp Leu Met Ser Tyr Leu
 340 345 350

Asn Tyr Gin lie Glu His His Leu Phe Pro Ser Cys Pro Gin Phe Arg
 355 360 365

Phe Pro Gly Tyr Val Ser Met Arg Val Arg Glu Phe Phe His Lys His
 370 375 380

Gly Leu Lys Tyr Asn Glu Val Gly Tyr Leu His Ala Leu Asn Leu Thr
 385 390 395 400

Phe ser Asn Leu Ala Ala val Ala lie val Glu
 405 410

<210> 9

<211> 777

<212> DNA

<213> Isochrysis galbana

<220>

<221> CDS

<222> (I).. (777)

<223> Delta-9- Elongase

<400> 9

atg gcc etc gca aac gac gcg gga gag cgc ate tgg gcg get gtg ace 48
 Met Ala Leu Ala Asn Asp Ala Gly Glu Arg lie Trp Ala Ala Val Thr
 1 5 10 15

gac ccg gaa ate etc att ggc ace ttc teg tac ttg eta etc aaa ccg 96
 Asp Pro Glu lie Leu lie Gly Thr Phe Ser Tyr Leu Leu Leu Lys Pro
 20 25 30

ctg etc cgc aat tec ggg ctg gtg gat gag aag aag ggc gca tac agg 144
 Leu Leu Arg Asn Ser Gly Leu Val Asp Glu Lys Lys Gly Ala Tyr Arg
 35 40 45

acg tec atg ate tgg tac aac gtt ctg ctg gcg etc ttc tct gcg ctg 192
 Thr Ser Met lie Trp Tyr Asn Val Leu Leu Ala Leu Phe Ser Ala Leu
 50 55 60

age ttc tac gtg acg gcg ace gcc etc ggc tgg gac tat ggt acg gcc 240
 Ser Phe Tyr Val Thr Ala Thr Ala Leu Gly Trp Asp Tyr Gly Thr Gly
 65 70 75 80

gcg tgg ctg cgc agg caa ace ggc gac aca ccg cag ccg etc ttc cag 288
 Ala Trp Leu Arg Arg Gin Thr Gly Asp Thr Pro Gin Pro Leu Phe Gin
 85 90 95

tgc ccg tec ccg gtt tgg gac teg aag etc ttc aca tgg ace gcc aag 336
 Cys Pro ser Pro val Trp Asp ser Lys Leu Phe Thr Trp Thr Ala Lys
 100 105 110

PhoenixTemp18528 .tmp .txt

gca ttc tat tac tec aag tac	gtg gag tac etc gac acg gcc tgg ctg	384
Ala Phe Tyr Tyr Ser Lys Tyr	VaI Glu Tyr Leu Asp Thr Ala Trp Leu	
115	120 125	
agg gtc tec ttt etc cag gcc ttc cac cac ttt ggc gcg ccg tgg gat	432	
Arg VaI Ser Phe Leu Gin Ala Phe His His Phe	Gly Ala Pro Trp Asp	
130	135 140	
gtg tac etc ggc att egg ctg cac aac gag ggc gta tgg ate ttc atg	480	
VaI Tyr Leu Gly H e Arg Leu His Asn Glu Gly VaI Trp H e Phe Met		
145	150 155 160	
ttt ttc aac teg ttc att cac ace ate atg tac ace tac tac ggc etc	528	
Phe Phe Asn Ser Phe lie His Thr lie Met Tyr Thr Tyr Tyr Gly Leu		
165	170 175	
ace gcc gcc ggg tat aag ttc aag gcc aag ccg etc ate ace gcg atg	576	
Thr Ala Ala Gly Tyr Lys Phe Lys Ala Lys Pro Leu lie Thr Ala Met		
180	185 190	
cag ate tgc cag ttc gtg ggc ggc ttc ctg ttg gtc tgg gac tac ate	624	
Gin lie Cys Gin Phe VaI Gly Gly Phe Leu Leu VaI Trp Asp Tyr lie		
195	200 205	
aac gtc ccc tgc ttc aac teg gac aaa ggg aag ttg ttc age tgg get	672	
Asn VaI Pro Cys Phe Asn Ser Asp Lys Gly Lys Leu Phe Ser Trp Ala		
210	215 220	
ttc aac tat gca tac gtc ggc teg gtc ttc ttg etc ttc tgc cac ttt	720	
Phe Asn Tyr Ala Tyr VaI Gly Ser VaI Phe Leu Leu Phe Cys His Phe		
225	230 235 240	
ttc tac cag gac aac ttg gca acg aag aaa teg gcc aag gcg ggc aag	768	
Phe Tyr Gin Asp Asn Leu Ala Thr Lys Lys Ser Ala Lys Ala Gly Lys		
245	250 255	
cag etc tag	777	
Gin Leu		

<210> 10

<211> 258

<212> PRT

<213> Isochrysis galbana

<400> 10

Met	Ala	Leu	Ala	Asn	Asp	Ala	Gly	Glu	Arg	lie	Trp	Ala	Ala	Val	Thr
1				5					10					15	
Asp	Pro	Glu	lie	Leu	lie	Gly	Thr	Phe	ser	Tyr	Leu	Leu	Leu	Lys	Pro
			20					25					30		
Leu	Leu	Arg	Asn	Ser	Gly	Leu	Val	Asp	Glu	Lys	Lys	Gly	Ala	Tyr	Arg
		35					40					45			
Thr	Ser	Met	lie	Trp	Tyr	Asn	Val	Leu	Leu	Ala	Leu	Phe	Ser	Ala	Leu
	50					55					60				

PhoenixTemp18528 .tmp .txt

Ser Phe Tyr Val Thr Ala Thr Ala Leu Gly Trp Asp Tyr Gly Thr Gly
65 70 75 80

Ala Trp Leu Arg Arg Gin Thr Gly Asp Thr Pro Gin Pro Leu Phe Gin
85 90 95

Cys Pro Ser Pro Val Trp Asp Ser Lys Leu Phe Thr Trp Thr Ala Lys
100 105 110

Ala Phe Tyr Tyr Ser Lys Tyr Val Glu Tyr Leu Asp Thr Ala Trp Leu
115 120 125

Arg val ser Phe Leu Gin Ala Phe His His Phe Gly Ala Pro Trp Asp
130 135 140

Val Tyr Leu Gly lie Arg Leu His Asn Glu Gly Val Trp lie Phe Met
145 150 155 160

Phe Phe Asn ser Phe lie His Thr lie Met Tyr Thr Tyr Tyr Gly Leu
165 170 175

Thr Ala Ala Gly Tyr Lys Phe Lys Ala Lys Pro Leu lie Thr Ala Met
180 185 190

Gin lie cys Gin Phe val Gly Gly Phe Leu Leu val Trp Asp Tyr lie
195 200 205

Asn Val Pro Cys Phe Asn Ser Asp Lys Gly Lys Leu Phe Ser Trp Ala
210 215 220

Phe Asn Tyr Ala Tyr Val Gly Ser Val Phe Leu Leu Phe Cys His Phe
225 230 235 240

Phe Tyr Gin Asp Asn Leu Ala Thr Lys Lys Ser Ala Lys Ala Gly Lys
245 250 255

Gin Leu

<210> 11

<211> 891

<212> DNA

<213> Acanthamoeba castellanii

<220>

<221> CDS

<222> (1)..(891)

<223> Delta-9-Elongase

PhoenixTemp18528 .tmp .txt

```

<400> 11
atg gcg get gcg acg gcg acg acg gca acg acg gcg gtg atg gag caa 48
Met Ala Ala Ala Thr Ala Thr Thr Ala Thr Thr Ala Val Met Glu Gin
1 5 10 15

gtg ccc att acg gag gcc ate ttc egg ccg gac etc tgg gtc gga egg 96
Val Pro lie Thr Glu Ala lie Phe Arg Pro Asp Leu Trp Val Gly Arg
20 25 30

gac cag tgg gag gcg aat gcc gtg age ttc gta tgg agg tac tgg tgg 144
Asp Gin Trp Glu Ala Asn Ala Val Ser Phe Val Trp Arg Tyr Trp Trp
35 40 45

ttc ttc ctg gtg atg gcc gtg gca tac ctg ccc ate ate ttc gcc etc 192
Phe Phe Leu Val Met Gly Val Ala Tyr Leu Pro lie lie Phe Gly Leu
50 55 60

aag tac tgg atg aag gat cgt ccg gcc ttc aac etc cgt egg ccg etc 240
Lys Tyr Trp Met Lys Asp Arg Pro Ala Phe Asn Leu Arg Arg Pro Leu
65 70 75 80

ate ttg tgg aat ate ttc atg gcg acg ttc teg ace gcc ggc ttc ctg 288
lie Leu Trp Asn lie Phe Met Ala Thr Phe Ser Thr Ala Gly Phe Leu
85 90 95

teg ate gtc tac ccc etc ate gag aac tgg gtc tac ccc ggc ggc ggc 336
Ser lie Val Tyr Pro Leu lie Glu Asn Trp Val Tyr Pro Gly Gly Gly
100 105 110

etc ace ccg cat gag ttc ate tgc teg gcc age tac tec tac aag ttt 384
Leu Thr Pro His Glu Phe lie Cys Ser Ala Ser Tyr Ser Tyr Lys Phe
115 120 125

ggt gat tgc gcc ate tgg gtg ttc etc ttc aac atg teg aag ate etc 432
Gly Asp Cys Ala lie Trp Val Phe Leu Phe Asn Met Ser Lys lie Leu
130 135 140

gag ttc gtc gac ace ate ttc ate gtc ccc agg aag ace cac etc ggc 480
Glu Phe val Asp Thr lie Phe lie val Pro Arg Lys Thr His Leu Gly
145 150 155 160

ttc etc cac tac tac cac cac ate ate ace tac tec ttc tgc etc tac 528
Phe Leu His Tyr Tyr His His lie lie Thr Tyr Ser Phe Cys Leu Tyr
165 170 175

gcc ggc cag tac atg cac cac tac aac tgt ggc ggc tat ttc ttc tgc 576
Ala Gly Gin Tyr Met His His Tyr Asn cys Gly Gly Tyr Phe Phe cys
180 185 190

etc atg aac ttc ttc gtc cac ggc ate atg tac ttc tac tac get etc 624
Leu Met Asn Phe Phe Val His Gly lie Met Tyr Phe Tyr Tyr Ala Leu
195 200 205

cgc tec atg ggc ttc cgt ccc tec ttc gat att ggc ate ace ttc etc 672
Arg ser Met Gly Phe Arg Pro ser Phe Asp lie Gly lie Thr Phe Leu
210 215 220

cag att ttg caa atg gtg etc ggc gtg gcc ate ate ace ate tec gcc 720
Gin lie Leu Gin Met Val Leu Gly Val Ala lie lie Thr lie Ser Ala
225 230 235 240

ggc tgc gag aag gtg gac ccc ate gga acg ace ttc ggc tac ttt att 768
Gly Cys Glu Lys Val Asp Pro lie Gly Thr Thr Phe Gly Tyr Phe lie
245 250 255

tat ttc teg ttc ttc gtc etc ttc tgc aag ttc ttc tac tac cgc tac 816

```

PhoenixTempl8528.tmp.txt

Tyr Phe Ser Phe Phe Val Leu Phe Cys Lys Phe Phe Tyr Tyr Arg Tyr
 260 265 270

ate gcc acg ccc gcc aag aag ccc gag gcc gcc gcc aag teg cca gcc 864
 lie Ala Thr Pro Ala Lys Lys Pro Glu Ala Ala Ala Lys Ser Pro Ala
 275 280 285

ace aag ccc aag agg aag cac gac taa 891
 Thr Lys Pro Lys Arg Lys His Asp
 290 295

<210> 12
 <211> 296
 <212> PRT
 <213> Acanthamoeba castellanii

<400> 12

Met Ala Ala Ala Thr Ala Thr Thr Ala Thr Thr Ala val Met Glu Gln
 1 5 10 15

Val Pro lie Thr Glu Ala lie Phe Arg Pro Asp Leu Trp Val Gly Arg
 20 25 30

Asp Gln Trp Glu Ala Asn Ala val ser Phe val Trp Arg Tyr Trp Trp
 35 40 45

Phe Phe Leu Val Met Gly Val Ala Tyr Leu Pro lie lie Phe Gly Leu
 50 55 60

Lys Tyr Trp Met Lys Asp Arg Pro Ala Phe Asn Leu Arg Arg Pro Leu
 65 70 75 80

lie Leu Trp Asn lie Phe Met Ala Thr Phe Ser Thr Ala Gly Phe Leu
 85 90 95

Ser lie Val Tyr Pro Leu lie Glu Asn Trp Val Tyr Pro Gly Gly Gly
 100 105 110

Leu Thr Pro His Glu Phe lie Cys Ser Ala Ser Tyr Ser Tyr Lys Phe
 115 120 125

Gly Asp Cys Ala lie Trp Val Phe Leu Phe Asn Met Ser Lys lie Leu
 130 135 140

Gl u Phe Val Asp Thr lie Phe lie Val Pro Arg Lys Thr His Leu Gly
 145 150 155 160

Phe Leu His Tyr Tyr His His lie lie Thr Tyr Ser Phe Cys Leu Tyr
 165 170 175

Ala Gly Gln Tyr Met His His Tyr Asn Cys Gly Gly Tyr Phe Phe Cys

Phoenixr empl8528.tmp.txt

180

185

190

Leu Met Asn Phe Phe Val His Gly lie Met Tyr Phe Tyr Tyr Ala Leu
 195 200 205

Arg ser Met Gly Phe Arg Pro ser Phe Asp lie Gly lie Thr Phe Leu
 210 215 220

Gln lie Leu Gln Met Val Leu Gly Val Ala lie lie Thr lie Ser Ala
 225 230 235 240

Gly Cys Glu Lys Val Asp Pro lie Gly Thr Thr Phe Gly Tyr Phe lie
 245 250 255

Tyr Phe Ser Phe Phe Val Leu Phe Cys Lys Phe Phe Tyr Tyr Arg Tyr
 260 265 270

lie Ala Thr Pro Ala Lys Lys Pro Glu Ala Ala Ala Lys Ser Pro Ala
 275 280 285

Thr Lys Pro Lys Arg Lys His Asp
 290 295

<210> 13

<211> 1320

<212> DNA

<213> Thraust rochyt ri urn

<220>

<221> cds

<222> (l) ..(1320)

<223> Del ta- 5- Desatu rase

<400> 13

atg ggc aag ggc age gag ggc cgc age gcg gcg cgc gag atg acg gcc 48
 Met Gly Lys Gly Ser Glu Gly Arg Ser Ala Ala Arg Glu Met Thr Ala
 1 5 10 15

gag gcg aac ggc gac aag egg aaa acg att ctg ate gag ggc gtc ctg 96
 Glu Ala Asn Gly Asp Lys Arg Lys Thr lie Leu lie Glu Gly val Leu
 20 25 30

tac gac gcg acg aac ttt aag cac ccg ggc ggt teg ate ate aac ttc 144
 Tyr Asp Ala Thr Asn Phe Lys His Pro Gly Gly Ser lie lie Asn Phe
 35 40 45

ttg ace gag ggc gag gcc ggc gtg gac gcg acg cag gcg tac cgc gag 192
 Leu Thr Glu Gly Glu Ala Gly Val Asp Ala Thr Gln Ala Tyr Arg Glu
 50 55 60

ttt cat cag egg tec ggc aag gcc gac aag tac etc aag teg ctg ccg 240

PhoenixTemp18528 .tmp .txt

Phe 65	His	Gin	Arg	Ser	Gly 70	Lys	Ala	Asp	Lys	Tyr 75	Leu	Lys	Ser	Leu	Pro 80	
aag	ctg	gat	gcg	tec	aag	gtg	gag	teg	egg	ttc	teg	gee	aaa	gag	cag	288
Lys	Leu	Asp	Ala	Ser 85	Lys	Val	Glu	Ser	Arg 90	Phe	Ser	Ala	Lys	Glu 95	Gin	
gcg	egg	cgc	gac	gcc	atg	acg	cgc	gac	tac	gcg	gcc	ttt	cgc	gag	gag	336
Ala	Arg	Arg	Asp 100	Ala	Met	Thr	Arg	Asp 105	Tyr	Ala	Ala	Phe	Arg 110	Glu	Glu	
etc	gtc	gcc	gag	ggg	tac	ttt	gac	ccg	teg	ate	ccg	cac	atg	att	tac	384
Leu	Val	Ala 115	Glu	Gly	Tyr	Phe	Asp 120	Pro	Ser	lie	Pro	His 125	Met	lie	Tyr	
cgc	gtc	gtg	gag	ate	gtg	gcg	etc	ttc	gcg	etc	teg	ttc	tgg	etc	atg	432
Arg	val 130	val	Glu	lie	val	Ala 135	Leu	Phe	Ala	Leu	ser 140	Phe	Trp	Leu	Met	
tec	aag	gcc	teg	ccc	ace	teg	etc	gtg	ctg	ggc	gtg	gtg	atg	aac	ggc	480
Ser 145	Lys	Ala	Ser	Pro 150	Thr	Ser	Leu	Val	Leu	Gly 155	Val	Val	Met	Asn	Gly 160	
att	gcg	cag	ggc	cgc	tgc	ggc	tgg	gtc	atg	cac	gag	atg	ggc	cac	ggg	528
lie	Ala	Gin	Gly 165	Arg	cys	Gly	Trp	val	Met 170	His	Glu	Met	Gly 175	His	Gly	
teg	ttc	acg	ggc	gtc	ate	tgg	etc	gac	gac	egg	atg	tgc	gag	ttc	ttc	576
Ser	Phe	Thr 180	Gly	Val	lie	Trp	Leu	Asp 185	Asp	Arg	Met	Cys	Glu 190	Phe	Phe	
tac	ggc	gtc	ggc	tgc	ggc	atg	age	ggg	cac	tac	tgg	aag	aac	cag	cac	624
Tyr	Gly 195	val	Gly	Cys	Gly	Met	ser 200	Gly	His	Tyr	Trp	Lys 205	Asn	Gin	His	
age	aag	cac	cac	gcc	gcg	ccc	aac	cgc	etc	gag	cac	gat	gtc	gat	etc	672
Ser 210	Lys	His	His	Ala	Ala	Pro 215	Asn	Arg	Leu	Glu	His 220	Asp	Val	Asp	Leu	
aac	acg	ctg	ccc	ctg	gtc	gcc	ttt	aac	gag	cgc	gtc	gtg	cgc	aag	gtc	720
Asn 225	Thr	Leu	Pro	Leu	Val 230	Ala	Phe	Asn	Glu	Arg 235	Val	Val	Arg	Lys	Val 240	
aag	ccg	gga	teg	ctg	ctg	gcg	etc	tgg	ctg	cgc	gtg	cag	gcg	tac	etc	768
Lys	Pro	Gly	Ser 245	Leu	Leu	Ala	Leu	Trp	Leu	Arg 250	Val	Gin	Ala	Tyr 255	Leu	
ttt	gcg	ccc	gtc	teg	tgc	ctg	etc	ate	ggc	ctt	ggc	tgg	acg	etc	tac	816
Phe	Ala	Pro 260	Val	Ser	Cys	Leu	Leu	lie 265	Gly	Leu	Gly	Trp	Thr 270	Leu	Tyr	
ctg	cac	ccg	cgc	tac	atg	ctg	cgc	ace	aag	egg	cac	atg	gag	ttc	gtc	864
Leu	His 275	Pro	Arg	Tyr	Met	Leu	Arg 280	Thr	Lys	Arg	His	Met 285	Glu	Phe	Val	
tgg	ate	ttc	gcg	cgc	tac	att	ggc	tgg	ttc	teg	etc	atg	ggc	get	etc	912
Trp	lie 290	Phe	Ala	Arg	Tyr	lie	Gly 295	Trp	Phe	Ser	Leu	Met 300	Gly	Ala	Leu	
ggc	tac	teg	ccg	ggc	ace	teg	gtc	ggg	atg	tac	ctg	tgc	teg	ttc	ggc	960
Gly 305	Tyr	Ser	Pro	Gly	Thr 310	Ser	Val	Gly	Met	Tyr 315	Leu	Cys	Ser	Phe	Gly 320	
etc	ggc	tgc	att	tac	att	ttc	ctg	cag	ttc	gcc	gtc	age	cac	acg	cac	1008
Leu	Gly	Cys 325	lie	Tyr	lie	Phe	Leu	Gin	Phe 330	Ala	Val	Ser	His 335	Thr	His	
ctg	ccg	gtg	ace	aac	ccg	gag	gac	cag	ctg	cac	tgg	etc	gag	tac	gcg	1056
Leu	Pro	Val	Thr	Asn	Pro	Glu	Asp	Gin	Leu	His	Trp	Leu	Glu	Tyr	Ala	

PhoenixTemp18528.tmp.txt

340	345	350	
gcc gac cac acg gtg aac att age ace aag tec tgg etc gtc acg tgg 1104	Ala Asp His Thr Val Asn lie Ser Thr Lys Ser Trp Leu Val Thr Trp	355	360
tgg atg teg aac ctg aac ttt cag ate gag cac cac etc ttc ccc acg 1152	Trp Met ser Asn Leu Asn Phe Gin lie Glu His His Leu Phe Pro Thr	370	375
gcg ccg cag ttc cgc ttc aag gaa ate agt cct cgc gtc gag gcc etc 1200	Ala Pro Gin Phe Arg Phe Lys Glu lie Ser Pro Arg Val Glu Ala Leu	385	390
ttc aag cgc cac aac etc ccg tac tac gac ctg ccc tac acg age gcg 1248	Phe Lys Arg His Asn Leu Pro Tyr Tyr Asp Leu Pro Tyr Thr Ser Ala	405	410
gtc teg ace ace ttt gcc aat ctt tat tec gtc ggc cac teg gtc ggc 1296	Val Ser Thr Thr Phe Ala Asn Leu Tyr Ser Val Gly His Ser Val Gly	420	425
gcc gac ace aag aag cag gac tga 1320	Ala Asp Thr Lys Lys Gin Asp	435	

<210> 14

<211> 439

<212> PRT

<213> Thraustrochytrium

<400> 14

Met Gly Lys Gly Ser Glu Gly Arg Ser Ala Ala Arg Glu Met Thr Ala	1	5	10	15
Glu Ala Asn Gly Asp Lys Arg Lys Thr lie Leu lie Glu Gly Val Leu	20	25	30	
Tyr Asp Ala Thr Asn Phe Lys His Pro Gly Gly Ser lie lie Asn Phe	35	40	45	
Leu Thr Glu Gly Glu Ala Gly Val Asp Ala Thr Gin Ala Tyr Arg Glu	50	55	60	
Phe His Gin Arg Ser Gly Lys Ala Asp Lys Tyr Leu Lys Ser Leu Pro	65	70	75	80
Lys Leu Asp Ala Ser Lys Val Glu Ser Arg Phe Ser Ala Lys Glu Gin	85	90	95	
Ala Arg Arg Asp Ala Met Thr Arg Asp Tyr Ala Ala Phe Arg Glu Glu	100	105	110	
Leu Val Ala Glu Gly Tyr Phe Asp Pro Ser lie Pro His Met lie Tyr	115	120	125	

PhoenixTtempl8528 .tmp .txt

Arg Val Val Glu lie Val Ala Leu Phe Ala Leu Ser Phe Trp Leu Met
 130 135 140

Ser Lys Ala Ser Pro Thr Ser Leu Val Leu Gly Val Val Met Asn Gly
 145 150 155 160

lie Ala Gin Gly Arg Cys Gly Trp Val Met His Glu Met Gly His Gly
 165 170 175

Ser Phe Thr Gly Val lie Trp Leu Asp Asp Arg Met Cys Glu Phe Phe
 180 185 190

Tyr Gly Val Gly Cys Gly Met Ser Gly His Tyr Trp Lys Asn Gin His
 195 200 205

Ser Lys His His Ala Ala Pro Asn Arg Leu Glu His Asp Val Asp Leu
 210 215 220

Asn Thr Leu Pro Leu Val Ala Phe Asn Glu Arg Val Val Arg Lys Val
 225 230 235 240

Lys Pro Gly Ser Leu Leu Ala Leu Trp Leu Arg Val Gin Ala Tyr Leu
 245 250 255

Phe Ala Pro Val Ser Cys Leu Leu lie Gly Leu Gly Trp Thr Leu Tyr
 260 265 270

Leu His Pro Arg Tyr Met Leu Arg Thr Lys Arg His Met Glu Phe Val
 275 280 285

Trp lie Phe Ala Arg Tyr lie Gly Trp Phe Ser Leu Met Gly Ala Leu
 290 295 300

Gly Tyr Ser Pro Gly Thr Ser Val Gly Met Tyr Leu Cys Ser Phe Gly
 305 310 315 320

Leu Gly Cys lie Tyr lie Phe Leu Gin Phe Ala Val Ser His Thr His
 325 330 335

Leu Pro Val Thr Asn Pro Glu Asp Gin Leu His Trp Leu Glu Tyr Ala
 340 345 350

Ala Asp His Thr Val Asn lie Ser Thr Lys Ser Trp Leu Val Thr Trp
 355 360 365

Trp Met Ser Asn Leu Asn Phe Gin lie Glu His His Leu Phe Pro Thr
 370 375 380

Ala Pro Gin Phe Arg Phe Lys Glu lie ser Pro Arg val Glu Ala Leu
 385 390 395 400

Phoenixr empl8528.tmp.txt

Phe Lys Arg His Asn Leu Pro Tyr Tyr Asp Leu Pro Tyr Thr Ser Ala
 405 410 415

Val Ser Thr Thr Phe Ala Asn Leu Tyr Ser Val Gly His Ser Val Gly
 420 425 430

Ala Asp Thr Lys Lys Gin Asp
 435

<210> 15

<211> 1353

<212> DNA

<213> Acanthamoeba casteliani

<220>

<221> cds

<222> (1) .. (1353)

<223> Delta-5-Desaturase

<400> 15

atg gcc ace gca tct gca tec aac gtt etc cgc ctg ccc gga gag gga 48
 Met Ala Thr Ala Ser Ala Ser Asn Val Leu Arg Leu Pro Gly Glu Gly
 1 5 10 15

etc gcg act ggc etc gag cag etc gag tgg gcc gaa gtg cag aag cac 96
 Leu Ala Thr Gly Leu Glu Gin Leu Glu Trp Ala Glu Val Gin Lys His
 20 25 30

aac acg cgc gag age teg tgg ctg gtg att aac gac cag gtg tac gac 144
 Asn Thr Arg Glu Ser Ser Trp Leu Val lie Asn Asp Gin Val Tyr Asp
 35 40 45

ate ace aac ttc ggc egg cgc cat ccc ggt ggc aag gta ate tac cac 192
 lie Thr Asn Phe Gly Arg Arg His Pro Gly Gly Lys Val lie Tyr His
 50 55 60

tac gcg ggt caa gat gcc acg gac teg ttt egg get ctt cac ccc gat 240
 Tyr Ala Gly Gin Asp Ala Thr Asp Ser Phe Arg Ala Leu His Pro Asp
 65 70 75 80

tec gcc ctg gtg atg aag tat etc aag ccc etc etc ate ggt caa gtg 288
 Ser Ala Leu Val Met Lys Tyr Leu Lys Pro Leu Leu lie Gly Gin Val
 85 90 95

gca ccc ggc tea tec ace gca gca teg att gtt gat ggc gcc cgc ccg 336
 Ala Pro Gly Ser Ser Thr Ala Ala Ser lie Val Asp Gly Ala Arg Pro
 100 105 110

gcg ccc teg gca ttc gta gag gaa ttc aga cag gtg cgc aaa gaa ttc 384
 Ala Pro Ser Ala Phe Val Glu Glu Phe Arg Gin Val Arg Lys Glu Phe
 115 120 125

gag gag cag ggc ctg ttc gag gcc age tgg tec ttc ttc ttc ggg atg 432
 Glu Glu Gin Gly Leu Phe Glu Ala Ser Trp Ser Phe Phe Phe Gly Met
 130 135 140

PhoenixTemp18528 .tmp .txt

ctg gcc cac ate ttc ctg etc gag get gcc gcc tac tac age ate aag	480
Leu Ala His lie Phe Leu Leu Glu Ala Ala Ala Tyr Tyr Ser lie Lys	
145 150 155 160	
ctg ctg ggc aac agt tgg ccc gtc tac etc etc gcc gtc ggc etc etc	528
Leu Leu Gly Asn Ser Trp Pro Val Tyr Leu Leu Ala Val Gly Leu Leu	
165 170 175	
gcc act gcc cag gca cag gcc ggc tgg etc cag cac gat tgt ggg cac	576
Ala Thr Ala Gin Ala Gin Ala Gly Trp Trp Leu Gin His Asp Cys Gly His	
180 185 190	
ttg tec gtg ttc aag aag teg aag tgg aac cat tgg atg cac tac ate	624
Leu Ser Val Phe Lys Lys Ser Lys Trp Asn His Trp Met His Tyr lie	
195 200 205	
gtc ate tgc cac ate aag ggc gcc teg cga gcc tgg tgg aac tgg cgt	672
Val lie Cys His lie Lys Gly Ala Ser Arg Ala Trp Trp Asn Trp Arg	
210 215 220	
cac ttt gag cac cac gca aag ccc aac gtg gtg cgc aag gac ccc gac	720
His Phe Glu His His Ala Lys Pro Asn Val Val Arg Lys Asp Pro Asp	
225 230 235 240	
ate ace ttc ccc aac etc ttc ctt etc ggc gac cac ctg acg cgc aag	768
lie Thr Phe Pro Asn Leu Phe Leu Leu Gly Asp His Leu Thr Arg Lys	
245 250 255	
tgg gcc aag gcc aag aag gga gtg atg ccc tac aac aag cag cac etc	816
Trp Ala Lys Ala Lys Lys Gly Val Met Pro Tyr Asn Lys Gin His Leu	
260 265 270	
tac tgg tgg get ttc ccc ccg etc ctg ctg ccc gtc tac ttc cac tac	864
Tyr Trp Trp Ala Phe Pro Pro Leu Leu Leu Pro Val Tyr Phe His Tyr	
275 280 285	
gac aac att cga tac gtc ttc cag cac aag cac tgg tgg gac etc ttc	912
Asp Asn lie Arg Tyr Val Phe Gin His Lys His Trp Trp Asp Leu Phe	
290 295 300	
tgg ate gcc acg ttc ttc gcg aag cac ttc acg etc tac ggc ccg ctg	960
Trp lie Ala Thr Phe Phe Ala Lys His Phe Thr Leu Tyr Gly Pro Leu	
305 310 315 320	
atg ggc ggc tgg ggc gcg ttc tgg ttc tac atg ctg gtg cgc acg gtc	1008
Met Gly Gly Trp Gly Ala Phe Trp Phe Tyr Met Leu Val Arg Thr Val	
325 330 335	
gag age cac tgg ttc aca tgg gtg ace cag atg aac cac ate ccc atg	1056
Glu Ser His Trp Phe Thr Trp Val Thr Gin Met Asn His H e Pro Met	
340 345 350	
cac gtc gac aac gac cgc gag ctg gac tgg ccc ace ctg cag ggt etc	1104
His Val Asp Asn Asp Arg Glu Leu Asp Trp Pro Thr Leu Gin Gly Leu	
355 360 365	
gcc acg tgc aac gtc gag ggc age etc ttc aac gac tgg ttc acg ggc	1152
Ala Thr Cys Asn Val Glu Gly Ser Leu Phe Asn Asp Trp Phe Thr Gly	
370 375 380	
cac etc aac tac cag ate gag cac cac etc ttc ccc ace atg ccc cgc	1200
His Leu Asn Tyr Gin lie Glu His His Leu Phe Pro Thr Met Pro Arg	
385 390 395 400	
cac aac tac gcg gtg gcc aac aag aag gtc cag gcc etc tac aag aag	1248
His Asn Tyr Ala val Ala Asn Lys Lys val Gin Ala Leu Tyr Lys Lys	
405 410 415	

PhoenixTemp18528 .tmp .txt

cac ggc gtg ccg atg cag ace aag ggc etc ate gaa gcc ttc gcc gac 1296
 His Gly Val Pro Met Gin Thr Lys Gly Leu lie Glu Ala Phe Ala Asp
 420 425 430

ate gtc aag teg etc gag cac tat ggt gag gtg tgg aag gag gcc tac 1344
 lie Val Lys Ser Leu Glu His Tyr Gly Glu Val Trp Lys Glu Ala Tyr
 435 440 445

tac ggc taa 1353
 Tyr Gly
 450

<210> 16

<211> 450

<212> PRT

<213> Acanthamoeba castellanii

<400> 16

Met Ala Thr Ala Ser Ala Ser Asn Val Leu Arg Leu Pro Gly Glu Gly
 1 5 10 15

Leu Ala Thr Gly Leu Glu Gin Leu Glu Trp Ala Glu Val Gin Lys His
 20 25 30

Asn Thr Arg Glu Ser Ser Trp Leu Val lie Asn Asp Gin Val Tyr Asp
 35 40 45

lie Thr Asn Phe Gly Arg Arg His Pro Gly Gly Lys Val lie Tyr His
 50 55 60

Tyr Ala Gly Gin Asp Ala Thr Asp ser Phe Arg Ala Leu His Pro Asp
 65 70 75 80

Ser Ala Leu Val Met Lys Tyr Leu Lys Pro Leu Leu lie Gly Gin Val
 85 90 95

Ala Pro Gly Ser ser Thr Ala Ala ser lie val Asp Gly Ala Arg Pro
 100 105 110

Ala Pro Ser Ala Phe Val Glu Glu Phe Arg Gin Val Arg Lys Glu Phe
 115 120 125

Gl u Gl u Gi n Gly Leu Phe Gl u Ala ser Trp ser Phe Phe Phe Gly Met
 130 135 140

Leu Ala His lie Phe Leu Leu Gl u Ala Ala Ala Tyr Tyr Ser lie Lys
 145 150 155 160

Leu Leu Gly Asn Ser Trp Pro Val Tyr Leu Leu Ala Val Gly Leu Leu
 165 170 175

Phoenix\templ8528 .tmp .txt

Ala Thr Ala Gin Ala Gin Ala Gly Trp Leu Gin His Asp Cys Gly His
 180 185 190

Leu Ser Val Phe Lys Lys Ser Lys Trp Asn His Trp Met His Tyr lie
 195 200 205

Val lie Cys His lie Lys Gly Ala Ser Arg Ala Trp Trp Asn Trp Arg
 210 215 220

His Phe Glu His His Ala Lys Pro Asn Val Val Arg Lys Asp Pro Asp
 225 230 235 240

lie Thr Phe Pro Asn Leu Phe Leu Leu Gly Asp His Leu Thr Arg Lys
 245 250 255

Trp Ala Lys Ala Lys Lys Gly Val Met Pro Tyr Asn Lys Gin His Leu
 260 265 270

Tyr Trp Trp Ala Phe Pro Pro Leu Leu Leu Pro val Tyr Phe His Tyr
 275 280 285

Asp Asn lie Arg Tyr Val Phe Gin His Lys His Trp Trp Asp Leu Phe
 290 295 300

Trp lie Ala Thr Phe Phe Ala Lys His Phe Thr Leu Tyr Gly Pro Leu
 305 310 315 320

Met Gly Gly Trp Gly Ala Phe Trp Phe Tyr Met Leu Val Arg Thr Val
 325 330 335

Glu Ser His Trp Phe Thr Trp Val Thr Gin Met Asn His lie Pro Met
 340 345 350

His Val Asp Asn Asp Arg Glu Leu Asp Trp Pro Thr Leu Gin Gly Leu
 355 360 365

Ala Thr Cys Asn Val Glu Gly Ser Leu Phe Asn Asp Trp Phe Thr Gly
 370 375 380

His Leu Asn Tyr Gin lie Glu His His Leu Phe Pro Thr Met Pro Arg
 385 390 395 400

His Asn Tyr Ala Val Ala Asn Lys Lys Val Gin Ala Leu Tyr Lys Lys
 405 410 415

His Gly Val Pro Met Gin Thr Lys Gly Leu lie Glu Ala Phe Ala Asp
 420 425 430

lie Val Lys Ser Leu Glu His Tyr Gly Glu Val Trp Lys Glu Ala Tyr
 435 440 445

Tyr Gly

PhoenixTemp18528 .tmp .txt

450

<210> 17

<211> 1374

<212> DNA

<213> Perkinsus marinus

<220>

<221> CDS

<222> (1)..(1374)

<223> Delta-5-Desaturase

<400> 17

atg act act tea ace act act gtg caa eta caa gaa gac ctg tea agt 48
Met Thr Thr Ser Thr Thr Val Gi n Leu Gi n Gl u Asp Leu Ser Ser
1 5

ggt gac cag aac gcc cac ccc agt cca age cga get act cct agt gtt 96
Gly Asp Gi n Asn Ala Hi s Pro Ser Pro Ser Arg Ala Thr Pro Ser Val
20 25 30

ggt gat act aag gag gat gcg agg gtt gtt ate aaa eta ttt ggt aca 144
Gly Asp Thr Lys Gl u Asp Ala Arg Val Val lie Lys Leu Phe Gly Thr
35 40 45

tgg gtt gat gtt aca get tgg ttg aat gac cat cct ggt ggt tct aaa 192
Trp Val Asp Val Thr Ala Trp Leu Asn Asp His Pro Gly Gly Ser Lys
50 55 60

gtg etc aga gca ttc aac aag aag gac gcg act gat get gtt atg gcc 240
Val Leu Arg Ala Phe Asn Lys Lys Asp Ala Thr Asp Ala val Met Ala
65 70 75 80

atg cac act gat gaa get ate aag cgc ate ate aga ttt tea aat gtg 288
Met His Thr Asp Glu Ala lie Lys Arg lie lie Arg Phe Ser Asn Val
85 90 95

gtc tec teg gcc ccc ate aac gcc tct att ggt gat gtc cag gtt att 336
val ser ser Ala Pro lie Asn Ala ser lie Gly Asp val Gin val lie
100 105 110

gag aaa tct eta teg aga gaa cag ttg atg tat tac aag etc cgc act 384
Glu Lys Ser Leu Ser Arg Glu Gin Leu Met Tyr Tyr Lys Leu Arg Thr
115 120 125

ctt get aga aac cag gcc tgg ttt caa age aat eta tta tac gaa gga 432
Leu Ala Arg Asn Gin Gly Trp Phe Gin ser Asn Leu Leu Tyr Glu Gly
130 135 140

gtg aaa gca atg ata gcc ttc ggt ttg etc ate ate ggg ttt get act 480
Val Lys Ala Met lie Ala Phe Gly Leu Leu lie lie Gly Phe Ala Thr
145 150 155 160

etc tac ttc gac tat ggt att tgg tea ace gca ctg ata ggt ttc get 528
Leu Tyr Phe Asp Tyr Gly lie Trp Ser Thr Ala Leu lie Gly Phe Ala
165 170 175

tgg ttt cag ctg ggg tgg ttg gga cat gac tgg tct cat cat aca get 576

Phoenix\templ8528 .tmp .txt

450

455

<210> 18

<211> 457

<212> PRT

<213> Perkinsus marinus

<400> 18

Met Thr Thr Ser Thr Thr Thr Val Gi n Leu Gi n Gl u Asp Leu Ser Ser
 1 5 10 15

Gly Asp Gl n Asn Ala His Pro Ser Pro Ser Arg Ala Thr Pro Ser Val
 20 25 30

Gly Asp Thr Lys Gl u Asp Ala Arg Val Val lie Lys Leu Phe Gly Thr
 35 40 45

Trp Val Asp Val Thr Ala Trp Leu Asn Asp His Pro Gly Gly Ser Lys
 50 55 60

Val Leu Arg Ala Phe Asn Lys Lys Asp Ala Thr Asp Ala Val Met Ala
 65 70 75 80

Met His Thr Asp Gl u Ala lie Lys Arg lie lie Arg Phe Ser Asn Val
 85 90 95

Val Ser Ser Ala Pro lie Asn Ala Ser lie Gly Asp Val Gi n Val lie
 100 105 110

Gl u Lys Ser Leu Ser Arg Gl u Gi n Leu Met Tyr Tyr Lys Leu Arg Thr
 115 120 125

Leu Ala Arg Asn Gi n Gly Trp Phe Gi n Ser Asn Leu Leu Tyr Gl u Gly
 130 135 140

Val Lys Ala Met lie Ala Phe Gly Leu Leu lie lie Gly Phe Ala Thr
 145 150 155 160

Leu Tyr Phe Asp Tyr Gly lie Trp Ser Thr Ala Leu lie Gly Phe Ala
 165 170 175

Trp Phe Gi n Leu Gly Trp Leu Gly His Asp Trp Ser His His Thr Ala
 180 185 190

Leu Pro Lys Ser Thr Thr Asn Cys Ala Asn Tyr Asn Asp Tyr Leu Gly
 195 200 205

Trp Leu Thr Gly Leu Ala Arg Gly Asn Thr Leu Leu Trp Trp Lys Leu
 210 215 220

PhoenixTtempl8528 .tmp .txt

Arg His Asn Thr His His Val Leu Thr Asn Gin Tyr Glu Asn Asp Pro
225 230 235 240

Asp lie Leu Thr Gin Pro Pro Leu His Phe Phe Glu Asp Phe Asp Val
245 250 255

Gly Asn Val Asn Arg Tyr Gin Ala Val Tyr Tyr Leu Pro Met Leu Thr
260 265 270

Leu Leu His Leu Phe Trp Leu Tyr Glu Ser Val Leu Val Cys Leu Arg
275 280 285

Gin Ser Lys Ser lie Asn Arg Tyr Asn Arg Met His Ala Arg Arg Asp
290 295 300

Thr Val Ala Leu Val Leu His lie Leu lie Val Gly lie lie Ser Tyr
305 310 315 320

Thr Ser Gly Lys Tyr Leu Leu lie Leu Leu Ala Tyr Met Leu Ser Gly
325 330 335

Phe Leu Thr Ala Val Val Phe Ala Ser His Tyr Asn Glu Pro Arg
340 345 350

Val Ala Ser Gly Glu Ser Leu Ser Leu Val Arg Gin Thr Leu Leu Thr
355 360 365

Thr lie Asn lie Gly Ser Phe Ser Asp Thr His Trp Glu Lys Lys Leu
370 375 380

Trp Phe Tyr Leu Thr Gly Gly Leu Asn Met Gin lie Glu His His Leu
385 390 395 400

Phe Pro Thr Met Pro Arg His Asn Leu Pro Lys Thr Thr Phe Leu Val
405 410 415

Lys Ser Leu Ala Gin Glu Leu Gly Leu Pro Tyr Lys Glu Thr Asn lie
420 425 430

Val Ser Leu Thr Lys Ala Ala Val Thr Thr Leu His His Asn Ala Leu
435 440 445

Arg Asn lie Glu Arg Leu Leu Ala Arg
450 455

<210> 19

<211> 1224

<212> DNA

<213> Acanthamoeba castellanii

PhoenixTemp18528 .tmp .txt

<220>

<221> CDS

<222> (1)..(1224)

<223> Delta-12/Delta-15-Desaturase

<400> 19

atg	act	att	act	ace	cag	ace	ttg	aac	cag	aag	get	get	aag	aag	48	
Met	Thr	Ile	Thr	Thr	Gln	Thr	Leu	Asn	Gln	Lys	Ala	Ala	Lys	Lys		
1			5					10					15			
gga	gga	aag	gag	agg	get	cca	att	att	cca	aag	gag	aac	get	cca	ttc	96
Gly	Gly	Lys	Glu	Arg	Ala	Pro	lie	lie	Pro	Lys	Glu	Asn	Ala	Pro	Phe	
			20					25					30			
act	ttg	gga	cag	ate	aag	gga	get	ate	cca	cct	cat	etc	ttc	aag	cac	144
Thr	Leu	Gly	Gln	lie	Lys	Gly	Ala	lie	Pro	Pro	His	Leu	Phe	Lys	His	
		35					40					45				
tec	atg	ttg	aag	tct	ttc	tec	tac	ttg	gga	gtg	gat	ttg	ttg	gag	tct	192
Ser	Met	Leu	Lys	Ser	Phe	Ser	Tyr	Leu	Gly	Val	Asp	Leu	Leu	Glu	Ser	
	50					55					60					
ace	ate	tgg	ttg	ttc	etc	ate	ttg	tac	ttg	gat	gga	etc	act	aag	gag	240
Thr	lie	Trp	Leu	Phe	Leu	lie	Leu	Tyr	Leu	Asp	Gly	Leu	Thr	Lys	Glu	
65					70					75					80	
aac	ace	ttg	ttg	aac	tgg	act	tgc	tgg	gtt	gca	tac	tgg	ttg	tac	caa	288
Asn	Thr	Leu	Leu	Asn	Trp	Thr	Cys	Trp	Val	Ala	Tyr	Trp	Leu	Tyr	Gln	
				85					90					95		
gga	ttg	act	tgg	act	gga	att	tgg	gtg	ttg	get	cat	gag	tgt	gga	cat	336
Gly	Leu	Thr	Trp	Thr	Gly	lie	Trp	Val	Leu	Ala	His	Glu	Cys	Gly	His	
			100					105					110			
gga	gga	ttc	gtt	get	caa	gag	tgg	ttg	aac	gat	ace	gtg	ggt	ttc	att	384
Gly	Gly	Phe	Val	Ala	Gln	Glu	Trp	Leu	Asn	Asp	Thr	Val	Gly	Phe	lie	
		115				120						125				
ttc	cat	ace	gtg	etc	tac	gtt	cca	tac	ttc	tec	tgg	aag	ttc	tct	cat	432
Phe	His	Thr	Val	Leu	Tyr	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Phe	Ser	His	
	130					135					140					
get	aag	cac	cat	cac	tac	ace	aac	cac	atg	act	aag	gat	gag	cca	ttc	480
Ala	Lys	His	His	His	Tyr	Thr	Asn	His	Met	Thr	Lys	Asp	Glu	Pro	Phe	
145					150				155						160	
gtg	cca	cat	aca	ate	act	cca	gag	caa	agg	get	aaa	gtg	gat	caa	gga	528
Val	Pro	His	Thr	lie	Thr	Pro	Glu	Gln	Arg	Ala	Lys	Val	Asp	Gln	Gly	
				165					170					175		
gag	ttg	cca	cat	cca	aac	aag	cca	tec	etc	ttc	get	ttc	tac	gag	aga	576
Glu	Leu	Pro	His	Pro	Asn	Lys	Pro	Ser	Leu	Phe	Ala	Phe	Tyr	Glu	Arg	
			180					185					190			
tgg	gtg	ate	cca	ttc	gtg	atg	ttg	ttc	ttg	gga	tgg	cca	etc	tac	ttg	624
Trp	Val	lie	Pro	Phe	Val	Met	Leu	Phe	Leu	Gly	Trp	Pro	Leu	Tyr	Leu	
		195				200						205				
tct	ate	aac	get	tct	gga	cca	cca	aag	aag	gag	ttg	gtt	tec	cac	tac	672
Ser	lie	Asn	Ala	Ser	Gly	Pro	Pro	Lys	Lys	Glu	Leu	Val	Ser	His	Tyr	
	210					215					220					

PhoenixT empl8528 .tmp .txt

gat cca aag get tec ate ttc aac aag aaa gat tgg tgg aag ate ttg 720
 Asp Pro Lys Ala Ser lie Phe Asn Lys Lys Asp Trp Trp Lys lie Leu
 225 230 235 240

etc tct gat ttg gga ttg gtt get tgg act ttg get ttg tgg aag ttg 768
 Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu
 245 250 255

gga gag act ttc gga ttc gga ttg gtg get get ctt tac att cca cca 816
 Gly Glu Thr Phe Gly Phe Gly Leu Val Ala Ala Leu Tyr lie Pro Pro
 260 265 270

gtg etc gtt ace aac tct tac ttg gtg get ate ace ttc ttg caa cac 864
 Val Leu Val Thr Asn Ser Tyr Leu Val Ala lie Thr Phe Leu Gin His
 275 280 285

ace gat gat ate etc cca cat tac gat get act gag tgg act tgg ttg 912
 Thr Asp Asp lie Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu
 290 295 300

aga gga get ttg tgc act gtg gat aga tct ttg gga tgg ttc gga gat 960
 Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp
 305 310 315 320

tac aag ace cat cac ate gtt gat act cat gtg ace cac cac ate ttc 1008
 Tyr Lys Thr His His lie Val Asp Thr His Val Thr His His lie Phe
 325 330 335

tct tac etc cca ttc tat aac get gag gag get act aag get att aag 1056
 Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala lie Lys
 340 345 350

cca gtg ttg aag gag tat cac tgc gag gat aag aga gga ttc ttc cac 1104
 Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His
 355 360 365

ttc tgg tac ttg ttc ttc aag ace get get gag aac tct gtt gtg gat 1152
 Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp
 370 375 380

aac gag ace aac aag tec cca gga ate ttc tac ttc ttc agg gag gag 1200
 Asn Glu Thr Asn Lys Ser Pro Gly lie Phe Tyr Phe Phe Arg Glu Glu
 385 390 395 400

att aag cac gga aag get cat tga 1224
 lie Lys His Gly Lys Ala His
 405

<210> 20

<211> 407

<212> PRT

<213> Acanthamoeba caste! lanii

<400> 20

Met Thr lie Thr Thr Thr G n Thr Leu Asn G n Lys Ala Al a Lys Lys
 1 5 10 15

Gly Gly Lys G u Arg Al a Pro lie lie Pro Lys G u Asn Al a Pro Phe
 20 25 30

PhoenixTempl8528 .tmp .txt

Thr Leu Gly Gln lie Lys Gly Ala lie Pro Pro His Leu Phe Lys His
 35 40 45

Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser
 50 55 60

Thr lie Trp Leu Phe Leu lie Leu Tyr Leu Asp Gly Leu Thr Lys Glu
 65 70 75 80

Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln
 85 90 95

Gly Leu Thr Trp Thr Gly He Trp Val Leu Ala His Glu Cys Gly His
 100 105 110

Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe lie
 115 120 125

Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His
 130 135 140

Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe
 145 150 155 160

Val Pro His Thr He Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly
 165 170 175

Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg
 180 185 190

Trp val lie Pro Phe val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu
 195 200 205

Ser lie Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr
 210 215 220

Asp Pro Lys Ala ser lie Phe Asn Lys Lys Asp Trp Trp Lys lie Leu
 225 230 235 240

Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu
 245 250 255

Gly Glu Thr Phe Gly Phe Gly Leu val Ala Ala Leu Tyr lie Pro Pro
 260 265 270

Val Leu Val Thr Asn Ser Tyr Leu Val Ala lie Thr Phe Leu Gln His
 275 280 285

Thr Asp Asp lie Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu
 290 295 300

PhoenixTempl8528 .tmp .txt

Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp
 305 310 315 320

Tyr Lys Thr His His lie Val Asp Thr His Val Thr His His H e Phe
 325 330 335

Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala lie Lys
 340 345 350

Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His
 355 360 365

Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn ser val val Asp
 370 375 380

Asn Glu Thr Asn Lys Ser Pro Gly lie Phe Tyr Phe Phe Arg Glu Glu
 385 390 395 400

lie Lys His Gly Lys Ala His
 405

<210> 21

<211> 1224

<212> DNA

<213> Acanthamoeba castellanii

<220>

<221> CDS

<222> (1)..(1224)

<223> Delta-12/Delta-15-Desaturase

<400> 21

atg acg ate acg acg acg cag aca ctg aat cag aag gca gcc aag aag 48
 Met Thr H e Thr Thr Thr Gin Thr Leu Asn Gin Lys Ala Ala Lys Lys
 1 5 10 15

ggc gga aag gag cgc get ccg ate att ccc aag gag aac gcc ccc ttc 96
 Gly Gly Lys Glu Arg Ala Pro lie lie Pro Lys Glu Asn Ala Pro Phe
 20 25 30

act ctg ggc cag ate aag ggc gcc att cct ccg cat etc ttc aag cac 144
 Thr Leu Gly Gin lie Lys Gly Ala lie Pro Pro His Leu Phe Lys His
 35 40 45

age atg etc aaa tec ttc age tat ctg ggc gtg gat ctg ctg gag age 192
 Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser
 50 55 60

ace ate tgg etc ttc etc ate etc tac etc gac ggc etc ace aag gag 240
 Thr lie Trp Leu Phe Leu lie Leu Tyr Leu Asp Gly Leu Thr Lys Glu
 65 70 75 80

PhoenixTemp18528 .tmp .txt																
aac	acg	etc	etc	aac	tgg	act	tgc	tgg	ggt	gcg	tac	tgg	etc	tac	cag	288
Asn	Thr	Leu	Leu	Asn	Trp	Thr	Cys	Trp	Val	Ala	Tyr	Trp	Leu	Tyr	Gln	
				85					90					95		
ggt	ctg	ace	tgg	act	ggc	att	tgg	gtg	ctg	gee	cac	gag	tgt	ggc	cat	336
Gly	Leu	Thr	Trp	Thr	Gly	lie	Trp	Val	Leu	Ala	His	Glu	Cys	Gly	His	
			100					105					110			
ggc	ggc	ttc	gtg	gcg	cag	gag	tgg	etc	aac	gac	acg	gtc	ggc	ttc	ate	384
Gly	Gly	Phe	Val	Ala	Gln	Glu	Trp	Leu	Asn	Asp	Thr	Val	Gly	Phe	He	
		115					120					125				
ttc	cac	ace	gtc	etc	tac	gtg	ccc	tac	ttc	teg	tgg	aag	ttc	tec	cac	432
Phe	His	Thr	Val	Leu	Tyr	Val	Pro	Tyr	Phe	Ser	Trp	Lys	Phe	Ser	His	
	130					135					140					
gee	aag	cac	cac	cac	tac	ace	aac	cac	atg	aca	aag	gac	gag	ccc	ttc	480
Ala	Lys	His	His	His	Tyr	Thr	Asn	His	Met	Thr	Lys	Asp	Glu	Pro	Phe	
145					150					155					160	
gtg	ccc	cac	ace	ate	ace	cct	gag	cag	agg	gcc	aag	gtc	gac	cag	ggc	528
Val	Pro	His	Thr	lie	Thr	Pro	Glu	Gln	Arg	Ala	Lys	Val	Asp	Gln	Gly	
				165					170					175		
gag	ctg	ccc	cac	ccc	aac	aag	ccc	tec	etc	ttc	gcc	ttc	tac	gaa	agg	576
Glu	Leu	Pro	His	Pro	Asn	Lys	Pro	Ser	Leu	Phe	Ala	Phe	Tyr	Glu	Arg	
			180					185					190			
tgg	gtc	ate	ccc	ttc	gtc	atg	etc	ttc	etc	ggc	tgg	ccg	etc	tac	ctg	624
Trp	Val	lie	Pro	Phe	Val	Met	Leu	Phe	Leu	Gly	Trp	Pro	Leu	Tyr	Leu	
		195				200						205				
tec	ate	aac	gcc	tct	ggc	cct	ccc	aag	aag	gag	ctt	gtg	tec	cac	tac	672
Ser	lie	Asn	Ala	Ser	Gly	Pro	Pro	Lys	Lys	Glu	Leu	Val	Ser	His	Tyr	
	210					215					220					
gac	ccc	aaa	gcc	age	ate	ttc	aac	aag	aag	gac	tgg	tgg	aag	ate	ctt	720
Asp	Pro	Lys	Ala	Ser	lie	Phe	Asn	Lys	Lys	Asp	Trp	Trp	Lys	lie	Leu	
225					230					235					240	
etc	tct	gac	etc	ggc	ctt	gtg	gcg	tgg	ace	ctg	gcc	etc	tgg	aag	ctg	768
Leu	ser	Asp	Leu	Gly	Leu	val	Ala	Trp	Thr	Leu	Ala	Leu	Trp	Lys	Leu	
			245						250					255		
ggc	gag	ace	ttc	ggc	ttc	ggt	etc	gtg	gcc	gcc	etc	tac	att	ccg	ccc	816
Gly	Glu	Thr	Phe	Gly	Phe	Gly	Leu	Val	Ala	Ala	Leu	Tyr	lie	Pro	Pro	
			260					265					270			
gtg	ctg	gtg	ace	aac	tec	tac	ctg	gtg	gcc	ate	ace	ttc	etc	cag	cac	864
val	Leu	val	Thr	Asn	ser	Tyr	Leu	val	Ala	lie	Thr	Phe	Leu	Gln	His	
		275					280					285				
ace	gac	gac	att	ctg	ccc	cac	tac	gac	gcc	ace	gag	tgg	ace	tgg	etc	912
Thr	Asp	Asp	lie	Leu	Pro	His	Tyr	Asp	Ala	Thr	Glu	Trp	Thr	Trp	Leu	
	290					295					300					
agg	ggt	get	etc	tgc	act	gtt	gat	cgt	teg	ctg	ggc	tgg	ttc	ggc	gac	960
Arg	Gly	Ala	Leu	cys	Thr	val	Asp	Arg	ser	Leu	Gly	Trp	Phe	Gly	Asp	
305					310					315					320	
tac	aag	acg	cac	cac	ate	gtc	gac	ace	cac	gtg	acg	cac	cac	ate	ttc	1008
Tyr	Lys	Thr	His	His	lie	Val	Asp	Thr	His	Val	Thr	His	His	lie	Phe	
				325					330					335		
teg	tac	ctg	ccg	ttc	tac	aac	gcc	gag	gag	gcc	ace	aag	gcc	ate	aag	1056
Ser	Tyr	Leu	Pro	Phe	Tyr	Asn	Ala	Glu	Glu	Ala	Thr	Lys	Ala	lie	Lys	
			340					345					350			
ccc	gtg	etc	aag	gag	tac	cac	tgc	gag	gac	aag	cgt	ggc	ttc	ttc	cac	1104

PhoenixTemp18528.tmp.txt
 Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His
 355 360 365
 ttc tgg tat ctg ttc ttc aag ace gcc gcc gag aac age gtt gtc gac 1152
 Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp
 370 375 380
 aac gag ace aac aag age ccc ggc ate ttc tac ttc ttc egg gag gag 1200
 Asn Glu Thr Asn Lys Ser Pro Gly lie Phe Tyr Phe Phe Arg Glu Glu
 385 390 395 400
 ate aag cac ggc aag gcc cac tag 1224
 lie Lys His Gly Lys Ala His
 405

<210> 22
 <211> 407
 <212> PRT
 <213> Acanthamoeba castellanii

<400> 22
 Met Thr lie Thr Thr Thr Gln Thr Leu Asn Gln Lys Ala Ala Lys Lys
 1 5 10 15
 Gly Gly Lys Glu Arg Ala Pro lie lie Pro Lys Glu Asn Ala Pro Phe
 20 25 30
 Thr Leu Gly Gln lie Lys Gly Ala lie Pro Pro His Leu Phe Lys His
 35 40 45
 Ser Met Leu Lys Ser Phe Ser Tyr Leu Gly Val Asp Leu Leu Glu Ser
 50 55 60
 Thr lie Trp Leu Phe Leu lie Leu Tyr Leu Asp Gly Leu Thr Lys Glu
 65 70 75 80
 Asn Thr Leu Leu Asn Trp Thr Cys Trp Val Ala Tyr Trp Leu Tyr Gln
 85 90 95
 Gly Leu Thr Trp Thr Gly lie Trp Val Leu Ala His Glu Cys Gly His
 100 105 110
 Gly Gly Phe Val Ala Gln Glu Trp Leu Asn Asp Thr Val Gly Phe lie
 115 120 125
 Phe His Thr Val Leu Tyr Val Pro Tyr Phe Ser Trp Lys Phe Ser His
 130 135 140
 Ala Lys His His His Tyr Thr Asn His Met Thr Lys Asp Glu Pro Phe
 145 150 155 160
 Val Pro His Thr lie Thr Pro Glu Gln Arg Ala Lys Val Asp Gln Gly

Phoenix\templ8528 .tmp .txt

165

170

175

Glu Leu Pro His Pro Asn Lys Pro Ser Leu Phe Ala Phe Tyr Glu Arg
180 185 190

Trp val lie Pro Phe val Met Leu Phe Leu Gly Trp Pro Leu Tyr Leu
195 200 205

Ser lie Asn Ala Ser Gly Pro Pro Lys Lys Glu Leu Val Ser His Tyr
210 215 220

Asp Pro Lys Ala Ser lie Phe Asn Lys Lys Asp Trp Trp Lys lie Leu
225 230 235 240

Leu Ser Asp Leu Gly Leu Val Ala Trp Thr Leu Ala Leu Trp Lys Leu
245 250 255

Gly Glu Thr Phe Gly Phe Gly Leu Val Ala Ala Leu Tyr lie Pro Pro
260 265 270

Val Leu Val Thr Asn Ser Tyr Leu Val Ala lie Thr Phe Leu Gin His
275 280 285

Thr Asp Asp lie Leu Pro His Tyr Asp Ala Thr Glu Trp Thr Trp Leu
290 295 300

Arg Gly Ala Leu Cys Thr Val Asp Arg Ser Leu Gly Trp Phe Gly Asp
305 310 315 320

Tyr Lys Thr His His lie Val Asp Thr His Val Thr His His H e Phe
325 330 335

Ser Tyr Leu Pro Phe Tyr Asn Ala Glu Glu Ala Thr Lys Ala lie Lys
340 345 350

Pro Val Leu Lys Glu Tyr His Cys Glu Asp Lys Arg Gly Phe Phe His
355 360 365

Phe Trp Tyr Leu Phe Phe Lys Thr Ala Ala Glu Asn Ser Val Val Asp
370 375 380

Asn Glu Thr Asn Lys Ser Pro Gly lie Phe Tyr Phe Phe Arg Glu Glu
385 390 395 400

H e Lys His Gly Lys Ala His
405

<210> 23

<211> 1254

<212> DNA

PhoenixTtempl8528 .tmp .txt

<213> Perkinsus marinus

<220>

<221> CDS

<222> (1).. (1254)

<223> Delta-12-Desaturase

<400> 23

atg ace caa act gag gtc caa gcc gga ccg tgt aga gat ggt agg aac 48
 Met Thr Gln Thr Glu val Gln Ala Gly Pro cys Arg Asp Gly Arg Asn
 1 5 10 15

etc aag agt gag get gat gtt aaa ggc ttc act gcg gag gag ttt act 96
 Leu Lys Ser Glu Ala Asp Val Lys Gly Phe Thr Ala Glu Glu Phe Thr
 20 25 30

aag gtt ggg ccg tct gtg tgt get ata caa tea get ate ccc atg cac 144
 Lys val Gly Pro ser val cys Ala lie Gln ser Ala lie Pro Met His
 35 40 45

tgt cgt gat agg age ctg tea agg tct gtc eta tgc gtc ate agg gat 192
 Cys Arg Asp Arg Ser Leu Ser Arg Ser Val Leu Cys Val lie Arg Asp
 50 55 60

etc etc tac ata aca gca tgt get get gtg cag tac tct ctg ttg gcg 240
 Leu Leu Tyr lie Thr Ala cys Ala Ala val Gln Tyr ser Leu Leu Ala
 65 70 75 80

tta gta ccc ccg gac tea ace etc ctg agg gca gtc etc tgg ggt gtt 288
 Leu Val Pro Pro Asp Ser Thr Leu Leu Arg Ala Val Leu Trp Gly Val
 85 90 95

tac att ttc tgg caa ggc gtc ttt ttt act ggt att tgg gtg atg ggc 336
 Tyr lie Phe Trp Gln Gly Val Phe Phe Thr Gly lie Trp Val Met Gly
 100 105 110

cac gag tgc ggc cat ggg get ttt tec cct tat tct atg ctg aac gat 384
 His Glu Cys Gly His Gly Ala Phe Ser Pro Tyr Ser Met Leu Asn Asp
 115 120 125

agt att ggt ttt gtc etc cac teg gcc etc ttg gta ccc tac ttc age 432
 Ser lie Gly Phe Val Leu His Ser Ala Leu Leu Val Pro Tyr Phe Ser
 130 135 140

tgg cag tac tec cat gcg agg cac cat aag ttc ace aac cac get act 480
 Trp Gln Tyr Ser His Ala Arg His His Lys Phe Thr Asn His Ala Thr
 145 150 155 160

aag ggt gag age cat gtc ccc age ctg gaa agt gag atg ggc gta ttc 528
 Lys Gly Glu Ser His Val Pro Ser Leu Glu Ser Glu Met Gly Val Phe
 165 170 175

agt cgt ata cag aag gcc ctg gag ggt tat ggt etc gat gat gtc ttc 576
 Ser Arg lie Gln Lys Ala Leu Glu Gly Tyr Gly Leu Asp Asp Val Phe
 180 185 190

cca gtc ttc cct ata gtg atg etc ctg gtt ggg tat cct gtg tat etc 624
 Pro Val Phe Pro lie Val Met Leu Leu Val Gly Tyr Pro Val Tyr Leu
 195 200 205

ttc tgg aat gca tea ggt ggg cgt gtg ggc tac gat cgc cgt ccg tac 672
 Phe Trp Asn Ala Ser Gly Gly Arg Val Gly Tyr Asp Arg Arg Pro Tyr

PhoenixTemp18528.tmp.txt

210		215		220		
age gac act aag cca tct cat ttc aat ccc aac ggt ggc ctt ttc cct						720
Ser Asp Thr Lys Pro Ser His Phe Asn Pro Asn Gly Gly Leu Phe Pro						
225		230		235		240
cct tat atg aga gag aaa gtc etc ctt agt gga gtt ggc tgt age ata						768
Pro Tyr Met Arg Glu Lys val Leu Leu ser Gly val Gly Cys ser lie						
		245		250		255
ace etc ctt att ttg gcc tat tgt get ggg agg gta ggc ctt age agt						816
Thr Leu Leu lie Leu Ala Tyr Cys Ala Gly Arg Val Gly Leu Ser Ser						
		260		265		270
gta ttg ttg tgg tat ggt tgt ccc tac ctt atg ace aac gcc tgg eta						864
Val Leu Leu Trp Tyr Gly Cys Pro Tyr Leu Met Thr Asn Ala Trp Leu						
		275		280		285
acg ctg tat ace tec eta cag cac acg cat gaa gga gtc ccc cat tat						912
Thr Leu Tyr Thr Ser Leu Gin His Thr His Glu Gly Val Pro His Tyr						
		290		295		300
ggc gat gag get ttc ace ttc ate aga ggt gcc tta get tct ate gat						960
Gly Asp Glu Ala Phe Thr Phe lie Arg Gly Ala Leu Ala Ser lie Asp						
		310		315		320
cgt cca ccg tat ggc att ttc tct acg cat ttt cac cac gaa att ggc						1008
Arg Pro Pro Tyr Gly lie Phe Ser Thr His Phe His His Glu lie Gly						
		325		330		335
ace act cat gtt ctg cac cac att gat tct agg ate ccc tgt tac cat						1056
Thr Thr His Val Leu His His lie Asp Ser Arg lie Pro Cys Tyr His						
		340		345		350
get aga gaa gcc act gat get ate aag cct att ctg ggg gat tac tat						1104
Ala Arg Glu Ala Thr Asp Ala lie Lys Pro lie Leu Gly Asp Tyr Tyr						
		355		360		365
agg gag gat ggt act cct ata gta aag gca ttt ttg aag gtc cac aga						1152
Arg Glu Asp Gly Thr Pro lie Val Lys Ala Phe Leu Lys Val His Arg						
		370		375		380
gag tgc aag ttc ate gga ggc etc aac ggc gtc cag ttt tac cgt cct						1200
Glu Cys Lys Phe lie Gly Gly Leu Asn Gly Val Gin Phe Tyr Arg Pro						
		385		390		400
ggg cag egg ccg cag cag cag ccc tgc ggc age aac get cgc act tct						1248
Gly Gin Arg Pro Gin Gin Gin Pro Cys Gly Ser Asn Ala Arg Thr Ser						
		405		410		415
cgt tag						1254
Arg						

<210> 24

<211> 417

<212> PRT

<213> Perkinsus marinus

<400> 24

Met Thr Gln Thr Glu Val Gln Ala Gly Pro Cys Arg Asp Gly Arg Asn
1 5 10 15

PhoenixTemp18528 .tmp .txt

Leu Lys Ser Glu Ala Asp Val Lys Gly Phe Thr Ala Glu Glu Phe Thr
 20 25 30

Lys Val Gly Pro Ser Val Cys Ala lie Gin Ser Ala lie Pro Met His
 35 40 45

Cys Arg Asp Arg Ser Leu Ser Arg Ser Val Leu Cys Val lie Arg Asp
 50 55 60

Leu Leu Tyr lie Thr Ala Cys Ala Ala Val Gin Tyr Ser Leu Leu Ala
 65 70 75 80

Leu Val Pro Pro Asp Ser Thr Leu Leu Arg Ala Val Leu Trp Gly Val
 85 90 95

Tyr lie Phe Trp Gin Gly Val Phe Phe Thr Gly lie Trp Val Met Gly
 100 105 110

His Glu Cys Gly His Gly Ala Phe Ser Pro Tyr Ser Met Leu Asn Asp
 115 120 125

Ser lie Gly Phe Val Leu His Ser Ala Leu Leu Val Pro Tyr Phe Ser
 130 135 140

Trp Gin Tyr Ser His Ala Arg His His Lys Phe Thr Asn His Ala Thr
 145 150 155 160

Lys Gly Glu Ser His Val Pro Ser Leu Glu Ser Glu Met Gly Val Phe
 165 170 175

Ser Arg lie Gin Lys Ala Leu Glu Gly Tyr Gly Leu Asp Asp Val Phe
 180 185 190

Pro Val Phe Pro lie Val Met Leu Leu Val Gly Tyr Pro Val Tyr Leu
 195 200 205

Phe Trp Asn Ala Ser Gly Gly Arg Val Gly Tyr Asp Arg Arg Pro Tyr
 210 215 220

Ser Asp Thr Lys Pro Ser His Phe Asn Pro Asn Gly Gly Leu Phe Pro
 225 230 235 240

Pro Tyr Met Arg Glu Lys Val Leu Leu Ser Gly Val Gly Cys Ser lie
 245 250 255

Thr Leu Leu lie Leu Ala Tyr Cys Ala Gly Arg Val Gly Leu Ser Ser
 260 265 270

Val Leu Leu Trp Tyr Gly Cys Pro Tyr Leu Met Thr Asn Ala Trp Leu
 275 280 285

Phoenixr empl8528 . tmp . txt

Thr Leu Tyr Thr Ser Leu Gln His Thr His Glu Gly Val Pro His Tyr
 290 295 300

Gly Asp Glu Ala Phe Thr Phe lie Arg Gly Ala Leu Ala Ser lie Asp
 305 310 315 320

Arg Pro Pro Tyr Gly lie Phe Ser Thr His Phe His His Glu lie Gly
 325 330 335

Thr Thr His Val Leu His His lie Asp Ser Arg lie Pro Cys Tyr His
 340 345 350

Ala Arg Glu Ala Thr Asp Ala lie Lys Pro lie Leu Gly Asp Tyr Tyr
 355 360 365

Arg Glu Asp Gly Thr Pro lie Val Lys Ala Phe Leu Lys Val His Arg
 370 375 380

Glu Cys Lys Phe lie Gly Gly Leu Asn Gly Val Gln Phe Tyr Arg Pro
 385 390 395 400

Gly Gln Arg Pro Gln Gln Gln Pro Cys Gly Ser Asn Ala Arg Thr Ser
 405 410 415

Arg

<210> 25

<211> 31

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(31)

<223>

<400> 25
ggtaccatgg cgatcacgac gacgcagaca c

31

<210> 26

<211> 34

PhoenixTemp18528 .tmp .txt

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(34)

<223>

<400> 26

gagctcctag tgggccttgc cgtgcttgat ctcc

34

<210> 27

<211> 30

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(30)

<223>

<400> 27

ggtaccatgg tctcacaac cccggcctc

30

<210> 28

<211> 29

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

PhoenixT empl8528 .tmp .txt

<222> (I).. (29)

<223>

<400> 28

ggagctctca gttctcagca cccatcttc

29

<210> 29

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 29

ggtaccatgg ccaccgcac tgcac

26

<210> 30

<211> 27

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(27)

<223>

<400> 30

ggagctttag ccgtagtag cctcctt

27

<210> 31

Phoenix\temp\18528 .tmp .txt

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 31

ggtaccatgg cggctgcgac ggcgac

26

<210> 32

<211> 27

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(27)

<223>

<400> 32

ggagctttag tcgtgcttcc tcttggg

27

<210> 33

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

Phoenix\templ8528 .tmp .txt

<221> misc_feature

<222> (1)..(26)

<223>

<400> 33

ggtaccatga cccaaactga ggtcca

26

<210> 34

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(26)

<223>

<400> 34

ggagctctaa cgagaagtgc gagcgt

26

<210> 35

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(26)

<223>

<400> 35

ggtaccatgt cttctcttac cctcta

26

PhoenixT empl8528 .tmp .txt

<210> 36

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 36

ggagctctat tccactatgg caacag

26

<210> 37

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 37

ggtaccatga ctacttcaac cactac

26

<210> 38

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

PhoenixTemp18528 .tmp .txt

<220>

<221> misc_feature

<222> (1)..(26)

<223>

<400> 38

ggagctctac ctagcaagca atctct

26

<210> 39

<211> 34

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(34)

<223>

<400> 39

ggatccacca tggcgatcac gacgacgcag acac

34

<210> 40

<211> 36

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(36)

<223>

<400> 40

ggtctagact agtgggcctt gccgtgcttg atctcc

36

PhoenixTemp18528 .tmp .txt

<210> 41

<211> 33

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(33)

<223>

<400> 41

ggatccagga tggctctcac aacccggcc etc

33

<210> 42

<211> 30

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(30)

<223>

<400> 42

ggcttagatc agttctcagc acccatcttc

30

<210> 43

<211> 26

<212> DNA

<213> unknown

<220>

Phoenix t empl8528 . tmp . txt

<223> artificial sequence

<220>

<221> misc_feature

<222> (1) .. (26)

<223>

<400> 43
ggatccatgg ccaccgcatc tgcac

26

<210> 44

<211> 29

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1) .. (29)

<223>

<400> 44
ggtctagatt agccgtagta ggcctcctt

29

<210> 45

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1) .. (26)

<223>

PhoenixT empl8528 .tmp .txt

<400> 45
ggatccatgg cggctgcgac ggcgac 26

<210> 46

<211> 29

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(29)

<223>

<400> 46
ggtctagatt agtcgtgctt cctottggg 29

<210> 47

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 47
ggatccatga cccaaactga ggtcca 26

<210> 48

<211> 28

<212> DNA

<213> unknown

PhoenixTemp18528 .tmp .txt

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(28)

<223>

<400> 48

ggctctagact aacgagaagt gcgagcgt

28

<210> 49

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(26)

<223>

<400> 49

ggatccatgt cttctcttac cctcta

26

<210> 50

<211> 28

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> misc_feature

<222> (1)..(28)

<223>

PhoenixT empl8528 .tmp .txt

<400> 50
ggcttagact attccactat ggcaacag 28

<210> 51

<211> 26

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(26)

<223>

<400> 51
ggatccatga ctacttcaac cactac 26

<210> 52

<211> 28

<212> DNA

<213> unknown

<220>

<223> artificial sequence

<220>

<221> mi sc_feature

<222> (1)..(28)

<223>

<400> 52
ggcttagact acctagcaag caatctct 28