



**First application of loop-mediated isothermal amplification (LAMP) assays for rapid identification of mating type in the heterothallic fungus *Aspergillus fumigatus***

Journal:	<i>Mycoses</i>
Manuscript ID	MYC-OA-2018-353.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	King, Kevin; Rothamsted Research, BCP Hawkins, Nichola ; Rothamsted Research, BCP Atkins, Sarah; Rothamsted Research, BCP Dyer, Paul; University of Nottingham, School of Life Sciences West, Jonathan ; Rothamsted Research, BCP Fraaije , Bart; Rothamsted Research, BCP
Keywords:	<i>Aspergillus fumigatus</i> , diagnostics, fungal pathogen, mating type, sexual reproduction
Area of Expertise:	Laboratory Diagnostics, Molecular Mycology
Additional Keywords:	

SCHOLARONE™  
Manuscripts

## First application of loop-mediated isothermal amplification (LAMP) assays for rapid identification of mating type in the heterothallic fungus *Aspergillus fumigatus*

K. M. King, N. J. Hawkins, S. Atkins, P. S. Dyer\*, J. S. West and B. A. Fraaije  
Rothamsted Research, Biointeractions and Crop Protection Department, Harpenden,  
Hertfordshire, AL5 2JQ, UK

\*School of Life Sciences, University of Nottingham, University Park, Nottingham NG7 2RD,  
UK

Corresponding author: Kevin King, Rothamsted Research, Harpenden, Hertfordshire, UK; +44  
(0)1582 763133; [Kevin.king@rothamsted.ac.uk](mailto:Kevin.king@rothamsted.ac.uk)

**Running title:** Fungal mating-type identification using LAMP

**Author contributions:** K.M.K., N.J.H., P.S.D., J.S.W., and B.A.F. conceived the ideas; K.M.K. and S.A. collected the data; K.M.K. analysed the data; K.M.K. led the writing; all authors critically reviewed the manuscript prior to submission.

### Summary

**Background** Loop-mediated isothermal amplification (LAMP) assays, which operate at a single temperature and require no post-reaction processing, have been described for rapid species-specific detection of numerous fungi. The technology has much less commonly been applied to identification of other key genetic traits such as fungicide resistance, and has not yet been applied to mating-type determination in any fungus.

**Objectives** To develop first LAMP assays for mating-type identification in a fungus, in this instance with the saprophytic mould and human opportunistic pathogen *Aspergillus fumigatus*, a heterothallic ascomycete requiring isolates of opposite mating type (*MAT1-1*, *MAT1-2*) for sexual reproduction.

**Methods** New LAMP primer sets, targeted to *MAT* gene sequences, were screened against 34 *A. fumigatus* isolates (of known mating type) from diverse clinical, environmental and geographic sources to establish if they could distinguish *MAT1-1* or *MAT1-2* genotypes.

**Results and conclusions** The new assays, operating at a single temperature of 65°C, correctly identified the mating-type of *A. fumigatus* isolates in <20 minutes, and thus have numerous research and practical applications. Similar *MAT* LAMP assays could now be developed for other fungi of agricultural, environmental, industrial and/or medical importance.

**Keywords:** *Aspergillus fumigatus*, diagnostics, fungal pathogen, mating type, sexual reproduction

### Introduction

The fungus *Aspergillus fumigatus* is a saprophytic mould commonly found on plant debris and in soil. It is also an opportunistic human pathogen causing allergic symptoms and life-threatening invasive infections. The incidence of invasive aspergillosis (IA) has been increasing in recent years largely due to increased numbers of immunocompromised individuals in the population unable to fight off infection.<sup>1</sup> For more than 145 years, *A. fumigatus* was only known to reproduce asexually, although several signatures of cryptic sexuality were present, e.g. presence and expression of mating (*MAT*) genes and evidence of gene recombination within natural populations.<sup>2</sup> However, the breakthrough 2009 discovery of

a functional sexual cycle<sup>3</sup> had several implications including: (a) potentially explaining high genotypic diversity observed in populations; (b) production of sexually-derived airborne ascospores potentially more resilient to unfavourable environmental conditions; and (c) generation of sexual progeny with potentially greater pathogenicity and/or reduced sensitivity to fungicides.<sup>4</sup> *Aspergillus fumigatus* possesses a heterothallic (obligate outbreeding) mating system, with highly dissimilar stretches of DNA, termed ‘idiomorphs’, present in isolates of opposite mating type as is characteristic for heterothallic ascomycete species.<sup>5</sup> Thus, *MATI-1* isolates contain an alpha-domain *MATI-1-1* gene whereas *MATI-2* isolates contain a high-mobility group *MATI-2-1* gene together with a recently described *MATI-2-4* gene.<sup>6</sup> A multiplex PCR-based assay for determination of mating type has previously been developed for *A. fumigatus*.<sup>2</sup>

More recently, loop-mediated isothermal amplification (LAMP) assays have become increasingly used for rapid species-specific detection of numerous fungi, including *A. fumigatus*.<sup>7</sup> LAMP technology, first described by in 2000<sup>8</sup>, typically involves 4-6 primers in each reaction and has several purported advantages over PCR-based diagnostics. These include faster reaction times, potentially improved sensitivity and specificity, increased tolerance of sample inhibitors, no requirement for additional post-reaction processing (e.g. resolving PCR products on agarose gels) and use of only a single constant reaction temperature thus raising the possibility of field-based detection. Despite these advantages, LAMP assays have much less commonly been applied to detection of other key genetic traits such as fungicide resistance, one recent example being an assay targeted to a 34 bp tandem repeat in the *cyp51A* gene that has been associated with azole resistance in *A. fumigatus*.<sup>9</sup> To date, however, LAMP assays have not been used for rapid detection of different mating types in fungi. The objective of the present study was therefore to develop and evaluate for the first time whether LAMP assays could be used for the rapid identification of mating type in a fungus, with a focus here on the human opportunistic pathogen *A. fumigatus*.

## Methods

### Ethics statement

The authors confirm that the ethical policies of the journal, as noted on the journal’s author guidelines page, have been adhered to. No ethical approval was required as the research in this article related to micro-organisms.

### Fungal isolates, DNA extraction and initial molecular characterization

Details of *A. fumigatus* isolates, including source material and geographic origin, are given in Table 1; all isolates are maintained as -80°C glycerol stocks at Rothamsted Research, UK. Genomic DNA was extracted from *A. fumigatus* spores, harvested from one-week old cultures grown on Sabouraud dextrose agar (Lab M Ltd, UK) at 37°C, using a MasterPure yeast DNA purification kit (Epicentre, USA) into a final volume of 100 µL TE buffer. DNA was quantified via nanodrop spectrophotometer and diluted to 10 ng/ µL using PCR grade water. The mating type of these isolates was first determined using the published multiplex PCR assay<sup>2</sup>. Amplicons were resolved on agarose gels, with 834 bp or 438 bp products amplified from *MATI-1* or *MATI-2* isolates, respectively.

### Design and validation of MAT LAMP assays

For the *MATI-1* LAMP assay, *MAT* idiomorph sequence was downloaded from GenBank (Accession: AY898661<sup>2</sup>), with LAMP primers targeted to the internal *MATI-1-1* gene. For the *MATI-2* LAMP assay, *MAT* idiomorph sequence was sourced from the *A. fumigatus* Ensembl genome (isolate AF293; gene ID: AFUA\_3G06170), with LAMP primers targeted to the internal *MATI-2-1* gene. LAMP primer sets were designed using the free online software package PrimerExplorer (v. 5) with default settings.

For screening isolates against each of the *MAT* LAMP assays, 15  $\mu$ L reactions contained 0.3  $\mu$ L BIP primer (final concentration 2  $\mu$ M), 0.3  $\mu$ L FIP primer (2  $\mu$ M), 0.15  $\mu$ L LOOPB primer (1  $\mu$ M), 0.15  $\mu$ L LOOPF primer (1  $\mu$ M), 0.3  $\mu$ L B3 primer (0.2  $\mu$ M), 0.3  $\mu$ L F3 primer (0.2  $\mu$ M) (Table 1),  $\mu$ L 7.5  $\mu$ L isothermal mastermix (ISO-001; Optigene, UK) and 1  $\mu$ L DNA template (10 ng total DNA). No-template (PCR-grade water) controls were included in each test run. LAMP assays were run at 65°C for 30 minutes (FAM fluorescence measured every 30 secs), followed by a final dissociation step at 95°C for 1 min; 55°C for 30 secs and 95°C for 30 secs. Assays were run with a MX3000p qPCR system (Agilent), with data analysed using inbuilt 7500 SDS software (v.1.4; Applied Biosystems). Dissociation curves were checked manually after each run to confirm the presence of a single peak.

## Results

### Development and validation of *MAT* LAMP assays

For all *A. fumigatus* isolates tested, identical *MAT* genotype results were obtained using the previously described multiplex PCR assay<sup>2</sup> (see Figure 1 for representative results) and the new *MAT*-specific LAMP assays developed in the present study (Figure 1, Table 2). The new *MATI-1* and *MATI-2* -specific LAMP assays gave positive results within 10 – 20 mins (i.e. clear amplification curves) only for isolates of corresponding *MATI-1* or *MATI-2* type, respectively (Fig 2A). Positive results obtained with each *MAT*-specific LAMP assays gave single dissociation curves of c. 89.5°C ( $\pm$ 0.3), indicating specific amplification of the targeted *MAT* gene regions (Fig 2B). No-template (water) controls tested negative, i.e. no amplification curves or dissociation plot peaks were observed (data not shown).

## Discussion

This study reports the first use of LAMP technology to rapidly (within 20 mins) establish the mating-type identity for a fungus, as demonstrated here for isolates of *A. fumigatus*. The *MATI-1* and *MATI-2* -specific LAMP assays appeared robust, being successfully applied to isolates of known opposite *MAT* type from a diverse range of clinical and environmental sources (air, food, plant and soil) and geographic locations (Africa, Asia, Europe and North and South America). These assays will be of use in research into the applied biology of this important human opportunistic pathogen. For example, they will allow the rapid set-up of sexual crosses with isolates of known opposite *MAT* type, subsequent analysis of the *MAT* type inheritance of the progeny, and through progeny analysis the determination of the genetic basis of traits such as antifungal resistance and virulence.

It should now be possible to develop similar LAMP assays targeting *MAT* gene sequences to allow rapid mating-type determination in other heterothallic fungi of medical [e.g. *Aspergillus lentulus* - another causal agent of human aspergillosis<sup>10</sup>], agricultural [e.g. *Zymoseptoria tritici* - cause of wheat Septoria leaf blotch<sup>11</sup>], environmental [e.g. *Hymenoscyphus fraxineus* - cause of ash dieback<sup>12</sup>] and industrial [e.g. *Penicillium chrysogenum* - used in penicillin production<sup>13</sup>] importance. Such assays could also provide a better understanding into the reproductive strategies of various fungal pathogens, providing

insight into their evolutionary potential and possible risk of breakdown of disease management strategies.<sup>14</sup> Furthermore, they could also be used to indirectly assess possible cryptic sexuality in fungi for which no sexual stage is yet known, given that frequency dependent selection operating on *MAT* genes generally, although not always, results in a 1:1 distribution of mating types.<sup>5,14</sup>

## Acknowledgements

These studies were partly supported by the Rothamsted Research Smart Crop Protection (SCP) strategic programme (BBS/OS/CP/000001) funded through the Biotechnology and Biological Sciences Research Council's Industrial Strategy Challenge Fund and by CropLife International through project 14209 'Investigating the sources and spread of azole resistant *Aspergillus fumigatus* strains in the environment'.

## Conflict of interest

No conflict of interest is declared.

## Figure legend

**Figure 1** Representative results from screening of *Aspergillus fumigatus* isolates with the multiplex PCR mating-type assay.<sup>2</sup> *MAT1-1* and *MAT1-2* type isolates are distinguished by amplicons of 834 bp and 438 bp respectively. 'L' indicates EasyLadder 1 (Bioline); NTC indicates no template control.

**Figure 2** Representative results from screening of *Aspergillus fumigatus* isolates of *MAT1-1* (18-C6-9) or *MAT1-2* (G2-2) type screened against the new *MAT*-specific LAMP assays. Shown are (A) amplification plots and (B) dissociation plots. See base of figure for explanatory legend. No-template (water) controls also gave negative results (data not shown).

## References

1. Dagenais TRT, Keller NP. Pathogenesis of *Aspergillus fumigatus* in invasive aspergillosis. *Clinical Microbiology Reviews* 2009; **22**: 447-65
2. Paoletti M, Rydholm C, Schwier EU *et al.* Evidence for sexuality in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Current Biology* 2005; **15**: 1242-248.
3. O'Gorman CM, Fuller HT, Dyer PS. Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature* 2009; **457**: 471-74.
4. Zhang J, Snelders E, Zwaan BJ *et al.* A novel environmental azole resistance mutation in *Aspergillus fumigatus* and a possible role of sexual reproduction in its emergence. *mBio* 2017; **8**: pp.e00791-17.
5. Dyer PS, Inderbitzin P, Debuchy R. Mating-type structure, function, regulation and evolution in the Pezizomycotina. In: Wendland J, ed. *Growth, Differentiation and Sexuality, The Mycota I*. 3rd ed. Switzerland: Springer International Publishing 2016; 351-385.
6. Yu Y, Amich J, Will C *et al.* The novel *Aspergillus fumigatus* *MAT1-2-4* mating-type gene is required for mating and cleistothecia formation. *Fungal Biology and Genetics* 2017; **108**, 1-12.
7. Tang Q, Shuguang T, Yu N *et al.* Development and evaluation of a loop-mediated isothermal amplification method for rapid detection of *Aspergillus fumigatus*. *Journal of Clinical Microbiology* 2016; **54**: 950-55.

8. Notomi T, Okayama H, Masubuchi H *et al.* Loop-mediated isothermal amplification of DNA. *Nucleic Acids Research* 2000; **28**: e63.
9. Yu L-S, Rodriguez-Manzano J, Malpartida-Cardenas K *et al.* Rapid and sensitive detection of azole-resistant *Aspergillus fumigatus* by tandem-repeat loop-mediated isothermal amplification. *The Journal of Molecular Diagnostics* 2018; doi: <https://doi.org/10.1016/j.jmoldx.2018.10.004>
10. Swilaiman SS, O’Gorman CM, Balajee SA *et al.* Discovery of a Sexual Cycle in *Aspergillus lentulus*, a close relative of *A. fumigatus*. *Eukaryotic Cell* 2013; **12**: 962-69.
11. Waalwijk C, Mendes O, Verstappen EC *et al.* Isolation and characterization of the mating-type idiomorphs from the wheat septoria leaf blotch fungus *Mycosphaerella graminicola*. *Fungal Genetics and Biology* 2002; **35**: 277-86.
12. Gross A, Zaffarano PL, Duo A *et al.* Reproductive mode and life cycle of the ash dieback pathogen *Hymenoscyphus pseudoalbidus*. *Fungal Genetics and Biology* 2012; **49**: 977-86.
13. Böhm J, Hoff B, O’Gorman CM *et al.* Sexual reproduction and mating-type-mediated strain development in the penicillin-producing fungus *Penicillium chrysogenum*. *Proceedings of the National Academy of Sciences* 2013; **110**: 1476-481.
14. McDonald BA, Linde CC. Pathogen population genetics, evolutionary potential, and durable resistance. *Annual Review of Phytopathology* 2002; **40**: 349-79.

**Table 1** Primer sets used in the present study

<b>Purpose / Primer name</b>	<b>Primer sequence (5' – 3')</b>	<b>Source</b>
<b>New <i>MAT1-1</i>-specific LAMP assay:</b>		
AFMAT1F3	CGGTTGGCGATATCGTGAA	Present study
AFMAT1B3	GCCATCTGTCTCTTCAGGAG	
AFMAT1FIP	CAGCGAAGGCCATTGTGGAAGTTACTGGCTACGTGTCTGAGA	
AFMAT1BIP	ACGGCATTGAGATCACTGGCGCCACTTCAGGAGTTGCGAA	
AFMAT1LOOPF	TTGGTCCGTTTCGTGTGGC	
AFMAT1LOOPB	ACGATGCCATTGTGACTGAC	
<b>New <i>MAT1-2</i>-specific LAMP assay:</b>		
AFMAT2F3	CCCGTCTTGGGTAAGTGTCT	Present study
AFMAT2B3	GTGCGAAGGACTCAGTTACG	
AFMAT2FIP	CAACAGGTGCGCCAATGAGTGAGAGTTCCTCCTGAGCTTGA	
AFMAT2BIP	GCTCTCCGTGTTATGCGTACCCAGCTTCACCGTGAGATGC	
AFMAT2LOOPF	CACTGTCATTCCGTGTTATCGG	
AFMAT2LOOPB	CAGCTTTTTCCGGAACAGCT	
<b>Multiplex PCR mating type assay:</b>		
AFM1	CCTTGACGCGATGGGGTGG	2
AFM2	CGCTCCTCATCAGAACAACCTCG	
AFM3	CGGAAATCTGATGTCGCCACG	



**Table 2** Validation of new *Aspergillus fumigatus* MAT LAMP assays by screening of isolates from diverse environmental sources and geographic localities

Isolate	Source	Origin	MAT type (multiplex PCR) <sup>a</sup>	LAMP detection time (min)	
				MAT1-1 assay	MAT1-2 assay
47-255	Clinical	Europe	MAT1-1	10-11	Negative
47-257	Clinical	Europe	MAT1-1	9-10	Negative
47-258	Clinical	Europe	MAT1-1	9-10	Negative
47-2	Clinical	North America	MAT1-1	9-10	Negative
C6-UT1	Food	Asia	MAT1-1	15-16	Negative
C1-2-UT3	Food	South America	MAT1-1	9-10	Negative
C3-UT1	Food	South America	MAT1-1	9-10	Negative
C3-UT3	Food	South-America	MAT1-1	9-10	Negative
O5-5	Plant	Africa	MAT1-1	9-10	Negative
18-C6-9	Plant	Europe	MAT1-1	9-10	Negative
18-C7-8	Plant	Europe	MAT1-1	8-9	Negative
O9-8	Plant	Europe	MAT1-1	12-13	Negative
T4-1	Plant	Europe	MAT1-1	15-16	Negative
G4-1	Plant	South America	MAT1-1	8-9	Negative
O10-1	Plant	South America	MAT1-1	19-20	Negative
1-2.2-B1	Soil	Europe	MAT1-1	9-10	Negative
1-2.2-B2	Soil	Europe	MAT1-1	9-10	Negative
STNL1-B1	Soil	Europe	MAT1-1	9-10	Negative
STNL1-A8	Soil	Europe	MAT1-1	9-10	Negative
SWG1-A9	Soil	Europe	MAT1-1	12-13	Negative
BKCb-1	Air	Europe	MAT1-2	Negative	9-10
47-246	Clinical	Europe	MAT1-2	Negative	8-9
Af65	Clinical	Europe	MAT1-2	Negative	10-11
Af293	Clinical	Europe	MAT1-2	Negative	9-10
C5-T8	Food	Africa	MAT1-2	Negative	9-10
15-37-1	Food	Asia	MAT1-2	Negative	9-10
C1-1-T3	Food	South America	MAT1-2	Negative	9-10
C7-T2	Food	South America	MAT1-2	Negative	9-10
C7-UT1	Food	South America	MAT1-2	Negative	8-9
G2-2	Plant	Europe	MAT1-2	Negative	7-8
SWF5-C6	Soil	Europe	MAT1-2	Negative	7-8
PG1-5	Soil	Europe	MAT1-2	Negative	9-10
WSN19-3	Soil	Europe	MAT1-2	Negative	9-10
SWUK5-A9	Soil	Europe	MAT1-2	Negative	8-9

<sup>a</sup> Determined by mating multiplex PCR assay.<sup>2</sup>






Figure 1 Representative results from screening of *Aspergillus fumigatus* isolates with the multiplex PCR mating-type assay.2 *MAT1-1* and *MAT1-2* type isolates are distinguished by amplicons of 834 bp and 438 bp respectively. 'L' indicates Easyladder 1 (Bioline); NTC indicates no template control.

169x34mm (220 x 220 DPI)




Figure 2 Representative results from screening of *Aspergillus fumigatus* isolates of MAT1-1 (18-C6-9) or MAT1-2 (G2-2) type screened against the new MAT-specific LAMP assays. Shown are (A) amplification plots and (B) dissociation plots. See base of figure for explanatory legend. No-template (water) controls also gave negative results (data not shown).

170x260mm (300 x 300 DPI)

## First application of loop-mediated isothermal amplification (LAMP) assays for rapid identification of mating type in the heterothallic fungus *Aspergillus fumigatus*

K. M. King, N. J. Hawkins, S. Atkins, P. S. Dyer\*, J. S. West and B. A. Fraaije  
Rothamsted Research, Biointeractions and Crop Protection Department, Harpenden,  
Hertfordshire, AL5 2JQ, UK

\*School of Life Sciences, University of Nottingham, University Park, Nottingham NG7 2RD,  
UK

Corresponding author: Kevin King, Rothamsted Research, Harpenden, Hertfordshire, UK; +44  
(0)1582 763133; [Kevin.king@rothamsted.ac.uk](mailto:Kevin.king@rothamsted.ac.uk)

**Running title:** Fungal mating-type identification using LAMP

**Author contributions:** K.M.K., N.J.H., P.S.D., J.S.W., and B.A.F. conceived the ideas; K.M.K. and S.A. collected the data; K.M.K. analysed the data; K.M.K. led the writing; all authors critically reviewed the manuscript prior to submission.

### Summary

**Background** Loop-mediated isothermal amplification (LAMP) assays, which operate at a single temperature and require no post-reaction processing, have been described for rapid species-specific detection of numerous fungi. The technology has much less commonly been applied to identification of other key genetic traits such as fungicide resistance, and has not yet been applied to mating-type determination in any fungus.

**Objectives** To develop first LAMP assays for mating-type identification in a fungus, in this instance with the saprophytic mould and human opportunistic pathogen *Aspergillus fumigatus*, a heterothallic ascomycete requiring isolates of opposite mating type (*MAT1-1*, *MAT1-2*) for sexual reproduction.

**Methods** New LAMP primer sets, targeted to *MAT* gene sequences, were screened against 34 *A. fumigatus* isolates (of known mating type) from diverse clinical, environmental and geographic sources to establish if they could distinguish *MAT1-1* or *MAT1-2* genotypes.

**Results and conclusions** The new [assays/diagnostics](#), operating at a single temperature of 65°C, correctly identified the mating-type of *A. fumigatus* isolates in <20 minutes, and thus have numerous research and practical applications. Similar *MAT* LAMP assays could now be developed for other fungi of agricultural, environmental, industrial and/or medical importance.

**Keywords:** *Aspergillus fumigatus*, diagnostics, fungal pathogen, mating type, sexual reproduction

### Introduction

The fungus *Aspergillus fumigatus* is a saprophytic mould commonly found on plant debris and in soil. It is also an opportunistic human pathogen causing allergic symptoms and life-threatening invasive infections. The incidence of invasive aspergillosis (IA) has been increasing in recent years largely due to increased numbers of immunocompromised individuals in the population unable to fight off infection.<sup>1</sup> For more than 145 years, *A. fumigatus* was only known to reproduce asexually, although several signatures of cryptic sexuality were present, e.g. presence and expression of mating (*MAT*) genes and evidence of gene recombination within natural populations.<sup>2</sup> However, the breakthrough 2009 discovery of

a functional sexual cycle<sup>3</sup> had several implications including: (a) potentially explaining high genotypic diversity observed in populations; (b) production of sexually-derived airborne ascospores potentially more resilient to unfavourable environmental conditions; and (c) generation of sexual progeny with potentially greater pathogenicity and/or reduced sensitivity to fungicides.<sup>4</sup> *Aspergillus fumigatus* possesses a heterothallic (obligate outbreeding) mating system, with highly dissimilar stretches of DNA, termed ‘idiomorphs’, present in isolates of opposite mating type as is characteristic for heterothallic ascomycete species.<sup>5</sup> Thus, *MATI-1* isolates contain an alpha-domain *MATI-1-1* gene whereas *MATI-2* isolates contain a high-mobility group *MATI-2-1* gene together with a recently described *MATI-2-4* gene.<sup>6</sup> A multiplex PCR-based assay for determination of mating type has previously been developed for *A. fumigatus*.<sup>2</sup>

More recently, loop-mediated isothermal amplification (LAMP) assays have become increasingly used for rapid species-specific detection of numerous fungi, including *A. fumigatus*.<sup>7</sup> LAMP technology, first described by in 2000<sup>8</sup>, typically involves 4-6 primers in each reaction and has several purported advantages over PCR-based diagnostics. These include faster reaction times, potentially improved sensitivity and specificity, increased tolerance of sample inhibitors, no requirement for additional post-reaction processing (e.g. resolving PCR products on agarose gels) and use of only a single constant reaction temperature thus raising the possibility of field-based detection. Despite these advantages, LAMP assays have much less commonly been applied to detection of other key genetic traits such as fungicide resistance, one recent example being an assay targeted to a 34 bp tandem repeat in the *cyp51A* gene that has been associated with azole resistance in *A. fumigatus*.<sup>9</sup> To date, however, LAMP assays have not been used for rapid detection of different mating types in fungi. The objective of the present study was therefore to develop and evaluate for the first time whether LAMP assays could be used for the rapid identification of mating type in a fungus, with a focus here on the human opportunistic pathogen *A. fumigatus*.

## Methods

### Ethics statement

The authors confirm that the ethical policies of the journal, as noted on the journal’s author guidelines page, have been adhered to. No ethical approval was required as the research in this article related to micro-organisms.

### Fungal isolates, DNA extraction and initial molecular characterization

Details of *A. fumigatus* isolates, including source material and geographic origin, are given in Table 1; all isolates are maintained as -80°C glycerol stocks at Rothamsted Research, UK. Genomic DNA was extracted from *A. fumigatus* spores, harvested from one-week old cultures grown on Sabouraud dextrose agar (Lab M Ltd, UK) at 37°C, using a MasterPure yeast DNA purification kit (Epicentre, USA) into a final volume of 100 µL TE buffer. DNA was quantified via nanodrop spectrophotometer and diluted to 10 ng/ µL using PCR grade water. The mating type of these isolates was first determined using the published multiplex PCR assay<sup>2</sup>. Amplicons were resolved on agarose gels, with 834 bp or 438 bp products amplified from *MATI-1* or *MATI-2* isolates, respectively.

### Design and validation of MAT LAMP assays

For the *MATI-1* LAMP assay, *MAT* idiomorph sequence was downloaded from GenBank (Accession: AY898661<sup>2</sup>), with LAMP primers targeted to the internal *MATI-1-1* gene. For the *MATI-2* LAMP assay, *MAT* idiomorph sequence was sourced from the *A. fumigatus* Ensembl genome (isolate AF293; gene ID: AFUA\_3G06170), with LAMP primers targeted to the internal *MATI-2-1* gene. LAMP primer sets were designed using the free online software package PrimerExplorer (v. 5) with default settings.

For screening isolates against each of the *MAT* LAMP assays, 15  $\mu$ L reactions contained 0.3  $\mu$ L BIP primer (final concentration 2  $\mu$ M), 0.3  $\mu$ L FIP primer (2  $\mu$ M), 0.15  $\mu$ L LOOPB primer (1  $\mu$ M), 0.15  $\mu$ L LOOPF primer (1  $\mu$ M), 0.3  $\mu$ L B3 primer (0.2  $\mu$ M), 0.3  $\mu$ L F3 primer (0.2  $\mu$ M) (Table 1),  $\mu$ L 7.5  $\mu$ L isothermal mastermix (ISO-001; Optigene, UK) and 1  $\mu$ L DNA template (10 ng total DNA). No-template (PCR-grade water) controls were included in each test run. LAMP assays were run at 65°C for 30 minutes (FAM fluorescence measured every 30 secs), followed by a final dissociation step at 95°C for 1 min; 55°C for 30 secs and 95°C for 30 secs. Assays were run with a MX3000p qPCR system (Agilent), with data analysed using inbuilt 7500 SDS software (v.1.4; Applied Biosystems). Dissociation curves were checked manually after each run to confirm the presence of a single peak.

## Results

### Development and validation of *MAT* LAMP assays

For all *A. fumigatus* isolates tested, identical *MAT* genotype results were obtained using the previously described multiplex PCR assay<sup>2</sup> (see Figure 1 for representative results) and the new *MAT*-specific LAMP assays developed in the present study (Figure 1, Table 2). The new *MATI-1* and *MATI-2* -specific LAMP assays gave positive results within 10 – 20 mins (i.e. clear amplification curves) only for isolates of corresponding *MATI-1* or *MATI-2* type, respectively (Fig 2A). Positive results obtained with each *MAT*-specific LAMP assays gave single dissociation curves of c. 89.5°C (+0.3), indicating specific amplification of the targeted *MAT* gene regions (Fig 2B). No-template (water) controls tested negative, i.e. no amplification curves or dissociation plot peaks were observed (data not shown).

~~For all *A. fumigatus* isolates tested, identical *MAT* genotype results were obtained using the previously described multiplex PCR assay<sup>2</sup> (see Figure 1 for representative results) and the new *MAT*-specific LAMP assays developed in the present study (Table 2). For the *MATI-1* and *MATI-2* -specific LAMP assays, conclusive results were obtained with no requirement for gel electrophoresis within 20 mins and 10 mins, respectively. Positive results obtained with each *MAT*-specific LAMP assays gave single dissociation curves of 89.5°C ( $\pm$ 0.3), indicating specific amplification of the targeted *MAT* gene regions.~~

## Discussion

This study reports the first use of LAMP technology to rapidly (within 20 mins) establish the mating-type identity for a fungus, as demonstrated here for isolates of *A. fumigatus*. The *MATI-1* and *MATI-2* -specific LAMP assays appeared robust, being successfully applied to isolates of known opposite *MAT* type from a diverse range of clinical and environmental sources (air, food, plant and soil) and geographic locations (Africa, Asia, Europe and North and South America). These assays will be of use in research into the applied biology of this important human opportunistic pathogen. For example, they will allow the rapid set-up of sexual crosses with isolates of known opposite *MAT* type, subsequent analysis of the *MAT* type inheritance of

the progeny, and through progeny analysis the determination of the genetic basis of traits such as antifungal resistance and virulence.

It should now be possible to develop similar LAMP assays targeting *MAT* gene sequences to allow rapid mating-type determination in other heterothallic fungi of medical [e.g. *Aspergillus lentulus* - another causal agent of human aspergillosis<sup>10</sup>], agricultural [e.g. *Zymoseptoria tritici* - cause of wheat Septoria leaf blotch<sup>11</sup>], environmental [e.g. *Hymenoscyphus fraxineus* - cause of ash dieback<sup>12</sup>] and industrial [e.g. *Penicillium chrysogenum* - used in penicillin production<sup>13</sup>] importance. Such assays could also provide a better understanding into the reproductive strategies of various fungal pathogens, providing insight into their evolutionary potential and possible risk of breakdown of disease management strategies.<sup>14</sup> Furthermore, they could also be used to indirectly assess possible cryptic sexuality in fungi for which no sexual stage is yet known, given that frequency dependent selection operating on *MAT* genes generally, although not always, results in a 1:1 distribution of mating types.<sup>5,14</sup>

### Acknowledgements

These studies were partly supported by the Rothamsted Research Smart Crop Protection (SCP) strategic programme (BBS/OS/CP/000001) funded through the Biotechnology and Biological Sciences Research Council's Industrial Strategy Challenge Fund and by CropLife International through project 14209 'Investigating the sources and spread of azole resistant *Aspergillus fumigatus* strains in the environment'.

### Conflict of interest

No conflict of interest is declared.

### Figure legend

**Figure 1** Representative results from screening of *Aspergillus fumigatus* isolates with the multiplex PCR mating-type assay.<sup>2</sup> *MAT1-1* and *MAT1-2* type isolates are distinguished by amplicons of 834 bp and 438 bp respectively. 'L' indicates EasyLadder 1 (Bioline); NTC indicates no template control.

**Figure 2** Representative results from screening of *Aspergillus fumigatus* isolates of *MAT1-1* (18-C6-9) or *MAT1-2* (G2-2) type screened against the new *MAT*-specific LAMP assays. Shown are (A) amplification plots and (B) dissociation plots. See base of figure for explanatory legend. No-template (water) controls also gave negative results (data not shown).

### References

1. Dagenais TRT, Keller NP. Pathogenesis of *Aspergillus fumigatus* in invasive aspergillosis. *Clinical Microbiology Reviews* 2009; **22**: 447-65
2. Paoletti M, Rydholm C, Schwier EU *et al.* Evidence for sexuality in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Current Biology* 2005; **15**: 1242-248.
3. O'Gorman CM, Fuller HT, Dyer PS. Discovery of a sexual cycle in the opportunistic fungal pathogen *Aspergillus fumigatus*. *Nature* 2009; **457**: 471-74.
4. Zhang J, Snelders E, Zwaan BJ *et al.* A novel environmental azole resistance mutation in *Aspergillus fumigatus* and a possible role of sexual reproduction in its emergence. *mBio* 2017; **8**: pp.e00791-17.

5. Dyer PS, Inderbitzin P, Debuchy R, Mating-type structure, function, regulation and evolution in the Pezizomycotina. In: Wendland J, ed. *Growth, Differentiation and Sexuality, The Mycota I*. 3rd ed. Switzerland: Springer International Publishing 2016; 351-385.
6. Yu Y, Amich J, Will C *et al*. The novel *Aspergillus fumigatus* MAT1-2-4 mating-type gene is required for mating and cleistothecia formation. *Fungal Biology and Genetics* 2017; **108**, 1-12.
7. Tang Q, Shuguang T, Yu N *et al*. Development and evaluation of a loop-mediated isothermal amplification method for rapid detection of *Aspergillus fumigatus*. *Journal of Clinical Microbiology* 2016; **54**: 950-55.
8. Notomi T, Okayama H, Masubuchi H *et al*. Loop-mediated isothermal amplification of DNA. *Nucleic Acids Research* 2000; **28**: e63.
9. Yu L-S, Rodriguez-Manzano J, Malpartida-Cardenas K *et al*. Rapid and sensitive detection of azole-resistant *Aspergillus fumigatus* by tandem-repeat loop-mediated isothermal amplification. *The Journal of Molecular Diagnostics* 2018; doi: <https://doi.org/10.1016/j.jmoldx.2018.10.004>
10. Swilaiman SS, O’Gorman CM, Balajee SA *et al*. Discovery of a Sexual Cycle in *Aspergillus lentulus*, a close relative of *A. fumigatus*. *Eukaryotic Cell* 2013; **12**: 962-69.
11. Waalwijk C, Mendes O, Verstappen EC *et al*. Isolation and characterization of the mating-type idiomorphs from the wheat septoria leaf blotch fungus *Mycosphaerella graminicola*. *Fungal Genetics and Biology* 2002; **35**: 277-86.
12. Gross A, Zaffarano PL, Duo A *et al*. Reproductive mode and life cycle of the ash dieback pathogen *Hymenoscyphus pseudoalbidus*. *Fungal Genetics and Biology* 2012; **49**: 977-86.
13. Böhm J, Hoff B, O’Gorman CM *et al*. Sexual reproduction and mating-type-mediated strain development in the penicillin-producing fungus *Penicillium chrysogenum*. *Proceedings of the National Academy of Sciences* 2013; **110**: 1476-481.
14. McDonald BA, Linde CC. Pathogen population genetics, evolutionary potential, and durable resistance. *Annual Review of Phytopathology* 2002; **40**: 349-79.