

Rothamsted Repository Download

A - Papers appearing in refereed journals

Bird, I. F., Cornelius, M. J., Keys, A. J. and Whittingham, C. P. 1972. Adenosine triphosphate synthesis and the natural electron acceptor for synthesis of serine from glycine in leaves (ATP). *Biochemical Journal*. 128 (1), pp. 191-192.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1042/bj1280191>

The output can be accessed at:

<https://repository.rothamsted.ac.uk/item/96z7x/adenosine-triphosphate-synthesis-and-the-natural-electron-acceptor-for-synthesis-of-serine-from-glycine-in-leaves-atp>.

© Please contact library@rothamsted.ac.uk for copyright queries.

Adenosine Triphosphate Synthesis and the Natural Electron Acceptor for Synthesis of Serine from Glycine in Leaves

By I. F. BIRD, M. J. CORNELIUS, A. J. KEYS and C. P. WHITTINGHAM
Rothamsted Experimental Station, Harpenden, Herts., U.K.

(Received 13 March 1972)

Mitochondria from the leaves of higher plants (Kisaki *et al.*, 1971; Bird *et al.*, 1971) and from rat and avian livers (Sanadi & Bennett, 1960; Kawasaki *et al.*, 1966) catalyse a reaction between two molecules of glycine to form one molecule each of serine, CO₂ and NH₃. Kisaki *et al.* (1971), using mitochondria from leaves of *Spinacia oleracea*, found that the reaction was faster in N₂ than in air and that NAD⁺ was necessary as the electron acceptor. By contrast, the reaction catalysed by mitochondria from leaves of *Nicotiana tabacum* required aerobic conditions, was accompanied by the synthesis of ATP and ADP, and used one atom of O₂ for each molecule of serine formed (Bird *et al.*, 1972). The affinity of the reacting system for O₂ was high (Fig. 1). The reaction was inhibited by inhibitors of the electron-transport system in mitochondria and was little affected by added NAD⁺. The immediate acceptor of electrons arising from conversion of glycine into serine is therefore linked to part of the electron-transport system of mitochondria.

The rate of reaction was measured by using [1-¹⁴C]glycine as the substrate. The ¹⁴CO₂ evolved was absorbed in 2M-NaOH and measured by liquid-scintillation counting. Reaction vessels were shaken throughout the reaction period. The ADP phosphorylated was measured by a modification (Nobel, 1967) of the method of Nielsen & Lehninger (1955). Leaves were obtained from tobacco plants (*Nicotiana tabacum* var. White Burley) kept in darkness for 24 h. Leaf lamina was ground with 4 vol. of medium (pH 7.7) containing 0.4M-sucrose, 20mM-sodium citrate, 0.2M-tris-HCl, 10mM-KH₂PO₄, 5mM-EDTA and 1% (w/v) of bovine serum albumin. Coarse debris was removed by squeezing through two layers of muslin and filtering through cottonwool. The filtrate was centrifuged (100g for 10min) and the pellet discarded. The supernatant liquid was centrifuged (38000g for 10min) and the pellet obtained was resuspended in an appropriate volume of medium (pH 7.7) containing 0.4M-sucrose, 20mM-sodium citrate, 4mM-MgCl₂, 10mM-KH₂PO₄ and 1% (w/v) of bovine serum albumin.

Ferricyanide could act as the final electron acceptor instead of O₂ (Table 1). Serine synthesis was strongly inhibited by antimycin A whether O₂ (Bird *et al.*, 1972) or ferricyanide (Table 1, Expt. A) was the final electron acceptor. Amytal (2mM) inhibited

serine synthesis by less than 20%. With liver mitochondria oxidation of substrates linked through NAD⁺ to the electron-transport chain is completely inhibited by 1.8mM-Amytal (Jailing *et al.*, 1955); measurements made with an oxygen electrode (Rank Bros.), with pyruvate as substrate, confirmed that this is also true for tobacco leaf mitochondria. Therefore oxidation associated with the synthesis of serine from glycine does not proceed through the internal pool of NAD⁺ (Lehninger, 1951) in mitochondria, and coupling site I of the electron-transport system is not involved. Ferricyanide accepts electrons between coupling sites II and III (Lee *et al.*, 1967) and should halve phosphorylation where oxidations involve only phosphorylation sites II and III. Table 1, Expt. B, shows that in the presence of ferricyanide




Fig. 1. Rate of serine synthesis at various oxygen concentrations

Reaction mixtures (pH 7.7; 0.66 ml) contained particles from 0.1 g of leaf, 1 μmol of NAD⁺, 0.1 μmol of pyridoxal phosphate, 2 μmol of ADP, 10 μmol (0.5 μCi) of [1-¹⁴C]glycine, 200 μmol of sucrose, 10 μmol of sodium citrate, 5 μmol of KH₂PO₄, 2 μmol of MgCl₂ and 5 mg of bovine serum albumin. The reaction vessels were evacuated and filled with measured volumes of O₂ and N₂ to give atmospheres with the O₂ content shown. During the reaction the vessels were kept on a mechanical shaker at 25°C.

Table 1. *Effects of Amytal, ferricyanide and antimycin A on serine synthesis from glycine and accompanying phosphorylation of ADP*

Reaction mixtures were as described in Fig. 1 except for the addition, where indicated, of antimycin A (6.67nmol), $K_3Fe(CN)_6$ (2 μ mol) and Amytal (1.33 μ mol) or the omission of [1- ^{14}C]glycine.

Atmosphere in the reaction vessel	Additions to reaction mixtures	Rate of synthesis (μ mol/h per g fresh wt. of leaf)		
		Serine	ATP	ATP increase caused by the presence of glycine
Expt. A				
Air	—	4.32	—	—
Air	$K_3Fe(CN)_6$	5.28	—	—
Air	Amytal	3.66	—	—
N ₂	—	0.50	—	—
N ₂	$K_3Fe(CN)_6$	5.80	—	—
N ₂	Amytal	0.60	—	—
N ₂	Amytal + $K_3Fe(CN)_6$	4.70	—	—
N ₂	Antimycin A	0.18	—	—
N ₂	Antimycin A + $K_3Fe(CN)_6$	0.46	—	—
Expt. B				
Air	Glycine absent	—	6.65	—
Air	—	4.56	12.21	5.56
N ₂	Glycine absent	—	3.26	—
N ₂	—	0.92	2.81	-0.45
N ₂	$K_3Fe(CN)_6$, glycine absent	—	5.51	—
N ₂	$K_3Fe(CN)_6$	4.18	8.50	2.99

in N₂ the increase in phosphorylation caused by the presence of glycine was only 54% of that where O₂ was the final electron acceptor.

The results confirm our view that NAD⁺ reduction is not important during serine synthesis from glycine in leaves, but that the electron-transport system of mitochondria coupled to phosphorylation sites II and III is responsible for the oxidation involved and that two molecules of ATP are incidentally synthesized for each molecule of serine formed.

Bird, I. F., Keys, A. J. & Whittingham, C. P. (1971) *Proc. Int. Congr. Photosynth. Res. 2nd.*, *Stresa* in the press
 Bird, I. F., Cornelius, M. J., Keys, A. J. & Whittingham, C. P. (1972) *Phytochemistry* **11**, 1587-1594

Jailing, O., Lindberg, O. & Ernster, L. (1955) *Acta Chem. Scand.* **9**, 198-199
 Kawasaki, H., Sato, T. & Kikuchi, G. (1966) *Biochem. Biophys. Res. Commun.* **23**, 227-233
 Kisaki, T., Imai, A. & Tolbert, N. E. (1971) *Plant Cell Physiol.* **12**, 267-273
 Lee, C. P., Sottocasa, G. L. & Ernster, L. (1967) *Methods Enzymol.* **10**, 33-37
 Lehninger, A. L. (1951) *Phosphorus Metabolism* (McElroy, W. D. & Glass, B., eds.), vol. 1, pp. 344-366, The Johns Hopkins Press, Baltimore
 Nielsen, S. O. & Lehninger, A. L. (1955) *J. Biol. Chem.* **215**, 555-570
 Nobel, P. S. (1967) *Plant Physiol.* **42**, 1389-1394
 Sanadi, D. R. & Bennett, M. J. (1960) *Biochim. Biophys. Acta* **39**, 367-369