

Rothamsted Repository Download

A - Papers appearing in refereed journals

Chen, G. Q., Van-Erp, H., Martin-Moreno, J., Johnson, K., Morales, E., Browse, J., Eastmond, P. J. and Lin, J-T. 2016. Expression of castor LPAT2 enhances ricinoleic acid content at the sn-2 position of triacylglycerols in lesquerella seed. *International Journal Of Molecular Sciences*. 17 (4), p. 507.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.3390/ijms17040507>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/8v21v>.

© 6 April 2016. Licensed under the Creative Commons CC BY.



Article

Expression of Castor LPAT2 Enhances Ricinoleic Acid Content at the *sn*-2 Position of Triacylglycerols in Lesquerella Seed

Grace Q. Chen ^{1,*}, Harrie van Erp ^{2,3,†}, Jose Martin-Moreno ³, Kumiko Johnson ¹, Eva Morales ¹, John Browse ², Peter J. Eastmond ³ and Jiann-Tsyh Lin ¹

¹ Western Regional Research Center, Agricultural Research Service, U.S. Department of Agriculture, 800 Buchanan St., Albany, CA 94710, USA; kumiko.johnson@ars.usda.gov (K.J.); evadesalta@gmail.com (E.M.); jiann.lin@ars.usda.gov (J.-T.L.)

² Institute of Biological Chemistry, Washington State University, Pullman, Washington, DC 99164, USA; harrie.van-erp@rothamsted.ac.uk (H.E.); jab@wsu.edu (J.B.)

³ Department of Plant Biology and Crop Science, Rothamsted Research, Harpenden, Hertfordshire AL5 2JQ, UK; jmartinm83@gmail.com (J.M.-M.); peter.eastmond@rothamsted.ac.uk (P.J.E.)

* Correspondence: grace.chen@ars.usda.gov; Tel.: +1-510-559-5627

† These authors contributed equally to this work.

Academic Editor: Ute Roessner

Received: 4 March 2016; Accepted: 30 March 2016; Published: 6 April 2016

Abstract: Lesquerella is a potential industrial oilseed crop that makes hydroxy fatty acid (HFA). Unlike castor its seeds are not poisonous but accumulate lesquerolic acid mostly at the *sn*-1 and *sn*-3 positions of triacylglycerol (TAG), whereas castor contains ricinoleic acid (18:1OH) at all three positions. To investigate whether lesquerella can be engineered to accumulate HFAs in the *sn*-2 position, multiple transgenic lines were made that express castor lysophosphatidic acid acyltransferase 2 (*RcLPAT2*) in the seed. *RcLPAT2* increased 18:1OH at the *sn*-2 position of TAGs from 2% to 14%–17%, which resulted in an increase of tri-HFA-TAGs from 5% to 13%–14%. Our result is the first example of using a LPAT to increase ricinoleic acid at the *sn*-2 position of seed TAG. This work provides insights to the mechanism of HFA-containing TAG assembly in lesquerella and directs future research to optimize this plant for HFA production.

Keywords: lesquerella; *Physaria fendleri*; lysophosphatidic acid acyltransferase; hydroxy fatty acid; seed triacylglycerol; metabolic engineering

1. Introduction

Lesquerella (*Physaria fendleri*, formerly *Lesquerella fendleri* (Gray) Wats.) (Brassicaceae) [1] is valued for the unusual hydroxy fatty acids (HFAs) in its seed oil. The major HFA in lesquerella is lesquerolic acid (20:1OH) [2], comprising 55%–60% of total seed fatty acids (FAs) [3–8]. The main source of HFAs is castor (*Ricinus communis*), 90% of its seed oil is ricinoleic acid (18:1OH). 18:1OH is used as a feedstock for the manufacture of a range of products, such as lubricants, plasticizers and surfactants [9]. However, castor cultivation is limited by the presence of the toxin ricin in its seed. Lesquerella seeds do not contain ricin and therefore its oil provides a safe source of HFAs. Efforts have been made to improve lesquerella through plant breeding [4]. Since lesquerella can be transformed by *Agrobacterium* [10], genetic engineering also provides an alternative means to improve this potential crop.

Seed oils or triacylglycerols (TAGs) are storage lipids in which FAs are esterified to each of the three hydroxy groups of a glycerol backbone. FAs are synthesized in the plastid, exported and converted to acyl-coenzyme As (acyl-CoAs) for the synthesis of TAG in the endoplasmic reticulum (ER) [11,12]. The synthesis of TAG requires the Kennedy pathway [12–15] (Figure 1). The first acylation reaction is

catalyzed by a glycerol-3-phosphate acyltransferase (GPAT) and yields lysophosphatidic acid (LPA), which in turn is acylated by a lysophosphatidic acid acyltransferase (LPAT) to produce phosphatidic acid (PA). PA is then converted to 1,2-*sn*-diacylglycerol (DAG) by PA phosphatase (PAP). Finally, a third FA is esterified to the *sn*-3 position of DAG by 1,2-*sn*-diacylglycerol acyltransferase (DGAT) to produce TAG. Alternatively, FAs can be directly incorporated into phosphatidylcholine (PC) by the reactions of the acyl editing cycle [16,17] (Figure 1). These reactions are catalyzed by the forward and reverse reactions of lyso-PC acyltransferase (LPCAT) to yield acyl-CoA, or by a phospholipase A-type activity to yield a free FA that then is activated to acyl-CoA. Because PC is the substrate for FA-modifying enzymes such as desaturases and hydroxylases, rapid de-acylation and re-acylation of PC causes the acyl-CoA pool to be enriched with modified FAs (mFA), which can subsequently be used for TAG synthesis. FAs can also be acylated to the *sn*-3 position of DAG by Phospholipid: DAG acyltransferase (PDAT) [18]. In addition, metabolic labeling experiments showed that there are two pools of DAG, *de novo* DAG and PC-derived DAG [19,20] (Figure 1). The *de novo* synthesis of DAG involves the Kennedy pathway, as described above. In the PC-derived DAG pathway, *de novo* DAG is used to synthesize PC. Removal of the head group of PC by PC:DAG cholinephosphotransferase (PDCT) can generate PC-derived DAG [21,22] (Figure 1). Alternatively, PC-derived DAG can be produced by the reverse action of CDP-choline: DAG cholinephosphotransferase (CPT) [23], or a lipase-based mechanism using phospholipase C, or phospholipase D plus PAP. As FAs in PC can be modified, the conversion of PC into DAG also provides a means to increase the amount of mFAs in TAG.




Figure 1. Proposed pathways for TAG biosynthesis in lesquerella seed. Blue arrows indicate reactions involved in Kennedy pathway. Purple arrows indicate reactions involved in acyl editing. Green arrows indicate reactions involved in PC-derived DAG synthesis. The rest of reactions are indicated by black arrows. Enzymes catalyzing these reactions are underlined. Fatty acid numerical symbols are described as in Table 2. Abbreviations: A, DAG production by the reverse reaction of CPT, phospholipase C or phospholipase D plus PAP; G3P, glycerol-3-phosphate; LPA, lysophosphatidic acid; PA, phosphatidic acid; LPC, lysophosphatidylcholine; PC, phosphatidylcholine; DAG, diacylglycerol; TAG, triacylglycerol; GPAT, glycerol 3-phosphate acyltransferase; LPAT, lysophosphatidic acid acyltransferase; PAP, phosphatidic acid phosphatase; FAH12, Δ^{12} oleic acid hydroxylase; FAD2, Δ^{12} oleic acid desaturase; FAD3, Δ^{15} (ω -3) linoleic acid desaturase; KCS3, 3-ketoacyl-CoA synthase 3; LPCAT, lysophosphatidylcholine acyltransferase; CPT, CDP-choline:DAG cholinephosphotransferase; PDCT, PC:DAG cholinephosphotransferase; DGAT, diacylglycerol acyltransferase.

Castor TAG contains 90% 18:10H, over 70% of the TAGs having all three *sn* positions esterified with 18:10H [24]. The 18:10H is synthesized by hydroxylation of 18:1, which is esterified to the *sn*-2 position of PC, efficiently released and activated to 18:10H-CoA, and then redistributed into the *sn*-1, *sn*-2, and *sn*-3 positions of TAG [25,26]. Biochemical evidence indicates that in lesquerella, 18:10H is initially synthesized by hydroxylation of 18:1 at the *sn*-2 position of PC, released and

activated to 18:1OH-CoA. Subsequently, 18:1OH-CoA is rapidly elongated to 20:1OH-CoA for TAG synthesis [27,28]. The lesquerella gene *PfKCS3* responsible for the elongation of 18:1OH-CoA to 20:1OH-CoA has been identified [29] (Figure 1). In addition to 18:1OH and 20:1OH, lesquerella seed accumulates a low level of densipolic acid (18:2OH) and auricollic acid (20:2OH), which were suggested to be synthesized by a microsomal Δ -15 desaturase [27,28].

Arabidopsis, which does not accumulate HFAs, has been used as a model to study genes associated with HFA synthesis [30]. A key gene encoding the oleate 12-hydroxylase (FAH12) was first isolated from castor (*CFAH12* or *RcFAH12*) [31], and then from lesquerella (*PfFAH12*) [32]. Expression of *RcFAH12* or *PfFAH12* in Arabidopsis led to accumulation of HFAs to no more than 17% in seed oil [32–36]. To boost HFA production, additional genes from castor have been co-expressed with *RcFAH12* in Arabidopsis, including *RcDGAT2* [37], *RcPDAT1A* [38], or *RcPDAT1-2* [39], and *RcPDCT* [21]. The overexpression of *RcDGAT2* and *RcPDAT1A* resulted in an increase in seed HFA content from 17% to 23%–28% [38]. Co-expression of castor DGAT, PDAT and PDCT, while simultaneously suppressing the expression of endogenous acyltransferases, may increase the flux of HFA into TAG through more efficient utilization of HFA-containing substrates, such as HFA-CoA, HFA-DAG, and HFA-PC, in their pathways [20,21,38–40].

Lesquerella TAGs contain ~60% 20:1OH and almost all of it is esterified to the *sn*-1 and *sn*-3 positions [3,7] (Figure 1). The absence of HFAs at the *sn*-2 position of TAG might be caused by the selectivity of LPATs in lesquerella for common FAs. This is supported by the fact that LPAT enzymes from plants are some of the most stringent acyltransferases regarding substrate discrimination [41]. Based on several studies of plant microsomal (putative ER-localized) LPATs, Frentzen (1998) proposed the presence of at least two distinct classes of LPATs, class-A and -B, sharing little sequence homology between them. The class-A LPATs had selectivity for unsaturated C₁₈ acyl groups, and they are ubiquitously expressed in higher plants for membrane lipid and TAG synthesis [41]. The class-B LPATs are present in *Limnanthes* [42] and *Cocos* [43]. Their seed oils are enriched with the unusual FAs (UFAs), erucic acid (22:1) and lauric acid (12:0), respectively, at the *sn*-2 position of TAGs. Their corresponding members of class-B LPATs possessed substrate preference for UFAs and were used to improve FA composition in industrial oilseeds by genetic engineering [41,44–46]. Interestingly, a yeast LPAT sharing little homology to plant LPATs also has preference for esterifying unusual very long chain FAs at the *sn*-2 position of seed TAGs [47]. Most plant species do not have a representative of the class-B gene, including Arabidopsis where five LPATs have been identified [48,49]. Arabidopsis LPAT2 (*AtLPAT2*) is the lone gene that encodes the ubiquitous and ER-located LPAT, and is thus suggested to be the key enzyme involved in membrane and seed oil biosynthesis [49]. Recently, a highly seed-expressed LPAT was identified from *Sterculia foetida* whose seeds accumulate up to 60% of desaturated cyclopropane FA [50]. *SfLPAT* showed strong homology to *AtLPAT2* and enabled the incorporation of the unusual cyclopropane FA at the *sn*-2 position of LPA in transgenic Arabidopsis [50]. In castor, whole genome data [51] allowed the identification and analysis of the LPAT multigene family [51–55]. Among them, two genes, castor lysophosphatidic acid acyltransferase 2 (*RcLPAT2*) and castor lysophosphatidic acid acyltransferase B (*RcLPATB*), representing class-A and -B respectively, were found to express highly in seed tissues and thus were characterized in more detail [55]. Spatial and temporal gene expression profiling indicated that *RcLPAT2* exhibited a generalized constitutive pattern typical to that of class-A members. *In vitro* enzyme assays indicated that *RcLPAT2* had a preference for 18:1OH-CoA when *sn*-1-ricinoleate-*sn*-glycerol-3-phosphate (18:1OH-LPA) was used as an acyl acceptor, whereas when *sn*-1-oleoyl-*sn*-glycerol-3-phosphate (18:1-LPA) was used as the acyl-acceptor, *RcLPAT2* preferred 18:1-CoA as a substrate [55]. These results suggest that *RcLPAT2* might participate in both TAG and membrane glycerolipid synthesis. However, unlike the LPATBs from *Limnanthes* and *Cocos*, the expression of *RcLPATB* was not restricted to seeds as it was also expressed at a significant level in leaf, stem and floral organs. However, *in vitro* assays indicated that *RcLPATB* had a low enzyme activity and a broad specificity for acyl-CoAs compared with that of *RcLPAT2*. *RcLPATB* catalyzed the incorporation of saturated FAs, 12:0–16:0, at rates surpassing those of monounsaturated FAs including

18:1-CoA and 18:1OH-CoA, independently of the acceptors 18:1-LPA or 18:1OH-LPA [55]. Therefore, it remains questionable if RcLPATB is involved in HFA-containing TAG synthesis. We note that before the completion of castor genome sequencing, expression of a castor gene (GenBank ID, EU391594) designated as a “putative castor *LPAT*” [37] did not change HFA composition in transgenic *Arabidopsis* expressing *RcFAH12*. The sequence (EU391594) shows no significant similarity to RcLPAT2 (GenBank ID, JQ796917) and is now annotated as a putative castor 1-acyl-*sn*-glycerol-3-phosphate acyltransferase zeta precursor [51] (GenBank ID, XP_002525812).

Castor oil has long been used by industry as a raw material for manufacturing numerous established products. If lesquerella oil can be reengineered to resemble castor oil, this would provide an alternative source of castor oil that is safe, cost-competitive, and readily adaptable by existing industrial technologies. Increasing the levels of tri-HFA-TAG and 18:1OH, are key steps towards achieving this goal. We therefore over-expressed RcLPAT2 in lesquerella seeds. Analysis of seed TAGs suggests that RcLPAT2 is able to acylate the *sn*-2 position of LPA with 18:1OH and as a result the amounts of 18:1OH and tri-HFA-TAGs were increased. Additional genetic modifications for engineering a castor-oil producing lesquerella crop are discussed. To our knowledge this is the first time that castor *RcLPAT2* has been expressed in another oilseed to modify HFA content and also the first example of metabolic engineering in lesquerella.

2. Results

2.1. Castor *Lysophosphatidic Acid Acyltransferase 2* (*RcLPAT2*) Increased 18:1OH Accumulation in Transgenic Seeds

We have produced selfed T₁ seeds from 16 independent primary transgenic plants (T₀); expressing *RcLPAT2* under the control of the seed-specific napin (*napA*) promoter. Analysis of transgene *RcLPAT2* copy number in T₀ plants indicated that 12 of them have one copy and four have two copies (Table 1). Among the lines, the FA composition exhibited some variations in each non-HFA (Table S1) as well as HFA (Table 1), which can be explained by the heterozygous nature of lesquerella, as it is nearly a wild species, and similar seed-to-seed variations were reported [5]. However, the average levels of each non-HFAs, palmitic (16:0), palmitoleic (16:1), stearic (18:0), oleic (18:1), linoleic (18:2), linolenic (18:3), and eicosenoic (20:1) acids, from transgenic lines are similar to that of wild type (wt) seeds (Table S1). Interestingly, all transgenic lines had increased 18:1OH content ranging from 1.35%–4.08% compared with that of wt at 1.20% (Table 1). Among them, four T₁ seed populations (lines 1, 3, 4 and 5) showed relative high 18:1OH contents from 3.79%–4.08%. The correlation between 18:1OH content and transgene *RcLPAT2* copy number is weak ($r = 0.1$, $p = 0.713$). About 63% (10/16) of the lines also showed an increase in 18:2OH (Table 1), despite the overall low contents of 18:2OH (0.15%–0.75%) in transgenic lines and wt (0.24%) (Table 1). The correlation between 18:1OH and 18:2OH contents was strong ($r = 0.91$, $p < 0.00001$). With the increase of 18:1OH, we observed a small decrease in 20:1OH levels showing an average of 50.03% compared to 51.50% in wt (Table 1). For 20:2OH, its average level was 3.22%, which is slightly higher than the 3.07% measured for wt (Table 1). Total HFA levels varied from 52.56% (line 8) to 60.44% (line 9) among transgenic lines with an average of 56.08% which is comparable to 56.18% of wt (Table 1). To examine the inheritance of the *RcLPAT2* transgene, we selected hygromycin resistant T₁ seedlings and examined *RcLPAT2* copy number in those relatively high 18:1OH T₁ populations. We identified two homozygous lines, 3-1 and 4-5, and produced selfed T₂ seeds for further analysis. For FA composition, we observed profiles similar to that of T₁ seeds (Table 2). We note that the 18:2OH content in wt was 0.24% in Table 1 and 0.02% in Table 2. The discrepancy was likely caused by the heterozygous nature of lesquerella seeds and by small sample size used for the analysis (10 seeds per sample for Table 1, and 15 seeds per sample for Table 2). The increased 18:1OH accumulation phenotype in lines 3-1 and 4-5 was inherited into the T₂ generation, although the levels of 18:1OH in homozygous lines increased only to 5.13% and 4.72%, respectively (Table 2). In general, there was no obvious alteration on growth phenotype in transgenic lines associated with the expression of *RcLPAT2*.

Table 1. RclPAT2 copy number in primary transgenic lines and hydroxy fatty acid (HFA) composition of T₁ seeds.

HFA Content		18:1OH	18:2OH	20:1OH	20:2OH	Total HFA
Wild type		1.20 ± 0.09	0.24 ± 0.06	51.74 ± 0.98	3.07 ± 0.39	56.18 ± 0.48
Transgenics ID	Copy number					
Line 1	2	3.79 ± 0.04 ***	0.53 ± 0.04 **	45.64 ± 1.32 *	3.11 ± 0.15	53.08 ± 1.46
Line 2	1	1.74 ± 0.08 *	0.30 ± 0.08	51.43 ± 0.50	3.45 ± 0.21	56.91 ± 0.33
Line 3	1	4.08 ± 0.31 ***	0.74 ± 0.12 *	48.99 ± 0.69	3.60 ± 0.11	57.81 ± 0.34
Line 4	2	3.81 ± 0.06 ***	0.60 ± 0.03 **	50.02 ± 0.48	3.38 ± 0.24	57.81 ± 0.34
Line 5	1	3.97 ± 0.13 ***	0.75 ± 0.20 *	48.35 ± 1.88	3.58 ± 0.31	56.64 ± 1.27
Line 7	1	3.29 ± 0.12 ***	0.34 ± 0.03	51.09 ± 0.22	2.84 ± 0.15	57.56 ± 0.21
Line 8	1	1.42 ± 0.08 **	0.15 ± 0.07	48.24 ± 1.41	2.75 ± 0.20	52.56 ± 1.26
Line 9	1	2.87 ± 0.21 ***	0.53 ± 0.07 *	53.51 ± 0.33	3.53 ± 0.31	60.44 ± 0.36 **
Line 10	1	2.66 ± 0.12	0.43 ± 0.06 *	48.98 ± 1.11	3.09 ± 0.06	55.17 ± 1.00
Line 11	1	1.42 ± 0.08	0.23 ± 0.04	49.57 ± 0.46	3.28 ± 0.09	54.50 ± 0.31 *
Line 12	1	1.35 ± 0.05	0.15 ± 0.03	51.40 ± 1.75	3.26 ± 0.26	56.15 ± 1.58
Line 13	1	2.08 ± 0.11 **	0.29 ± 0.08	47.43 ± 2.07	3.20 ± 0.49	53.00 ± 2.07
Line 14	1	1.97 ± 0.07 **	0.14 ± 0.07	53.09 ± 0.82	2.00 ± 0.13	57.19 ± 0.75
Line 15	1	1.99 ± 0.10 **	0.24 ± 0.13	52.62 ± 0.70	2.93 ± 0.26	57.77 ± 0.47
Line 16	2	1.48 ± 0.04	0.28 ± 0.03	48.96 ± 1.07	3.94 ± 0.10	54.66 ± 1.12
Line 17	2	1.42 ± 0.04	0.23 ± 0.04	51.18 ± 0.59	3.61 ± 0.26	56.44 ± 0.41
Average among lines		2.46 ± 0.26	0.37 ± 0.05	50.03 ± 0.54	3.22 ± 0.11	56.08 ± 0.54

Each HFA composition was measured as percentage of total fatty acids. Triplicates of 10-seed samples were measured for wild type and each independent transgenic line. All data are averages of three measurements ± SE. Fatty acid legend: 18:1OH is ricinoleic; 18:2OH is densipolic; 20:1OH is lesquerolic; and 20:2OH is auricolic acid. Two-tailed student's *t*-test. * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001.

Table 2. Fatty acid (FA) composition of homozygous T₂ seeds.

FA	Wild Type	Line 3-1	Line 4-5
16:0	2.75 ± 0.08	3.51 ± 0.28	2.87 ± 0.07
16:1	1.16 ± 0.10	1.02 ± 0.09	1.00 ± 0.17
18:0	3.42 ± 0.83	3.31 ± 0.17	3.43 ± 0.20
18:1	15.19 ± 0.04	14.62 ± 0.30	14.96 ± 0.24
18:2	8.07 ± 0.28	8.43 ± 0.35	8.78 ± 0.48
18:3	13.90 ± 0.05	11.56 ± 0.65	11.40 ± 0.64
20:1	0.88 ± 0.12	0.37 ± 0.03 *	0.68 ± 0.21
18:1OH	1.43 ± 0.07	4.27 ± 0.10 ***	4.23 ± 0.41 **
18:2OH	0.02 ± 0.01	0.86 ± 0.14 *	0.49 ± 0.25
20:1OH	50.35 ± 0.59	47.82 ± 0.89	47.20 ± 0.62 *
20:2OH	2.63 ± 0.18	3.17 ± 0.12	3.50 ± 0.18 *

Each FA was measured as percentage of total FA. Triplicates of 15-seed sample were measured for wild type and transgenic lines. All data are averages of three measurements ± SE. Fatty acid legend: 16:0 is palmitic; 16:1 is palmitoleic; 18:0 is stearic; 18:1 is oleic; 18:2 is linoleic; 18:3 is linolenic; 20:1 is eicosenoic acid; 18:1OH is ricinoleic; 18:2OH densipolic; 20:1OH is lesquerolic; 20:2OH is auricolic acid. Two-tailed student's *t*-test. * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001.

2.2. Overexpression of RclPAT2 Altered the Composition of Hydroxy Fatty Acid (HFA)-Containing Triacylglycerol (TAG) Species and Increased the Percentage of 18:1OH at the sn-2 Position

There are four possible molecular species of TAG based on the number of HFA chains esterified to the glycerol backbone: 0-, 1-, 2-, and 3-HFA-TAG. To investigate if there is any change in TAG species composition in the transgenic seeds, we determined the relative amount of each TAG species as a percentage of total TAGs in wt and transgenic lines 3-1 and 4-5. In wt, the relative amount of 0-, 1-, 2-, and 3-HFA-TAG was 4.79%, 12.17%, 79.13% and 4.79% respectively (Figure 2), which are comparable to reported levels [3,56]. In both transgenic lines, there are increases in 0-, 1-, and 3-HFA-TAGs showing 6.13%, 26.06%, and 14.38% in line 3-1 and 8.01%, 22.64%, and 12.93% in line 4-5, respectively (Figure 2). However, we observed that 2-HFA-TAG decreased 29%–33%, showing 53.43% and 56.41% in lines

3-1 and 4-5, respectively (Figure 2). Regiochemical analysis was performed for lines 3-1 and 4-5 to determine whether overexpression of *RcLPAT2* increases the amount of HFAs at the *sn*-2 position. As shown in Figure 3, *sn*-2 18:1OH content increased from 2.19% to 17.29% or 14.85% in lines 3-1 and 4-5 respectively. Only very small increases were observed for *sn*-2 20:1OH from 0.93% to 1.64% or 1.53% in lines 3-1 and 4-5 respectively.




Figure 2. Molecular species composition of TAG. TAG molecular species in seeds of wild type (wt) (open bars) and transgenic line 3-1 (black bars), line 4-5 (grey bars) are measured as a weight percent of total TAG. 0-, 1-, 2-, and 3-HFA represent TAG molecular species with zero, one, two, or three HFAs, respectively (no stereochemistry implied). Triplicates of 50-seed sample were measured for wild type and transgenic lines. The HFAs represent the sum of 18:1OH and 20:1OH. FA numerical symbols are described as in Table 2. The data represent averages of three replicates \pm SE. Two-tailed Student's *t* test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.




Figure 3. HFA content at the *sn*-2 position of TAG. Ricinoleic acid (18:1OH, open bars) and lesquerolic acid (20:1OH, black bars) at the *sn*-2 position of TAG from wild type (wt) and transgenic line 3-1 and line 4-5 are measured in weight percent of total FA. Triplicates of 50-seed sample were measured for wild type and transgenic lines. The data represent averages of three replicates \pm SE. Two-tailed Student's *t* test. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.


3. Discussion

To evaluate the utility of *RcLPAT2* for the engineering of a lesquerella crop producing castor oil, we have produced transgenic lesquerella expressing *RcLPAT2* under the control of a seed specific promoter. Our results indicate that *RcLPAT2* enables the incorporation of 18:1OH at *sn*-2 position of LPA which increases the accumulation of 18:1OH and also tri-HFA-TAGs in lesquerella. This is the first successful step in the engineering of lesquerella seeds in order to produce castor oil. However, the overexpression of *RcLPAT2* did not lead to an overall increase in total HFAs (Tables 1 and 2, Figure 2). In fact, we observed small decreases in 20:1OH (Tables 1 and 2), indicating that in part, the increase in 18:1OH was at the expense of 20:1OH. Since *RcLPAT2* and *PfKCS3* utilize the same substrate 18:1OH-CoA, it is possible that *RcLPAT2* is competing with *PfKCS3* for 18:1OH-CoA substrate, limiting the availability of 18:1OH-CoA for the elongation reaction. Therefore, silencing *PfKCS3* would block the formation of 20:1OH and allow more 18:1OH-CoA to be directly incorporated into TAGs. In addition, the endogenous *PfFAH12* is a bifunctional oleate 12 hydroxylase:desaturase [32], that might not be strong enough to keep up with the demand of 18:1OH-CoA during the seed development. If this is the case, co-expression of a strict 12-hydroxylase gene from castor (*RcFAH12*) [31] or from *Physaria lindheimeri* (*PIFAH12*) [57] might be necessary to boost 18:1OH-CoA availability for incorporation into seed TAGs. On the other hand, lesquerella produces the polyunsaturated FA 18:2 at 8%–9% (Table S1, Table 2), which indicates that it must have an active desaturase *PfFAD2* [58] that converts 18:1 to 18:2. Since 18:1 is also the substrate for *FAH12* to synthesize 18:1OH, suppressing *PfFAD2* activity might increase the availability of 18:1 for *FAH12* to produce 18:1OH. Moreover, a microsomal Δ -15 desaturase (*PfFAD3*) could be silenced to further suppress the conversion of 18:1OH to 18:2OH. Our transgenic lines expressing *RcLPAT2* can be used as genetic backgrounds for studying the effect of additional transgenes such as *RcFAH12* or *PIFAH12* over-expression and/or *PfKCS3/PfFAD2/PfFAD3* silencing in order to develop a castor oil-producing lesquerella.

In transgenic lesquerella expressing *RcLPAT2*, we observed an increase in 0-, 1-, and 3-HFA-TAG levels and a reduction in 2-HFA-TAG species (Figure 2). Since the total HFA content remained unchanged, it is likely that HFAs were redistributed from *sn*-1 and/or *sn*-3 to the *sn*-2 position of TAG due to the limitation of the HFA supply. Regiochemical analysis showed that *sn*-2 18:1OH or 20:1OH increased 6–7-fold or 1.5-fold, respectively (Figure 3). This indicates that *RcLPAT2* allows for a more efficient acylation of 18:1OH than 20:1OH to the *sn*-2 position of TAG *in vivo*.

Although a number of mechanisms may facilitate accumulation of 18:1OH at the *sn*-2 position of TAG, the Kennedy pathway can be perceived as a major route for the channeling of 18:1OH into the *sn*-2 position of LPA by *RcLPAT2*. Figure 4 summarizes how *RcLPAT2* might influence TAG biosynthesis through the Kennedy pathway in transgenic lesquerella seeds. A lesquerella LPAT (*PfLPAT*) could acylate LPA with HFAs at a low level since we detected about 2% of 18:1OH and 1% of 20:1OH at the *sn*-2 position of wt TAGs (Figures 3 and 4). We have identified a *PfLPAT2* with 92% sequence identity to *AtLPAT2* [59]. Gene expression analysis showed that *PfLPAT2* is ubiquitously expressed in leaf, stem, flower bud, and developing seeds, suggesting it has a house-keeping role in membrane and storage lipid biosynthesis throughout plant life. In lines 3-1 and 4-5, small increase of 2% of 20:1OH at the *sn*-2 position of TAG were detected, however, significant increases of 18:1OH up to 15%–17% were detected (Figures 3 and 4). Thus unlike *PfLPAT2*, *RcLPAT2* has evolved to acylate 18:1OH to the *sn*-2 position of LPA in seeds. Our transgenic experiments demonstrate that *RcLPAT2* can be used to produce tri-HFA-TAGs in transgenic plants. In order to further increase the HFA level at the *sn*-2 position, the expression of *PfLPAT2* could be silenced, since substrate competition between endogenous and transgenic acyltransferases can limit the accumulation of UFAs [40,60].

A. Kennedy pathway in wild type lesquerella



B. Kennedy pathway in transgenic lesquerella line 3-1 expressing RcLPAT2




Figure 4. Simplified figure showing the effect of R_cLPAT2 on the TAG biosynthetic pathway in lesquerella focusing on the Kennedy pathway. In the Kennedy pathway sequential acylation of a glycerol backbone takes place, by the GPAT, LPAT and DGAT enzymes in order to generate TAG. (A) In lesquerella the GPAT and DGAT enzymes have high selectivity for 20:1-OH leading to acylation of the *sn*-1 and *sn*-3 positions of the glycerol backbone with 20:1-OH. LPAT has specificity for non-hydroxylated FAs, preventing the incorporation of significant amounts of HFA (~2% 18:1-OH and ~1% 20:1-OH) at the *sn*-2 position of the glycerol backbone (Figure 3); (B) In transgenic lesquerella seeds expressing R_cLPAT2, e.g., line 3-1, approximately 19% of FAs at the *sn*-2 position of TAG consist of HFAs (~17% 18:1-OH and ~2% 20:1-OH) (Figure 3). However, the total amount of HFAs in the seed oil is not increased, which leads to a decrease in acylation of 20:1-OH the *sn*-1 and *sn*-3 positions (Tables 1 and 2). This indicates that R_cLPAT2 has a high specificity for acylating 18:1-OH to the *sn*-2 position of LPA in lesquerella, which results in an increased amount of tri-HFA-TAGs in transgenic seed oil (Figure 2). 18:n means 18:1, 18:2, or 18:3, described as in Table 2. Other abbreviations are as in Figure 1. Red words show the effect of R_cLPAT2.

In lesquerella, acyl editing may provide 18:1OH for TAG assembly. Recent studies have suggested that LPCATs are responsible for incorporation of newly synthesized FA into PC (forward reaction), and transferring mFA such as PUFA or HFA produced on PC back to the acyl-CoA pool (reverse reaction) [61–64] (Figure 1). When the reverse reactions were measured, a ricinoleoyl group at the *sn*-2 position of PC was removed three- to six-fold faster than an oleoyl group by seven LPCATs from five species tested, including lesquerella [64]. Our previous results showed that there were no HFAs detectable in the PC fraction of total lipids isolated from developing lesquerella seeds [6], it is likely that acyl editing facilitates 18:1OH to be immediately released from PC and then utilized by R_cLPAT2 in these transgenic lesquerella lines.

We cannot exclude that the PC-derived DAG pathway may also contributed to the accumulation of 18:1OH at the *sn*-2 position of TAG (Figure 1). There is strong evidence that plants enriched with PUFAs in seed TAG may utilize the PC-derived pathway [62]. Lesquerella seed TAGs contain about 22% PUFAs (18:2 and 18:3) (Table 2), Therefore, it is likely that PC-derived DAGs are utilized in TAG assembly in lesquerella. Overexpression of R_cLPAT2 in developing seeds of lesquerella could enrich the *de novo* DAG pool with *sn*-2 18:1OH-DAGs. The 18:1OH on the *sn*-2 position of the *de novo* DAG could end up at the *sn*-2 position of PC by the action of CPT, generating *sn*-2 18:1OH-PC. If a significant FA flux through PC-derived DAG exists in lesquerella, some of the *sn*-2 18:1OH-PC could be converted by PDCT to *sn*-2 18:1OH-DAG for TAG assembly. The generation of 18:1OH-PC could also provide an opportunity for a Δ -15 desaturase to desaturate 18:1OH-PC to form 18:2OH-PC, resulting in observed small increases of 18:2OH in the transgenic lines.

In conclusion, this is the first demonstration that RcLPAT2 can be used to acylate 18:1OH at the *sn*-2 position of 20:1OH-LPA to generate tri-HFA-TAG, a valuable property for the engineering of a new castor oil-producing crop, such as lesquerella. This work also represents the first successful attempt in genetic engineering of lipid metabolic pathways in lesquerella.

4. Materials and Methods

4.1. Construction of *pGPro4-PnapA-RcLPAT2* Binary Vector

The map and sequence of the *pGPro4* vector was described [65]. The promoter sequence of *napA* was cloned into the *pGPro4* vector, resulting in *pGPro4-PnapA*. The activity of the *napA* promoter is seed-specific in lesquerella [66]. *RcLPAT2* was PCR amplified from castor cDNA with KOD polymerase (Novagen, Billerica, MA, USA) (Forward: 5'CACCATGGCTGTTGCAGCTGTAGC3', Reverse: 5'CTAGTCTGTTGTTTTCTG3') and cloned in the *pENTR-D-TOPO* vector (Life Technologies Ltd., Paisley, UK). Sequencing of *RcLPAT2* showed that there was one nucleotide change from A to G at position 1154 in the cloned *RcLPAT2* in comparison to the *RcLPAT2* sequence (GenBank ID: JQ796917) from the genome sequence in NCBI. This nucleotide change led to an amino acid change (Arg→Gly). This difference is most likely due to the different ecotypes used. To generate *pGPro4-PnapA-RcLPAT2*, *RcLPAT2* in *pENTR-D-TOPO* was PCR-amplified with gene-specific primers containing *Sal* I and *BstE* II restriction enzyme sites (Forward: 5'ACTGAGTCGACCGATCCATGGCTGTTGCAGCTGTAGC3', Reverse: 5'ACGTGTCTAGAGGTGACCCTAGTCTGTTGTTTTCTGC3') using Roche High fidelity DNA polymerase (Roche, Indianapolis, IN, USA). The PCR product was ligated in the *pGPro4-PnapA* vector digested with *Sal* I and *BstE* II.

4.2. Plant Transformation, and Estimation of Transgene Copy Number

The lesquerella seeds, WCL-LY2 [67], were kindly provided by Dave Dierig (USDA-ARS, Arid-Land Agricultural Research Center, Maricopa, AZ, USA). Plant transformation was performed using the *Agrobacterium tumefaciens* strain AGL1 [68] carrying the binary vector *pGPro4-PnapA-RcLPAT2*. Tissue culture and plant growth conditions were as described before [10]. We have produced a total of 17 primary transgenic plants (T_0), plant #6 was lost before maturity. In the greenhouse, we generated selfed T_1 seeds by hand-pollination between different flowers from the same plant. Transgenic T_1 plants were obtained by germinating T_1 seeds in basal medium supplemented with 50 mg/L hygromycin for two weeks and healthy T_1 seedlings were transplanted into soil for T_2 seed production. Transgene copy numbers was estimated for T_0 plants, and T_1 lines 3-1 and 4-5 using a qPCR method described before [69]. Optimized primer sequences for *RcPLAT2* are 5'TAGGCTGGTCCATGTGGTTTT3' (forward) and 5'TTCATCCTTGCCAGCTT3' (reverse). For internal control *KCS4/5*, the primer sets are the same as previously reported [69].

4.3. Lipid Extraction, TAG Species, Regiochemical and GC Analysis

The GC analysis for Table 1 and Table S1 was performed as described [38]. For Table 2, GC analysis was performed in the following way. One milliliter of hexanes and one mL of methanol containing 2.5% sulfuric acid (*v/v*) were added to seeds and incubated for 20 min at 100 °C. The top layer was extracted, dried under nitrogen and dissolved in 100 μ L hexanes. GC was performed on a Restek RTX-2330 column (30 m \times 0.25 mm inner diameter \times 0.2 μ m) using the following GC parameters: 120 °C for 1 min, a ramp to 180 °C at 40 °C \cdot min⁻¹, a ramp to 190 °C at 7 °C \cdot min⁻¹ with a 1 min hold, followed by a ramp to 250 °C at 30 °C \cdot min⁻¹ with a 6 min hold. Lipid extractions, analysis of TAG species and regiochemical analysis were performed as described [38]. However, the solvent system used for separating MAGs and 1-HFA-MAGs was based on Bates and Browse, 2011 [20]. In addition, regiochemical analysis was performed on total TAGs instead of separate TAG species as previously described [38]. In brief, TAG species analysis was performed in the following way. 1.5 mg of total lipids from wild type and transgenic lines were run in triplicate on TLC plates (Z292974-1PAK,

Sigma-Aldrich Company, Dorset, UK) alongside castor oil and flax oil standards. The TLC plates were first developed for 12 cm in CHCl₃:methanol:HOAc (93:3:0.5, v/v/v), dried for 15 min, followed by a full development in CHCl₃:methanol:HOAc (99:0.5:0.5, v/v/v). Lipids were visualized under UV after staining with the fluorescent dye primuline. Bands representing 3-HFA-, 2-HFA-, 1-HFA- and 0-HFA-TAGs were scraped from the TLC plates. The identity of the TAG species in each band was confirmed by measuring the percentage HFAs in each of these TAG fractions by GC (100%, 66%, 33%, and 0% HFAs in the 3-HFA-, 2-HFA-, 1-HFA- and 0-HFA-TAG fractions respectively). The amount of FAs in the 3-HFA-, 2-HFA-, 1-HFA- and 0-HFA-TAG fractions was calculated based on a 15:0 TAG internal standard added to each sample before sample preparation for GC. The relative percentage of each TAG species was calculated in the following way: $\mu\text{g FAs in the individual 3-HFA-, 2-HFA-, 1-HFA- or 0-HFA-TAG fractions} / \mu\text{g FAs in the combined 3-HFA-, 2-HFA-, 1-HFA- and 0-HFA-TAG fractions} = \text{relative \% of each TAG species as a fraction of total TAG}$. GC analysis was performed by transmethylation of FAs in 1 N methanolic HCl (Sigma-Aldrich Company) for two hours at 85 °C. FAMES were extracted by adding 500 μL of hexane, and 1 mL KCl, vortexing and centrifugation to separate the hexane layer from the aqueous phase. The hexane layer was collected and dried down under nitrogen and 200 μL BSTFA + TMCS (BSTFA + TMCS 99:1 (Sylon BFT), Sigma-Aldrich Company) was added and incubated at 80 °C for two hours to silylate the FAMES. The BSTFA + TMCS was evaporated and the silylated FAMES were dissolved in hexane. Quantification was performed by GC with flame ionization detection on a HP-1MS column (30 m \times 0.25 mm i.d. \times 0.25 μm , Agilent Technologies LDA UK Limited, Stockport, Cheshire, UK) using the following GC parameters: 50 °C for 1 min, a ramp to 200 °C at 50 °C \cdot min⁻¹, followed by a ramp to 325 °C at 3 °C \cdot min⁻¹. The identity of each silylated FAME was determined by GC-MS. In contrast to the GC results obtained for Tables 1 and 2 and Table S1, no 18:2OH and 20:2OH FAs could be detected in the analysis performed for Figures 2 and 3 using GC-MS. Probably the 18:1/18:2-OHs and 20:1/20:2-OHs cannot be resolved using the GC methods we used for these experiments.

Supplementary Materials: Supplementary materials can be found at <http://www.mdpi.com/1422-0067/17/4/507/s1>.

Acknowledgments: This study was supported by the U.S. Department of Agriculture Current Research Information System Project 2030-21410-020-00D (to Grace Q. Chen), the U.S. National Science Foundation grant IOS-1339385 (to John Browse) and the UK Biotechnology and Biological Sciences Research Council through Grant BBS/E/C/00005207 (to Peter Eastmond). USDA is an equal opportunity provider and employer. Mention of a specific product name by the U.S. Department of Agriculture does not constitute an endorsement and does not imply a recommendation over other suitable products.

Author Contributions: Grace Q. Chen and Harrie van Erp conceived and supervised the study, and also designed experiments and wrote the manuscript; Jose Martin-Moreno, Kumiko Johnson, Eva Morales, Grace Q. Chen and Harrie Van Erp performed experiments; and John Browse, Peter J. Eastmond and Jiann-Tsyh Lin made critical manuscript revisions.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Al-Shehbaz, I.A.; O’Kane, S.L., Jr. *Lesquerella* is united with *Physaria* (Brassicaceae). *Novon* **2002**, *12*, 319–329. [[CrossRef](#)]
2. Smith, C.R., Jr.; Wilson, T.L.; Miwa, T.K.; Zobel, H.; Lohmar, R.L.; Wolff, I.A. Lesquerolic acid. A new hydroxy acid from *Lesquerella* seed oil. *J. Org. Chem.* **1961**, *26*, 2903–2905. [[CrossRef](#)]
3. Hayes, D.G.; Kleiman, R. A detailed triglyceride analysis of *Lesquerella fendleri* oil: Column chromatographic fractionation followed by supercritical fluid chromatography. *J. Am. Oil Chem. Soc.* **1996**, *73*, 267–269. [[CrossRef](#)]
4. Dierig, D.A.; Wang, G.; McCloskey, W.B.; Thorp, K.R.; Isbell, T.A.; Ray, D.T.; Foster, M.A. *Lesquerella*: New crop development and commercialization in the U.S. *Ind. Crops Prod.* **2011**, *34*, 1381–1385. [[CrossRef](#)]
5. Isbell, T.A.; Mund, M.S.; Evangelista, R.L.; Dierig, D.A. Method for analysis of fatty acid distribution and oil content on a single *Lesquerella fendleri* seed. *Ind. Crops Prod.* **2008**, *28*, 231–236. [[CrossRef](#)]

6. Chen, G.Q.; Lin, J.T.; Lu, C. Hydroxy fatty acid synthesis and lipid gene expression during seed development in *Lesquerella fendleri*. *Ind. Crops Prod.* **2010**, *34*, 1286–1292. [[CrossRef](#)]
7. Hayes, D.G.; Kleiman, R. Recovery of hydroxy fatty acids from *Lesquerella* oil with lipases. *J. Am. Oil Chem. Soc.* **1992**, *69*, 982–985. [[CrossRef](#)]
8. Lin, J.T.; Fagerquist, C.K.; Chen, G.Q. Ratios of regioisomers of the molecular species of triacylglycerols in *Lesquerella* (*Physaria fendleri*) oil estimated by mass spectrometry. *J. Am. Oil Chem. Soc.* **2016**, *93*, 183–191. [[CrossRef](#)]
9. Caupin, H.J. *Products from Castor Oil: Past, Present and Future*; Gunstone, F.D., Padley, F.B., Eds.; Marcel Dekker: New York, NY, USA, 1997; pp. 787–795.
10. Chen, G.Q. Effective reduction of chimeric tissue in transgenics for the stable genetic transformation of *Lesquerella fendleri*. *HortScience* **2011**, *46*, 86–90.
11. Chapman, K.D.; Ohlrogge, J.B. Compartmentation of triacylglycerol accumulation in plants. *J. Biol. Chem.* **2012**, *287*, 2288–2294. [[CrossRef](#)] [[PubMed](#)]
12. Bates, P.D.; Stymne, S.; Ohlrogge, J. Biochemical pathways in seed oil synthesis. *Curr. Opin. Plant Biol.* **2013**, *16*, 358–364. [[CrossRef](#)] [[PubMed](#)]
13. Kennedy, E.P. Biosynthesis of complex lipids. *Fed. Proc.* **1961**, *20*, 934–940. [[PubMed](#)]
14. Weiss, S.B.; Kennedy, E.P.; Kiyasu, J.Y. The enzymatic synthesis of triglycerides. *J. Biol. Chem.* **1960**, *235*, 40–44. [[CrossRef](#)] [[PubMed](#)]
15. Barron, E.J.; Stumpf, P.K. Fat metabolism in higher plants XIX. The biosynthesis of triglycerides by avocado-mesocarp enzymes. *Biochim. Biophys. Acta* **1962**, *60*, 329–337. [[CrossRef](#)]
16. Lands, W.E. Lipid metabolism. *Annu. Rev. Biochem.* **1965**, *34*, 313–346. [[CrossRef](#)] [[PubMed](#)]
17. Li-Beisson, Y.; Shorrosh, B.; Beisson, F.; Andersson, M.X.; Arondel, V.; Bates, P.D.; Baud, S.; Bird, D.; DeBono, A.; Durrett, T.P.; *et al.*. Acyl-lipid metabolism. *Arabidopsis Book* **2013**, *11*, e0161. [[CrossRef](#)] [[PubMed](#)]
18. Dahlqvist, A.; Stahl, U.; Lenman, M.; Banas, A.; Lee, M.; Sandager, L.; Ronne, H.; Stymne, S. Phospholipid:diacylglycerol acyltransferase: An enzyme that catalyzes the acyl-CoA-independent formation of triacylglycerol in yeast and plants. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 6487–6492. [[CrossRef](#)] [[PubMed](#)]
19. Bates, P.D.; Durrett, T.P.; Ohlrogge, J.B.; Pollard, M. Analysis of acyl fluxes through multiple pathways of triacylglycerol synthesis in developing soybean embryos. *Plant Physiol.* **2009**, *150*, 55–72. [[CrossRef](#)] [[PubMed](#)]
20. Bates, P.D.; Browse, J. The pathway of triacylglycerol synthesis through phosphatidylcholine in *Arabidopsis* produces a bottleneck for the accumulation of unusual fatty acids in transgenic seeds. *Plant J.* **2011**, *68*, 387–399. [[CrossRef](#)] [[PubMed](#)]
21. Hu, Z.; Ren, Z.; Lu, C. The phosphatidylcholine diacylglycerol cholinephosphotransferase is required for efficient hydroxy fatty acid accumulation in transgenic *Arabidopsis*. *Plant Physiol.* **2012**, *158*, 1944–1954. [[CrossRef](#)] [[PubMed](#)]
22. Lu, C.; Xin, Z.; Ren, Z.; Miquel, M.; Browse, J. An enzyme regulating triacylglycerol composition is encoded by the *ROD1* gene of *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 18837–18842. [[CrossRef](#)] [[PubMed](#)]
23. Slack, C.R.; Campbell, L.C.; Browse, J.A.; Roughan, P.G. Some evidence for the reversibility of the cholinephosphotransferase-catalysed reaction in developing linseed cotyledons *in vivo*. *Biochim. Biophys. Acta* **1983**, *754*, 10–20. [[CrossRef](#)]
24. Lin, J.T.; Turner, C.; Liao, L.P.; McKeon, T.A. Identification and quantification of the molecular species of acylglycerols in castor oil by HPLC using ELSD. *J. Liquid Chromatogr. Relat. Technol.* **2003**, *26*, 773–780. [[CrossRef](#)]
25. Moreau, R.A.; Stumpf, P.K. Recent studies of the enzymic synthesis of ricinoleic acid by developing castor beans. *Plant Physiol.* **1981**, *67*, 672–676. [[CrossRef](#)] [[PubMed](#)]
26. Bafor, M.; Smith, M.A.; Jonsson, L.; Stobart, K.; Stymne, S. Ricinoleic acid biosynthesis and triacylglycerol assembly in microsomal preparations from developing castor-bean (*Ricinus communis*) endosperm. *Biochem. J.* **1991**, *280*, 507–514. [[CrossRef](#)] [[PubMed](#)]
27. Engeseth, N.; Stymne, S. Desaturation of oxygenated fatty acids in *Lesquerella* and other oil seeds. *Planta* **1996**, *198*, 238–245. [[CrossRef](#)]
28. Reed, D.W.; Taylor, D.C.; Covello, P.S. Metabolism of hydroxy fatty acids in developing seeds in the genera *Lesquerella* (Brassicaceae) and *Linum* (Linaceae). *Plant Physiol.* **1997**, *114*, 63–68. [[PubMed](#)]

29. Moon, H.; Smith, M.A.; Kunst, L. A condensing enzyme from the seeds of *Lesquerella fendleri* that specifically elongates hydroxy fatty acids. *Plant Physiol.* **2001**, *127*, 1635–1643. [[CrossRef](#)] [[PubMed](#)]
30. Lee, K.-R.; Chen, G.Q.; Kim, H.U. Current progress towards the metabolic engineering of plant seed oil for hydroxy fatty acids production. *Plant Cell Rep.* **2015**, *34*, 603–615. [[CrossRef](#)] [[PubMed](#)]
31. Van de Loo, F.J.; Broun, P.; Turner, S.; Somerville, C. An oleate 12-hydroxylase from *Ricinus communis* L. is a fatty acyl desaturase homolog. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 6743–6747. [[CrossRef](#)] [[PubMed](#)]
32. Broun, P.; Boddupalli, S.; Somerville, C. A bifunctional oleate 12-hydroxylase: Desaturase from *Lesquerella fendleri*. *Plant J.* **1998**, *13*, 201–210. [[CrossRef](#)] [[PubMed](#)]
33. Broun, P.; Somerville, C. Accumulation of ricinoleic, lesquerolic, and densipolic acids in seeds of transgenic arabidopsis plants that express a fatty acyl hydroxylase cDNA from castor bean. *Plant Physiol.* **1997**, *113*, 933–942. [[CrossRef](#)] [[PubMed](#)]
34. Smith, M.A.; Moon, H.; Chowrira, G.; Kunst, L. Heterologous expression of a fatty acid hydroxylase gene in developing seeds of *Arabidopsis thaliana*. *Planta* **2003**, *217*, 507–516. [[CrossRef](#)] [[PubMed](#)]
35. Kumar, R.; Wallis, J.G.; Skidmore, C.; Browse, J. A mutation in Arabidopsis cytochrome b5 reductase identified by high-throughput screening differentially affects hydroxylation and desaturation. *Plant J.* **2006**, *48*, 920–932. [[CrossRef](#)] [[PubMed](#)]
36. Lu, C.; Fulda, M.; Wallis, J.G.; Browse, J. A high-throughput screen for genes from castor that boost hydroxy fatty acid accumulation in seed oils of transgenic Arabidopsis. *Plant J.* **2006**, *45*, 847–856. [[CrossRef](#)] [[PubMed](#)]
37. Burgal, J.; Shockey, J.; Lu, C.; Dyer, J.; Larson, T.; Graham, I.; Browse, J. Metabolic engineering of hydroxy fatty acid production in plants: RcDGAT2 drives dramatic increases in ricinoleate levels in seed oil. *Plant Biotechnol. J.* **2008**, *6*, 819–831. [[CrossRef](#)] [[PubMed](#)]
38. Van Erp, H.; Bates, P.D.; Burgal, J.; Shockey, J.; Browse, J. Castor phospholipid:diacylglycerol acyltransferase facilitates efficient metabolism of hydroxy fatty acids in transgenic Arabidopsis. *Plant Physiol.* **2011**, *155*, 683–693. [[CrossRef](#)] [[PubMed](#)]
39. Kim, H.U.; Lee, K.R.; Go, Y.S.; Jung, J.H.; Suh, M.C.; Kim, J.B. Endoplasmic reticulum-located PDAT1-2 from castor bean enhances hydroxy fatty acid accumulation in transgenic plants. *Plant Cell Physiol.* **2011**, *52*, 983–993. [[CrossRef](#)] [[PubMed](#)]
40. Van Erp, H.; Shockey, J.; Zhang, M.; Adhikari, N.D.; Browse, J. Reducing isozyme competition increases target fatty acid accumulation in seed triacylglycerols of transgenic Arabidopsis. *Plant Physiol.* **2015**, *168*, 36–46. [[CrossRef](#)] [[PubMed](#)]
41. Frentzen, M. Acyltransferases from basic science to modified seed oils. *Lipid/Fett* **1998**, *100*, 161–166. [[CrossRef](#)]
42. Phillips, B.E.; Smith, C.R., Jr.; Tallent, W.H. Glycerides of *Limnanthes douglasii* seed oil. *Lipids* **1971**, *6*, 93–99. [[CrossRef](#)]
43. Balachandran, C.; Arumughan, C.; Mathew, A.G. Distribution of major chemical constituents and fatty acids in different regions of coconut endosperm. *J. Am. Oil Chem. Soc.* **1985**, *62*, 1583–1586. [[CrossRef](#)]
44. Brough, C.L.; Coventry, J.M.; Christie, W.W.; Kroon, J.T.M.; Brown, A.P.; Barsby, T.L.; Slabas, A.R. Towards the genetic engineering of triacylglycerols of defined fatty acid composition: Major changes in erucic acid content at the *sn*-2 position affected by the introduction of a 1-acyl-*sn*-glycerol-3-phosphate acyltransferase from *Limnanthes douglasii* into oil seed rape. *Mol. Breed.* **1996**, *2*, 133–142.
45. Knutzon, D.S.; Hayes, T.R.; Wyrick, A.; Xiong, H.; Davies, H.M.; Voelker, T.A. Lysophosphatidic acid acyltransferase from coconut endosperm mediates the insertion of laurate at the *sn*-2 position of triacylglycerols in lauric rapeseed oil and can increase total laurate levels. *Plant Physiol.* **1999**, *120*, 739–746. [[CrossRef](#)] [[PubMed](#)]
46. Lassner, M.W.; Levering, C.K.; Davies, H.M.; Knutzon, D.S. Lysophosphatidic acid acyltransferase from meadowfoam mediates insertion of erucic acid at the *sn*-2 position of triacylglycerol in transgenic rapeseed oil. *Plant Physiol.* **1995**, *109*, 1389–1394. [[CrossRef](#)] [[PubMed](#)]
47. Zou, J.; Katavic, V.; Giblin, E.M.; Barton, D.L.; MacKenzie, S.L.; Keller, W.A.; Hu, X.; Taylor, D.C. Modification of seed oil content and acyl composition in the brassicaceae by expression of a yeast *sn*-2 acyltransferase gene. *Plant Cell* **1997**, *9*, 909–923. [[CrossRef](#)] [[PubMed](#)]
48. Kim, H.U.; Huang, A.H.C. Plastid lysophosphatidyl acyltransferase is essential for embryo development in Arabidopsis. *Plant Physiol.* **2004**, *134*, 1206–1216. [[CrossRef](#)] [[PubMed](#)]

49. Kim, H.U.; Li, Y.; Huang, A.H. Ubiquitous and endoplasmic reticulum-located lysophosphatidyl acyltransferase, LPAT2, is essential for female but not male gametophyte development in Arabidopsis. *Plant Cell* **2005**, *17*, 1073–1089. [[CrossRef](#)] [[PubMed](#)]
50. Yu, X.H.; Prakash, R.R.; Sweet, M.; Shanklin, J. Coexpressing *Escherichia coli* cyclopropane synthase with *Sterculia foetida* Lysophosphatidic acid acyltransferase enhances cyclopropane fatty acid accumulation. *Plant Physiol.* **2014**, *164*, 455–465. [[CrossRef](#)] [[PubMed](#)]
51. Chan, A.P.; Crabtree, J.; Zhao, Q.; Lorenzi, H.; Orvis, J.; Puiu, D.; Melake-Berhan, A.; Jones, K.M.; Redman, J.; Chen, G.; *et al.* Draft genome sequence of the oilseed species *Ricinus communis*. *Nat. Biotechnol.* **2010**, *28*, 951–956. [[CrossRef](#)] [[PubMed](#)]
52. Cagliari, A.; Margis-Pinheiro, M.; Loss, G.; Mastroberti, A.A.; de Araujo Mariath, J.E.; Margis, R. Identification and expression analysis of castor bean (*Ricinus communis*) genes encoding enzymes from the triacylglycerol biosynthesis pathway. *Plant Sci.* **2010**, *179*, 499–509. [[CrossRef](#)] [[PubMed](#)]
53. Troncoso-Ponce, M.A.; Kilaru, A.; Cao, X.; Durrett, T.P.; Fan, J.; Jensen, J.K.; Thrower, N.A.; Pauly, M.; Wilkerson, C.; Ohlrogge, J.B. Comparative deep transcriptional profiling of four developing oilseeds. *Plant J.* **2011**, *68*, 1014–1027. [[CrossRef](#)] [[PubMed](#)]
54. Brown, A.P.; Kroon, J.T.; Swarbreck, D.; Febrer, M.; Larson, T.R.; Graham, I.A.; Caccamo, M.; Slabas, A.R. Tissue-specific whole transcriptome sequencing in castor, directed at understanding triacylglycerol lipid biosynthetic pathways. *PLoS ONE* **2012**, *7*, e30100. [[CrossRef](#)] [[PubMed](#)]
55. Arroyo-Caro, J.M.; Chileh, T.; Kazachkov, M.; Zou, J.; Alonso, D.L.; Garcia-Maroto, F. The multigene family of lysophosphatidate acyltransferase (LPAT)-related enzymes in *Ricinus communis*: Cloning and molecular characterization of two LPAT genes that are expressed in castor seeds. *Plant Sci.* **2013**, *199–200*, 29–40. [[CrossRef](#)] [[PubMed](#)]
56. Lin, J.T.; Chen, G.Q. Identification of TAG and DAG and their FA constituents in Lesquerella (*Physaria fendleri*) oil by HPLC and MS. *J. Am. Oil Chem. Soc.* **2013**, *90*, 1819–1829. [[CrossRef](#)]
57. Dauk, M.; Lam, P.; Kunst, L.; Smith, M.A. A FAD2 homologue from *Lesquerella lindheimeri* has predominantly fatty acid hydroxylase activity. *Plant Sci.* **2007**, *173*, 43–49. [[CrossRef](#)]
58. Lozinsky, S.; Yang, H.; Forseille, L.; Cook, G.R.; Ramirez-Erosa, I.; Smith, M.A. Characterization of an oleate 12-desaturase from *Physaria fendleri* and identification of 5'UTR introns in divergent FAD2 family genes. *Plant Physiol. Biochem. PPB/Soc. Fr. Physiol. Veg.* **2014**, *75*, 114–122. [[CrossRef](#)] [[PubMed](#)]
59. Kim, H.U.; Chen, G.Q. Identification of hydroxy fatty acid and triacylglycerol metabolism-related genes in Lesquerella through seed transcriptome analysis. *BMC Genom.* **2015**, *16*. [[CrossRef](#)] [[PubMed](#)]
60. Liu, J.; Rice, A.; McGlew, K.; Shaw, V.; Park, H.; Clemente, T.; Pollard, M.; Ohlrogge, J.; Durrett, T.P. Metabolic engineering of oilseed crops to produce high levels of novel acetyl glyceride oils with reduced viscosity, freezing point and calorific value. *Plant Biotechnol. J.* **2015**, *13*, 858–865. [[CrossRef](#)] [[PubMed](#)]
61. Xu, J.; Carlsson, A.S.; Francis, T.; Zhang, M.; Hoffman, T.; Giblin, M.E.; Taylor, D.C. Triacylglycerol synthesis by PDAT1 in the absence of DGAT1 activity is dependent on re-acylation of LPC by LPCAT2. *BMC Plant Biol.* **2012**, *12*. [[CrossRef](#)] [[PubMed](#)]
62. Bates, P.D.; Browse, J. The significance of different diacylglycerol synthesis pathways on plant oil composition and bioengineering. *Front. Plant Sci.* **2012**, *3*. [[CrossRef](#)] [[PubMed](#)]
63. Wang, L.; Shen, W.; Kazachkov, M.; Chen, G.; Chen, Q.; Carlsson, A.S.; Stymne, S.; Weselake, R.J.; Zou, J. Metabolic interactions between the lands cycle and the Kennedy pathway of glycerolipid synthesis in Arabidopsis developing seeds. *Plant Cell* **2012**, *24*, 4652–4669. [[CrossRef](#)] [[PubMed](#)]
64. Lager, I.; Yilmaz, J.L.; Zhou, X.R.; Jasieniecka, K.; Kazachkov, M.; Wang, P.; Zou, J.; Weselake, R.; Smith, M.A.; Bayon, S.; *et al.* Plant acyl-CoA:lysophosphatidylcholine acyltransferases (LPCATs) have different specificities in their forward and reverse reactions. *J. Biol. Chem.* **2013**, *288*, 36902–36914. [[CrossRef](#)] [[PubMed](#)]
65. Chen, G.Q.; Thilmony, R.; Lin, J.-T. Transformation of *Lesquerella fendleri* with the new binary vector pGPro4-35s. *Online J. Biol. Sci.* **2011**, *11*, 90–95.
66. Chen, G.Q.; Lin, J.-T. A napin promoter activates gene expression in developing seeds of *Lesquerella fendleri*. *Online J. Biol. Sci.* **2012**, *12*, 113–117. [[CrossRef](#)]
67. Dierig, D.A.; Tomasi, P.M.; Dahlquist, G.H. Registration of WCL-LY2 high oil *Lesquerella fendleri* germplasm. *Crop Sci.* **2001**, *41*, 604–605. [[CrossRef](#)]

68. Lazo, G.R.; Stein, P.A.; Ludwig, R.A. A DNA transformation-competent Arabidopsis genomic library in Agrobacterium. *Nat. Biotechnol.* **1991**, *9*, 963–967. [[CrossRef](#)]
69. Chen, G.Q.; Lin, J.-T. Use of quantitative polymerase chain reaction for determining copy numbers of transgenes in *Lesquerella fendleri*. *Am. J. Agric. Biol. Sci.* **2010**, *5*, 415–421. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons by Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).