

# Rothamsted Repository Download

## A - Papers appearing in refereed journals

Evens, N. P., Buchner, P. H., Williams, L. E. and Hawkesford, M. J. 2017. The role of ZIP transporters and group F bZIP transcription factors in the Zn-deficiency response of wheat (*Triticum aestivum*). *The Plant Journal*. 92 (2), pp. 291-304.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1111/tpj.13655>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/84561>.

© 3 August 2017, Rothamsted Research. Licensed under the Creative Commons CC BY.

# The role of ZIP transporters and group F bZIP transcription factors in the Zn-deficiency response of wheat (*Triticum aestivum*)

Nicholas P. Evens<sup>1,2</sup>, Peter Buchner<sup>1</sup>, Lorraine E. Williams<sup>2,\*</sup> and Malcolm J. Hawkesford<sup>1,\*</sup>

<sup>1</sup>Rothamsted Research, West Common, Harpenden, Hertfordshire AL5 2JQ, UK, and

<sup>2</sup>Biological Science, University of Southampton, Life Sciences Building 85, Highfield Campus, Southampton SO17 1BJ, UK

Received 6 June 2017; revised 25 July 2017; accepted 28 July 2017; published online 3 August 2017.

\*For correspondence (e-mails malcolm.hawkesford@rothamsted.ac.uk; l.e.williams@soton.ac.uk).

## SUMMARY

Understanding the molecular basis of zinc (Zn) uptake and transport in staple cereal crops is critical for improving both Zn content and tolerance to low-Zn soils. This study demonstrates the importance of group F bZIP transcription factors and ZIP transporters in responses to Zn deficiency in wheat (*Triticum aestivum*). Seven group F TabZIP genes and 14 ZIPs with homeologs were identified in hexaploid wheat. Promoter analysis revealed the presence of Zn-deficiency-response elements (ZDREs) in a number of the ZIPs. Functional complementation of the *zrt1/zrt2* yeast mutant by TaZIP3, -6, -7, -9 and -13 supported an ability to transport Zn. Group F TabZIPs contain the group-defining cysteine–histidine-rich motifs, which are the predicted binding site of Zn<sup>2+</sup> in the Zn-deficiency response. Conservation of these motifs varied between the TabZIPs suggesting that individual TabZIPs may have specific roles in the wheat Zn-homeostatic network. Increased expression in response to low Zn levels was observed for several of the wheat ZIPs and bZIPs; this varied temporally and spatially suggesting specific functions in the response mechanism. The ability of the group F TabZIPs to bind to specific ZDREs in the promoters of TaZIPs indicates a conserved mechanism in monocots and dicots in responding to Zn deficiency. In support of this, TabZIPF1-7DL and TabZIPF4-7AL afforded a strong level of rescue to the Arabidopsis hypersensitive *bzip19 bzip23* double mutant under Zn deficiency. These results provide a greater understanding of Zn-homeostatic mechanisms in wheat, demonstrating an expanded repertoire of group F bZIP transcription factors, adding to the complexity of Zn homeostasis.

**Keywords:** zinc, micronutrient, biofortification, ZIP transporter, membrane transport, bZIP, wheat (*Triticum aestivum*), transcription factor.

## INTRODUCTION

Micronutrient deficiency in humans is an issue of global concern. Enhancing the micronutrient content of staple crops is therefore an important objective for modern agriculture. Cereals such as wheat (*Triticum aestivum*) are relatively low in essential micronutrients such as zinc (Zn) and iron in their edible tissues. This presents a major problem when cereals form the main part of the diet and is a particular issue in developing countries (Kumssa *et al.*, 2015). Crop yield is also detrimentally affected when plants are deficient in micronutrients as they are required throughout plant development (Brown *et al.*, 1993). Zn has essential roles in plant growth, phytohormone activity, enzyme activation and modification of gene expression (Broadley *et al.*, 2007). The yield reductions from reduced Zn availability are ultimately associated with damage to cell

proteins, lipids and DNA (Cakmak, 2000). The development of crops that maintain growth and yield under low Zn availability would have clear benefits, and for that an understanding of the homeostatic network that determines Zn efficiency is required. Wheat cultivation occupies the largest area of any crop, therefore this study addresses mechanisms contributing to adaptation to low Zn in this monocot cereal.

Plants have developed sophisticated sensing and response mechanisms allowing them to adapt to variations in micronutrient availability. The complex nature of these homeostatic mechanisms is starting to be resolved in the model plant Arabidopsis. In this dicot, two bZIP (basic-leucine zipper domain) transcription factors, bZIP19 and -23, are involved in adapting to Zn deficiency by

inducing the expression of particular family members of membrane transporters, the ZIPs (ZRT, IRT-related proteins) (Assunção *et al.*, 2010). These have emerged as a key membrane transporter family in the journey of Zn from soil to seed (Palmgren *et al.*, 2008). In Arabidopsis, eight members of the ZIP family have ability to transport Zn (AtIRT1, AtZIP1, -2, -3, -7, -10, -11 and -12) (Grotz *et al.*, 1998; Milner *et al.*, 2013), and several have been shown to be induced in response to Zn deficiency. ZIPs have also been characterised in cereals such as rice (*Oryza sativa*) and barley (*Hordeum vulgare*). In rice, OsZIP3, -4, -5 and -8 are functional Zn transporters (Ishimaru *et al.*, 2005; Yang *et al.*, 2009b; Lee *et al.*, 2010a) while OsIRT1 and -2 as well as OsZIP6 and -7 can transport Fe but not Zn (Bugchio *et al.*, 2002; Ishimaru *et al.*, 2006; Kavitha *et al.*, 2015). The expression of *OsZIP4*, -5 and -8 is induced by Zn deficiency in both the root and shoot (Suzuki *et al.*, 2012). In barley HvIRT1 and HvZIP3, -5 and -8 have all been shown to rescue the *zrt1/zrt2* Zn mutant yeast strain to a varying degree, indicating the ability to transport Zn (Pedas *et al.*, 2008, 2009). Some of the rice and barley ZIPs are also induced by Zn deficiency, including *OsZIP1*, -3, -4, -5 and -8 (Ramesh *et al.*, 2003; Ishimaru *et al.*, 2005; Lee *et al.*, 2010b) and *HvZIP3*, -5, -7, -8, -10 and -13 (Pedas *et al.*, 2009; Tiong *et al.*, 2013, 2015). In wheat little is known about this family except for *TdZIP1* from wild emmer wheat (*Triticum turgidum* ssp. *dicoccoides*), a Zn transporter with higher expression under Zn deficiency (Durmaz *et al.*, 2011). In this study we have identified Zn-responsive ZIPs in wheat (*T. aestivum*) and confirmed their Zn transport capability. A particular step forward was to investigate the mechanism that leads to changes in the expression of these transporters in response to Zn deficiency.

In Arabidopsis, increases observed in expression of particular ZIPs are proposed to occur via binding of AtbZIP19 and AtbZIP23 to Zn-deficiency-response elements (ZDREs) in ZIP promoters (Assunção *et al.*, 2010). This operates as an adaptive mechanism to increase uptake and distribution of Zn in response to Zn deficiency. Evidence for this was provided by the extreme hypersensitivity of a *bzip19 bzip23* double mutant to low-Zn conditions and the lack of Zn-induced expression of key Zn transporters, including *AtZIP1*, -3, -4, -5, -9 and -10 (Assunção *et al.*, 2010). These genes contain one or more copies of the ZDRE in their promoter and direct binding by AtbZIP19 to this motif in *AtZIP4* was demonstrated previously (Assunção *et al.*, 2010).

Further analysis of the *bzip19* and *bzip23* single mutants indicates that while they may overlap in their targets, *AtbZIP19* and *AtbZIP23* may also operate to alter the expression of a specific subset of genes (Inaba *et al.*, 2015). The mechanism whereby low Zn is sensed is still to be elucidated, but it is suggested that under normal cellular Zn

concentrations a Zn<sup>2+</sup> ion binds to the cysteine–histidine-rich motifs present in the group F bZIPs, making them inactive. Upon a reduction in cellular Zn concentration, Zn<sup>2+</sup> dissociates, thereby activating the bZIP dimer which in turn binds to the ZDRE motif and brings about an increase in transcription of Zn-responsive genes (Assunção *et al.*, 2013). We show that wheat contains an additional level of complexity in adapting to Zn deficiency with an expanded number of group F bZIP transcription factors, allowing further modulation of the response to variations in Zn growth conditions.

## RESULTS

### A multi-gene family of ZIPs is present in wheat

A comprehensive analysis of the genomic information recently available for hexaploid wheat was performed. From this analysis, 14 *TaZIP* genes were identified (Figure 1). Modern bread wheat (*T. aestivum*) is hexaploid, containing three genomes – A, B and D; homeologous genes are present on all three genomes. Full homeolog complements were found for all of the *TaZIPs* in each of the A, B and D genomes (Figure S1 in the Supporting Information). A previous phylogenetic analysis of *TaZIPs* identified 11 homologs (Tiong *et al.*, 2015); *TaZIP2*, -8 and -9 have not formerly been identified. The *TaZIP* homeologs group closely to one another (Figure S1) and for 12 of the *TaZIPs* the most closely related homolog was that from barley, followed by *Brachypodium* and rice. *TaZIP9* and *TaZIP13* showed high relatedness: *TaZIP9-2DS* [*TaZIP9* present on the short arm (S) of chromosome 2D] and *TaZIP13-2DL* [*TaZIP13* present on the long arm (L) of chromosome 2D] had 83% nucleotide sequence similarity (Table S1 and Figure S2), and both are located on chromosome 2, suggesting a wheat-specific gene duplication. This is further supported by there being only one barley gene in this group.

The phylogeny of ZIPs displays three distinct clades (Figure 1). All clades contain both monocot and dicot ZIP homologs, suggesting that ZIPs were present before the divergence of monocots and dicots. *AtPutZnT*, *AtIAR1* and *AtZTP29* and their cereal homologs (*TaZIP11*, *TaZIP14* and *TaZIP16*) are situated in a subclade of clade 3 and are more distantly related to the other ZIPs included in the phylogenetic analysis. Each sub-clade also contains monocot and dicot members, but there is also evidence suggesting species-specific ZIP expansion with possible functional divergence.


### Wheat ZIPs functionally complement *zrt1/zrt2*, a Zn-sensitive yeast mutant

A functional role in Zn homeostasis for a selection of the wheat ZIPs (*TaZIP3*, -6, -7, -9 and -13) was tested using yeast complementation. These *TaZIPs* were selected due to



compared with the empty pYES2 vector control, consistent with them functioning as Zn transporters. None of the *TaZIPs* heterologously expressed in the *fet3/fet4* Fe-uptake

mutant strain were able to rescue the Fe-deficient phenotype (Figure 2e–g), suggesting that none of the *TaZIPs* investigated are functional Fe transporters.



**Figure 2.** Complementation of Zn and Fe yeast uptake mutants with *TaZIP* genes. The yeast Zn uptake mutant *zrt1/zrt2* (a–d) and the Fe uptake mutant *fet3/fet4* (e–g) were transformed with five wheat *TaZIP* genes, shown here with empty vector controls (e.v.) in both the mutant and wild-type yeast strains. Each spot is a dilution of the culture starting on the left of each plate (undiluted, 1:2, 1:10, 1:100, 1:1000) with the contents of selective media described above each plate.

### TaZIP expression is induced by Zn deficiency

Relative gene expression levels of *TaZIP3*, -5, -6, -7 and -13 were determined in root and shoot tissue obtained during a 1-week hydroponic Zn-starvation period. Under these conditions there were no observable differences in wheat plants after 1 week and they showed similar root and shoot fresh weight at the end of this period (Figure S3). However, Zn levels were clearly reduced in both roots and shoots (Figure S3). Other elements were measured, but they showed little change over this period (Figure S4). There was a small but significant increase in root Mn levels in -Zn grown plants but no corresponding change in shoots (Figure S4). Thus the major effect was on the Zn content in both roots and shoots. Having established a system for testing the effects of Zn deficiency, we determined the transcriptional changes of a selected number of ZIPs. In roots, *TaZIP3*, -5, -7 and -13 showed increased expression under the -Zn treatment, with only *TaZIP6* expression remaining fairly stable (Figure 3). *TaZIP3*, -5 and -13 showed significantly higher expression levels from the first time point (day 1); *TaZIP7* expression was significantly increased from day 3. In the shoot material, expression of all five *TaZIPs* increased under -Zn conditions but the timing varied between individual genes. *TaZIP6* was the slowest to respond, remaining constant until day 5. As in the roots, *TaZIP3* showed significantly higher expression from day 1; however, *TaZIP5* and -13 were slower to respond in the shoot compared with the root. The magnitude of gene-expression changes varied between the *TaZIPs* as well as between roots and shoots. *TaZIP5* expression increased to a greater extent in the shoots compared with roots, whereas for *TaZIP13* the opposite was observed. *TaZIP* induction under Zn-deficient conditions was also observed over a longer starvation period (Figure S5). In this case, plants were only grown for 7 days rather than 14 days prior to the Zn starvation. Slight differences were observed, indicating that the developmental stage at which Zn starvation was implemented could have an effect, but overall the results were similar. In this case an induction was seen in *TaZIP6* expression in the root but again it was not as high as the induction observed for the other genes at day seven.

### Wheat contains seven group F *TabZIPs* with a full set of homeologs identified in the wheat genome

Searching the wheat genome revealed seven group F *TabZIPs* with homeologs identified on the A, B and D genomes. The previous genome-wide identification of *TabZIPs* in the wheat genome (Li *et al.*, 2015) reported 11 group F *TabZIPs*. However, this was inaccurate as closer examination showed that a considerable proportion of these are homeologs of the same gene (Table S2). Our analysis identified seven *TabZIPFs*, each with three homeologs. The phylogenetic analysis presented in Figure 4 shows that


these have a related barley homolog. *TabZIPF3a* and *TabZIPF3b* are closely related to *TabZIPF3a-7AL* and *TabZIPF3b-7AL*, sharing 88% sequence similarity. Their sequence similarity, the location of each on chromosome seven and the fact that each has a barley homolog suggests they have evolved through gene duplication, which is likely to have occurred before the speciation of wheat and barley. *TabZIPF1*, -2 and -4 also form a clade more closely related to *TabZIPF3a* and -3b than *TabZIPF5* and -6. *TabZIPF1*, -2, -3a, -3b and -4 cluster more closely to the two group F Arabidopsis *bZIPs* shown to be involved in the Zn-deficiency response (*AtbZIP19* and *AtbZIP23*) rather than to *AtbZIP24*, which has a reported function in salt tolerance (Yang *et al.*, 2009a).

### Group F *bZIP* expression is induced by low-Zn conditions

The relative gene expression levels of *TabZIPF1*, -3a, -3b and -4 were determined in root and shoot material from a 1-week period of Zn starvation. The four *TabZIPFs* were all induced under Zn-deficient conditions in both roots and shoots but to rather different extents (Figure 5). *TabZIPF3a*, -3b and -4 showed the greatest induction in both roots and shoots. In contrast, *TabZIPF1* was slower to respond to the -Zn treatment, taking until day 5 in the root and day 3 in the shoot to be expressed to a significantly higher level. The magnitude of induction of *TabZIPF1* was less than that of the other *TabZIPFs* examined.

### Group F *bZIPs* rescue the Zn-deficiency phenotype of an Arabidopsis *bzip19-4 bzip23-2* mutant indicating conserved homeostatic mechanisms

*TabZIPF1-7DL*, *TabZIPF3b-7BL*, *TabZIPF4-7AL* and *TabZIPF4-7DL* were investigated further due to variations in the conservation of their group F motif 1. They were cloned into the Arabidopsis expression vector pMDC32 (Curtis and Grossniklaus, 2003), which contains a dual 35S CaMV constitutive promoter. The sequences of the four cloned *TabZIPFs* are shown in Figure S6. All four cloned *TabZIPs* show high similarity to *AtbZIP19* and *AtbZIP23* in both the *bZIP* domain and group F motif 2; however, the group F motif 1 is less conserved. *TabZIPF3b-7BL* and *TabZIPF4-7DL* both contain a truncated group F motif 1 compared with the other group F *bZIPs* as well as their homeologs. *TabZIPF1-7DL* and *TabZIPF4-7AL* have a group F motif 1 with higher similarity to *AtbZIP19* and *AtbZIP23*. To test their importance in the Zn-deficiency response mechanism, the ability of these four *TabZIPs* to rescue the Zn hypersensitivity of the Arabidopsis *bzip19 bzip23* double mutant was tested. An alternative double mutant Arabidopsis line, *bzip19-4 bzip23-2*, was created by Nazri *et al.* (2017) that demonstrates the same Zn hypersensitivity as the Arabidopsis *bzip19-1 bzip23-1* double mutant previously characterised (Assunção *et al.*, 2010). The *bzip19-4 bzip23-2* double mutant was transformed with the four



**Figure 3.** *TaZIP* gene expression analysis in wheat material throughout a 1-week Zn starvation period. Relative expression levels (normalised relative quantification, NRQ) of five wheat *ZIP* transporter gene transcripts in root and shoot material throughout 1 week of Zn starvation. NRQ values are normalised to *TaActin3* and *TaSuccDH* expression and means of three biological replicates are given ( $\pm$ SEM). Bars within individual graphs displaying an asterisk show a significant difference between treatment means at a given time point. Significance ( $P < 0.05$ ) was tested *post hoc*, using Fisher's least significant difference test on  $\log_2(1/\text{NRQ})$ -transformed data. +Zn = 8  $\mu\text{M}$  Zn (grey bars), -Zn = 0  $\mu\text{M}$  Zn (white bars).


*TabZIP*s and two independent transgenic lines were characterised in detail. These were compared with the wild type and the non-transformed double mutants on half MS media with (15  $\mu\text{M}$ ) and without (0  $\mu\text{M}$ ) Zn. Expression of both *TabZIPF1-7DL* and *TabZIPF4-7AL* significantly improved the performance of the *bzip19-4 bzip23-2* double mutant under low-Zn conditions and plants displayed a significantly higher fresh weight than the mutant (Figures 6 and S7). *TabZIPF3b-7BL* expression slightly increased the growth of the double mutant in one line only under low Zn, though to a lesser extent than both *TabZIPF1-7DL* and *TabZIPF4-7AL*. *TabZIPF4-7DL* provided no rescue to the double mutant (Figure 6).

#### Ability of *TabZIP* to bind to ZDRE motifs identified in *TaZIP* promoters

The rescue of the *bzip19 bzip23* double mutant by the wheat *bZIP*s suggested a conserved mechanism of action

for these group F *bZIP* transcription factors. In *Arabidopsis* *bZIP19* functions in binding to ZDRE motifs in Zn-responsive genes (Assunção *et al.*, 2010). Therefore, the presence of ZDREs in the promoters of *TaZIP* genes was determined. Regions upstream of the start codon were analysed at up to 2000 bp where possible. Motifs were regarded as potential ZDREs if they had no more than one mismatch to the ZDRE consensus (RTGTCGACAY) reported by Assunção *et al.* (2010). ZDREs were present in the promoter regions of a number of the *ZIP* genes; these are shown in Table S3 with their location relative to the predicted start codon. *TaZIP*s with a completely conserved ZDRE domain include *TaZIP5*, -7, -8 and -10; additionally *TaZIP1*, -3, -6, -9 and -13 contain at least one ZDRE motif with one mismatch to the consensus in their promoter sequence.

To directly test the regulatory link between the *TabZIP*s and *TaZIP*s, the binding ability of *in vitro* synthesised *bZIP*s to putative ZDRE motifs discovered in the promoter




**Figure 4.** Phylogenetic tree of group F bZIP proteins from cereals and Arabidopsis.

A neighbour-joining tree was generated for Arabidopsis (At), rice, *Oryza sativa* (Os), *Brachypodium distachyon* (Bd), barley, *Hordeum vulgare* (Hv), and wheat, *Triticum aestivum* (Ta) group F bZIP coding sequence translations. The Muscle algorithm (Edgar, 2004) was used for the alignment of sequences and the phylogenetic tree was created using MEGA (v.5.2) software. Evolutionary distances were computed using the *p*-distance method and are in units of the number of amino acid differences per site. A thousand bootstrap replicates were used and bootstrap values are shown as percentages. Gene nomenclature for Arabidopsis is from Jakoby *et al.* (2002), *Brachypodium* is from Liu and Chu (2015), rice is from Corrêa *et al.* (2008) and barley is from Pourabed *et al.* (2015). Full accession information for the wheat genes is provided in Table S2.

regions of *TaZIPs* was determined (Figure 7). *AtbZIP19*, used as a control, created a band shift (indicative of binding) when two or three copies of the *AtZIP4* ZDRE were present (probe names *Ass2Z* and *Ass3Z*), but not with a mutated three-copy *AtZIP4* ZDRE probe (*Ass3Zmut*). *AtbZIP19* also produced band shifts with the previously untested *TaZIP* ZDRE probes 3, 5, 13 and 7. No shift was observed with the *TaZIP6* ZDRE probe, suggesting *AtbZIP19* does not interact with the putative ZDRE identified in the promoter of *TaZIP6* (Figure 7a).

*TabZIPF1-7DL* created a band shift with *Ass3z* and *Ass2z*, which are based on *AtZIP4* ZDREs, but not the mutated version, *Ass3Zmut*. *TaZIP* ZDRE probes 3, 5, 13 and 7 all produced a band shift indicating an interaction of *TabZIPF1-7DL* with these ZDREs. As with *AtbZIP19*, however, *TaZIP6* ZDRE did not produce a band shift (Figure 7b). *TabZIPF3b-7BL* was only able to bind to the probe containing three *AtZDREs* (*Ass3Z*) and none of the *TaZIP* ZDRE probes (Figure 7c). *Ass3Z* is a modification of the ZDREs present in the promoter of *AtZIP4*. It has three



**Figure 5.** *TabZIP* gene expression analysis in wheat material throughout a 1-week Zn starvation period. Relative expression levels (normalised relative quantification, NRQ) of five wheat *bZIP* transcription factor gene transcripts in root and shoot material throughout 1 week of Zn starvation. NRQ values are normalised to *TaActin3* and *TaSuccDH* expression and means of three biological replicates are given ( $\pm$ SEM). Bars within individual graphs displaying an asterisk show a significant difference between treatment means at a given time point. Significance ( $P < 0.05$ ) was tested *post hoc*, using Fisher's least significant difference test on  $\log_2(1/\text{NRQ})$ -transformed data. +Zn = 8  $\mu\text{M}$  Zn (grey bars), -Zn = 0  $\mu\text{M}$  Zn (white bars).

ZDREs adjacent to each other whereas the promoter has two ZDREs spaced 128-bp apart (Figure 7). Neither *TabZIPF4-7AL* nor *TabZIPF4-7DL* bound to any of the ZIP ZDRE probes tested (Figure 7d, e).

## DISCUSSION


### TaZIPs have an important role in the transport of Zn

Analysis of the wheat genome revealed 14 *TaZIPs*, each with a homeolog on the A, B and D genomes. This is an increase in the number in the previous phylogenetic analysis of *TaZIPs* (Tiong *et al.*, 2015) where 12 *TaZIPs* were identified and homeologs were not reported. The lengths ranged from 277 to 578 amino acids with much of the variation coming from the more divergent homologs *TaZIP11*, -14 and -16 (homologs of *AtPutZnT*, *AtIAR* and *AtZTP29*, respectively). *TaZIP11*, -14 and -16 were more distantly related to the other *TaZIPs* identified, suggesting a different role or cellular localisation in the plant. This is supported in knockout studies of *AtIAR* in Arabidopsis and heterologous expression of *AtZTP29* in yeast (Lasswell *et al.*, 2000; Wang *et al.*, 2010).

Functional characterisation using the yeast *zrt1/zrt2* mutant shows that *TaZIP3*, -6, -7, -9 and -13 rescue the Zn

hypersensitivity of this mutant. When considered alongside the Zn-induced gene expression patterns of the *TaZIPs* determined in this study, this is strong evidence that ZIP members from this clade are important Zn transporters in wheat. Although not tested in yeast, *TaZIP5* was induced under Zn-deficient conditions, suggesting a role in the Zn homeostatic framework. Barley homologs of three *TaZIPs* investigated in this study have shown Zn-responsive expression (*TaZIP3/HvZIP3*, *TaZIP5/HvZIP5* and *TaZIP7/HvZIP7*) (Pedas *et al.*, 2009; Tiong *et al.*, 2015). Previously, from yeast studies, Pedas *et al.* (2009) indicated that *HvIRT1* and *HvZIP3*, -5 and -8 could transport Zn. Additionally, work in yeast suggests that the wild emmer wheat (*T. turgidum* ssp. *dicoccoides*) *TdZIP1* (*TaZIP3* homolog) may also function as a Zn transporter (Durmaz *et al.*, 2011). The five *TaZIPs* characterised in this study were able to complement the yeast *zrt1/zrt2* mutant but were unable to rescue the Fe-uptake mutant *fet3/fet4*, suggesting there is substrate selectivity for these particular ZIPs.

The barley *TaZIP6* homolog *HvZIP6* is unresponsive to Zn status in the shoot but expression has been shown to increase in the root (Tiong *et al.*, 2015). Additionally, *OsZIP6* expression is induced in both the roots and the shoots of Zn-deficient rice plants (Kavitha *et al.*, 2015). In




**Figure 6.** Functional complementation of the Arabidopsis *bzip19-4 bzip23-2* mutant with group F TabZIPs. Complementation analysis with (a) TabZIPF1-7DL, (b) TabZIPF3b-7BL, (c) TabZIPF4-7AL and (d) TabZIPF4-7DL. Fresh weight analysis results shown are mean average fresh weights per seedling  $\pm$  SEM from six plates with four seedlings per plate, per line ( $n = 24$ ). Bars with different letters indicate a significant difference ( $P < 0.05$ ) tested on log-transformed data using Fisher's least significant difference test. +Zn = 15  $\mu$ M Zn (grey bars) and -Zn = 0  $\mu$ M Zn (white bars). Representative plates at both the +Zn (15  $\mu$ M Zn) and -Zn (0  $\mu$ M Zn) media concentrations are shown. wt indicates the wild-type line and dm indicates the untransformed *bzip19-4 bzip23-2* line. #1 and #2 are two corresponding, independent TabZIP transformed *bzip19-4 bzip23-2* lines. All plates illustrated are 18 days post-germination.

this study we show that *TaZIP6* expression was not induced in the root during 1 week of Zn-deficient growth but was significantly induced in the shoot after 5 days. The specific spatiotemporal expression patterns indicate that individual *TaZIPs* may have precise roles within the plant. Feasibly the Zn requirement of different regions in the root and shoot varies and may explain the differential pattern of increased gene expression initiation observed and its magnitude.

### Specific group F bZIPs induce the transcription of key Zn transporters, indicating a conserved homeostatic mechanism

The analysis presented here discovered seven distinct sets of group F bZIPs present in the wheat genome. Amongst the *TabZIPs* identified (and between homeologs), conservation of the cysteine-histidine-rich group F motif 1 varied. The group F motif 2 was widely conserved, as was the



**Figure 7.** Electrophoretic mobility shift assays determining the binding ability of AtbZIP19 and TabZIPs to Zn-deficiency-response element (ZDRE)-containing probes.

Electrophoretic mobility shift assays were used to investigate the binding ability of AtbZIP19 (a) and five TabZIPs [TabZIPF1-7DL (b), TabZIPF3b-7BL (c), TabZIPF4-7AL (d) and TabZIPF4-7DL (e)] to Arabidopsis ZDRE-containing probes and wheat ZDRE-containing probes. Lanes C1–C3 are control lanes: C1 is TabZIPF protein only, no probe. C2 is probe (Ass3Z), no protein. C3 is probe (Ass3Z) with TNT wheat germ mix (no template). All other lanes contain *in vitro* synthesised TabZIPs (as labelled) with the following probes: lane A3 is Ass3Z, a3 is Ass3Zmut (mutated Ass3Z) and A2 is Ass2Z which all contain Arabidopsis ZDREs. Z3–Z13 are TaZIP3ZDRE–TaZIP13ZDRE, which all contain wheat ZDREs. Asterisks indicate a band shift. Probe sequences are provided in Table S7.

general bZIP domain. Previously, the two cysteine–histidine-rich motifs present in group F bZIPs were hypothesised to be the binding site for  $Zn^{2+}$  ions (Assunção *et al.*, 2010, 2013). As a small and efficient electron acceptor,  $Zn^{2+}$  is able to form tetrahedral complexes with sulphur and nitrogen ligands found in the side chains of cysteine and histidine (Tauris *et al.*, 2009; Pace and Weerapana, 2014).

The role of four *TabZIPFs* in the Zn homeostatic mechanism was investigated by expression in the Arabidopsis *bzip19-4 bzip23-2* double mutant line. *TabZIPF1-7DL* and *TabZIPF4-7AL* provided partial complementation of this mutant, resulting in improved growth under the  $-Zn$  treatment compared with the untransformed *bzip19-4 bzip23-2* line. This indicates that the role of these genes in wheat is likely to be in the Zn-deficiency response, and furthermore that they play a similar role in wheat and Arabidopsis. They did not completely rescue the *bzip19-4 bzip23-2* mutant. This may be due to slight differences in  $Zn^{2+}$  affinity of the TabZIPs, or a reduced binding ability to ZDREs in the promoter regions of Zn-responsive Arabidopsis genes. *TabZIPF1-7DL* and *TabZIPF4-7AL* show divergence from AtbZIP19 and AtbZIP23 in the 3' end of group F motif 1 (Figure S6), this may increase the  $Zn^{2+}$  affinity of these TabZIPs, although this requires further study.

Binding of AtbZIP19 and *TabZIPF1-7DL* to ZDREs found in the promoter of *AtZIP4* as well as *TaZIP3*, *-5*, *-7* and *-13* demonstrated that an analogous Zn-homeostatic mechanism exists in the two species. *TabZIPF4-7AL* partially rescued the *bzip19-4 bzip23-2* line, but did not bind to the

ZDRE-containing probes *in vitro*. This suggests this bZIP functions in the Zn-deficiency response of wheat but the lack of binding indicates that either it acts through a different binding mechanism or that, potentially, binding is not observed here due to a requirement for native ZIP promoter flanking sequence or for a narrow and specific Zn concentration range not simulated in the electrophoretic mobility shift assay (EMSA).

*TabZIPF3b-7BL* and *TabZIPF4-7DL* were unable to restore the Zn-deficiency response of the *bzip19-4 bzip23-2* mutant. Neither of these *TabZIPFs* contains a well-conserved group F motif 1, which could be a causal factor for their lack of rescue ability. Despite being homeologs with high sequence similarity (84%), *TabZIPF4-7AL* and *TabZIPF4-7DL* showed different rescue abilities. This is most likely due to their different levels of group F motif 1 conservation. Neither *TabZIPF3b-7BL* nor *TabZIPF4-7DL* were able to bind to TaZDRE-containing probes *in vitro*. Their presence in the wheat genome and expression *in planta* could be the result of genetic redundancy due to the hexaploidy of wheat, although an alternative hypothesis is a role in the fine tuning of the Zn homeostatic mechanism in wheat. The truncated motif in these *TabZIPFs* may facilitate the increased expression of specific Zn homeostatic genes at precise plant tissue locations. This may permit the plant to alter expression of particular genes in response to certain conditions or developmental stages without changing the expression of an entire suite of genes throughout the whole plant at a single given point in the Zn concentration continuum.

The expression profiles of *TabZIPF1*, *-3a*, *-3b* and *-4* show they were induced, although to different degrees, under Zn-deficient conditions. The current Zn-homeostatic model for adapting to Zn deficiency suggests that dissociation of  $Zn^{2+}$  from group F bZIPs results in their binding to ZDRE motifs in the promoters of Zn-responsive genes to increase their expression, seen particularly in the ZIPs (Assunção *et al.*, 2010). Transcriptional regulation of the bZIPs themselves was not included as part of the model. Modifications may include a further level of intricacy, revealed here in wheat, whereby induction of bZIPs would allow for a sustained adaptive response under prolonged deficiency. Due to the differential effects observed across the *TabZIPs* in response to Zn deficiency, it is possible that certain *bZIPs*, notably the members with lower induction due to Zn deficiency, are 'master regulators' of other *bZIPs*. *TabZIPF1* may exert a global induction of *ZIPs* at critical levels of Zn deficiency while also influencing other *bZIPs* that then contribute to further adaptive responses. As such, *TabZIPF1* may be a promising target for future breeding or transgenic strategies.

In summary, our findings demonstrate that wheat contains a complex system for regulating Zn homeostasis. It has an expanded number of group F bZIP transcription factors that serve in altering the expression of ZIPs by binding to ZDREs in their promoters. These genes show differences in spatial and temporal expression patterns, indicating that there is a homeostatic network allowing adaptation to low Zn availability by contributing to Zn uptake and distribution under changing nutrient availability. The variation in conservation of cysteine-histidine-rich motifs throughout the wheat group F bZIPs may provide additional refinement to the homeostatic mechanism.

## EXPERIMENTAL PROCEDURES

### Bioinformatics analysis

Gene and coding sequences of wheat *ZIPs* and *bZIPs* were identified using a Blast analysis of the IWGSC wheat survey sequence database and the TGACv1 assembly (<http://www.plants.ensemble.org>) with rice, barley and *Brachypodium* homologs. Phylogenetic analysis was performed by multiple protein sequence alignment using GENIEOUS v.8.1.3 (<http://www.geneious.com>; Kearse *et al.*, 2012) with the Muscle algorithm (Edgar, 2004). MEGA5 (Tamura *et al.*, 2011) was used for calculation of phylogenetic trees using the neighbour-joining method with evolutionary distances computed using the  $p$ -distance method and 1000 bootstrap replicates conducted for each phylogeny.

### Plant growth (*T. aestivum* and *Arabidopsis thaliana*)

**Hydroponic culture 1-week starvation experiment.** *Triticum aestivum* cv. Paragon seedlings were germinated on sterile water-soaked soft paper tissue for 7 days before transfer to single-plant hydroponic aerated culture. Plants were cultivated for the first 3 days on half-strength before changing to full-strength modified Letcombe liquid nutrient solution [1.5 mM  $Ca(NO_3)_2$ , 5 mM  $KNO_3$ , 2 mM  $NaNO_3$ , 1 mM  $MgSO_4$ , 0.5 mM  $KH_2PO_4$ , 25  $\mu M$  FeEDTA,

0.2  $\mu M$   $CuCl_2 \cdot 2H_2O$ , 1  $\mu M$   $H_3BO_3$ , 0.6  $\mu M$   $MnCl_2 \cdot 4H_2O$ , 0.1  $\mu M$   $Na_2MoO_4 \cdot 2H_2O$ , 5  $\mu M$  KCl, 8  $\mu M$   $ZnCl_2$ , 2.56 mM 2-(*N*-morpholine)-ethanesulphonic acid (MES) buffer, pH = 5.8; Drew and Saker, 1984] including the chelator 75.5  $\mu M$  HEDTA. The solution was changed three times a week. Two weeks after germination, plants (apart from the +Zn control plants) were Zn-starved by omitting the  $ZnCl_2$  from the culture solution (-Zn). Plants were grown in a growth chamber with 16-h day conditions of 20°C/70% humidity and 500  $\mu mol m^{-2} sec^{-1}$  light and night conditions of 16°C/80% humidity. Samples were taken at day 0, 1, 3, 5 and 7 of Zn starvation. Whole roots (washed with deionised water and dried briefly on soft paper towels) and whole shoots were sampled, weighed and immediately frozen in liquid nitrogen and stored at -80°C.

**Hydroponic culture 3-week starvation experiment.** Conditions as above except for a reduction to 7 days rather than 14 days of hydroponic culture in +Zn solution prior to Zn-starving the plants.

*Arabidopsis thaliana* plants were grown in growth chambers with 16-h day conditions, 23°C and 120  $\mu mol m^{-2} sec^{-1}$  light, night conditions 18°C. Soil used was autoclaved and comprised equal proportions of vermiculite, Levingtons M2 and John Innes No. 2 compost in 8-cm pots supplemented with 0.28 g  $L^{-1}$  Intercept insecticide (Bayer, <https://www.bayer.com/>).

### Plant total RNA isolation

All wheat material was homogenised using a SPEX freezer mill (SPEX CertiPrep Ltd, <http://www.spexcsp.com/>) in liquid nitrogen, aliquotted into 2-ml micro-tubes and stored at -80°C. Total RNA was isolated by a modified method based on Verwoerd *et al.* (1989) with additional phenol-chloroform-isoamyl alcohol extractions. Possible genomic DNA contamination was removed by RNase-free DNase treatment. The final air-dried pellet was dissolved in an appropriate volume of RNase-free water.

### Cloning full-length *TabZIPs* and *TaZIPs*

First-strand cDNA synthesis was performed from 2  $\mu g$  total RNA and a dT-adapter primer (Invitrogen Superscript III, <http://www.invitrogen.com/>; standard protocol, 2 h synthesis time). Full-length *TabZIPs* were amplified using the oligonucleotide primers specified in Table S4 with *Pfu* DNA polymerase (Promega, <http://www.promega.com/>). *TabZIP* amplicons were subsequently topoisomerase-cloned into pENTR/D-TOPO (Invitrogen) and transformed into *Escherichia coli*. *TabZIP* plant expression vectors were created by Gateway (Invitrogen) recombination of pENTR:*TabZIP* vectors into pMDC32 (Curtis and Grossniklaus, 2003).

Full-length *TaZIPs* were amplified using Q5 High-fidelity DNA polymerase (New England Biolabs, <https://www.neb.com/>). *TaZIP* amplicons were cloned into pGEM-T Easy vector (Promega) and subcloned into a pYES2 yeast expression vector using *EcoRI* digestion and subsequent ligation. All vectors were sequenced before *Arabidopsis* or yeast transformation.

### Functional complementation in yeast

The following strains of the yeast *Saccharomyces cerevisiae* were used in this study: wild-type DY1457 (MATa, *ade1*+/ *can1*, *his3*, *leu2*, *trp1*, *ura3*), *zrt1/zrt2* (DY1457 + *zrt1::LEU2*, *zrt2::HIS3*) and *fet3/fet4* (DY1457 + *fet3-2::HIS3*, *fet3-1::LEU2*). Yeast strains were transformed with pYES2*TaZIP* vectors using the S.c. EasyComp Transformation Kit (Invitrogen) according to the manufacturer's instructions. Following transformation, PCR-confirmed pYES2*TaZIP* containing *zrt1/zrt2* colonies was inoculated in 10 ml of SC-glucose minus uracil

(pH 5.3) overnight (30°C with shaking at 200 r.p.m.). Inoculums were pelleted (3 min, 1300 *g*), suspended in SC-galactose minus uracil (pH 5.3) and incubated for 4 h (30°C with shaking at 200 r.p.m.) to allow gene induction. Inoculums were then pelleted and washed with SC-galactose minus uracil minus Zn (pH 5.3) to remove excess Zn from the pellet. Inoculums were diluted to OD<sub>600</sub> = 0.4 using SC-glucose minus uracil minus Zn (pH 5.3) and serial dilutions made (1:2, 1:10, 1:100 and 1:1000). Seven microlitres of the dilutions was plated onto SC-galactose minus uracil plates (pH 5.3) containing either 200 μM ZnSO<sub>4</sub> (+Zn), 0 Zn + 0 mM, 2 mM, 5 mM and 7.5 mM EGTA, incubated at 30°C and photographed after 8 days.

Following *fet3/fet4* transformation, positive colonies were confirmed using PCR and inoculated in 10 ml SC-glucose minus uracil + 10 μM FeCl<sub>3</sub> (pH 4.0) overnight (30°C with shaking at 200 r.p.m.). Inoculums were pelleted (3 min, 1300 *g*), suspended in SC-galactose minus uracil + 10 μM FeCl<sub>3</sub> (pH 4.0) and incubated for 4 h (30°C with shaking at 200 r.p.m.) to allow gene induction. Inoculums were diluted to OD<sub>600</sub> = 0.4 using SC-glucose minus uracil + 10 μM FeCl<sub>3</sub> (pH 4.0) and serial dilutions made (1:2, 1:10, 1:100 and 1:1000). Seven microlitres of the dilutions was plated onto SC-galactose minus uracil minus Fe (pH 4.0) plates supplemented with 0 μM, 0.74 μM or 10 μM FeCl<sub>3</sub>, incubated at 30°C and photographed after 3 days.

### Expression analysis of TaZIPs and TabZIPs

Real-time PCR was performed using the Applied Biosystems 7500 Real Time PCR System and the SYBR<sup>®</sup> Green Jumpstart<sup>™</sup> Taq ReadyMix<sup>™</sup> (Sigma-Aldrich, <http://www.sigmaaldrich.com>). The 25-μl reactions contained 1 μl of cDNA and 250 nM of each primer. The primers used are given in Table S5 and were designed to cover gene expression of the homeologous genes from all three wheat genomes. Mean primer efficiencies were estimated using the linear phase of all individual reaction amplification curves (Ramakers *et al.*, 2003) calculated using the LinRegPCR package (Ruijter *et al.*, 2009). The stable, constitutive genes Actin 3 and Succinate Dehydrogenase were used to determine the normalised quantification of expression. The normalised relative quantity (NRQ) of expression was calculated in relation to the Ct values and the primer efficiency (*E*) of both the target gene (*X*) and the mean of the two normalisation reference genes (*N*) as normalised relative expression (NRE) based on Rieu and Powers (2009):

$$\text{NRE} = (E_X)^{-\text{Ct}_X} / (E_N)^{-\text{Ct}_N}$$

Statistically significant changes in relation to Zn treatment were calculated using a two-way analysis of variance (ANOVA) on log<sub>2</sub>(1/NRQ)-transformed data followed by a *post hoc* Fisher's least significant difference test (LSD) at the 5% significance level. All statistical analyses were done in Genstat 17th edition (VSN International, <https://www.vsn.co.uk/>).

### Complementation of the Arabidopsis *bzip19-4 bzip23-2* mutant line with group F TabZIPs

pMDC32TabZIP plasmids were transformed into *Agrobacterium tumefaciens* GV3850 by electroporation. *Arabidopsis thaliana* (Col-8) *bzip19-4 bzip23-2* double mutants (Nazri *et al.*, 2017) were transformed using the floral dip method (Clough and Bent, 1998) with an additional 3-h pre-induction of *vir* genes by the addition of 100 μM acetosyringone to the culture prior to floral dipping. Homozygous T<sub>3</sub> plants were used in subsequent phenotype assays.

Growth assays were conducted using two independent T<sub>3</sub> homozygous lines from each pMDC32-TabZIP transformation. Four sterilised seeds of each T<sub>3</sub> line, the Col-8 wild type and the

untransformed *bzip19-4 bzip23-2* were plated onto six square plates of 0.5 × MS medium (Murashige and Skoog, 1962) containing 1% (w/v) sucrose and 0.8% (w/v) agarose containing 15 μM ZnSO<sub>4</sub> (+Zn) and 0 μM ZnSO<sub>4</sub> (-Zn). Following 18 days of growth, plates were photographed and root and shoot fresh weights were measured. Individual fresh weights were obtained by calculating the average from the combined weight of four seedlings prior to statistical analysis. A two-way ANOVA was performed on log-transformed data for each complementation experiment (the four TabZIPs investigated) followed by a *post hoc* Fisher's LSD at the 5% significance level.

### Electrophoretic mobility shift assay

Full-length coding sequences of bZIPs were PCR-amplified from sequenced vectors using primer pairs containing an SP6 promoter and Kozak region (forward) and a poly-A tail (reverse) (Table S6). bZIP proteins were *in vitro* translated using the T<sub>N</sub>T SP6 High-Yield Wheat Germ Protein Expression Kit (Promega) following the manufacturer's instructions with 690 ng of purified PCR product added to 18 μl of wheat germ master mix. ZDREs containing biotinylated oligonucleotides and nonbiotin-labelled complementary oligonucleotides were synthesised (Table S7) (Eurofins, <https://www.eurofins.com/>) and annealed at a concentration of 1 pmol μl<sup>-1</sup>, 10 mM 2-amino-2-(hydroxymethyl)-1,3-propanediol (TRIS), 1 mM EDTA, 50 mM NaCl for 5 min at 95°C and slowly cooling to room temperature (20°C) overnight. *In vitro* translated proteins (3 μl of the 30-μl reaction) were incubated with 4 μl of the annealed oligonucleotide solution in a 20-μl binding reaction containing 20 mM TRIS-HCl (pH 7.5), 10 mM KCl, 1 mM EDTA, 0.25 μg μl<sup>-1</sup> BSA, 1 mM DTT and 0.25 μg μl<sup>-1</sup> salmon sperm DNA at 28°C for 30 min. The DNA-protein complex was analysed on a 6% native PAGE DNA retardation gel (Thermo Fisher Scientific, <https://www.thermofisher.com/>) using 0.5 × TRIS borate-EDTA at 100 V for 75 min. After electrophoresis, the gel was blotted to Amersham Hybond-N<sup>+</sup> membrane (GE Healthcare, [http://www3.gehealthcare.com/en/global\\_gateway](http://www3.gehealthcare.com/en/global_gateway)) using the XCell II<sup>™</sup> Blot module (Thermo Fisher Scientific). The membrane was crosslinked using a Stratalinker<sup>®</sup> UV crosslinker and the signal detected using Chemiluminescent Nucleic Acid Detection Module (Thermo Fisher Scientific) according to manufacturer's instructions using the Odyssey<sup>®</sup> FC imaging system (Li-Cor, <https://www.licor.com/>), with a 2-min exposure.

### Determination of the Zn content in plant material

Cryogenically milled roots and shoots of hydroponically grown *T. aestivum* cv. Paragon were freeze dried for 2 days, then digested in 5 ml of nitric acid:perchloric acid (85:15, v/v; 70% concentration, trace analysis grade; Fisher Scientific) for a minimum of 2 h at room temperature followed by a 5-h programmed thermoblock cycle. Five millilitres of 25% (v/v) nitric acid was added to the solution and the tubes were reheated for 1 h at 80°C. Ultra-pure water (>18 MΩ) was added to to make up to 9 ml, mixed well and re-warmed for a further 30 min at 80°C. After cooling, the solutions were made up to final volumes of 10 ml with deionised H<sub>2</sub>O. Inductively coupled plasma optical emission spectrometry analysis was carried out using an Optima inductively coupled plasma-optical emission spectrometer (Perkin Elmer Life and Analytical Sciences, <http://www.perkinelmer.com/>).

### ACKNOWLEDGEMENTS

This study was supported by a University of Southampton and Rothamsted Research BBSRC DTP studentship (BB/J014451/1) and the 20:20 Wheat<sup>®</sup> programme (BBS/E/C/00005202) at Rothamsted

Research. Rothamsted Research receives strategic funding from the BBSRC.

## CONFLICTS OF INTEREST

The authors confirm that they have no conflicts of interest to declare.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

**Figure S1.** Phylogenetic tree of *Triticum aestivum* and barley ZIP proteins showing homeolog grouping.

**Figure S2.** Multiple sequence alignment of TaZIPs.

**Figure S3.** Phenotypic effect of Zn starvation throughout a 3-week growth period.

**Figure S4.** Mineral concentration analysis of wheat root and shoot samples during a 1-week Zn- starvation period.

**Figure S5.** Gene expression analysis of TaZIPs in wheat root and shoot material throughout an extended Zn starvation period of 3 weeks.

**Figure S6.** Multiple sequence alignment of cloned group F TabZIPs, AtbZIP19 and AtbZIP23.

**Figure S7.** Confirmatory PCR of 35S::TabZIP-transformed Arabidopsis lines.

**Table S1.** TaZIP identity matrix.

**Table S2.** Wheat TaZIP and TabZIP gene identification details.

**Table S3.** Overview of ZDREs present in promoters of TaZIPs.

**Table S4.** Oligonucleotide primer sequences used for cloning of full length TaZIPs and TabZIPs.

**Table S5.** Oligonucleotide primer sequences used for SYBR Green real time RT-PCR expression analysis.

**Table S6.** Oligonucleotide primer sequences used for PCR-amplification of TabZIPs. With SP6 promoters and Poly-A tails prior to in vitro transcription translation.

**Table S7.** Complementary oligonucleotides used in EMSAs.

## REFERENCES

- Assunção, A.G.L., Herrero, E., Lin, Y., Huettel, B., Talukdar, S. and Smaczniak, C. (2010) *Arabidopsis thaliana* transcription factors bZIP19 and bZIP23 regulate the adaptation to zinc deficiency. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 10296–10301.
- Assunção, A.G.L., Persson, D.P., Husted, S., Schjørring, J.K., Alexander, R.D. and Aarts, M.G.M. (2013) Model of how plants sense zinc deficiency. *Metallomics*, **5**, 1110–1116.
- Broadley, M.R., White, P.J., Hammond, J.P., Zelko, I. and Lux, A. (2007) Zinc in plants. *New Phytol.* **173**, 677–702.
- Brown, P., Cakmak, I. and Zhang, Q. (1993) Form and function of zinc in plants. In *Zinc in Soils and Plants* (Robson, A.D. ed). Dordrecht, the Netherlands: K. A. Publishers, pp. 93–106.
- Bughio, N., Yamaguchi, H., Nishizawa, N.K., Nakanishi, H. and Mori, S. (2002) Cloning an iron-regulated metal transporter from rice. *J. Exp. Bot.* **53**, 1677–1682.
- Cakmak, I. (2000) Tansley Review No.111: possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *New Phytol.* **146**, 185–205.
- Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735–743.
- Corréa, L.G.G., Riaño-Pachón, D.M., Schrago, C.G., dos Santos, R.V., Mueller-Roeber, B. and Vincenz, M. (2008) The role of bZIP transcription factors in green plant evolution: adaptive features emerging from four founder genes. *PLoS ONE*, **3**, e2944.
- Curtis, M.D. and Grossniklaus, U. (2003) A Gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant Physiol.* **133**, 462–469.
- Drew, M.C. and Saker, L.R. (1984) Uptake and long-distance transport of phosphate, potassium and chloride in relation to internal ion concentrations in barley; evidence of non-allosteric regulation. *Planta*, **160**, 500–507.
- Durmaz, E., Coruh, C., Dinler, G. et al. (2011) Expression and cellular localization of ZIP1 transporter under zinc deficiency in wild emmer wheat. *Plant Mol. Biol.* **29**, 582–596.
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797.
- Grotz, N., Fox, T., Connolly, E., Park, W., Guerinot, M.L. and Eide, D. (1998) Identification of a family of zinc transporter genes from *Arabidopsis* that respond to zinc deficiency. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 7220–7224.
- Inaba, S., Kurata, R., Kobayashi, M., Yamagishi, Y., Mori, I., Ogata, Y. and Fukao, Y. (2015) Identification of putative target genes of bZIP19, a transcription factor essential for *Arabidopsis* adaptation to Zn deficiency in roots. *Plant J.* **84**, 323–334.
- Ishimaru, Y., Suzuki, M., Kobayashi, T., Takahashi, M., Nakanishi, H., Mori, S. and Nishizawa, N.K. (2005) OsZIP4, a novel zinc-regulated zinc transporter in rice. *J. Exp. Bot.* **56**, 3207–3214.
- Ishimaru, Y., Suzuki, M., Tsukamoto, T. et al. (2006) Rice plants take up iron as an Fe<sup>3+</sup>-phytosiderophore and as Fe<sup>2+</sup>. *Plant J.* **45**, 335–346.
- Jakoby, M., Weisshaar, B., Dröge-Laser, W., Vicente-Carbajosa, J., Tiedemann, J., Kroj, T. and Parcy, F. (2002) bZIP transcription factors in *Arabidopsis*. *Trends Plant Sci.* **7**, 106–111.
- Kavitha, P., Kuruvilla, S. and Mathew, M. (2015) Functional characterization of a transition metal ion transporter, OsZIP6 from rice (*Oryza sativa* L.). *Plant Physiol. Biochem.* **97**, 165–174.
- Keare, M., Moir, R., Wilson, A. et al. (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, **28**, 1647–1649.
- Kumssa, D.B., Joy, E.J.M., Ander, E.L., Watts, M.J., Young, S.D., Walker, S. and Broadley, M.R. (2015) Dietary calcium and zinc deficiency risks are decreasing but remain prevalent. *Sci. Rep.* **5**, 1–11.
- Lasswell, J., Rogg, L.E., Nelson, D.C., Rongey, C. and Bartel, B. (2000) Cloning and characterization of IAR1, a gene required for auxin conjugate sensitivity in *Arabidopsis*. *Plant Cell*, **12**, 2395–2408.
- Lee, S., Jeong, H.J., Kim, S.A., Lee, J., Guerinot, M.L. and An, G. (2010a) OsZIP5 is a plasma membrane zinc transporter in rice. *Plant Mol. Biol.* **73**, 507–517.
- Lee, S., Kim, S.A., Lee, J., Guerinot, M.L. and An, G. (2010b) Zinc deficiency-inducible OsZIP8 encodes a plasma membrane-localized zinc transporter in rice. *Mol. Cells*, **29**, 551–558.
- Li, X., Gao, S., Tang, Y., Li, L., Zhang, F., Feng, B., Fang, Z., Ma, L. and Zhao, C. (2015) Genome-wide identification and evolutionary analyses of bZIP transcription factors in wheat and its relatives and expression profiles of anther development related TabZIP genes. *BMC Genom.* **16**, 1–21.
- Liu, X. and Chu, Z. (2015) Genome-wide evolutionary characterization and analysis of bZIP transcription factors and their expression profiles in response to multiple abiotic stresses in *Brachypodium distachyon*. *BMC Genom.* **16**, 1–15.
- Milner, M., Seamon, J., Craft, E. and Kochain, L. (2013) Transport properties of members of the ZIP family in plants and their role in Zn and Mn homeostasis. *J. Exp. Bot.* **64**, 369–381.
- Murashige, T. and Skoog, F. (1962) A revised medium for rapid growth and bio assays with Tobacco tissue cultures. *Physiol. Plant.* **15**, 473–497.
- Nazri, A.Z., Griffin, J.H.C., Peaston, K.A., Alexander-Webber, D.G.A. and Williams, L.E. (2017) F-group bZIPs in barley - a role in Zn deficiencies. *Plant Cell Environ.* <https://doi.org/10.1111/pce.13045>.
- Pace, N. and Weerapana, E. (2014) Zinc-binding cysteines: diverse functions and structural motifs. *Biomolecules*, **4**, 419–434.
- Palmgren, M.G., Clemens, S., Williams, L.E., Krämer, U., Borg, S., Schjørring, J.K. and Sanders, D. (2008) Zinc biofortification of cereals: problems and solutions. *Trends Plant Sci.* **13**, 464–473.
- Pedás, P., Ytting, C.K., Fuglsang, A.T., Jahn, T.P., Schjørring, J.K. and Husted, S. (2008) Manganese efficiency in barley: identification and characterization of the metal ion transporter HvIRT1. *Plant Physiol.* **148**, 455–466.

- Pedas, P., Schjoerring, J.K. and Husted, S. (2009) Identification and characterization of zinc-starvation-induced ZIP transporters from barley roots. *Plant Physiol. Biochem.* **47**, 377–383.
- Pourabed, E., Ghane Golmohamadi, F., Soleymani Monfared, P., Razavi, S.M. and Shobbar, Z.S. (2015) Basic leucine zipper family in barley: genome-wide characterization of members and expression analysis. *Mol. Biotechnol.* **57**, 12–26.
- Ramakers, C., Ruijter, J.M., Lekanne Deprez, R.H. and Moorman, A.F.M. (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* **339**, 62–66.
- Ramesh, S.A., Shin, R., Eide, D.J. and Schachtman, D.P. (2003) Differential metal selectivity and gene expression of two zinc transporters from rice. *Plant Physiol.* **133**, 126–134.
- Rieu, I. and Powers, S.J. (2009) Real-time quantitative RT-PCR: design, calculations, and statistics. *Plant Cell*, **21**, 1031–1033.
- Ruijter, J.M., Ramakers, C., Hoogaars, W.M.H., Karlen, Y., Bakker, O., Hoff, M.J.B., Van Den Hoff, M.J. and Moorman, A.F.M. (2009) Amplification efficiency: linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Res.* **37**, e45.
- Suzuki, M., Bashir, K., Inoue, H., Takahashi, M., Nakanishi, H. and Nishizawa, N.K. (2012) Accumulation of starch in Zn-deficient rice. *Rice*, **5**, 9.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739.
- Tauris, B., Borg, S., Gregersen, P.L. and Holm, P.B. (2009) A roadmap for zinc trafficking in the developing barley grain based on laser capture microdissection and gene expression profiling. *J. Exp. Bot.* **60**, 1333–1347.
- Tiong, J., McDonald, G.K., Genc, Y., Pedas, P., Hayes, J.E., Toubia, J., Langridge, P. and Huang, C.Y. (2013) HvZIP7 mediates zinc accumulation in barley (*Hordeum vulgare*) at moderately high zinc supply. *New Phytol.* **201**, 131–143.
- Tiong, J., McDonald, G., Genc, Y., Shirley, N., Langridge, P. and Huang, C.Y. (2015) Increased expression of six ZIP family genes by zinc (Zn) deficiency is associated with enhanced uptake and root-to-shoot translocation of Zn in barley (*Hordeum vulgare*). *New Phytol.* **207**, 1097–1109.
- Verwoerd, T., Dekker, B. and Hoekema, A. (1989) A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Res.* **17**, 2362.
- Wang, M., Xu, Q., Yu, J. and Yuan, M. (2010) The putative *Arabidopsis* zinc transporter ZTP29 is involved in the response to salt stress. *Plant Mol. Biol.* **73**, 467–479.
- Yang, O., Popova, O.V., Süthoff, U., Lüking, I., Dietz, K.-J. and Goldack, D. (2009a) The *Arabidopsis* basic leucine zipper transcription factor AtbZIP24 regulates complex transcriptional networks involved in abiotic stress resistance. *Gene*, **436**, 45–55.
- Yang, X., Huang, J., Jiang, Y. and Zhang, H.S. (2009b) Cloning and functional identification of two members of the ZIP (Zrt, Irt-like protein) gene family in rice (*Oryza sativa* L.). *Mol. Biol. Rep.* **36**, 281–287.
- Zhao, H. and Eide, D. (1996a) The yeast ZRT1 gene encodes the zinc transporter protein of a high-affinity uptake system induced by zinc limitation. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 2454–2458.
- Zhao, H. and Eide, D. (1996b) The ZRT2 gene encodes the low affinity zinc transporter in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **271**, 23203–23210.