

# Rothamsted Repository Download

## A - Papers appearing in refereed journals

Comber, L., Harris, P., Lu, Y., Wu, L. and Atkinson, P. M. 2019. The forgotten semantics of regression modelling in Geography. *Geographical Analysis* . pp. 1-22.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1111/gean.12199>
- <https://doi.org/10.1111/gean.12199>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/8w9v8>.

© 29 May 2019, Please contact [library@rothamsted.ac.uk](mailto:library@rothamsted.ac.uk) for copyright queries.

# The Forgotten Semantics of Regression Modeling in Geography

Alexis John Comber<sup>1</sup> , Paul Harris<sup>2</sup> , Yihe Lü<sup>3</sup>,  
Lianhai Wu<sup>2</sup> , Peter M. Atkinson<sup>3,4,5,6</sup> 

<sup>1</sup>School of Geography, University of Leeds, Leeds, UK, <sup>2</sup>Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, UK, <sup>3</sup>Chinese Academy of Sciences, Beijing, 100085, China, <sup>4</sup>Faculty of Science and Technology, Lancaster University, Lancaster, UK, <sup>5</sup>School of Geography and Environmental Science, University of Southampton, Highfield, Southampton, SO17 1BJ, UK, <sup>6</sup>School of Natural and Built Environment, Queen's University, Belfast, BT7 1NN, UK

*This article is concerned with the semantics associated with the statistical analysis of spatial data. It takes the simplest case of the prediction of variable  $y$  as a function of covariate(s)  $x$ , in which predicted  $y$  is always an approximation of  $y$  and only ever a function of  $x$ , thus, inheriting many of the spatial characteristics of  $x$ , and illustrates several core issues using “synthetic” remote sensing and “real” soils case studies. The outputs of regression models and, therefore, the meaning of predicted  $y$ , are shown to vary due to (1) choices about data: the specification of  $x$  (which covariates to include), the support of  $x$  (measurement scales and granularity), the measurement of  $x$  and the error of  $x$ , and (2) choices about the model including its functional form and the method of model identification. Some of these issues are more widely recognized than others. Thus, the study provides definition to the multiple ways in which regression prediction and inference are affected by data and model choices. The article invites researchers to pause and consider the semantic meaning of predicted  $y$ , which is often nothing more than a scaled version of covariate(s)  $x$ , and argues that it is naïve to ignore this.*

## Introduction

There is a long and varied literature on the semantics associated with geographical information. Primarily this is concerned with how spatial phenomena are conceptualized, represented, and encoded in spatial data (e.g., Smith and Varzi 2000; Harvey 2000; Robbins 2001; Smith 2001; Mark and Turk 2003; Smith and Mark 2003; Comber, Fisher, and Wadsworth 2005; Pires 2005; Turk, Mark, and Stea 2011; Derungs et al. 2013). Much less attention has been paid to the semantics associated with the analysis of spatial data, that is, what we *do* with spatial data.

In this article, we describe how the specification of spatial models, and choices made therein, have an impact on the outcomes of a spatial analysis. For example, taking the simplest case of the prediction of  $y$  as a function of  $x$  with error  $\epsilon$ :

Correspondence: Peter M. Atkinson, Faculty of Science and Technology, Lancaster University, Lancaster, UK  
e-mail: pma@lancaster.ac.uk

Submitted: October 18, 2018. Revised version accepted: March 4, 2019

$$y = f(x) + \epsilon$$

$$y^* = f(x)$$

and noting that the predicted  $y$ , that is,  $y^*$ , will only ever approximate  $y$ , it is evident that  $y^*$  in Equation (1) is always, and can only ever be, a function of  $x$ . Thus, only  $f(x)$  is ever found and never  $y$ . This statement is axiomatic but often overlooked. It can lead to implicit or explicit claims about regression-based spatial outputs that are indefensible. For example, maps of disease incidence have been produced that are in effect simple transforms of covariates such as vegetation greenness (e.g., Beck, Lobitz, and Wood 2000; Anyamba, Linthicum, and Tucker 2001; Hay et al. 2005; Lui et al. 2015). In these, the disease map is actually a map of vegetation greenness scaled by additive and multiplicative factors. The map inherits all the spatial characteristics of vegetation greenness and only those of disease incidence that relate indirectly through greenness. Thus, the key points that this article makes are that  $y^*$  can only ever be a function of  $x$ ; that only  $f(x)$  is ever found and never  $y$ ; and that models built on these paradigms, including those using spatial data, may result in naïve interpretations of the results and rather surprising consequences in some extreme cases.

The semantic and characteristic variation in  $y^*$ , as a function of how  $y$  is predicted, will arise due to core decisions about the data and the model. Decisions relating to the data include their specification, support, measurement, and errors:

- The *specification of  $x$* : the  **$x$  issue**, on which covariates to include in the model.
- The *support of  $x$* : the  **$v$  issue** in which support effects are induced through the choice of measurement scales and granularity of  $x$ .
- The *measurement of  $x$* : the  **$m$  issue**, that is, how the desired  $x$  is measured.
- The *error of  $x$* : the  **$e$  issue** in which the accuracy with which we measure  $x$ , affects  $y^*$ .

Decisions relating to the model include its functional form, the method of model identification and the specification of  $y$  itself (a combined data and model issue).

- The *functional form of the model*: the  **$f$  issue**, in which the specification of the model itself has a semantic effect on  $y^*$ .
- The *method of model identification*: the  **$i$  issue**, the statistical method of estimating the model parameters.
- The *specification of  $y$* : the  **$y$  issue**. For any model incorporating spatial effects, semantic and characteristic variation in  $y^*$  will also arise as a function of the specification of  $y$ . The definition and sampling framework of  $y$  itself has a semantic effect on  $y^*$  at locations other than  $y$ .

This study illustrates the effects of data and model decisions through the  **$x$ ,  $v$ ,  $m$ ,  $e$ ,  $f$ , and  $i$  issues** and their impacts on the semantics of the resultant spatial operations. Focus is placed on the imparted effects on  $y^*$  through these issues (i.e., model prediction), and much less so the imparted effects on associated inferences (i.e., parameter uncertainty). The  **$y$  issue** is acknowledged, but not investigated, as its illustration is not straightforward and could not be captured adequately here (see Wang et al. (2012) and references therein). Although the stated issues are inherent to many statistical operations, we focus on regression, where the issues are illustrated through ordinary least squares (OLS) multiple regression, geographically weighted regression (GWR) (Brunsdon, Fotheringham, and Charlton 1996), and mixed models whose parameters

are estimated via restricted maximum likelihood (REML) (e.g., Welham et al. 2004) or via maximum likelihood (ML). The next section describes the six issues in more detail before they are illustrated using a “synthetic” case study of Landsat 5 remote sensing imagery. The analysis is repeated using a “real” case study of a soils data set for the Loess Plateau, China. Finally, we conclude the study with key outputs and discussion points.

## Semantic issues in detail

Regression seeks to model changes in  $y$ , the target variable, with changes in  $x$ , the covariates, where  $x$  is always (and naively) assumed free of measurement error. For this study,  $y$  is always fixed in fitting the regression, meaning that regardless of the issues described, the aim is always toward the same  $y$ , producing a  $y^*$  with associated residuals ( $y - y^*$ ). As we are not presenting the **issue**  $y$ , only *in-sample* prediction of  $y$  is conducted and not *out-of-sample*.

### The $x$ issue

The  **$x$  issue** as relates to the number  $k_x$  and choice  $c_x$  of potential covariates to present to the regression. Covariate selection has a limiting effect on the possible value and semantic meaning of  $y^*$ . Covariate *selection* can be a difficult task. For example, two or more covariates displaying similar predictive ability in relation to  $y$  can commonly be retained to increase the accuracy of  $y^*$ , but at the same time can compromise model interpretability due to their collinearity. (Note that presence of such covariance among covariates can confound the semantics of any model due to uncertainty about whether the observed response is a result of  $x_1$  or of  $x_2$ , imposing a further level of intra- $x$  semantics on the model.) This duality can result in an  **$i$  issue** of whether to fit the model using, say, some penalized regression or not (e.g., Zou and Hastie 2005). Covariate *availability*, links to the  **$f$  issue** (and indirectly to the  **$i$  issue**), as missing covariates in a spatial study are often reflected in spatial effects (e.g., non-stationary relationships and/or autocorrelated residuals), that would not be present if the regression were fully-specified (i.e., with a full set of covariates, such that a multiple regression fitted using OLS suffices) (see Cressie and Chan 1989).

### The $v$ issue

The support  $v$  of  $x$  is an important concept in geostatistics and quantitative geography. The support is the space on which an observation is defined, or on which a measurement is made and is defined by parameters, such as size, shape, orientation, and position, as well as the dimension of interest (e.g., 2D, 3D, and 2D plus time). The support represents the lower limit of what we can know about the real world. In contrast, the spatial extent of the sample set defines the space of interest, while the sampling scheme (e.g., random, systematic grid) defines how that space is represented. Nothing can be known *beyond* the extent of the sample set and nothing can be known *within* the support (Atkinson and Tate 2000). Indeed, all that can be known exists in the relations *between* the observations of the specific sampling scheme. As a result, data are always a function of the real world and the sampling framework, as well as conceptual and semantic choices defining measurable properties (such as “biomass per unit area”). The support has an important effect on the semantic meaning of any regression prediction: as the support increases, some of the potential variation in the property of interest is lost to within the support (i.e., it is integrated out) and all that is left is the variation between supports. Because of this, the measured variance collapses as the support size increases.

Where spatial structure exists, neighboring observations are commonly similar to each other (positive spatial correlation) or more rarely dissimilar (negative spatial correlation), entailing in both instances that decrease in variance is smaller as the support increases. This means that less variance is lost to within the support. This is also one aspect of the modifiable areal unit problem (Openshaw 1984), where, as the areal size changes (up- or down-scaling), the correlation between a given pair of variables changes, conditional upon the scales of spatial variation that exist in the same variables. Gotway and Young (2002), Zhang, Atkinson, and Goodchild (2014) and Murakami and Tsutsumi (2015) describe the effects of varying support of the observation units.

In terms of the *v* issue,  $y^*$  and  $(y-y^*)$  are a function, either directly or indirectly, of the sampling framework of  $x$ , specifically, the size, geometry, and orientation of the support over which the variables were measured. If the supports of the covariates  $x$  change, then the predictions  $y^*$  will change and, thus, the semantics of the model or its predicted state will change. Most obviously, the variance of  $y^*$  will change.

### **The *m* issue**

The *m* issue can manifest itself in several ways. Measurements of a property will vary according to different options for how they are measured analytically. If these variables are then used separately as covariates to predict  $y$ , then clearly  $y^*$  will vary according to which form of the covariate  $x$  is specified. Thus, the specific impact of any covariate  $x$  is dependent on its method of measurement, and each individual covariate may be affected in this manner in different ways.

An interesting *m* issue consideration arises in remote sensing. The pixel values in a remotely sensed image are generally accepted to be integrals (e.g., of radiance or brightness) over the measurement support, which is approximated as a pixel. The measured values are integrals across different variables because the angle of view varies across the support. This may matter less for certain satellite-based observations, but it may be of greater consequence for airborne, drone, or ground-based measurements. For example, when measuring the radiance of a crop using a ground-based instrument, a single measurement is likely to include the tops of the canopy at nadir, but also the sides of the canopy at the edge of the support. Thus, a single value is realized as the integral of fundamentally different variables. This semantic mixing in  $x$  will impact on  $y^*$ .

### **The *e* issue**

The *e* issue is concerned with measurement error and how the accuracy with which we measure  $x$ , affects  $y^*$ . The  $x$ 's can never be known perfectly and while they *might* be error free, we can never know that. Thus, the error  $e_x$  in  $x$  will propagate through the model transform to  $y^*$ . Here  $e_x$  may include systematic and random contributions of given magnitudes, will have its own distributional form, and may have a spatio-temporal character (e.g., autocovariance). It may also be related directly to  $x$  (i.e., heteroscedasticity). All of these properties will impact on  $y^*$ , and regression models to account for them exist (e.g., Christensen 2011). Most fundamentally, the larger the error  $e_x$  relative to the signal in  $x$  (i.e., the smaller the signal-to-noise ratio in  $x$ ), the more degraded is the semantic information content of  $y^*$ . Observe that the error  $e_x$  is different to the error or residual of  $y^*$ .

### **The *f* issue**

The *f* issue is concerned with model specification or choice. For regression modeling this can include models that deal with non-linearity or heteroskedasticity, relationship non-stationarity (e.g., GWR) and autocorrelated residuals (e.g., mixed models). For each model category there

exist alternative models (or modeling paradigms) with broadly the same objective. For example, weighted regression can be used to deal with non-linearity or heteroskedasticity (e.g., Carroll and Rupert 1988), an expansion model can be used to model non-stationary relationships (Cassetti 1972), and a simultaneous autoregressive model can be used to counter autocorrelated residuals (e.g., Anselin 1990; Cressie 1993).

Model specification frequently includes secondary choices within the chosen model. These include the kernel function in GWR (e.g., Gaussian, bi-square, etc.) or the variogram model in a mixed model (e.g., exponential, spherical, etc.). All these choices can be directed based on model fitting, but only where the options have been explored by the investigator. This is analogous to the case for the data (the ***x* issue**), whereby the data selected by the model fitting can be chosen only from the set provided by the investigator: the chosen model can only be from the set of models investigated.

### The *i* issue

Model identification concerns the statistical estimation method used for fitting a chosen model. Examples include weighted least squares (WLS) for addressing non-linearity or heteroskedasticity and partial least squares (PLS) for addressing collinearity (e.g., Frank and Friedman 1993). The method of fitting can also determine the number  $k_x$  and set  $c_x$  of covariates to include in the model and the use of an intercept term (e.g., with stepwise, LASSO or elastic net approaches; see Zou and Hastie 2005). The choice of adopting a Bayesian inference framework also falls mostly into this category, for example, through fitting options such as Markov chain Monte Carlo (MCMC) or Integrated Nested Laplace Approximation (INLA) methods (Rue, Martino, and Chopin 2009). The choice of estimation method will determine the identification of the model and parameters specified and, thus, this choice will have an effect on  $y^*$ , its semantic meaning and character.

The method of identification strongly links with model specification. However, we do not attempt to illustrate more complex problems in which it can be difficult to isolate non-linearity, non-stationarity, and autocorrelation effects, for a given spatial data set (e.g., Anselin 2010; Basile et al. 2014; Harris 2018) or illustrate those that additionally consider scale-dependent processes (e.g., Dong et al. 2015). Instead, we choose to illustrate routine methods of identification in mixed models with REML and ML, both of which separate trend (first-order effects) from autocorrelation (second-order) effects (see also, Armstrong 1984).

### Summary

It is clear that the semantic meaning of the regression-based prediction of  $y^*$  is a function of the number  $k_x$  and choice  $c_x$  of covariates  $x$  (the ***x* issue**), the given supports (the ***v* issue**), the way that they are measured (the ***m* issue**), together with their (unknown) errors (the ***e* issue**). The fitted regression in terms of its specification (the ***f* issue**) and its identified parameters (the ***i* issue**) will also impact on  $y^*$ . These issues and, in particular, their impact on the prediction of  $y^*$  and model inference, are illustrated through two case studies with *non-spatial* multiple regression models and GWR, and mixed models as *spatial* regression models. GWR and the mixed model represent the ***f* issue**, and the ***i* issue** is represented by OLS for multiple regression, together with REML/ML for the mixed models. All study regression models are described in the Appendix to this article.

## Synthetic case study: Landsat 5 imagery

In the first case study, remote sensing imagery is used to illustrate the semantic impacts of regression modeling on  $y^*$  and model inference due to the  $x$ ,  $v$ ,  $m$ ,  $e$ ,  $f$ , and  $i$  issues. Cloud-free Landsat 5 imagery from 2011 was obtained for the East of England (path 201, row 023, 30<sup>th</sup> October 2011) with a spatial resolution of 30 m. Three of the Landsat 5 spectral bands were used in the regression analyses, with the Blue band (0.45–0.52  $\mu\text{m}$ ) as the target variable ( $y$ ), and the Red and Green bands (0.63–0.69  $\mu\text{m}$  and 0.52–0.60  $\mu\text{m}$ , respectively) as the covariates  $x_1$  and  $x_2$ , respectively. A  $50 \times 50$  pixel area was selected randomly and extracted for use (Fig. 1).

A total of eight regression analyses were undertaken to illustrate the semantic effects of the six issues, where in each case the aim was to predict the Blue band ( $y$ ) using the Red band ( $x_1$ ) and in three instances, the Green band ( $x_2$ ) also. The eight regression analyses were as follows:

- a **Reference:**  $y = f(x_1)$  with multiple regression;
- b  **$x$  issue:**  $y = f(x_1 + x_2)$  with multiple regression;
- c  **$v$  issue:**  $y = f(x_1)$  with multiple regression, where the Red band ( $x_1$ ) is aggregated to 60 m spatial resolution using a nearest neighbor algorithm;
- d  **$m$  issue:**  $y = f(x_1)$  with multiple regression, where the Red band ( $x_1$ ) is altered slightly by taking its square root;
- e  **$e$  issue:**  $y = f(x_1)$  with multiple regression, where a relatively small amount of random noise is added to the Red band ( $x_1$ );
- f **first  $f$  issue:**  $y = f(x_1)$  with GWR;
- g **second  $f$  issue:**  $y = f(x_1 + x_2)$  with a mixed model fitted by REML;
- h  **$i$  issue:**  $y = f(x_1 + x_2)$  with a mixed model fitted by ML.


Fig. 2 maps the actual  $y$ 's (all the same), the different predicted  $y^*$ 's, the covariates  $x_1$  and  $x_2$ , and the residuals ( $y - y^*$ ) arising from the eight regression fits and the six semantic issues  $x$ ,  $v$ ,



**Figure 1.** The  $50 \times 50$  pixel area of 2011 Landsat 5 imagery used in the study, showing bands 3, 2, and 1 through the Red, Green, and Blue channels.

*m*, *e*, *f*, and *i*. The distributions of  $y^*$  are also shown via boxplots in Fig. 3, along with the corresponding residuals ( $y-y^*$ ) in Fig. 4. It is also possible to recompare the residuals ( $y-y^*$ ) as in Fig. 5, together with  $y^*$ , with the covariate  $x_1$  through  $(x_1-y^*)$  as in Fig. 6, where maps from both figures use the same legend and class breaks. Model inference for each of the eight regression models is summarized in Table 1.

From Fig. 2, it is clear that for regression models (a, c, d, e, and f) the  $y^*$  are much closer to  $x_1$  than to  $y$ . This is particularly true for regressions (c and e), the *v* and *e* issues, respectively, where changes in the support of  $x_1$  and the error associated with  $x_1$  are very clearly reflected in  $y^*$ . For regression (d), the *m* issue, the  $y^*$  have a similar spatial pattern to  $x_1$  as that found in the reference regression (a), but the  $y^*$  has reduced variability. This effect similarly results for regression (f), the first *f* issue with GWR, in comparison to the reference regression (a); but in this case, the  $y^*$  show the least spatial similarity to  $x_1$ , (at least for regressions using only a single covariate). Thus, accounting for spatial effects (the *f* issue) reduces the similarity between the  $y^*$  and  $x_1$ , as might be expected.



**Figure 2.** Maps of the original and same target variable ( $y$ ), different  $y^*$ , covariates ( $x_1$  and  $x_2$ ), and model residuals ( $y-y^*$ ) arising from the six issues and eight regressions (a–h).




Figure 2. Continued






Figure 3. Boxplot distributions of  $y^*$  arising from the six issues and eight regressions (a–h) (outliers given with a small transparency term).

When an additional covariate,  $x_2$  is added for the **x issue** (in regressions b, g and h), the  $y^*$  are no longer as close to  $x_1$ , but are now a reflection of  $x_1$  and  $x_2$ , combined. Variability in  $y^*$  reduces when viewing the second **f issue** (a mixed model), in comparison to its reference regression (b);



**Figure 4.** Boxplot distributions of the regression residuals ( $y-y^*$ ) from the six issues and eight regressions (a–h). Outliers are given with a small transparency term.




**Figure 5.** The residuals ( $y-y^*$ ) arising from the six issues and eight regressions (a–h).

and this is similarly true with the first **f issue** (GWR), in comparison to its reference regression (a). On viewing only regressions (g and h), the chosen **i issue**, appears to have little to no effect on  $y^*$ , which was anticipated.

From Fig. 3, the  $y^*$  from single covariate regressions (c, d, e, and f, for issues **v**, **m**, **e**, and **f**, respectively) all display clear differences to that found with the reference regression (a). Similarly, the double covariate regression (b, for the **x issue**) provides clear differences in its  $y^*$  to that of the single covariate reference regression (a). Similarly, the double covariate regressions (g and h, for issues **f** and **i**, respectively), both display differences to that found with their reference regression (b), but not with each other. The results for Fig. 4, mimic those given for Fig. 3, but in terms of the residuals ( $y-y^*$ ) rather than  $y^*$ .

From Fig. 5, it is evident that regressions for five of the six issues (**x**, **v**, **m**, **e**, and **f**) all provide residuals that are different in spatial pattern to the reference regression and also to any alternative reference regression (i.e., the second **f issue** and the **i issue** with the **x issue** as their reference). For regressions representing the **v** and **e issues**, this effect can be quite striking. Fig. 6,



**Figure 6.** The differences between  $x_1$  and the  $y^*$  values ( $x_1 - y^*$ ) from the six issues and eight regressions (a–h).

**Table 1.** Model summaries for the eight regression analyses applied to the image case study

Semantic issue and regression form	Covariate	Estimate	<i>t</i> -value	Pr. ( $> t $ )	
<b>Reference:</b> $y = f(x_1)$ , multiple regression (MR)	$x_1$	0.943	548.022	0.000	
<b><i>x</i> issue:</b> $y = f(x_1 + x_2)$ , MR	$x_1$	1.193	126.506	0.000	
	$x_2$	-0.283	-26.889	0.000	
<b><i>v</i> issue:</b> $y = f(x_1)$ , MR, $x_1$ aggregated to 60 m	$x_1$	0.927	259.293	0.000	
<b><i>m</i> issue:</b> $y = f(x_1)$ , MR, square root of $x_1$	$x_1$	0.919	566.906	0.000	
<b><i>e</i> issue:</b> $y = f(x_1)$ , MR, random noise added to $x_1$	$x_1$	0.930	321.496	0.000	
<b>first <i>f</i> issue:</b> $y = f(x_1)$ , <i>Min.</i> GWR*	–	0.919	298.183	0.000	
	<i>1<sup>st</sup> Quartile</i>	–	0.935	431.140	0.000
	<i>Median</i>	$x_1$	0.943	484.059	0.000
	<i>3<sup>rd</sup> Quartile</i>	–	0.951	523.779	0.000
	<i>Max.</i>	–	0.992	572.124	0.000
<b>second <i>f</i> issue:</b> $y = f(x_1 + x_2)$ , mixed model REML	$x_1$	0.524	28.783	0.000	
	$x_2$	0.193	12.000	0.000	
<b><i>i</i> issue:</b> $y = f(x_1 + x_2)$ , mixed model ML	$x_1$	0.524	28.789	0.000	
	$x_2$	0.193	11.997	0.000	


\*The GWR *t*-values are not corrected for multiple hypothesis tests (see Gollini et al., 2015).

depicts how the spatial characteristics of  $y^*$  move closer to or further away from the spatial characteristics of the covariate  $x_1$ , depending on the illustrated semantic issue. Here, the regression for the *m* issue appears to provide  $y^*$  values that are the most similar to the covariate  $x_1$ , while the regressions for the *v* and *e* issues provide  $y^*$  values that are least similar.

The model inference summaries (Table 1) are only given for completeness and are largely uninteresting given that this is a synthetic case study. However, none of the six semantic issues result in a change of significance for any regression coefficients. The coefficient estimates can change, however; for example, the covariate  $x_2$ , negatively relates to  $y$  in the *x* issue, whereas the same relationship is a positive relationship in the second *f* issue.

## Real case study: soils in the Loess Plateau, China

In this second case study, the predictive and inferential impacts of the *x*, *v*, *m*, *f*, and *i* issues are illustrated for a soils data set in the Loess Plateau, China. This area is dominated by thick loess deposits and suffers from intense erosion. It contains several fragile ecosystems (Chen et al. 2007; Wang et al. 2008) and has been the subject of much research, particularly focused on reducing the impacts of soil erosion by vegetation restoration, re-greening, and sustainable agriculture initiatives (Hu et al. 2017; Li et al. 2017; Fu et al. 2018; Luo et al. 2018). Soil moisture is driven by limited precipitation, most of which falls between June and September. Understanding hydrological processes, and particularly soil conductivity, is critical in the Loess Plateau as these underpin the effectiveness of vegetation restoration activities. Soil data were collected at 243 locations in the Loess Plateau, China (Fig. 7) and analyzed in the laboratory to generate



**Figure 7.** The 243 sample locations in the Loess Plateau, China, with a slight shading transparency to show sites that overlap.


measurements of several soil characteristics including soil physical properties (soil hydraulic conductivity,  $K_s$ , in  $\text{cm min}^{-1}$  and saturated soil water content, SSWC), and soil composition (% Clay, % Silt, and % Sand). All soil measures were recorded at three different depths of 0–10 cm, 10–20 cm, and 20–40 cm.

In this study,  $K_s$  at 0–10 cm depth is considered as the target variable,  $y$ , with covariates SSWC at 0–10 cm and % Clay and % Silt both at 0–10 cm (% Sand at 0–10 cm is dropped to deal in a simple way with the compositional nature of the data). Additional covariates in terms of landscape (terrain) indices were obtained for each sample location from a series of DEMs at spatial resolutions of 30 m and 90 m. The “terrain” function in the R “raster” package (v2.6-7, Hijmans and van Etten 2017) was used to generate measures of aspect (0–360°) and stream flow direction (encoded in powers of 2 to indicate compass direction). The aspect variable was transformed to East-ness by taking the cosine of the angles, and flow direction was converted to a measure of Flow Northeast-ness in a similar way using the sine function only.


The soil and terrain covariates allow us to demonstrate the  $x$  issue, and as the two terrain covariates (East-ness and Flow Northeast-ness) are calculated over different supports of 30 and 90 m, the  $v$  issue, also. The  $m$  issue is not present in this data, where it would have been ideal to have covariate data from different methods of measuring the exact same property using three competing analytical techniques. However, we do have measurements of the *same* soil property collected at three different depths and, therefore, we use the SSWC at 20–40 cm, as alternative measurements of SSWC at 0–10 cm. This serves for illustration. If we had chosen to integrate the depth-dependent soils data, for example, from 0 to 20 cm or 0 to 40 cm then we would be provided with an alternative  $v$  issue.

The nature of the alternative covariates for the study of the  $v$  issue and  $m$  issue are shown in Figs. 8 and 9, respectively. The prediction accuracy of  $K_s$  at 0–10 cm should intuitively reduce for a regression calibrated with terrain variables over the larger supports ( $v$  issue), and similarly, reduce for a regression calibrated with the deeper values of % SWCC ( $m$  issue). These covariate decisions illustrate the potential for variation in regression outputs (predictions, coefficients, and uncertainties) due to decisions over choice of support (Fig. 8) and of measurement (Fig. 9). The final data-driven semantic issue, the  $e$  issue, is not illustrated.

To illustrate the two model-driven semantic issues (the  $f$  and  $i$  issues), multiple regression, GWR, and mixed models fitted using REML and ML were again specified. Thus, again two  $f$  issues are illustrated, where the  $i$  issue corresponds directly to the second  $f$  issue (via the mixed models). Again, the second  $f$  issue and its  $i$  issue are also given in the form used for the  $x$  issue



**Figure 8.** The  $v$  issue—variation in the terrain indices derived from different DEMs (probability densities). Note that the variance should decrease as sample support size increases.



**Figure 9.** The contrived *m* issue—variation in the distributions of the soil property variables (% Clay, % Silt, and % SSWC) sampled at the top and bottom depths (0–10 cm and 20–40 cm). The *m* issue is illustrated only using % SSWC and not % Clay and % Silt, also.

and not the reference regression. The resultant seven regression analyses chosen to illustrate the *x*, *v*, *m*, *f*, and *i* issues are as follows (with the differences to the reference regression highlighted in bold):

- Reference:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, Flow-NE and East-ness both at 30 m) with multiple regression;
- x* issue:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, Flow-NE and East-ness both at 30 m; **Clay and Silt both at 0–10 cm**) with multiple regression;
- v* issue:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, **Flow-NE and East-ness both at 90 m**) with multiple regression;
- m* issue:**  $K_s$  at 0–10 cm =  $f$ (**SSWC at 20–40 cm**, Flow-NE and East-ness both at 30 m) with multiple regression;
- first ***f* issue:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, Flow-NE and East-ness both at 30 m) with **GWR**;
- second ***f* issue:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, Flow-NE and East-ness both at 30 m; **Clay and Silt both at 0–10 cm**) with **mixed model fitted by REML**;
- i* issue:**  $K_s$  at 0–10 cm =  $f$ (SSWC at 0–10 cm, Flow-NE and East-ness both at 30 m; **Clay and Silt both at 0–10 cm**) with **mixed model fitted by ML**.

Fig. 10 maps the  $y^*$ 's. Model inference outputs for each of the seven regressions are summarized in Table 2.

From Fig. 10, it is useful to start with regression models (a, c, d, and e) where the  $K_s$  (at 0–10 cm) predictions  $y^*$  are informed by SSWC, Flow-NE and East-ness covariates, and relate to the *v*, *m*, and (first) *f* issues. The regressions for (c, d, and e) provide  $K_s$  prediction maps with a broadly similar spatial pattern to that found with the reference regression (a). However, the *v* issue (regression c) results in a reduced range of  $y^*$  and a likely reduction in its accuracy, and the possibility of missing areas of high or low  $K_s$ , that may be vital to the understanding of the soils process and its erosion. The *m* issue (regression d) appears to result in an increase in lower values of  $y^*$ , suggesting an under-prediction bias for  $K_s$ , especially toward the edges of the sampled area. Again, clear spatial interpretation issues for  $K_s$  may result. For the *f* issue care must be taken, as unlike the *v* and *m* issues, where likely inaccuracies are expected (including their direction), it is not known which of multiple regression and GWR is the most accurate. GWR




Figure 10. Maps of  $y^*$  arising from the five issues and seven regressions (a–g).

**Table 2.** Model summaries from the five issues and the seven regression analyses applied to the soils case study (changes from the reference regression highlighted in bold)

Regression form	Covariate	Estimate	<i>t</i> -value	Pr. ( $> t $ )
<b>Reference</b> (multiple regression)	SSWC at 0–10 cm	0.013	22.292	0.000
	Flow-NE at 30 m	0.085	2.660	0.008
	East-ness at 30 m	-0.082	-2.398	0.017
<b><i>x</i> issue</b> (multiple regression)	SSWC at 0–10 cm	0.038	13.497	0.000
	Flow-NE at 30 m	0.058	2.144	0.033
	East-ness at 30 m	-0.074	-2.571	0.011
	<b>Clay at 0–10 cm</b>	0.047	6.755	0.000
	<b>Silt both at 0–10 cm</b>	-0.029	-9.837	0.000
<b><i>v</i> issue</b> (multiple regression)	SSWC at 0–10 cm	0.013	22.116	0.000
	<b>Flow-NE at 90 m</b>	0.093	2.810	0.005
	<b>East-ness at 90 m</b>	0.026	0.691	0.490
<b><i>m</i> issue</b> (multiple regression)	<b>SSWC at 20–40 cm</b>	0.014	19.194	0.000
	Flow-NE at 30 m	0.096	2.733	0.007
	East-ness at 30 m	-0.088	-2.345	0.020
first <b><i>f</i> issue (GWR*</b> , showing IQRs only)	SSWC at 0–10 cm	0.002	2.799	0.000
	Flow-NE at 30 m	0.095	1.798	0.503
	East-ness at 30 m	0.092	1.783	0.354
second <b><i>f</i> issue (mixed model REML)</b>	SSWC at 0–10 cm	0.038	13.430	0.000
	Flow-NE at 30 m	0.060	2.260	0.025
	East-ness at 30 m	-0.080	-2.831	0.005
	<b>Clay at 0–10 cm</b>	0.044	6.185	0.000
	<b>Silt both at 0–10 cm</b>	-0.029	-9.488	0.000
<b><i>i</i> issue (mixed model ML)</b>	SSWC at 0–10 cm	0.038	13.430	0.000
	Flow-NE at 30 m	0.060	2.254	0.025
	East-ness at 30 m	-0.080	-2.815	0.005
	<b>Clay at 0–10 cm</b>	0.044	6.221	0.000
	<b>Silt both at 0–10 cm</b>	-0.029	-9.507	0.000

\*The GWR *t*-values are not corrected for multiple hypothesis tests (see Gollini et al., 2015).

provides a cluster of unusually high *K*<sub>s</sub> values to the mid-south of the sample area. If, as is likely, GWR predicts more accurately than multiple regression, then the mis-identification of such an area of high *K*<sub>s</sub> values is likely to be problematic, especially for this region's soil health (a region of approximately 100 km by 200 km).

Conversely the maps of  $y^*$  are highly similar for regressions (b, f, and g) where  $y^*$  are additionally informed by Clay and Silt covariates, encompassing the  $x$ , (second)  $f$ , and  $i$  issues. This suggests a multiple regression for predicting Ks to be as worthy as a mixed model (i.e., the  $f$  issue), regardless of its identification method (i.e., the  $i$  issue). For the  $x$  issue, the  $y^*$  from regression (a) need to be compared to  $y^*$  from regression (b), where the addition of the Clay and Silt covariates has an obvious influence on  $y^*$ . As previously discussed, this outcome may relate to where missing covariates are reflected in spatial effects (in this case the non-stationary relationships of regression e), that may not be present if the regression were better-specified (i.e., with the extra covariates). However, this would need to be tested via a second GWR fit using all five covariates.

The model inference summaries are given in Table 2, and similarly indicate little difference in regressions (b, f, and g), but clear differences for regressions (a, c, d, and e). For the latter, this includes changes in coefficient sign (for East-ness) and changes in coefficient significance (for East-ness with regression c, and Flow-NE and East-ness with regression e). Again, the  $x$  issue in covariate choice, appears hugely important to the interpretation of this study's soils process.

## Conclusions

This study illustrated how the semantics associated with regression prediction and inference are functions of choices over covariate data and regression models. The choice of covariates, the nature of their supports, the way that they are measured and their (unknown) errors were all shown to have an impact on the resultant prediction, and thereby its semantics. Prediction was also shown to be affected by model choices, specifically model specification and the method of parameter identification. The implications of each of the issues identified are summarized in Table 3 and, of course, in any analysis several issues may be present and may interact. Many of these semantic effects are known but they are infrequently articulated and almost never routinely described in reports and scientific articles despite being the subject of wider study. Given the ubiquity of the use of regression within quantitative geography and the profound nature of the semantic impacts, this study presents an opportunity for researchers in quantitative geography to pause and update their thinking. In the extreme case of simple regression with one covariate  $x_1$ , the prediction is nothing more than a scaled version of that covariate, inheriting the same (scaled) spatial correlation and it is fundamentally naive to ignore this. Even small differences in data and model decisions can result in profound differences in outcomes including predictions, residuals, coefficient estimates, and the relationship between covariates and predictions. There is a wider and potentially more pressing context for these observations. Some advocates of machine learning have suggested that most of the issues identified in this article can be ignored. However, as Marcus (2018, p. 15) states "deep learning is just a statistical technique, and all statistical techniques suffer from deviation from their assumptions."

The inclusion of GWR as a model choice raised some interesting issues in terms of the spatial structure of the predictions and residuals. For multiple regression, the impact on the predictions is a simple linear additive and multiplicative factor applied to the covariates. In GWR, spatial correlation is induced in the predictions that is not present in the covariates. In a mixed model, the impact includes spatial smoothing that is similarly not present in the covariates. More profoundly, as the nature of model information and model fit changes from location to location with GWR, this suggests that the semantics of  $y^*$  vary locally, also. This is evident in extensions to GWR, such as the geographically weighted LASSO (Wheeler 2009) and geographically

**Table 3.** Implications of the different semantic issues

Issue	Implications
<b><i>x</i> issue</b> (which covariates to include in the model)	This limits the possibility space of $y^*$ , which in turn, restricts its semantic meaning
<b><i>v</i> issue</b> (support effects/measurement scales)	Choices (e.g., size, geometry, orientation, and position) can result in different integrals $x_v$ on the supports $v$ which, in turn, can have a key effect on $y^*$
<b><i>m</i> issue</b> (how $x$ is measured)	The impact of any covariate $x$ is dependent on its method of measurement and $y^*$ will vary according to which form of $x$ is specified
<b><i>e</i> issue</b> (accuracy of measures of $x$ )	The error $e_x$ in $x$ will propagate through the model transform to the prediction of $y$ (related to the measurement of $x$ above) and will have its own distribution and spatio-temporal character
<b><i>f</i> issue</b> (specification of the model)	A semantic effect on $y^*$ through the resultant linear weighted combinations of $x$ 's. Models for spatial data can vary greatly (e.g., linear vs. non-linear, stationary vs. non-stationary). The nature of the model parameters effect $y^*$
<b><i>i</i> issue</b> (the statistical method of estimating the model parameters)	A semantic effect on $y^*$ will be evident in the linear weights for combining the $x$ 's. Although often small, the choice of estimation or inference method for identifying the model may affect $y^*$
<b><i>y</i> issue</b> (variation in $y^*$ due to the specification of $y$ )	The definition and sampling framework of $y$ itself has a semantic effect on $y^*$ at locations other than $y$ (i.e., predictions at un-sampled locations)

weighted elastic net logistic regression (Comber and Harris 2018), where covariate selection is local; locally compensated ridge GWR, where the method of estimation is local (Brunsdon, Charlton, and Harris 2012); and autoregressive GWR (Geniaux and Martinetti 2017; Brunsdon, Fotheringham, and Charlton 1998), where the method of identification is local.

In summary, the semantic issues highlighted in this article have clear relevance to the many geographical studies that routinely use regression in one form or another, many of which inform the core challenges that the world is facing today, such as the global disease burden, global mortality, poverty, food insecurity, losses in biodiversity, and increasing pollution through urbanization. It is important that researchers think carefully about the semantics associated with any interpretation of predictions and the potential impact of semantic issues, given the large number of ways in which regression prediction may be affected by data and model choices. It is hoped that this study will raise awareness among quantitative geographers, challenge naive application of regression, lead to proper consideration of the semantic interpretation of regression predictions and introduce the semantic lens to data and model choices in regression modeling. This study supports this purpose by providing a clear conceptual framework that defines the multiple ways in which data and model choices can impact on prediction and inference. The uncertainty in

prediction arising from these choices was deliberately omitted in establishing the core message of the article but will be the subject of a future study.

## Acknowledgements

This research was supported by the China-UK bilateral collaborative research on critical zone science (the Natural Environment Research Council Newton Fund NE/N007433/1, the National Natural Science Foundation of China NO. 41571130083), and the National Key Research and Development Program of China (No. 2016YFC0501601). All of the data preparation, analyses and mappings were undertaken in R 3.5.1, the open source software. The code and data used in this analysis are available from <https://github.com/lexcomber/SemanticsOfStats>. The DOI for the data and code are at <https://zenodo.org/badge/latestdoi/123798087>.

## Appendix: Regression models

### Multiple regression

The multiple regression model can be fitted using OLS, and has the form:

$$y_i = \beta_0 + \sum_{k=1}^m \beta_k x_{ik} + \varepsilon_i \quad (A1)$$

where for observations indexed by  $i = 1, \dots, n$ ,  $y_i$  is the target variable,  $x_{ik}$  is the value of the  $k^{th}$  covariate,  $m$  is the number of covariates,  $\beta_0$  is the intercept term,  $\beta_k$  is the regression coefficient for the  $k^{th}$  covariate, and  $\varepsilon_i$  is the random error term. In matrix terms, the coefficients of OLS regression are estimated from:

$$\hat{\beta}_{OLS} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X} \quad (A2)$$

where,  $\mathbf{X}$  is a  $(n \times (m + 1))$  covariate data matrix and  $\mathbf{y}$  is a  $(n \times 1)$  target data vector. The multiple regression prediction of  $y$  at any observed (and un-observed) spatial location  $\mathbf{s}$  is:

$$y_{MR}^*(\mathbf{s}) = \mathbf{x}(\mathbf{s})^T \hat{\beta}_{OLS} \quad (A3)$$

### GWR

Multiple regression fits a linear model with a single set of coefficient estimates that describes the global relationships between the target variable and the covariates. GWR fits a series of localized linear regressions using subsets of spatially weighted nearby data falling under a moving window or kernel. In this way, it provides a means to explore spatial heterogeneity in data relationships, by mapping the resultant coefficients. The basic form of GWR is similar to that given in Equation (A1), but with locations associated with the coefficient terms:

$$y_i = \beta_0(\mathbf{s}) + \sum_{k=1}^m \beta_k(\mathbf{s}) x_{ik} + \varepsilon_i \quad (A4)$$

where  $\mathbf{s}$  is now the spatial location of the  $i^{th}$  observation. In matrix terms, the coefficients of GWR are estimated from:

$$\hat{\beta}(\mathbf{s}) = (\mathbf{X}^T \mathbf{W}(\mathbf{s}) \mathbf{X})^{-1} \mathbf{X}^T \mathbf{W}(\mathbf{s}) \mathbf{y} \quad (A5)$$

where  $\mathbf{W}(\mathbf{s})$  is a  $(n \times n)$  diagonal spatial weighting matrix determined from a pre-specified kernel function. The GWR prediction of  $y$  at any location  $\mathbf{s}$  is:

$$y_{GWR}^*(\mathbf{s}) = \mathbf{x}(\mathbf{s})^T \hat{\boldsymbol{\beta}}(\mathbf{s}) \quad (\text{A6})$$

The critical issue in GWR is to determine the kernel bandwidth (i.e., how many data to include in each of the localized subsets). Optimum bandwidths can be found by minimizing a model fit diagnostic, usually via a leave-one-out cross-validation score (Brunsdon et al., 1996) or via an AIC approach which penalizes for model complexity (Hurvich, Simonoff, and Tsai 1998). In this article's "synthetic" case study, a user-specified bandwidth was used and in the "real" case study, it was found optimally by leave-one-out cross-validation. The "weighted" part of GWR concerns the choice of a distance decay kernel such that data points further away from the kernel center contribute less to each local regression. For both case studies, a Gaussian kernel was specified with a fixed distance bandwidth, where for each data point, a weight  $w_{i,j}$  was calculated based on its distance to the kernel center as follows:

$$w_{i,j} = \exp\left(-\frac{1}{2} \left(\frac{d_{ij}}{b}\right)^2\right) \quad (\text{A7})$$

where  $d_{i,j}$  is the distance from the kernel center at  $i$  to an observation at point  $j$  and  $b$  is the bandwidth. Details of bandwidth specification for GWR together with further kernel schemes can be found in Gollini et al. (2015). Observe that as the bandwidth  $b$  tends to infinity, GWR tends to the corresponding multiple regression.

### Mixed models

A mixed model has a similar form to that given in Equation (A1), but spatial autocorrelation in the error term is accounted for, if present. In matrix terms, the coefficients of a mixed model fitted by REML are unbiasedly estimated from:

$$\hat{\boldsymbol{\beta}}_{REML} = \left(\mathbf{X}^T [\boldsymbol{\Sigma}_\theta]^{-1} \mathbf{X}\right)^{-1} \mathbf{X}^T [\boldsymbol{\Sigma}_\theta]^{-1} \mathbf{y} \quad (\text{A8})$$

where  $[\boldsymbol{\Sigma}_\theta]^{-1}$  represents unbiased variogram information of the (spatially-autocorrelated) residual process,  $\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}_{REML}$ . In this study, an exponential variogram model was chosen for this purpose, which is defined as:

$$\gamma(h) = c_0 + c_1 \left(1 - \exp\left(-\frac{h}{a}\right)\right) \quad (\text{A9})$$

where the parameters  $c_0$  and  $c_1$  represent the partial sills of the variogram and  $a$  is the correlation range;  $h$  is distance, assuming isotropy. Details for specifying variograms for a mixed model can be found in Chilés and Delfiner (1999). The mixed model prediction of  $y$  at an observed location  $\mathbf{s}$  is:

$$y_{MM}^*(\mathbf{s}) = \mathbf{x}(\mathbf{s})^T \hat{\boldsymbol{\beta}}_{REML} \quad (\text{A10})$$

Observe that if prediction at unobserved locations is required then the mixed model extends to its corresponding unbiased (regression) kriging model, where Equation (A10) also extends accordingly (Chilés and Delfiner 1999). In this mixed model, the model is fitted by maximizing the *restricted* log-likelihood. However, to illustrate the *i* issue (method of identification), a mixed model was also fitted by ML, where for ML parameter estimation, the log-likelihood is maximized (and Equations (A8)–(A10), follow accordingly).

## References

- Anselin, L. (1990). "Spatial Dependence and Spatial Structural Instability in Applied Regression Analysis." *Journal of Regional Science* 30(2), 185–207.
- Anselin, L. (2010). "Thirty Years of Spatial Econometrics." *Papers in Regional Science* 89(1), 3–25.
- Anyamba, A., K. J. Linthicum, and C. J. Tucker. (2001). "Climate-Disease Connections: Rift Valley Fever in Kenya." *Cadernos de saude publica* 17, S133–40.
- Armstrong, M. (1984). "Problems with Universal Kriging." *Mathematical Geology* 16, 101–8.
- Atkinson, P. M., and N. J. Tate. (2000). "Spatial Scale Problems and Geostatistical Solutions: A Review." *Professional Geographer* 52, 607–23.
- Basile, R., M. Durban, R. Minguez, J. M. Montero, and J. Mur. (2014). "Modeling Regional Economic Dynamics: Spatial Dependence, Spatial Heterogeneity and Non-Linearities." *Journal of Economic Dynamics and Control* 48, 229–45.
- Beck, L. R., B. M. Lobitz, and B. L. Wood. (2000). "Remote Sensing and Human Health: New Sensors and New Opportunities." *Emerging Infectious Diseases* 6(3), 217–27.
- Brunsdon, C., M. Charlton, and P. Harris. (2012). "Living with Collinearity in Local Regression Models." Proceedings of the 10th International Symposium on Spatial Accuracy Assessment in Natural Resources and Environmental Sciences, Brazil.
- Brunsdon, C., A. S. Fotheringham, and M. Charlton. (1996). "Geographically Weighted Regression: A Method for Exploring Spatial Nonstationarity." *Geographical Analysis* 28, 281–98.
- Brunsdon, C., A. S. Fotheringham, and M. Charlton. (1998). "Spatial Nonstationarity and Autoregressive Models." *Environment and Planning A* 30(6), 957–93.
- Carroll, R. J., and D. Rupert. (1988). *Transformation and Weighting in Regression*. New York, NY: Chapman and Hall.
- Casetti, E. (1972). "Generating Models by the Expansion Method: Applications to Geographic Research." *Geographical Analysis* 4, 81–91.
- Chen, L. D., W. Wei, B. J. Fu, and Y. H. Lu. (2007). "Soil and Water Conservation on the Loess Plateau in China: Review and Perspective." *Progress in Physical Geography* 31, 389–403.
- Chilès, J. P., and P. Delfiner. (1999). *Geostatistics—Modelling Spatial Uncertainty*. New York, NY: Wiley.
- Christensen, W. F. (2011). "Filtered Kriging for Spatial Data with Heterogeneous Measurement Error Variances." *Biometrika* 67(3), 947–57.
- Comber, A. J., P. F. Fisher, and R. A. Wadsworth. (2005). "What is land cover?" *Environment and Planning B: Planning and Design* 32, 199–209.
- Comber, A., and P. Harris. (2018). "Geographically Weighted Elastic Net Logistic Regression." *Journal of Geographical Systems* 20(4), 317–41.
- Cressie, N. (1993). *Statistics for Spatial Data*. New York, NY: John Wiley and Sons.
- Cressie, N., and N. H. Chan. (1989). "Spatial Modeling of Regional Variables." *Journal of the American Statistical Association* 84, 393–401.
- Derungs, C., F. Wartmann, R. S. Purves, and D. M. Mark. (2013). "The Meanings of the Generic Parts of Toponyms: Use and Limitations of Gazetteers in Studies of Landscape Terms." In *Spatial Information Theory. COSIT 2013. Lecture Notes in Computer Science*, vol. 8116, 261–78, edited by T. Tenbrink, J. Stell, A. Galton, and Z. Wood. Cham: Springer.
- Dong, G., R. Harris, K. Jones, and J. Yu. (2015). "Multilevel Modelling with Spatial Interaction Effects with Application to an Emerging Land Market in Beijing, China." *PLoS ONE* 10(6), 1–18.
- Frank, I. E., and J. H. Friedman. (1993). "A Statistical View of Some Chemometrics Regression Tools." *Technometrics* 35(2), 109–35.
- Fu, W., Y. Lü, P. Harris, A. Comber, and L. Wu. (2018). "Peri-Urbanization may Vary with Vegetation Restoration: A Large Scale Regional Analysis." *Urban Forestry and Urban Greening* 29, 77–87.
- Geniaux, G., and D. Martinetti. (2017). "A New Method for Dealing Simultaneously with Spatial Autocorrelation and Spatial Heterogeneity in Regression Models." *Regional Science and Urban Economics* 72, 74–85. <https://doi.org/10.1016/j.regsciurbeco.2017.04.001>.
- Gollini, I., B. Lu, M. Charlton, C. Brunsdon, and P. Harris. (2015). "GWmodel: an R Package for Exploring Spatial Heterogeneity using Geographically Weighted Models." *Journal of Statistical Software* 63, 1–50.

- Gotway, C. A., and L. J. Young. (2002). "Combining Incompatible Spatial Data." *Journal of the American Statistical Association* 97(458), 632–48.
- Harris, P. (2018). "A Simulation Study on Specifying a Regression Model for Spatial Data: Choosing between Autocorrelation and Heterogeneity Effects." *Geographical Analysis* 51(2), 151–81.
- Harvey, F. (2000). "The Social Construction of Geographical Information Systems." *International Journal of Geographical Information Science* 14(8), 711–3.
- Hay, S. I., C. A. Guerra, A. J. Tatem, P. M. Atkinson, and R. W. Snow. (2005). "Opinion—Tropical Infectious Diseases: Urbanization, Malaria Transmission and Disease Burden in Africa." *Nature Reviews Microbiology* 3(1), 81–90.
- Hijmans, R. J., and J. van Etten. (2017). raster: Geographic Data Analysis and Modeling. R package version, 2.6-7. <https://cran.r-project.org/web/packages/raster/raster.pdf>.
- Hu, J., Y. Lü, B. Fu, A. J. Comber, and P. Harris. (2017). "Quantifying the Effect of Ecological Restoration on Runoff and Sediment Yields: A Meta-Analysis for the Loess Plateau." *Progress in Physical Geography* 41(6), 753–74. <https://doi.org/10.1177/0309133317738710>.
- Hurvich, C. M., J. S. Simonoff, and C. L. Tsai. (1998). "Smoothing Parameter Selection in Nonpara-Metric Regression Using an Improved Akaike Information Criterion." *Journal of the Royal Statistical Society B* 60(2), 271–93.
- Li, T., Y. Lü, B. Fu, A. J. Comber, P. Harris, and L. Wu. (2017). "Gauging Policy-Driven Large-Scale Vegetation Restoration Programmes Under a Changing Environment: Their Effectiveness and Socioeconomic Relationships." *Science of the Total Environment* 607–608, 911–9. <https://doi.org/10.1016/j.scitotenv.2017.07.044>.
- Liu, Y., J. Hu, I. Snell-Feikema, M. S. Bommel, A. Lamsal, and M. C. Wimberly. (2015). "Software to Facilitate Remote Sensing Data Access for Disease Early Warning Systems." *Environmental Modelling & Software* 74, 247–57.
- Luo, Y., Y. Lü, B. Fu, P. Harris, L. Wu, and A. Comber. (2018). "When Multi-Functional Landscape Meets Critical Zone Science: Advancing Multi-Disciplinary Research for Sustainable Human Well-Being." *National Science Review* 6(2), 349–58. <https://doi.org/10.1093/nsr/nwy003>.
- Marcus, G. (2018). Deep Learning: A Critical Appraisal. arXiv preprint arXiv:1801.00631.
- Mark, D. M., and A. G. Turk. (2003). "Landscape Categories in Yindjibarndi: Ontology, Environment, and Language." In *Spatial Information Theory. Foundations of Geographic Information Science. COSIT 2003. Lecture Notes in Computer Science*, vol. 2825, 28–45, edited by W. Kuhn, M. F. Worboys, and S. Timpf. Berlin: Springer.
- Murakami, D., and M. Tsutsumi. (2015). "Area-to-Point Parameter Estimation with Geographically Weighted Regression." *Journal of Geographical Systems* 17(3), 207–25.
- Openshaw, S. (1984). *The Modifiable Areal Unit Problem*. Norwich, England: Geobooks.
- Pires, P. (2005). "Geospatial Conceptualisation: A Cross-Cultural Analysis on Portuguese and American Geographical Categorisations." In Spaccapietra S, editor. *Journal on Data Semantics III*, LNCS 3534, 196–212.
- Robbins, P. (2001). "Fixed Categories in a Portable Landscape: The Causes and Consequences of Land-Cover Categorization." *Environment and Planning A: Economy and Space* 33, 161–79.
- Rue, H., S. Martino, and N. Chopin. (2009). "Approximate Bayesian Inference for Latent Gaussian Models by Using Integrated Nested Laplace Approximations." *Journal of the Royal Statistical Society B* 71(2), 319–92.
- Smith, B. (2001). "Fiat objects." *Topoi* 20(2), 131–48.
- Smith, B., and D. M. Mark. (2003). "Do Mountains Exist? Towards an Ontology of Landforms." *Environment and Planning B: Planning and Design* 30, 411–27.
- Smith, B., and A. C. Varzi. (2000). "Fiat and Bona Fide Boundaries." *Philosophical and Phenomenological Research* 60(2), 401–20.
- Turk, A. G., D. M. Mark, and D. Stea. (2011). "Ethnophysiology." In *Landscape in Language Transdisciplinary Perspectives*, 25–45, edited by D. M. Mark, A. G. Turk, N. Burenhult, and D. Stea. Amsterdam, the Netherlands: John Benjamins Publishing.
- Wang, J., A. Stein, B. Gao, and Y. Ge. (2012). "A Review of Spatial Sampling." *Spat Stat* 2, 1–14.
- Wang, L., Q. Wang, S. Wei, M. A. Shao, Y. Li. (2008). "Soil Desiccation for Loess Soils on Natural and Regrown Areas." *Forest Ecology and Management* 255, 2467–77.

## Geographical Analysis

- Welham, S., B. Cullis, B. Gogel, A. Gilmour, and R. Thompson. (2004). "Prediction in Linear Mixed Models." *Australian and New Zealand Journal of Statistics* 44, 571–86.
- Wheeler, D. (2009). "Simultaneous Coefficient Penalization and Model Selection in Geographically Weighted Regression: the Geographically Weighted Lasso." *Environment and Planning A* 41(3), 722–42.
- Zhang, J., P. M. Atkinson, and M. Goodchild. (2014). *Scale in Spatial Information and Analysis*. Boca Raton, FL: CRC Press.
- Zou, H., and T. Hastie. (2005). "Regularization and Variable Selection via the Elastic Net." *Journal of the Royal Statistical Society, Series B* 67(2), 301–20.