

Rothamsted Repository Download

C2 - Non-edited contributions to conferences

Blackwell, M. S. A., Turner, B., Granger, S. J., Hooper, A. M., Darch, T., Hawkins, J. M. B., Yuan, H. and McGrath, S. P. 2015. Using ³¹P-NMR to investigate dynamics of soil phosphorus compounds in the Rothamsted Long Term Experiments. EGU.

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/8v1v6/using-31p-nmr-to-investigate-dynamics-of-soil-phosphorus-compounds-in-the-rothamsted-long-term-experiments>.

© Please contact library@rothamsted.ac.uk for copyright queries.



Using ^{31}P -NMR to investigate dynamics of soil phosphorus compounds in the Rothamsted Long Term Experiments

Martin Blackwell (1), Ben Turner (2), Steve Granger (1), Tony Hooper (3), Tegan Darch (1), Jane Hawkins (1), Huimin Yuan (4), and Steve McGrath (3)

(1) Rothamsted Research, North Wyke, Okehampton, Devon, EX20 2SB, UK. , (2) Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Republic of Panama., (3) Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, UK., (4) College of Resources and Environmental Sciences, China Agricultural University, No. 2 Yuanmingyuan West Road, Haidian, Beijing 100193, China

The technique of ^{31}P -NMR spectroscopy has done more to advance the knowledge of phosphorus forms (especially organic phosphorus) in environmental samples than any other method. The technique has advanced such that specific compounds can be identified where previously only broad categories such as orthophosphate monoesters and diesters were distinguishable. The Soil Archive and Long Term Experiments at Rothamsted Research, UK, potentially provides an unequalled opportunity to use this technique to observe changes in soil phosphorus compounds with time and under different treatments, thereby enhancing our understanding of phosphorus cycling and use by plants. Some of the earliest work using this technique on soils was carried out by Hawkes et al. in 1984 and this used soils from two of the oldest Rothamsted Long Term Experiments, namely Highfield and Park Grass. Here we revisit the samples studied in this early work and reanalyse them using current methodology to demonstrate how the ^{31}P -NMR technique has advanced. We also present results from a study on the phosphorus chemistry in soils along the Hoosfield acid strip (Rothamsted, UK), where a pH gradient from 3.7 to 7.8 occurs in a single soil with little variation in total phosphorus (mean \pm standard deviation $399 \pm 27 \text{ mg P kg}^{-1}$). Soil pH was found to be an important factor in determining the proportion of phosphomonoesters and phosphodiester in the soil organic phosphorus, although total organic phosphorus concentrations were a relatively consistent proportion of the total soil phosphorus ($36 \pm 2\%$) irrespective of soil pH.

Key words.

^{31}P -NMR, soil organic phosphorus, long term experiments, Hoosfield acid strip