

Rothamsted Repository Download

A - Papers appearing in refereed journals

Fan, J., Urban, M., Parker, J. E., Brewer, H. C., Kelly, S. L., Hammond-Kosack, K. E., Fraaije, B. A., Liu, X. and Cools, H. J. 2013.
Characterization of the sterol 14alpha-demethylases of *Fusarium graminearum* identifies a novel genus-specific CYP51 function. *New Phytologist*. 198 (3), pp. 821-835.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1111/nph.12193>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/8qw19>.

© Rothamsted Research. Licensed under the Creative Commons CC BY.

Characterization of the sterol 14 α -demethylases of *Fusarium graminearum* identifies a novel genus-specific CYP51 function

Jieru Fan^{1,2}, Martin Urban³, Josie E. Parker⁴, Helen C. Brewer³, Steven L. Kelly⁴, Kim E. Hammond-Kosack³, Bart A. Fraaije², Xili Liu¹ and Hans J. Cools²

¹Department of Plant Pathology, China Agricultural University, Beijing, 100193, China; ²Department of Biological Chemistry and Crop Protection, Rothamsted Research, Harpenden, AL5 2JQ, UK; ³Department of Plant Biology and Crop Science, Rothamsted Research, Harpenden, AL5 2JQ, UK; ⁴Institute of Life Science and College of Medicine, Swansea University, Swansea, SA2 8PP, UK

Summary

Author for correspondence:
Hans J. Cools
Tel: +44 1582 763 133
Email: hans.cools@rothamsted.ac.uk

Received: 19 November 2012
Accepted: 15 January 2013

New Phytologist (2013) 198: 821–835
doi: 10.1111/nph.12193

Key words: azole target, CYP51 function, cytochrome P450 evolution, deoxynivalenol production, fungicide resistance, *Fusarium* head scab/blight, sterol biosynthesis, wheat infection.

- *CYP51* encodes the cytochrome P450 sterol 14 α -demethylase, an enzyme essential for sterol biosynthesis and the target of azole fungicides. In *Fusarium* species, including pathogens of humans and plants, three *CYP51* paralogues have been identified with one unique to the genus. Currently, the functions of these three genes and the rationale for their conservation within the genus *Fusarium* are unknown.
- Three *Fusarium graminearum* *CYP51*s (*FgCYP51*s) were heterologously expressed in *Saccharomyces cerevisiae*. Single and double *FgCYP51* deletion mutants were generated and the functions of the *FgCYP51*s were characterized *in vitro* and *in planta*.
- *FgCYP51A* and *FgCYP51B* can complement yeast *CYP51* function, whereas *FgCYP51C* cannot. *FgCYP51A* deletion increases the sensitivity of *F. graminearum* to the tested azoles. In $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ mutants, ascospore formation is blocked, and eburicol and two additional 14-methylated sterols accumulate. *FgCYP51C* deletion reduces virulence on host wheat ears.
- *FgCYP51B* encodes the enzyme primarily responsible for sterol 14 α -demethylation, and plays an essential role in ascospore formation. *FgCYP51A* encodes an additional sterol 14 α -demethylase, induced on ergosterol depletion and responsible for the intrinsic variation in azole sensitivity. *FgCYP51C* does not encode a sterol 14 α -demethylase, but is required for full virulence on host wheat ears. This is the first example of the functional diversification of a fungal *CYP51*.

Introduction

The cytochrome P450 sterol 14 α -demethylase (*CYP51*, syn. ERG11) is an essential enzyme in the biosynthesis of sterols, critical components of cell membranes of all eukaryotic organisms required for the regulation of membrane fluidity and permeability (Parks *et al.*, 1995; Lepesheva & Waterman, 2007). Of the many sterols identified in fungi, ergosterol is the most common and is required for fungal growth (Rodriguez *et al.*, 1985; Weete *et al.*, 2010). Consequently, *CYP51* is a widely exploited target for the control of fungal pathogens of humans and plants, with the azole (imidazole and triazole) fungicides being the leading class of antifungals for over three decades (Sheehan *et al.*, 1999). Surprisingly, despite their widespread long-term use and single-site mode of action, incidences of resistance to azole fungicides are rare. More commonly, reductions in sensitivity are reported with cross-resistance within the azole class often incomplete. To date, three mechanisms of resistance to azoles predominate in filamentous fungi: mutations in the

CYP51 gene encoding amino acid alterations, resulting in decreased affinity of the protein for inhibitors (Loffler *et al.*, 1997; Sanglard *et al.*, 1998; Wyand & Brown, 2005; Cools *et al.*, 2010); over-expression of the target *CYP51* gene, most frequently caused by insertions in the predicted promoter regions (Mellado *et al.*, 2007; Cools *et al.*, 2012); and over-expression of genes encoding efflux pumps (Sanglard *et al.*, 1995; Kretschmer *et al.*, 2009). These mechanisms can combine, and therefore resistance levels in fungal strains are often determined by combinations of *CYP51* amino acid alterations, *CYP51* gene over-expression and/or enhanced efflux (Perea *et al.*, 2001; Mellado *et al.*, 2007; Cools *et al.*, 2012).

Until recently, *CYP51* was thought to exist as a single gene in all phyla. Mammalian genomes, for example, contain one *CYP51*, with some nonfunctional pseudogenes identified (Rozman *et al.*, 1996). However, increasing genome sequence information has shown that this is not the case in all kingdoms. To date, multiple *CYP51* genes have been found in plants, including rice (12), oats (two), tobacco (two) and *Arabidopsis*

thaliana (two) (Lepesheva & Waterman, 2007). Filamentous fungi, particularly Ascomycetes, often possess two or more *CYP51* paralogues, for example in *Penicillium digitatum* (two), *Aspergillus fumigatus* (two), *A. nidulans* (two), *A. flavus* (three), *Magnaporthe oryzae* (two) and species of *Fusarium*, including *F. verticillioides*, *F. oxysporum* f. sp. *lycopersici* and *F. graminearum* (three). Molecular phylogenetic analysis has shown that the *CYP51* genes of fungi within the subphylum Pezizomycotina fall into three clades, designated A, B and C (Becher *et al.*, 2011). All species possess a *CYP51* in clade B (*CYP51B*). Species with multiple paralogues carry an additional *CYP51* in clade A (*CYP51A*), with duplications of *CYP51A* or *CYP51B* generating the third paralogue in some species, for example *A. flavus* and *A. terreus*, respectively. Uniquely, the third *CYP51* paralogue in *Fusarium* species forms a distinct clade, *CYP51C*. The *CYP51C* gene is found exclusively in *Fusarium* species, and is ubiquitous across the genus, as demonstrated by its use as a reliable phylogenetic marker (Fernández-Ortuño *et al.*, 2010).

Fungi with multiple *CYP51s* are intrinsically resistant to some azoles, although some remain effective. For example, *A. fumigatus* is well controlled by itraconazole and voriconazole, whereas fluconazole is ineffective. Species of *Fusarium*, for example *F. solani*, are resistant to commonly used medical azoles, including fluconazole, voriconazole and the recently introduced posaconazole (Nucci & Anaissie, 2007). Deletion of *CYP51A* increases the intrinsic sensitivity to some azoles in *M. oryzae* (e.g. tebuconazole and prochloraz), *A. fumigatus* (e.g. fluconazole and ketocozazole) and *F. graminearum* (e.g. tebuconazole and prochloraz) (Mellado *et al.*, 2005; Liu *et al.*, 2011; Yan *et al.*, 2011). Furthermore, resistance to effective azoles in fungi with multiple *CYP51s* is most frequently mediated by changes in the *CYP51A* paralogue. For example, over-expression of *CYP51A* has been reported for resistant isolates of *P. digitatum* (Hamamoto *et al.*, 2000; Ghosop *et al.*, 2007), and mutation of *AfCYP51A* is the most common mechanism of resistance in *A. fumigatus* isolates (Diaz-Guerra *et al.*, 2003; Mellado *et al.*, 2005), which, when combined with *AfCYP51A* over-expression, confers a multi-azole-resistant phenotype (Mellado *et al.*, 2007; Snelders *et al.*, 2008).

The control of *F. graminearum*, the most important pathogen causing *Fusarium* head blight (FHB) or head scab disease on wheat and barley, is primarily dependent on effective azole fungicides (e.g. tebuconazole, metconazole and prothioconazole). Effective application of azoles reduces the content of the harmful trichothecene mycotoxin deoxynivalenol (DON) in wheat grains (Beyer *et al.*, 2006; Paul *et al.*, 2008). However, an increased level of DON has been detected after treatment with sublethal doses of prothioconazole both *in vitro* and *in planta* (Audenaert *et al.*, 2010), and trichothecenes accumulate in grain samples after treatment with tebuconazole (Kulik *et al.*, 2012). In addition, a study using the enhanced green fluorescent protein gene (*egfp*) as a reporter demonstrated activation of the *F. graminearum* trichothecene biosynthetic *TRI5* gene by sublethal concentrations of tebuconazole (Ochiai *et al.*, 2007); both *TRI4* and *TRI5* transcript levels were higher after tebuconazole treatment in culture (Kulik *et al.*, 2012). Currently, although stress responses and

enhanced secondary metabolism have been proposed, the mechanism(s) responsible for altered mycotoxin production after azole treatment is unknown.

Functional analysis of multiple *CYP51s* has identified roles additional to primary sterol biosynthesis. In oats, a *CYP51* homologue, *AsCyp51H10*, is dispensable for sterol biosynthesis, but is required for the synthesis of avenacins, antimicrobial compounds unique to the genus *Avena* (Qi *et al.*, 2006). Heterologous expression in yeast identified *AsCyp51H10* as the first *CYP51* not classified as a sterol demethylase (Kunii *et al.*, 2012). Recent analysis of the 12 *CYP51s* of rice has differentiated some genes as *CYP51H*, a group likely to have functions beyond sterol biosynthesis (Inagaki *et al.*, 2011). Previous work on the *CYP51* genes of *F. graminearum* (*FgCYP51*) has demonstrated that the deletion of individual *FgCYP51* genes can reduce conidiation, but otherwise causes no changes in *in vitro* morphology, mycelial growth rate or ergosterol content (Liu *et al.*, 2011).

In this study, we have determined the competence of the three paralogous *FgCYP51* genes to act as sterol 14 α -demethylases by heterologous expression in *S. cerevisiae* strain YUG37:*erg11*, which carries a regulatable promoter controlling native *CYP51* expression, and found that *FgCYP51C* cannot complement the *CYP51* function of the yeast gene. We generated single ($\Delta FgCYP51A$, $\Delta FgCYP51B$ and $\Delta FgCYP51C$) and combined ($\Delta FgCYP51AC$ and $\Delta FgCYP51BC$) *CYP51* deletion mutants and characterized their function *in vitro* and *in planta*. We report distinct roles of the three *FgCYP51* genes, with the *FgCYP51C* gene specifically required for full virulence on host wheat ears, but not on *Arabidopsis* floral tissue or the fruits of apple and tomato.

Materials and Methods

Fungal strains, growth assays and sporulation tests

Fusarium graminearum isolates (Table 1) used in this study were routinely cultured on synthetic nutrient-poor agar (SNA) plates, as described previously (Urban *et al.*, 2002). Conidiation assays were performed on SNA plates, and the spores were germinated in YPS medium (0.3% (w/v) yeast extract, 0.3% (w/v) bacto-peptone and 20% (w/v) sucrose) (Cuzick *et al.*, 2008). Colony morphology was assayed on potato dextrose agar (PDA) plates in the dark at 22°C. Growth rate tests were conducted in potato dextrose broth (PDB) and Czapek Dox liquid medium (CZD) using an amended 96-well microtitre plate method. Absorbance measurements were made every 8 h for 96 h at 22°C (Fraaije *et al.*, 2007). Carrot agar plates were used for perithecia formation and ascospore production (Ehrenshaft *et al.*, 1995). Approximately 50 asci per strain/mutant were scraped from the plates and physically broken to determine ascospore formation.

Heterologous expression of *FgCYP51* genes in *S. cerevisiae* YUG37::*erg11*

Heterologous expression of *FgCYP51* genes in *S. cerevisiae* YUG37::*erg11* followed the method described by Cools *et al.* (2010). *FgCYP51* genes were amplified from cDNA of

Table 1 Fungal strains/mutants used in this study

| Strain | Description | Reference |
|---------------------|------------------------------|---------------------------------|
| Fg1955 | Wild-type, Finland origin | Nicolaisen <i>et al.</i> (2009) |
| PH-1 | Wild-type, USA origin | Cuomo <i>et al.</i> (2007) |
| $\Delta FgCYP51A4$ | CYP51A gene deletion mutant | This study |
| $\Delta FgCYP51A5$ | CYP51A gene deletion mutant | This study |
| $\Delta FgCYP51B3$ | CYP51B gene deletion mutant | This study |
| $\Delta FgCYP51B4$ | CYP51B gene deletion mutant | This study |
| $\Delta FgCYP51C2$ | CYP51C gene deletion mutant | This study |
| $\Delta FgCYP51C3$ | CYP51C gene deletion mutant | This study |
| $\Delta FgCYP51AC1$ | CYP51AC gene deletion mutant | This study |
| $\Delta FgCYP51AC3$ | CYP51AC gene deletion mutant | This study |
| $\Delta FgCYP51BC2$ | CYP51BC gene deletion mutant | This study |
| $\Delta FgCYP51BC3$ | CYP51BC gene deletion mutant | This study |
| $\Delta FgCYP51BC4$ | CYP51BC gene deletion mutant | This study |

F. graminearum isolate Fg1955 (Nicolaisen *et al.*, 2009) with the primers shown in Supporting Information Table S1 designed and cloned into the yeast expression vector pYES3/CT (Invitrogen). Plasmids were transformed into *S. cerevisiae* strain YUG37::*erg11* (Revankar *et al.*, 2004) using an S.c. EasyComp transformation kit (Invitrogen) with the pYES3/CT vector as a negative control. The complementation efficiency of two independent transformants per construct was screened on Synthetic Drop-out (SD) galactose (GAL) + raffinose (RAF) medium containing $3 \mu\text{g ml}^{-1}$ doxycycline (Sigma-Aldrich) by droplet inoculation with $5 \mu\text{l}$ of cell suspensions (six five-fold dilutions of a starting concentration of 1×10^6 cells ml^{-1}) for each transformant and incubation for 6 d at 30°C . The growth rate of YUG37::*erg11* transformants was determined by inoculation of liquid SD GAL + RAF medium containing $6 \mu\text{g ml}^{-1}$ doxycycline with $100 \mu\text{l}$ of a cell suspension of 1×10^6 cells ml^{-1} (grown for 24 h at 30°C). Absorbance was measured every 4 h for 96 h at 30°C and the growth rate was calculated at the linear growth stage.

Generation of gene deletion mutants

FgCYP51 gene deletion mutants were generated in *F. graminearum* strain PH-1 (NRRL 31084) (Cuomo *et al.*, 2007), the genome of which has been sequenced, using polyethyleneglycol (PEG)-mediated protoplast transformation of split-marker PCR fragments (Catlett *et al.*, 2003). The primers used to amplify the flanking sequences for each gene are listed in Table S1. PCR products were transformed into protoplasts of wild-type PH-1 to delete single *FgCYP51* genes. Transformants were selected with $75 \mu\text{g ml}^{-1}$ hygromycin B or geneticin. The split hygromycin markers of the *FgCYP51A* and *FgCYP51B* genes were transformed into the *FgCYP51C* gene deletion mutants carrying a geneticin selectable marker to generate the double deletion mutants, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$. Candidate transformants were screened using 5' flank and 3' flank PCRs to confirm the targeted deletion of the genes (Fig. S1). An additional PCR to amplify the 5' flank and 3' flank of the nontargeted *FgCYP51* gene confirmed that the remaining *FgCYP51* genes were not disrupted. Double deletion of *FgCYP51A* and *FgCYP51B* was considered to be lethal.

Fungicide sensitivity testing

Fungicide sensitivities were tested using 96-well microtitre plate sensitivity assays (Fraaije *et al.*, 2007). Briefly, spore suspensions of *Fusarium* isolates with 2×10^4 spores ml^{-1} were added to PDB amended with serial diluted concentrations of fungicides (Table S2). For *S. cerevisiae* transformants, SD GAL + RAF medium containing $6 \mu\text{g ml}^{-1}$ doxycycline (Sigma-Aldrich) and amended with the serial diluted concentrations of fungicides was inoculated with 1×10^6 cells ml^{-1} (Table S2) (Cools *et al.*, 2010). The absorbance was measured at 630 nm using a FLUOstar OPTIMA microplate reader (BMG Labtech GmbH, Offenburg, Germany) after 96 h of incubation at room temperature for *F. graminearum* and at 30°C for *S. cerevisiae* transformants. Fungicide sensitivities were determined as the 50% effective concentration (EC_{50}) using a dose–response relationship according to the BMG Labtech Optima Software. Three independent replicates were performed for statistical analysis.

Gene expression analysis

Total RNA was extracted from freeze–dried samples with TRIzol reagent (Invitrogen), followed by a 4 M lithium chloride purification and DNase I treatment. For *in vitro* *FgCYP51* expression analysis, RNA was extracted from mycelia of isolate Fg1955 grown in PDB for 24, 48, 72 and 96 h in the absence of azole, or after treatment with tebuconazole, epoxiconazole or prochloraz at the calculated EC_{50} at 24 h of incubation. *In vitro* gene expression of $\Delta FgCYP51$ mutants used RNA extracted after 48 h of growth in PDB, compared with expression in wild-type parent strain PH-1. For *in planta* expression analysis, RNA was extracted from inoculated wheat ears harvested at 7 d post-inoculation (dpi). First-strand cDNA was prepared using the SuperScript III first-strand synthesis system with oligo(dT) (Invitrogen). Real-time PCR amplifications were performed on an ABI 7500 sequence detection system (Applied Biosystems, Foster City, CA, USA) using SYBR Green I fluorescent dye detection. The relative quantities (RQs) were calculated using the $2^{-\Delta\Delta C_t}$ method with *F. graminearum actin* and β -*tubulin* as reference genes

and the statistical analysis of three biological replicates. The primers used for quantitative PCR are listed in Table S1.

Sterol extraction and analysis

Fusarium graminearum strains PH-1 and $\Delta FgCYP51$ mutants were collected after growth on PDB medium at 23°C for 48 h. Nonsaponifiable lipids were extracted as described previously (Kelly *et al.*, 1995). Briefly, pellets were resuspended in 2.5 ml of methanol, followed by the addition of 1.5 ml of 60% (w/v) KOH and 1 ml of 5 mg ml⁻¹ pyrogallol (in methanol). Samples were refluxed at 90°C for 2 h, and then extracted twice with hexane and dried in a rotary evaporator (Heto, Allerod, Denmark). Samples were derivatized by the addition of 100 µl of 90% bis(trimethylsilyl)-trifluoroacetamide (BSTFA) – 10% trimethylsilyl (TMS) (Sigma-Aldrich) and 50 µl of anhydrous pyridine (Sigma-Aldrich) and heating for 2 h at 80°C. Gas chromatography-mass spectrometry was performed using a VG12-250 mass spectrometer (VG Biotech, Mumbai, India) with splitless injection and chemical ionization. Data were analysed using an MSD Enhanced ChemStation (Agilent Technologies), and TMS esters of individual sterols were identified by reference to relative retention times, mass ions and fragmentation patterns. Data present the relative composition in total sterol in each sample. Three biological replicates were performed.

Plant disease assays

Wheat ear infection, gene expression and DON concentration measurement Wheat (*Triticum aestivum* L., cv Bobwhite) plants were grown and infected by point inoculation as described previously (Urban *et al.*, 2003). The 11th and 12th flowering spikelets counted from the stem were inoculated with 5 µl of 5 × 10⁴ spores per spikelet or mock inoculated with sterile water. The number of infected spikelets was recorded at 3, 7, 11, 16 and 21 dpi. For measurements of DON concentration and *in planta* gene expression assays, 14 flowering spikelets were inoculated per ear. The infected spikelets were harvested at 7 dpi for gene expression analyses and at 10 dpi for DON measurements. Samples were lyophilized and infected spikelets were ground to a fine powder in liquid nitrogen using a mortar and pestle. The powder was resuspended in water as described previously (Urban *et al.*, 2003). After incubation at 30°C for 30 min with shaking at 80 rpm, the solutions were centrifuged and the supernatants were analysed quantitatively for DON content using an EZ-Quant Vomitoxin (DON) plate kit (Diagnostix, Ontario, Canada), according to the manufacturer's instructions. Three wheat ears of different wheat plants were inoculated for each strain/mutant in each replicate experiment. Three independent replicate experiments (total of nine ears per strain/mutant inoculation) were carried out and analysed statistically.

Arabidopsis floral inoculations *Arabidopsis* plants (ecotype Landsberg) were grown and inoculated as described previously (Cuzick *et al.*, 2008). Five-week-old flowering plants were spray inoculated with c. 0.5 ml of conidial suspensions of 1 × 10⁵ spores ml⁻¹ of wild-type PH-1 or $\Delta FgCYP51$ strains,

and incubated in a 100% relative humidity box. Each isolate was sprayed onto three plants per experiment, with two experimental replicates performed (total of six plants per strain/mutant inoculation), with sterile water as a negative control. Disease symptoms on apical flowers and siliques were assessed using the *Fusarium*–*Arabidopsis* disease (FAD) scoring system (Urban *et al.*, 2002), with the modification of combining old and new silique scores. Floral and silique scores were added to give the total FAD score shown in Table S3.

Infection assays on apple fruit sections The ability of $\Delta FgCYP51$ mutants to colonize apple fruit sections relative to strain PH-1 was tested on cv Braeburn. After surface sterilization with 70% ethanol, apples were sliced into 8-mm-thick sections and inoculated in three discrete areas with a 5-µl droplet of a spore suspension containing 5 × 10⁵ ml⁻¹ conidia of wild-type isolate PH-1, sterile water control or a $\Delta FgCYP51$ gene mutant. In each experiment, three apple slices were inoculated for each mutant. Three independent experiments were carried out. Apple slices were sealed in a 9-cm Petri dish and incubated at 28°C for 6 d.

Infection assays on tomato fruits To assay the colonization of ripe tomato fruits (cultivar Encore) by $\Delta FgCYP51$ mutants, three wounds were made with a hyperdermic needle (BD microlance 23G 10.6 × 25) on 70% ethanol surface-sterilized tomatoes. Fruits were inoculated with 3 × 5 µl of 5 × 10⁵ ml⁻¹ conidia suspensions and incubated at 28°C in a sealed plastic container at 100% humidity for 6 d (Urban *et al.*, 2003). For each strain/mutant, three tomatoes were inoculated in each of three independent experiments. Mock inoculations were with sterile distilled water.

Statistical analysis

Data were analysed from two or three independent replicates using ANOVA. Following a significant *F*-test result ($P < 0.0001$ for plant infection, $P < 0.05$ for fungicide sensitivity tests and *in vitro/in planta* gene expression), means of independent replicates were compared using least-significant differences (LSD). No transformation of the data was required. $P < 0.05$ was considered to be significant. Greek letters were used to mark statistically significant differences ($P < 0.05$).

Results

FgCYP51A expression is induced by azole fungicides *in vitro*

In the absence of fungicide, transcript levels of *FgCYP51A* and *FgCYP51B* were highest at 72 h of incubation (Fig. 1). *FgCYP51C* expression did not change significantly across the time course. Twenty-four hours after treatment (48 h after inoculation) with subinhibitory concentrations of azoles, the expression of *FgCYP51A* was induced up to 30-fold (prochloraz and tebuconazole) or 100-fold (epoxiconazole) relative to the untreated control. *FgCYP51B* and *FgCYP51C* transcript levels did not change significantly after azole treatment.




Fig. 1 Time course of *in vitro* constitutive and induced *FgCYP51* expression. Growth of *Fusarium graminearum* was in potato dextrose broth (PDB) liquid medium. Treatment with prochloraz, tebuconazole or epoxiconazole at EC₅₀ concentration in the linear growth stage at 24 h of incubation, followed by sample collection at 24, 48, 72 and 96 h of incubation. The relative quantification (RQ) of the *FgCYP51A*, *FgCYP51B* and *FgCYP51C* genes was analysed using the $2^{-\Delta\Delta C_t}$ method with *actin* as the reference gene and untreated 24 h of incubation as a calibrator. The error bars show \pm SE of three biological replicates.




Fig. 2 Complementation of *Saccharomyces cerevisiae* strain YUG37::*erg11* with *FgCYP51A*, *FgCYP51B* or *FgCYP51C*. The growth of cells (six five-fold dilutions of a starting concentration of 1×10^6 cells ml⁻¹) in the absence (-DOX) and presence (+DOX) of doxycycline is shown. Yeast expression vector pYES3/CT without the *FgCYP51* gene was transformed into the yeast strain as a negative control.

Heterologous expression of *FgCYP51* genes in *S. cerevisiae* strain YUG37::*erg11*

FgCYP51A, *FgCYP51B* and *FgCYP51C* were expressed in *S. cerevisiae* strain YUG37::*erg11*, in which native *CYP51* gene (*ScCYP51*) expression is under the control of a doxycycline-repressible promoter (Revankar *et al.*, 2004). *FgCYP51A* and *FgCYP51B* genes were able to complement *ScCYP51* function; however, no growth was observed with transformants expressing *FgCYP51C* on doxycycline amended medium, a phenotype similar to the empty vector control (Fig. 2). Yeast transformants expressing *FgCYP51A* grew more slowly than those expressing *FgCYP51B* in the presence of doxycycline (Table 2). Yeast transformants expressing *FgCYP51A* were less sensitive to epoxiconazole than to tebuconazole and prochloraz. However, the sensitivity of *FgCYP51B* transformants was similar to all three azoles (Table 3). There was no difference in cycloheximide sensitivity, a non-azole fungicide, of yeast transformants expressing *FgCYP51A* relative to *FgCYP51B*.

In vitro growth characteristics of *FgCYP51* deletion strains

There were no differences in growth rate between mutants $\Delta FgCYP51A$, $\Delta FgCYP51B$, $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ (Table 4) and the wild-type parental strain PH-1 in PDB or CZD (Table 4). Nor were there morphological

Table 2 Growth rate of yeast transformants expressing *FgCYP51* genes of *Fusarium graminearum*

| Transformants | Growth rate maximum ^a | |
|---------------|----------------------------------|----------------------------|
| | -DOX ^b | +DOX |
| pYES-Fg51B | 8.6 \pm 0.2 | 3.5 \pm 0.1 ^c |
| pYES-Fg51A | 8.7 \pm 0.2 | 1.6 \pm 0.1 ^d |
| pYES-Fg51C | 11.4 \pm 0.0 | 0.1 \pm 0.0 ^d |
| pYES (Vector) | 10.7 \pm 1.4 | 0.0 \pm 0.0 ^d |

^aValues represent the greatest increase in the optical density at 600 nm (OD₆₀₀) in the absence or presence of doxycycline over a 4-h period measurement, and are means of two transformants from two independent replicates \pm SD.

^bCalculation at linear growth period (16–28 h).

^cCalculation at linear growth period (26–52 h).

^dCalculation at linear growth period (40–76 h).

differences on SNA plates under UV light (data not shown), or for single *FgCYP51* deletion strains on PDA incubated in the dark (Fig. 3a), confirming the findings of Liu *et al.* (2011). However, the morphology of double gene deletion mutants uniquely generated in this study ($\Delta FgCYP51AC$ and $\Delta FgCYP51BC$) on PDA incubated in the dark (Fig. 3a) was different. Growth was slower and the colony pigment was yellow rather than the pink of wild-type and single gene deletion strains. Although $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ strains produced around half the number of conidia of the PH-1 strain, conidial morphologies (data not shown) and germination rates were normal (Table 4).

FgCYP51B is essential for ascospore production

Perithecia were produced by all strains with no morphological differences relative to PH-1. However, 30 d after inoculation, no ascospores oozed from perithecia of $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ and, when perithecia were physically broken, no ascospores were found (Fig. 3b). Ascospores produced by $\Delta FgCYP51A$, $\Delta FgCYP51C$ and $\Delta FgCYP51AC$ showed wild-type morphology and germinated normally (data not shown).

Table 3 Azole sensitivity of *Fusarium graminearum* isolates and *Saccharomyces cerevisiae* YUG37:erg11 transformants

| Isolate/yeast transformant | EC ₅₀ (µg ml ⁻¹) ^a | | | |
|----------------------------|--|-----------------|-----------------|-----------------|
| | Prochloraz | Tebuconazole | Epoxiconazole | Cycloheximide |
| Fg1955 ^b | 0.0264 ± 0.0025 | 0.3410 ± 0.0010 | 1.4550 ± 0.1050 | – |
| pYES-Fg51A ^c | 0.0043 ± 0.0003 | 0.0063 ± 0.0003 | 0.0261 ± 0.0089 | 0.0537 ± 0.0005 |
| pYES-Fg51B ^c | 0.0160 ± 0.0031 | 0.0351 ± 0.0054 | 0.0103 ± 0.0044 | 0.0590 ± 0.0005 |

^aEC₅₀ values (µg ml⁻¹) are means of EC₅₀ from two independent replicates ± SD.

^b*Fusarium graminearum* isolate.

^c*Saccharomyces cerevisiae* YUG37:erg11 transformants expressing *FgCYP51* genes.

–, means not measured.




Fig. 3 *In vitro* characteristics of *Fusarium graminearum* wild-type PH-1 and *FgCYP51* gene deletion mutants. (a) Hyphal morphology after incubation on potato dextrose agar (PDA) in the dark. (b) Ascospore formation in perithecia of *F. graminearum* wild-type PH-1 and Δ*FgCYP51B* and Δ*FgCYP51BC* gene deletion mutants. Arrows show: a, asci with eight ascospores; b, hyphae forming the black pigmented perithecia.

Table 4 Phenotype characteristics of *FgCYP51* gene deletion mutants

| Strain/mutant | Growth rate ^a | | Conidiation ^b (10 ⁵ spores per plate) | Conidia germination ^c (%) |
|--------------------|-------------------------------|------------------------------|---|--------------------------------------|
| | CZD medium | PDB medium | | |
| PH-1 | 0.0145 ± 0.0005 ^{α*} | 0.3664 ± 0.0046 ^α | 38.2 ± 5.6 ^α | 94.7 ± 0.8 ^α |
| Δ <i>FgCYP51B</i> | 0.0135 ± 0.0019 ^α | 0.3499 ± 0.0705 ^α | 27.5 ± 4.8 ^β | 94.4 ± 1.1 ^α |
| Δ <i>FgCYP51A</i> | 0.0124 ± 0.0011 ^α | 0.3560 ± 0.0141 ^α | 33.8 ± 7.4 ^{αβ} | 94.6 ± 0.4 ^α |
| Δ <i>FgCYP51C</i> | 0.0112 ± 0.0002 ^α | 0.3304 ± 0.0032 ^α | 28.7 ± 3.8 ^{αβ} | 94.7 ± 0.5 ^α |
| Δ <i>FgCYP51AC</i> | 0.0114 ± 0.0001 ^α | 0.3627 ± 0.0060 ^α | 18.1 ± 3.2 ^γ | 94.4 ± 0.6 ^α |
| Δ <i>FgCYP51BC</i> | 0.0124 ± 0.0002 ^α | 0.3301 ± 0.0024 ^α | 19.4 ± 2.5 ^γ | 94.8 ± 0.7 ^α |

^aValues represent the greatest increase in the optical density at 600 nm (OD₆₀₀) in Czapek Dox medium (CZD) or potato dextrose broth (PDB) medium over an 8-h period and are means for two transformants from two independent replicates ± SD. The growth rate in CZD medium was calculated at the linear growth period (36–72 h), and the growth rate in PDB medium was calculated at the linear growth period (24–60 h).

^bConidia produced from 7 d of incubation on synthetic nutrient-poor agar (SNA) plate with 10⁴ conidia inoculation.

^cPercentage of germinated conidia was measured after 16 h of incubation in YPS medium at room temperature.

*Data were analysed using ANOVA ($P < 0.05$, F test). Different Greek letters are used to mark statistically significant differences between strains/mutants ($P < 0.05$, least-significant difference (LSD)).

Contribution of *FgCYP51A* and *FgCYP51B* to intrinsic differences in *F. graminearum* azole sensitivity

According to the sensitivity of Δ*FgCYP51A* and Δ*FgCYP51B* mutants, the seven azole fungicides tested can be divided into two groups (Table 5). Sensitivity to group 1, which includes tebuconazole, epoxiconazole, propiconazole and imazalil, is only affected significantly by *FgCYP51A* deletion. For example, Δ*FgCYP51A* strains are > 30-fold more sensitive to epoxiconazole

than are wild-type and Δ*FgCYP51B* strains. Group 2 includes metconazole, prochloraz and difenoconazole. Deletions of both *FgCYP51A* and *FgCYP51B* increased the sensitivity to these compounds. There was no significant difference in the sensitivity of single *FgCYP51C* gene deletion mutants to all the azoles tested relative to the wild-type PH-1, in contrast with previous reports (Liu *et al.*, 2011). In addition, there were no significant differences in azole sensitivity between Δ*FgCYP51A* and Δ*FgCYP51AC*, confirming that *FgCYP51C* has no effect on azole sensitivity.

Table 5 Azole sensitivity of *Fusarium graminearum* wild-type isolates PH-1 and *FgCYP51* gene deletion mutants^a

| Strain/mutant | Metconazole | | Prochloraz | | Difenoconazole | | Tebuconazole | | Epoxiconazole | | Propiconazole | | Imazalil | |
|-------------------|---|-----|---|-----|---|-----|---|-----|---|------|---|-----|---|-----|
| | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF |
| PH-1 | 0.0284 ± 0.0022 ^{**} | 1 | 0.0250 ± 0.0026 ^z | 1 | 0.6350 ± 0.0600 ^z | 1 | 0.2044 ± 0.0220 ^z | 1 | 0.4327 ± 0.0422 ^β | 1 | 0.4629 ± 0.0536 ^β | 1 | 0.4963 ± 0.0415 ^z | 1 |
| <i>ΔFgCYP51A</i> | 0.0132 ± 0.0029 ^β | 0.5 | 0.0099 ± 0.0020 ^γ | 0.4 | 0.2855 ± 0.0400 ^γ | 0.5 | 0.0668 ± 0.0137 ^β | 0.3 | 0.0139 ± 0.0036 ^γ | 0.03 | 0.0908 ± 0.0031 ^γ | 0.2 | 0.0668 ± 0.0249 ^β | 0.1 |
| <i>ΔFgCYP51B</i> | 0.0083 ± 0.0009 ^γ | 0.3 | 0.0116 ± 0.0017 ^γ | 0.5 | 0.4528 ± 0.0513 ^β | 0.7 | 0.1783 ± 0.0623 ^z | 0.8 | 0.3893 ± 0.0188 ^β | 0.9 | 0.4273 ± 0.0391 ^β | 0.9 | 0.5710 ± 0.1160 ^z | 1.1 |
| <i>ΔFgCYP51C</i> | 0.0298 ± 0.0024 ^z | 1.0 | 0.0240 ± 0.0039 ^z | 1.0 | 0.5348 ± 0.0784 ^z | 0.8 | 0.2284 ± 0.0420 ^z | 1.1 | 0.3923 ± 0.0320 ^β | 0.9 | 0.4492 ± 0.0792 ^β | 1.0 | 0.5770 ± 0.0790 ^z | 1.1 |
| <i>ΔFgCYP51AC</i> | 0.0158 ± 0.0006 ^β | 0.6 | 0.0136 ± 0.0012 ^γ | 0.5 | 0.2960 ± 0.0423 ^γ | 0.5 | 0.1153 ± 0.0084 ^β | 0.6 | 0.0268 ± 0.0013 ^γ | 0.06 | 0.0613 ± 0.0070 ^γ | 0.1 | 0.0406 ± 0.0131 ^β | 0.1 |
| <i>ΔFgCYP51BC</i> | 0.0140 ± 0.0037 ^β | 0.5 | 0.0190 ± 0.0034 ^β | 0.8 | 0.3683 ± 0.0359 ^{βγ} | 0.6 | 0.2053 ± 0.0394 ^z | 1.0 | 0.5470 ± 0.0351 ^z | 1.3 | 0.5703 ± 0.0526 ^z | 1.2 | 0.5066 ± 0.0688 ^z | 1.0 |

^aEC₅₀ (µg ml⁻¹) values are the means of two transformants from three independent replicates ± SD. Mean resistance factor (RF), representing the sensitivity change of each isolate, was calculated as the fold change in EC₅₀ compared with wild-type PH-1.

^bData were analysed using ANOVA ($P < 0.05$, F test). Different Greek letters are used to mark statistically significant differences between strains/mutants ($P < 0.05$, least-significant difference (LSD)).

Deletion of *FgCYP51B* and *FgCYP51C* decreases sensitivity to fenhexamid, an erg27 inhibitor

Testing the sensitivity to other ergosterol biosynthesis inhibitors (EBIs, Table 6) revealed no differences in amphotericin B sensitivity, an inhibitor that binds to ergosterol (Gray *et al.*, 2012), between PH-1 and the *FgCYP51* deletion mutants, suggesting no differences in ergosterol content between strains. *ΔFgCYP51B* and *ΔFgCYP51C* mutants were significantly less sensitive to fenhexamid, an erg27 inhibitor, relative to the wild-type strain. The *ΔFgCYP51BC* double mutants were least sensitive. There was no effect of *FgCYP51A* deletion on fenhexamid sensitivity. For all single *FgCYP51* deletion mutants, there was no difference in sensitivity to the erg1 inhibitor terbinafine relative to the wild-type strain PH-1. However, the double gene deletion mutants, *ΔFgCYP51AC* and *ΔFgCYP51BC*, were significantly less sensitive to terbinafine. There were no differences in sensitivity between mutant strains and the wild-type to the other EBIs tested, or to the β-tubulin assembly inhibitor carbendazim.

Sterol profiles of PH-1 and the *ΔFgCYP51* strains

Fourteen sterols were detected in the mycelia of the wild-type *F. graminearum* isolate PH-1 (Table 7). The most abundant sterol was ergosterol (72.98%), followed by 4,4-dimethyl ergosta-8,24(28)-dienol (5.24%), eburicol (3.46%), lanosterol (2.15%), 4-methyl ergosta-8,24-dienol (4.01%), brassicasterol (2.46%) and other minor sterols (< 2%). A proposed ergosterol biosynthesis pathway, based on the compounds detected in *FgCYP51* deletion mutants and the wild-type PH-1 (Table 7), and the reported sterol biosynthesis pathways of *Candida albicans*, *A. fumigatus* and *Paracoccidioides brasiliensis*, is shown in Fig. 4 (Visbal *et al.*, 2003; Alcazar-Fuoli *et al.*, 2008; Martel *et al.*, 2010a; Weete *et al.*, 2010).

The *ΔFgCYP51B* and *ΔFgCYP51BC* mutants accumulated significant amounts of eburicol, the CYP51 substrate commonly seen in filamentous fungi (Bean *et al.*, 2009). There was no difference in eburicol content in the *ΔFgCYP51A* mutant relative to PH-1 and, in the *ΔFgCYP51C* and *ΔFgCYP51AC* mutants, the eburicol content was reduced relative to PH-1. The product of CYP51, 4,4-dimethyl ergosta-8,14,24(28)-trienol, was less abundant in all *ΔFgCYP51* mutants relative to the wild-type, with the greatest reduction detected in *ΔFgCYP51BC*. Two additional 14-methylated sterols (4,4,14-trimethyl ergosta-trienol and 4,4,14-trimethyl ergosta-dienol) were only detected in *ΔFgCYP51B* and *ΔFgCYP51BC*. These are likely to have arisen from accumulated eburicol, through the desaturation of bonds in the sterol backbone by ERG4 (C-24 reductase) and/or ERG5 (C-22 desaturase) or, possibly, these actions combined with a rearrangement of double bonds by ERG2 (Δ⁸⁻⁷ isomerase). In addition, the content of episterol was less in both *ΔFgCYP51B* and *ΔFgCYP51BC*, although higher in *ΔFgCYP51A*, *ΔFgCYP51C* and *ΔFgCYP51AC* mutants. Other 14-demethylated sterols more abundant in *ΔFgCYP51C* and *ΔFgCYP51AC* mutants relative to PH-1 included episterol, ergosta-5,7,24(28)-trienol and ergosta-

Table 6 Ergosterol biosynthesis inhibitor (EBI) sensitivity of *Fusarium graminearum* wild-type isolate PH-1 and *FgCYP51* gene deletion mutants^a

| Strain/mutant | Amphotericin B | | Terbinafine | | Fenhexamid | | Fenpropimorph | | Carbendazim | |
|-------------------|---|-----|---|-----|---|-----|---|-----|---|-----|
| | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF | EC ₅₀ (µg ml ⁻¹) | RF |
| PH-1 | 0.1034 ± 0.0125 ^{α*} | 1 | 0.3650 ± 0.0323 ^α | 1 | 2.7029 ± 0.5793 ^α | 1 | 63.9457 ± 3.7561 ^α | 1 | 0.8249 ± 0.0062 ^α | 1 |
| <i>ΔFgCYP51A</i> | 0.0915 ± 0.0131 ^α | 0.9 | 0.3808 ± 0.0749 ^α | 1.0 | 2.4775 ± 0.4657 ^α | 0.9 | 61.7000 ± 5.6303 ^α | 1.0 | 0.8213 ± 0.0173 ^α | 1.0 |
| <i>ΔFgCYP51B</i> | 0.0921 ± 0.0206 ^α | 0.9 | 0.3263 ± 0.0223 ^α | 0.9 | 3.9425 ± 0.3495 ^β | 1.5 | 55.4000 ± 3.1633 ^α | 0.9 | 0.8000 ± 0.0085 ^α | 1.0 |
| <i>ΔFgCYP51C</i> | 0.0930 ± 0.0143 ^α | 0.9 | 0.4339 ± 0.1127 ^α | 1.2 | 3.5325 ± 0.6481 ^β | 1.3 | 56.6333 ± 9.0779 ^α | 0.9 | 0.8318 ± 0.0113 ^α | 1.0 |
| <i>ΔFgCYP51AC</i> | 0.0826 ± 0.0124 ^α | 0.8 | 0.5830 ± 0.1422 ^β | 1.6 | 3.3625 ± 0.8714 ^β | 1.2 | 59.6000 ± 9.1597 ^α | 0.9 | 0.8213 ± 0.0217 ^α | 1.0 |
| <i>ΔFgCYP51BC</i> | 0.0919 ± 0.0114 ^α | 0.9 | 0.5574 ± 0.0740 ^β | 1.5 | 5.4333 ± 0.7164 ^γ | 2.0 | 69.2333 ± 16.5599 ^α | 1.1 | 0.7977 ± 0.0146 ^α | 1.0 |

^aEC₅₀ values (µg ml⁻¹) are means of EC₅₀ of two transformants from three independent replicates ± SD. Mean resistance factor (RF), representing the sensitivity change of each isolate, was calculated as the fold change in EC₅₀ compared with wild-type PH-1.

*Data were analysed using ANOVA ($P < 0.05$, F test). Different Greek letters are used to mark statistically significant differences between strains/mutants ($P < 0.05$, least-significant difference (LSD)).

Table 7 Relative composition (%) of sterols of *Fusarium graminearum* wild-type PH-1 and *FgCYP51* gene deletion mutants

| Compound | | | Relative composition (%) ^a in the different <i>ΔFgCYP51</i> mutants | | | | | |
|--|----------------|----------------------|--|------------------|---------------------|------------------|-------------------|---------------------|
| Systematic name | Common name | Retention time (min) | PH-1 | <i>ΔFgCYP51A</i> | <i>ΔFgCYP51B</i> | <i>ΔFgCYP51C</i> | <i>ΔFgCYP51AC</i> | <i>ΔFgCYP51BC</i> |
| 14-methylated sterols | | | | | | | | |
| 4,4,14-trimethyl cholesta-8,24-dienol | Lanosterol | 36.0 | 2.15 ± 0.26 | 1.99 ± 0.10 | 1.76 ± 0.17 | 1.51 ± 0.02 | 1.58 ± 0.07 | 1.43 ± 0.04 |
| 4,4,14-trimethyl ergosta-8,24-dienol | Eburicol | 38.8 | 3.46 ± 0.65 | 3.33 ± 0.39 | 12.22 ± 1.59 | 1.05 ± 0.15 | 1.24 ± 0.035 | 14.26 ± 2.78 |
| 4,4,14-trimethyl ergosta-trienol | | 37.8 | ND | ND | 0.10 ± 0.02 | ND | ND | 0.05 ± 0.09 |
| 4,4,14-trimethyl ergosta-dienol | | 38.1 | ND | ND | 0.08 ± 0.07 | ND | ND | 0.04 ± 0.06 |
| 14-demethylated sterols | | | | | | | | |
| 4,4-dimethyl ergosta-8,14,24(28)-trienol | | 39.7 | 0.51 ± 0.11 | 0.32 ± 0.02 | 0.31 ± 0.01 | 0.23 ± 0.05 | 0.25 ± 0.04 | 0.09 ± 0.16 |
| 4,4-dimethyl ergosta-8,24(28)-dienol | | 40.2 | 5.24 ± 1.32 | 3.55 ± 0.38 | 3.76 ± 0.39 | 2.86 ± 0.74 | 2.45 ± 0.10 | 1.79 ± 0.62 |
| 4-methyl ergosta-8,24-dienol | | 36.7 | 4.01 ± 0.77 | 2.04 ± 0.27 | 3.08 ± 0.32 | 3.18 ± 0.73 | 2.34 ± 0.11 | 1.18 ± 0.99 |
| Ergosta-8,24-dienol | Fecosterol | 32.8 | 0.54 ± 0.06 | 0.56 ± 0.01 | 0.35 ± 0.01 | 0.67 ± 0.14 | 0.68 ± 0.05 | 0.13 ± 0.23 |
| Ergosta-7,24(28)-dienol | Episterol | 34.5 | 1.62 ± 0.05 | 2.87 ± 0.39 | 0.93 ± 0.10 | 2.75 ± 0.72 | 2.52 ± 0.30 | 0.98 ± 0.36 |
| Ergosta-5,7,24(28)-trienol | | 33.6 | 1.53 ± 0.39 | 0.05 ± 0.09 | 1.44 ± 0.05 | 2.41 ± 0.13 | 2.09 ± 0.64 | 0.56 ± 0.48 |
| Ergosta-5,7-dienol | | 33.9 | 1.73 ± 0.13 | 1.78 ± 0.25 | 0.99 ± 0.12 | 2.41 ± 0.15 | 2.63 ± 0.24 | 1.44 ± 0.38 |
| Ergosta-5,7,22,24(28)-tetraenol | | 30.3 | 0.88 ± 0.74 | 1.44 ± 0.33 | 0.83 ± 0.23 | 0.94 ± 0.13 | 0.78 ± 0.09 | 0.56 ± 0.34 |
| Ergosta-5,8,22,24(28)-tetraenol | | 32.3 | 0.97 ± 0.06 | 1.05 ± 0.05 | 0.85 ± 0.02 | 1.15 ± 0.07 | 1.12 ± 0.04 | 0.30 ± 0.29 |
| Ergosta-5,22,24(28)-trienol | | 30.0 | 1.53 ± 0.02 | 1.61 ± 0.02 | 1.77 ± 0.01 | 1.56 ± 0.09 | 1.41 ± 0.15 | 1.47 ± 0.18 |
| Ergosta-5,22-dienol | Brassicasterol | 29.7 | 2.46 ± 0.01 | 2.43 ± 0.08 | 2.72 ± 0.18 | 3.31 ± 0.30 | 2.74 ± 0.48 | 2.57 ± 0.25 |
| Ergosta-5,7,22-trienol | Ergosterol | 31.7 | 72.98 ± 2.53 | 74.74 ± 0.80 | 68.49 ± 0.91 | 75.66 ± 2.35 | 77.45 ± 2.12 | 72.85 ± 6.00 |

^aRelative compositions of sterols are means of the percentage of the total sterol mass for the isolates from three independent replicates ± SD. ND, not detected.

Accumulated 14-methylated sterols are shown in bold.

5,7-dienol. Although the composition of sterol intermediates differed between the *ΔFgCYP51* mutants and PH-1, there were no substantial differences in ergosterol content, a finding consistent with the similar *in vitro* growth rates of the various *ΔFgCYP51* mutants and PH-1.

Expression of genes in the ergosterol biosynthesis pathway

The relative expression of genes involved in the ergosterol biosynthesis pathway was measured after 48 h of incubation in PDB (Fig. 4a). *FgCYP51A* gene expression was almost 10-fold




Fig. 4 *FgCYP51* expression in wild-type PH-1 and *FgCYP51* gene deletion mutants *in vitro* and *in planta*. (a) *FgCYP51* gene expression *in vitro*. Mycelia were collected after 48 h of incubation in potato dextrose broth (PDB) medium inoculated with 1×10^4 spores ml^{-1} . (b) *FgCYP51* gene expression *in planta*. Fourteen spikelets were inoculated on each ear and collected at 7 d post-inoculation (dpi). The relative expression levels of *FgCYP51A* (black bars), *FgCYP51B* (light grey bars) and *FgCYP51C* (dark grey bars) genes were analysed with the $2^{-\Delta\Delta C_t}$ method with *actin* as the reference gene and PH-1 as a calibrator. The error bars show + SE between three biological replicates.

higher in $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ mutants relative to PH-1. When present, the expression of *FgCYP51B* and *FgCYP51C* was unchanged in the $\Delta FgCYP51$ mutants relative to PH-1. Of the other genes involved in ergosterol biosynthesis, there were no significant differences in the expression of *ERG2*, *ERG3A* or *ERG4* in any mutants relative to PH-1 (Fig. S2). *ERG7* and *ERG27* expression was significantly higher in the $\Delta FgCYP51A$ and $\Delta FgCYP51BC$ mutants. The expression of *ERG1*, encoding squalene epoxidase, the first enzyme in ergosterol biosynthesis, was significantly lower in the single $\Delta FgCYP51A$ and $\Delta FgCYP51C$, and double $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants, which are less sensitive to terbinafine, an *erg1* inhibitor (Table 6). *ERG3B* expression, encoding a C-5 (6) desaturase, was lower in $\Delta CYP51AC$ and $\Delta CYP51BC$ mutants. There were no significant differences in *actin* or β -*tubulin* gene expression (Fig. S3).

FgCYP51C is required for full virulence on wheat ears

In wheat ear infection assays, four of the *FgCYP51* gene deletion mutants showed reduced virulence relative to wild-type PH-1 (Fig. 5). There were no significant differences ($P < 0.05$, LSD) in $\Delta FgCYP51A$ infection over the time course relative to the wild-type. This suggests that *FgCYP51A* is not essential for wheat ear infection. The percentage ear infection caused by $\Delta FgCYP51B$ was significantly less than that of the wild-type at 16 dpi ($P < 0.05$, LSD), but infection levels were similar at 21 dpi, indicating that *FgCYP51B* gene deletion delays infection. Deletion of *FgCYP51C* decreased significantly ear infection at both 16 dpi and 21 dpi ($P < 0.05$, LSD), suggesting that *FgCYP51C* is required for full virulence on wheat ears. The virulence of $\Delta FgCYP51AC$ was similar to $\Delta FgCYP51C$, supporting the assertion that *FgCYP51A* has no effect on infection. The strongest reduction in wheat ear




Fig. 5 Infection of wheat ears with wild-type PH-1 and *FgCYP51* gene deletion mutants. The 11th and 12th spikelets (shown by the black dots) from the bottom of each ear were droplet inoculated with conidia. The number of infected spikelets was recorded at 3, 7, 11, 16 and 21 d post-inoculation (dpi). Mock inoculation with distilled water was the negative control. Three wheat ears of different wheat plants were inoculated per strain in each replicate experiment. Data were analysed from three independent experiments using ANOVA ($P < 0.001$, *F* test). Greek letters show the statistically significant differences between different strains on the same day ($P < 0.05$, least-significant difference (LSD)). Error bars show \pm SE. (a) Wheat ear infection over the time course. (b) Infected wheat ears at 21 dpi.




Fig. 6 Quantification of deoxynivalenol (DON) content in wheat spikelets infected with *Fusarium graminearum* PH-1 and *FgCYP51* gene deletion mutants at 10 d post-inoculation (dpi). Fourteen spikelets of each ear were inoculated and harvested after 10 d. The concentration of DON (ppm) per inoculated dry weight was measured by an EZ-Quant Vomitoxin (DON) plate kit (Diagnostix). Mock inoculated was used as a negative control. Error bars show \pm SE between three biological replicates.

infection was measured for $\Delta FgCYP51BC$ mutants, demonstrating a strong combined effect of the absence of both *FgCYP51C* and *FgCYP51B* on virulence. In addition, wheat spikelets inoculated with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants contained five-fold less DON at 10 dpi than those inoculated with PH-1 (Fig. 7). There was no significant difference in the amount of DON in wheat ears inoculated with $\Delta FgCYP51A$ and $\Delta FgCYP51B$ (Fig. 6).

The grains in wild-type PH-1, $\Delta FgCYP51A$ and $\Delta FgCYP51B$ infected wheat ears were shrivelled and bleached (Fig. 7c). Not all the grains of ears inoculated with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ were infected, although some spikelets were bleached (Fig. 7a). Rachises infected with PH-1, $\Delta FgCYP51A$ and $\Delta FgCYP51B$ were all bleached, whereas those infected with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ were a dark brown colour (Fig. 7b). In addition, grains in $\Delta FgCYP51C$,




Fig. 8 Pathogenicity of PH-1 and *FgCYP51* gene deletion mutants on *Arabidopsis* floral tissue. The apical inflorescences of 5-wk-old *Arabidopsis* plants (ecotype Landsberg) were spray inoculated with $c. 5 \times 10^4$ *Fusarium graminearum* spores per plant. *Arabidopsis*-*Fusarium* disease (FAD) scores for each plant were recorded at 4, 8, 11 and 15 d post-inoculation (dpi). Three plants were inoculated per *F. graminearum* strain/mutant. Data were analysed from two experimental replicates using ANOVA ($P < 0.001$, F test). Greek letters show the statistically significant differences among different strains on the same day ($P < 0.05$, least-significant difference (LSD)). Error bars show \pm SE between biological replicates.

$\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ infected ears had a pink pigment. This was absent from ears infected with PH-1 and the $\Delta FgCYP51A$ and $\Delta FgCYP51B$ mutants (Fig. 7c).

Gene expression in infected wheat ears

Similar to the *in vitro* assay, *FgCYP51A* expression was highest in wheat ears inoculated with $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ (Fig. 4b). Single *FgCYP51A* and *FgCYP51C* gene deletions did not affect the expression of *FgCYP51A* and *FgCYP51B* *in planta* (Fig. 4b). The expression of *TRI* genes, involved in DON production, was measured in inoculated wheat ears (Fig. S4). Although the RQs of *TRI4*, *TRI5* and *TRI15* transcripts varied between $\Delta FgCYP51$ mutants, these differences were not significant. There were no significant differences in *ERG1*, *ERG7* and




Fig. 7 Dissection of wheat ears infected with *Fusarium graminearum* PH-1 and *FgCYP51* gene deletion mutants. The 11th and 12th spikelets from the bottom of each ear were inoculated with conidia or sterile water as a control. Inoculated ears were dissected at 21 d post-inoculation (dpi). (a) Dissection of infected wheat ears infected with PH-1 and $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants. (b) Infected rachis. (c) Infected grains.

ERG27 expression between wild-type PH-1 and $\Delta FgCYP51$ mutants in infected wheat ears (Fig. S4).

Virulence on dicotyledonous hosts: *Arabidopsis*, tomato fruit and apple fruit sections

In contrast with wheat ear infection, there were no significant differences in disease values between the wild-type and $\Delta FgCYP51A$, $\Delta FgCYP51B$ and $\Delta FgCYP51C$ on *Arabidopsis* plants, a host which does not require DON mycotoxin for colonization (Cuzick *et al.*, 2008). Both $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ were significantly less virulent than wild-type PH-1 on *Arabidopsis* (Fig. 8), consistent with the virulence of these mutants on wheat ears. In addition, 8 d after inoculation, less aerial mycelia were visible and more green siliques were present on *Arabidopsis* plants infected with $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ in comparison with PH-1 and $\Delta FgCYP51A$, $\Delta FgCYP51B$ and $\Delta FgCYP51C$ (Fig. S5). All mutants infected tomato fruits and apple fruit sections, with phenotypes similar to that of wild-type PH-1 (Figs S6, S7).

Discussion

The recent increase in genome sequence information has revealed that many fungi, particularly ascomycetes, carry more than one copy of the azole fungicide target encoding gene, *CYP51*. In the Pezizomycotina, a subphylum of the ascomycetes that includes important pathogens of animals and plants, such as *Aspergillus* spp., *M. oryzae*, *Mycosphaerella graminicola* and *Fusarium* spp., *CYP51* paralogues have been classified into three phylogenetic clades, designated A, B and C (Becher *et al.*, 2011), with the *CYP51C* paralogue only found in *Fusarium* species. The *CYP51C* clade is ubiquitous across the genus *Fusarium* and, consequently, can be used as a reliable phylogenetic marker for the identification of different species (Fernández-Ortuño *et al.*, 2010). A previous study has reported that the deletion of individual *F. graminearum* *CYP51* genes (*FgCYP51A*, *FgCYP51B* or *FgCYP51C*) has no effect on colony morphology, vegetative growth rate or ergosterol content, although conidiation is reduced in all mutants and deletion of the *FgCYP51A* and *FgCYP51C* genes increases azole sensitivity (Liu *et al.*, 2011). In this study, by heterologous expression in yeast and systematic characterization of the impact of individual (*FgCYP51A*, *FgCYP51B*, *FgCYP51C*) and double (*FgCYP51AC* and *FgCYP51BC*) gene deletions on *in vitro* growth, fungicide sensitivity, total sterol composition and virulence on wheat ears and other plants, we describe distinct roles for the *FgCYP51* paralogues of *F. graminearum*.

FgCYP51B encodes the primary sterol 14 α -demethylase and is essential for ascospore production

The three *FgCYP51* genes of *F. graminearum* isolate Fg1955 were heterologously expressed in *S. cerevisiae* strain YUG37::*erg11*, which has been used previously to assess the impact of *CYP51* mutations on *Mycosphaerella graminicola* azole sensitivity and enzyme function (Cools *et al.*, 2010, 2011) and to analyse the

role of *AfCYP51A* and *AfCYP51B* genes in *A. fumigatus* (Martel *et al.*, 2010b). *FgCYP51A* and *FgCYP51B* were able to substitute *ScCYP51* function, whereas *FgCYP51C* could not (Fig. 2). In addition, transformants expressing *FgCYP51B* grew faster than those expressing *FgCYP51A* (Table 2). This suggests that the *FgCYP51B* protein is a more effective sterol 14 α -demethylase than *FgCYP51A* in yeast.

FgCYP51 gene deletion did not impact on ergosterol content, in accordance with Liu *et al.* (2011). However, the abundance of intermediate sterols was different in all mutants relative to the wild-type. The specific accumulation of eburicol in $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ suggests that the overall eburicol demethylation activity is perturbed in mutants lacking *FgCYP51B*. This is in contrast with those deficient in *FgCYP51A* or *FgCYP51C* activity, although the product of *CYP51*, 4,4-dimethyl ergosta-8,14,24(28)-trienol, was significantly less abundant in all $\Delta FgCYP51$ mutants relative to the wild-type. These data are consistent with studies of *A. fumigatus*. Deletions of either *AfCYP51A* or *AfCYP51B* blocked C14-demethylation, but far more eburicol accumulated in the *AfCYP51B* mutant than the *AfCYP51A* mutant (Alcazar-Fuoli *et al.*, 2008). Furthermore, substrate binding studies of *AfCYP51* proteins expressed in *Escherichia coli* detected strong binding with purified *AfCYP51B* using eburicol and lanosterol, in contrast with *AfCYP51A* (Warrilow *et al.*, 2010). As a consequence of accumulated eburicol, two additional novel 14-methylated sterol intermediates (4,4,14-trimethyl ergosta-trienol and 4,4,14-trimethyl ergosta-dienol) were detected in $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ (Fig. S8). These data suggest that Pezizomycotina *CYP51B*, including *FgCYP51B*, is central to effective sterol C14-demethylation.

Ascospores forcibly ejected from mature perithecia, formed by the overwintering fungus on field debris, are the primary source of inoculum for *F. graminearum* epidemics (Parry *et al.*, 1995; Trail *et al.*, 2005). In this study, no ascospores were formed in $\Delta FgCYP51B$ and $\Delta FgCYP51BC$ mutants (Fig. 3b), although all the *FgCYP51* gene deletion mutants produced superficially normal perithecia. This finding demonstrates that *FgCYP51B* is specifically required in the development of the sexual stage, a role that cannot be fulfilled by the up-regulation of *FgCYP51A*. Similarly, the delayed colonization of wheat ears by $\Delta FgCYP51B$ suggests that *FgCYP51A* cannot fully complement *FgCYP51B* function during wheat infection.

FgCYP51A encodes an inducible sterol 14 α -demethylase that determines azole sensitivity

In the absence of fungicide treatment, the relative transcript quantities of *FgCYP51A* and *FgCYP51B* were highest at 72 h of incubation, decreasing at 96 h during growth in rich medium (PDB). This pattern of expression is coincident with rapid fungal growth, which is linear between 24 and 60 h of incubation, and into stationary phase after 72 h of incubation (data not shown). The relative transcript quantities of *FgCYP51A* changed most over this time course. Previous studies have shown an increase in *CYP51A* gene expression after azole treatment and *CYP51B* deletion *in vitro* in *F. graminearum* (Liu *et al.*, 2010; Becher *et al.*,

2011) and *M. oryzae* (Yan *et al.*, 2011). We report similar results, with *FgCYP51A* expression induced over 100-fold by azoles *in vitro*, and *c.* 10-fold by *FgCYP51B* deletion both *in vitro* and *in planta*. The enhanced transcription of *FgCYP51A* on exposure to azoles suggests that this gene is not only responsive to chemical or genetic perturbation of *FgCYP51B* activity, but also other stresses induced by fungicide treatment. For example, in *S. cerevisiae*, *ScCYP51* expression is higher during growth on glucose, in the presence of haem, under oxygen-limiting growth conditions and during exposure to anaerobic conditions (Turi & Loper, 1992).

Amino acid substitutions and *CYP51* over-expression are the most common mechanisms of resistance to azoles in filamentous ascomycetes. In fungi with multiple *CYP51s*, it is the *CYP51A* paralogue that is most commonly altered, for example in *A. fumigatus* and *P. digitatum* (Ghosoph *et al.*, 2007; Mellado *et al.*, 2007). In addition, single gene deletion has confirmed that *AfCYP51A* is involved in intrinsic azole resistance, for example to fluconazole in *A. fumigatus* (Mellado *et al.*, 2005), and azole affinity studies have shown that the *AfCYP51A* protein has lower affinity than *AfCYP51B* for a wide range of azoles (Warrilow *et al.*, 2010). In *F. graminearum*, it has been suggested previously that different azole fungicides target different *FgCYP51s* (Liu *et al.*, 2011). In this study, deletion of *FgCYP51C*, either alone or with *FgCYP51A* ($\Delta FgCYP51AC$), had no effect on azole sensitivity. Deletion of *FgCYP51B* caused an increase in sensitivity to some azoles, with sensitivities to metconazole and prochloraz particularly affected. These data conflict with those of Liu *et al.* (2011), who reported increased sensitivity to tebuconazole and prochloraz of $\Delta FgCYP51C$ mutants, with *FgCYP51B* deletion having no impact on azole sensitivity. The reason for this discrepancy is unclear, although the different methodologies used for fungicide sensitivity testing and the different origins of the strains used in gene deletion studies may have contributed. However, clearly, in this study, an interaction of azole fungicides with *FgCYP51B* is consistent with the assertion that this paralogue is the primary sterol 14 α -demethylase in *F. graminearum*. In both studies, however, the deletion of *FgCYP51A* increased the sensitivity to all azoles tested and, particularly, epoxiconazole. Considering that *F. graminearum* isolate Fg1955 is least sensitive to epoxiconazole, *FgCYP51A* expression is inducible on azole exposure and yeast transformants expressing *FgCYP51A* are least sensitive to epoxiconazole (Table 3), it can be concluded that the intrinsically lower sensitivity to some azoles in *F. graminearum* is primarily determined by *FgCYP51A*.

FgCYP51C is a novel genus-specific *CYP51* gene required for full virulence on wheat ears

In the *S. cerevisiae* heterologous expression system, *FgCYP51A* and *FgCYP51B* were able to substitute for *ScCYP51* function. By contrast, *FgCYP51C* could not complement *ScCYP51*. This suggests that *FgCYP51C* cannot function as a sterol 14 α -demethylase. This loss or diversification of function is probably caused by substitutions in conserved putative substrate recognition sites (SRs) of *FgCYP51C*. The predicted *FgCYP51* amino acid

sequences are sufficiently identical (over 40%) to be considered as members of the same P450 family (Liu *et al.*, 2011). Analysis of residues conserved in eukaryotic *CYP51s* (Fig. S9, Lepesheva & Waterman, 2011) identified two residues (N304 and T305) unique to *FgCYP51C*. Although the importance of these residues in the function of *FgCYP51C* is unknown, substitutions T315N or S316T of rat *CYP51*, equivalent to N304 and T305, caused significant reductions in lanosterol demethylase activity (Nitahara *et al.*, 2001). Deletion of *FgCYP51C* had no impact on *in vitro* fungal morphology, growth rate, conidiation and spore germination at almost all vegetative stages, perithecia production, ascospore formation or azole sensitivity. In addition, there was no difference in eburicol or ergosterol content in $\Delta FgCYP51C$ mutants. However, $\Delta FgCYP51C$ mutants had less 4,4-dimethyl ergosta-8,14,24(28)-trienol, the product of *CYP51*, and accumulated the sterol intermediates episterol, ergosta-5,7,24(28)-trienol and ergosta-5,7-dienol, products of *ERG2*, *ERG3* and *ERG4* activity, respectively (Fig. S8). This suggests that *FgCYP51C* can impact indirectly on sterol 14 α -demethylation, *ERG2*, *ERG3* and *ERG4* activity. There was no difference in *ERG2*, *ERG3A*, *ERG3B* and *ERG4* gene expression in single $\Delta FgCYP51C$ mutants relative to wild-type PH-1 *in vitro*. However, the *ERG3B* gene was expressed less in both $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants, which grew more slowly and produced less aerial mycelia on rich medium in the dark and when inoculated on wheat ears and *Arabidopsis* floral tissues. The *CYP51* gene is required for aerobic viability in *S. cerevisiae*, *C. albicans* and *C. glabrata*. In a *CYP51*-deficient mutant, aerobic growth can be restored by null mutation or deletion of *ERG3* (Bard *et al.*, 1993; Kelly *et al.*, 1993, 1995; Geber *et al.*, 1995). However, to date, there is no evidence for *ERG3*-mediated azole resistance in filamentous fungi.

After inoculation with the $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants, the number of infected spikelets per wheat ear was reduced significantly relative to inoculations with wild-type PH-1, $\Delta FgCYP51A$ and $\Delta FgCYP51B$. In addition, not all the grain in bleached spikelets had a rough, shrivelled appearance, although infected grains were pink in wheat ears inoculated with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$, rather than the grey in plants inoculated with PH-1, $\Delta FgCYP51A$ and $\Delta FgCYP51B$ (Fig. 7c). The rachises of wheat heads infected with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ were dark brown, in contrast with the bleached rachises of plants inoculated with PH-1, $\Delta FgCYP51A$ and $\Delta FgCYP51B$. The blocking of fungal growth from inoculated spikelets to adjacent spikelets is correlated with an unidentified brown substance deposited in the rachis node in the additional wheat line CS-7EL, which carries resistance to FHB on the long arm of chromosome 7E(7EL) (Miller *et al.*, 2011). These data suggest that hyphal development of $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ during wheat ear infection is impaired.

In contrast with wheat ear infection, there were no differences in virulence between wild-type PH-1 and $\Delta FgCYP51C$ on *Arabidopsis*, in which the trichothecene mycotoxin DON is not required for fungal infection (Cuzick *et al.*, 2008), although colonization by the double $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants was impaired. This suggests that altered DON production is

responsible for the decreased virulence of the $\Delta FgCYP51C$ mutant on wheat ears. However, unlike mutants unable to produce DON, for example *TRI5* gene mutants, which cause only discrete eye-shaped lesions on spikelets and fail to infect the rachis (Cuzick *et al.*, 2008), $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ mutants are able to infect beyond the inoculated spikelet. In addition, *TRI4* and *TRI5* gene expression is not altered significantly in wheat ears inoculated with $\Delta FgCYP51C$, $\Delta FgCYP51AC$ and $\Delta FgCYP51BC$ relative to the wild-type. Therefore, rather than a biosynthetic requirement for DON biosynthesis, FgCYP51C is likely to have an indirect effect. A relationship between the sterol and trichothecene biosynthesis pathways has been reported previously. They share a common precursor, farnesol pyrophosphate, and the global regulator *TRI6*, located in the core *TRI* gene cluster. For example *ERG9* (FGSG_09381), encoding squalene synthase, the first step of sterol biosynthesis, was down-regulated in the $\Delta TRI6$ strain under nitrogen-deprived conditions, and *ERG25* (FGSG_10666) was up-regulated, although there was no impact on *FgCYP51C* expression (Nasmith *et al.*, 2011).

Conclusion

We have identified distinct functions of the three *CYP51* paralogues of *F. graminearum*. *FgCYP51B*, as the most conserved *CYP51* gene in all fungi, encodes the enzyme primarily responsible for sterol 14 α -demethylation, a role essential for ascospore formation. *FgCYP51A*, found in many human and agricultural pathogens, is induced by azoles and environmental stress, encodes a sterol 14 α -demethylase with the capacity to compensate for disruption of *FgCYP51B* function, and is responsible for intrinsic variation in sensitivity to different azoles. *FgCYP51C*, a *Fusarium*-specific *CYP51* gene, no longer functions as a sterol 14 α -demethylase, but rather is specifically required for full virulence on host wheat ears. This is the first example of functional diversification of a fungal *CYP51* gene.

Acknowledgements

The authors thank Nichola Hawkins and John Lucas (Rothamsted Research) for comments on the manuscript, and Mogens Nicolaisen (Aarhus University, Denmark) for supplying the isolate Fg1955. At Rothamsted Research, all experiments involving transgenic strains of *F. graminearum* were conducted in biological containment facilities under FERA licence number PHSI 181/6695 and with appropriate approval from the UK Health and Safety Executive. Jieru Fan was supported by an award of a Postgraduate Study Abroad Program Sponsored by the China Scholarship Council. Rothamsted Research receives grant-aided support from the Biotechnology and Biological Sciences Research Council (BBSRC) of the UK.

References

Alcazar-Fuoli L, Mellado E, Garcia-Effron G, Lopez JF, Grimalt JO, Cuenca-Estrella JM, Rodriguez-Tudela JL. 2008. Ergosterol biosynthesis pathway in *Aspergillus fumigatus*. *Steroids* 73: 339–347.

- Audenaert K, Callewaert E, Hofte M, De Saeger S, Haesaert G. 2010. Hydrogen peroxide induced by the fungicide prothioconazole triggers deoxynivalenol (DON) production by *Fusarium graminearum*. *BMC Microbiology* 10: 112.
- Bard M, Lees ND, Turi T, Craft D, Cofrin L, Barbuch R, Koegel C, Loper JC. 1993. Sterol synthesis and viability of *Erg11* (cytochrome-P450 lanosterol demethylase) mutations in *Saccharomyces cerevisiae* and *Candida albicans*. *Lipids* 28: 963–967.
- Bean TP, Cools HJ, Lucas JA, Hawkins ND, Ward JL, Shaw MW, Fraaije BA. 2009. Sterol content analysis suggests altered eburicol 14 α -demethylase (CYP51) activity in isolates of *Mycosphaerella graminicola* adapted to azole fungicides. *FEMS Microbiology Letters* 296: 266–273.
- Becher R, Weihmann F, Deising H, Wirsal S. 2011. Development of a novel multiplex DNA microarray for *Fusarium graminearum* and analysis of azole fungicide responses. *BMC Genomics* 12: 52.
- Beyer M, Klis MB, Klink H, Verreet JA. 2006. Quantifying the effects of previous crop, tillage, cultivar and triazole fungicides on the deoxynivalenol content of wheat grain – a review. *Journal of Plant Diseases and Protection* 113: 241–246.
- Catlett NL, Lee B-N, Yoder OC, Turgeon BG. 2003. Split-marker recombination for efficient targeted deletion of fungal genes. *Fungal Genetics Newsletter* 50: 9–11.
- Cools HJ, Bayon C, Atkins S, Lucas JA, Fraaije BA. 2012. Overexpression of the sterol 14 alpha-demethylase gene (*MgCYP51*) in *Mycosphaerella graminicola* isolates confers a novel azole fungicide sensitivity phenotype. *Pest Management Science* 68: 1034–1040.
- Cools HJ, Mullins JGL, Fraaije BA, Parker JE, Kelly DE, Lucas JA, Kelly SL. 2011. Impact of recently emerged sterol 14 α -demethylase (CYP51) variants of *Mycosphaerella graminicola* on azole fungicide sensitivity. *Applied and Environmental Microbiology* 77: 3830–3837.
- Cools HJ, Parker JE, Kelly DE, Lucas JA, Fraaije BA, Kelly SL. 2010. Heterologous expression of mutated eburicol 14 α -demethylase (CYP51) proteins of *Mycosphaerella graminicola* to assess effects on azole fungicide sensitivity and intrinsic protein function. *Applied and Environmental Microbiology* 76: 2866–2872.
- Cuomo CA, Gueldener U, Xu JR, Trail F, Turgeon BG, Di Pietro A, Walton JD, Ma LJ, Baker SE, Rep M *et al.* 2007. The *Fusarium graminearum* genome reveals a link between localized polymorphism and pathogen specialization. *Science* 317: 1400–1402.
- Cuzick A, Urban M, Hammond-Kosack K. 2008. *Fusarium graminearum* gene deletion mutants *map1* and *tri5* reveal similarities and differences in the pathogenicity requirements to cause disease on *Arabidopsis* and wheat floral tissue. *New Phytologist* 177: 990–1000.
- Diaz-Guerra TM, Mellado E, Cuenca-Estrella M, Rodriguez-Tudela JL. 2003. A point mutation in the 14 α -sterol demethylase gene *cyp51A* contributes to itraconazole resistance in *Aspergillus fumigatus*. *Antimicrobial Agents and Chemotherapy* 47: 1120–1124.
- Ehrenshaft M, Jenns AE, Daub ME. 1995. Targeted gene disruption of carotenoid biosynthesis in *Cercospora nicotianae* reveals no role for carotenoids in photosensitizer resistance. *Molecular Plant-Microbe Interactions* 8: 593–601.
- Fernández-Ortuño D, Loza-Reyes E, Atkins SL, Fraaije BA. 2010. The *CYP51C* gene, a reliable marker to resolve interspecific phylogenetic relationships within the *Fusarium* species complex and a novel target for species-specific PCR. *International Journal of Food Microbiology* 144: 301–309.
- Fraaije BA, Cools HJ, Kim SH, Motteram J, Clark WS, Lucas JA. 2007. A novel substitution I381V in the sterol 14 α -demethylase (CYP51) of *Mycosphaerella graminicola* is differentially selected by azole fungicides. *Molecular Plant Pathology* 8: 245–254.
- Geber A, Hitchcock CA, Swartz JE, Pullen FS, Marsden KE, Kwon-Chung KJ, Bennett JE. 1995. Deletion of the *Candida glabrata* *ERG3* and *ERG11* genes: effect on cell viability, cell growth, sterol composition, and antifungal susceptibility. *Antimicrobial Agents and Chemotherapy* 39: 2708–2717.
- Ghosh JM, Schmidt LS, Margosan DA, Smilanick JL. 2007. Imazalil resistance linked to a unique insertion sequence in the *PdCYP51* promoter region of *Penicillium digitatum*. *Postharvest Biology and Technology* 44: 9–18.

- Gray KC, Palacios DS, Dailey I, Endo MM, Uno BE, Wilcock BC, Burke MD. 2012. Amphotericin primarily kills yeast by simply binding ergosterol. *Proceedings of the National Academy of Sciences, USA* 109: 2234–2239.
- Hamamoto H, Hasegawa K, Nakaune R, Lee YJ, Makizumi Y, Akutsu K, Hibi T. 2000. Tandem repeat of a transcriptional enhancer upstream of the sterol 14 α -demethylase gene (*CYP51*) in *Penicillium digitatum*. *Applied and Environmental Microbiology* 66: 3421–3426.
- Inagaki YS, Etherington G, Geisler K, Field B, Dokarry M, Ikeda K, Mutsukado Y, Dicks J, Osbourn A. 2011. Investigation of the potential for triterpene synthesis in rice through genome mining and metabolic engineering. *New Phytologist* 191: 432–448.
- Kelly SL, Arnoldi A, Kelly DE. 1993. Molecular-genetic analysis of azole antifungal mode of action. *Biochemical Society Transactions* 21: 1034–1038.
- Kelly SL, Lamb DC, Corran AJ, Baldwin BC, Kelly DE. 1995. Mode of action and resistance to azole antifungals associated with the formation of 14 α -methylergosta-8,24(28)-dien-3 β ,6 α -diol. *Biochemical and Biophysical Research Communications* 207: 910–915.
- Kretschmer M, Leroch M, Mosbach A, Walker AS, Fillingner S, Mernke D, Schoonbeek HJ, Pradier JB, Leroux P, De Waard MA *et al.* 2009. Fungicide-driven evolution and molecular basis of multidrug resistance in field populations of the grey mould fungus *Botrytis cinerea*. *PLoS Pathogens* 5: e1000696.
- Kulik T, Lojko M, Jestoi M, Perkowski J. 2012. Sublethal concentrations of azoles induce *tri* transcript levels and trichothecene production in *Fusarium graminearum*. *FEMS Microbiology Letters* 335: 58–67.
- Kunii M, Kitahama Y, Fukushima EO, Seki H, Muranaka T, Yoshida Y, Aoyama Y. 2012. β -Amyrin oxidation by oat *CYP51H10* expressed heterologously in yeast cells: the first example of CYP51-dependent metabolism other than the 14-demethylation of sterol precursors. *Biological and Pharmaceutical Bulletin* 35: 801–804.
- Lepesheva GI, Waterman MR. 2007. Sterol 14 α -demethylase cytochrome P450 (*CYP51*), a P450 in all biological kingdoms. *Biochimica et Biophysica Acta (BBA) – General Subjects* 1770: 467–477.
- Lepesheva GI, Waterman MR. 2011. Structural basis for conservation in the *CYP51* family. *Biochimica et Biophysica Acta (BBA) – Proteins and Proteomics* 1814: 88–93.
- Liu X, Jiang JH, Shao JF, Yin YN, Ma ZH. 2010. Gene transcription profiling of *Fusarium graminearum* treated with an azole fungicide tebuconazole. *Applied Microbiology and Biotechnology* 85: 1105–1114.
- Liu X, Yu F, Schnabel G, Wu J, Wang Z, Ma Z. 2011. Paralogous *cyp51* genes in *Fusarium graminearum* mediate differential sensitivity to sterol demethylation inhibitors. *Fungal Genetics and Biology* 48: 113–123.
- Loffler J, Kelly SL, Hebart H, Schumacher U, LassFlörl C, Einsele H. 1997. Molecular analysis of *cyp51* from fluconazole-resistant *Candida albicans* strains. *FEMS Microbiology Letters* 151: 263–268.
- Martel CM, Parker JE, Bader O, Weig M, Gross U, Warrilow AGS, Rolley N, Kelly DE, Kelly SL. 2010a. Identification and characterization of four azole-resistant *erg3* mutants of *Candida albicans*. *Antimicrobial Agents and Chemotherapy* 54: 4527–4533.
- Martel CM, Parker JE, Warrilow AGS, Rolley NJ, Kelly SL, Kelly DE. 2010b. Complementation of a *Saccharomyces cerevisiae* ERG11/*CYP51* (sterol 14 α -demethylase) doxycycline-regulated mutant and screening of the azole sensitivity of *Aspergillus fumigatus* isoenzymes CYP51A and CYP51B. *Antimicrobial Agents and Chemotherapy* 54: 4920–4923.
- Mellado E, Garcia-Effron G, Alcázar-Fuoli L, Melchers WJG, Verweij PE, Cuenca-Estrella M, Rodríguez-Tudela JL. 2007. A new *Aspergillus fumigatus* resistance mechanism conferring *in vitro* cross-resistance to azole antifungals involves a combination of *cyp51A* alterations. *Antimicrobial Agents and Chemotherapy* 51: 1897–1904.
- Mellado E, Garcia-Effron G, Buitrago MJ, Alcázar-Fuoli L, Cuenca-Estrella M, Rodríguez-Tudela JL. 2005. Targeted gene disruption of the 14- α sterol demethylase (*cyp51a*) in *Aspergillus fumigatus* and its role in azole drug susceptibility. *Antimicrobial Agents and Chemotherapy* 49: 2536–2538.
- Miller SS, Watson EM, Lazebnik J, Gulden S, Balcerzak M, Fedak G, Ouellet T. 2011. Characterization of an alien source of resistance to Fusarium head blight transferred to Chinese Spring wheat 11. *Botany-Botanique* 89: 301–311.
- Nasmith CG, Walkowiak S, Wang L, Leung WWY, Gong YC, Johnston A, Harris LJ, Guttman DS, Subramaniam R. 2011. *Tri6* is a global transcription regulator in the phytopathogen *Fusarium graminearum*. *PLoS Pathogens* 7: e1002266.
- Nicolaisen M, Supronienė S, Nielsen LK, Lazzaro I, Spliid NH, Justesen AF. 2009. Real-time PCR for quantification of eleven individual *Fusarium* species in cereals. *Journal of Microbiological Methods* 76: 234–240.
- Nitahara Y, Kishimoto K, Yabusaki Y, Gotoh O, Yoshida Y, Horiuchi T, Aoyama Y. 2001. The amino acid residues affecting the activity and azole susceptibility of rat CYP51 (sterol 14-demethylase p450). *Journal of Biochemistry* 129: 761–768.
- Nucci M, Anaissie E. 2007. Fusarium infections in immunocompromised patients. *Clinical Microbiology Reviews* 20: 695–704.
- Ochiai N, Tokai T, Takahashi-Ando N, Fujimura M, Kimura M. 2007. Genetically engineered *Fusarium* as a tool to evaluate the effects of environmental factors on initiation of trichothecene biosynthesis. *FEMS Microbiology Letters* 275: 53–61.
- Parks L, Smith S, Crowley J. 1995. Biochemical and physiological effects of sterol alterations in yeast – a review. *Lipids* 30: 227–230.
- Parry DW, Jenkinson P, McLeod L. 1995. Fusarium ear blight (scab) in small-grain cereals – a review. *Plant Pathology* 44: 207–238.
- Paul PA, Lipps PE, Hershman DE, McMullen MP, Draper MA, Madden LV. 2008. Efficacy of triazole-based fungicides for Fusarium head blight and deoxynivalenol control in wheat: a multivariate meta-analysis. *Phytopathology* 98: 999–1011.
- Perea S, Lopez-Ribot JL, Kirkpatrick WR, McAtee RK, Santillan RA, Martinez M, Calabrese D, Sanglard D, Patterson TF. 2001. Prevalence of molecular mechanisms of resistance to azole antifungal agents in *Candida albicans* strains displaying high-level fluconazole resistance isolated from human immunodeficiency virus-infected patients. *Antimicrobial Agents and Chemotherapy* 45: 2676–2684.
- Qi X, Bakht S, Qin B, Leggett M, Hemmings A, Mellon F, Eagles J, Werck-Reichhart D, Schaller H, Lesot A *et al.* 2006. A different function for a member of an ancient and highly conserved cytochrome P450 family: from essential sterols to plant defense. *Proceedings of the National Academy of Sciences, USA* 103: 18848–18853.
- Revankar SG, Fu J, Rinaldi MG, Kelly SL, Kelly DE, Lamb DC, Keller SM, Wickes BL. 2004. Cloning and characterization of the lanosterol 14 α -demethylase (*ERG11*) gene in *Cryptococcus neoformans*. *Biochemical and Biophysical Research Communications* 324: 719–728.
- Rodríguez RJ, Low C, Bottema CDK, Parks LW. 1985. Multiple functions for sterols in *Saccharomyces cerevisiae*. *Biochimica et Biophysica Acta (BBA) – Lipids and Lipid Metabolism* 837: 336–343.
- Rozman D, Strömstedt M, Waterman MR. 1996. The three human cytochrome P450 lanosterol 14 α -demethylase (*CYP51*) genes reside on chromosomes 3, 7, and 13: structure of the two retrotransposed pseudogenes, association with a line-1 element, and evolution of the human *CYP51* family. *Archives of Biochemistry and Biophysics* 333: 466–474.
- Sanglard D, Ischer F, Koymans L, Bille J. 1998. Amino acid substitutions in the cytochrome P-450 lanosterol 14 α -demethylase (*CYP51A1*) from azole-resistant *Candida albicans* clinical isolates contribute to resistance to azole antifungal agents. *Antimicrobial Agents and Chemotherapy* 42: 241–253.
- Sanglard D, Kuchler K, Ischer F, Pagani JL, Monod M, Bille J. 1995. Mechanisms of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients involve specific multidrug transporters. *Antimicrobial Agents and Chemotherapy* 39: 2378–2386.
- Sheehan DJ, Hitchcock CA, Sibley CM. 1999. Current and emerging azole antifungal agents. *Clinical Microbiology Reviews* 12: 40–79.
- Snelders E, van der Lee HAL, Kuijpers J, Rijs AJMM, Varga J, Samson RA, Mellado E, Donders ART, Melchers WJG, Verweij PE. 2008. Emergence of azole resistance in *Aspergillus fumigatus* and spread of a single resistance mechanism. *PLoS Medicine* 5: e219.
- Trail F, Gaffoor I, Vogel S. 2005. Ejection mechanics and trajectory of the ascospores of *Gibberella zeae* (anamorph *Fusarium graminearum*). *Fungal Genetics and Biology* 42: 528–533.
- Turi TG, Loper JC. 1992. Multiple regulatory elements control expression of the gene encoding the *Saccharomyces cerevisiae* cytochrome-P450, lanosterol-

- 14- α -demethylase (Erg11). *Journal of Biological Chemistry* 267: 2046–2056.
- Urban M, Daniels S, Mott E, Hammond-Kosack K. 2002. *Arabidopsis* is susceptible to the cereal ear blight fungal pathogens *Fusarium graminearum* and *Fusarium culmorum*. *Plant Journal* 32: 961–973.
- Urban M, Mott E, Farley T, Hammond-Kosack K. 2003. The *Fusarium graminearum* MAP1 gene is essential for pathogenicity and development of perithecia. *Molecular Plant Pathology* 4: 347–359.
- Visbal G, Alvarez A, Moreno B, San-Blas G. 2003. S-adenosyl-L-methionine inhibitors Delta(24)-sterol methyltransferase and delta(24(28))-sterol methylreductase as possible agents against *Paracoccidioides brasiliensis*. *Antimicrobial Agents and Chemotherapy* 47: 2966–2970.
- Warrilow AGS, Melo N, Martel CM, Parker JE, Nes WD, Kelly SL, Kelly DE. 2010. Expression, purification, and characterization of *Aspergillus fumigatus* sterol 14- α demethylase (CYP51) isoenzymes A and B. *Antimicrobial Agents and Chemotherapy* 54: 4225–4234.
- Weete JD, Abril M, Blackwell M. 2010. Phylogenetic distribution of fungal sterols. *PLoS ONE* 5: e10899.
- Wyand RA, Brown JKM. 2005. Sequence variation in the CYP51 gene of *Blumeria graminis* associated with resistance to sterol demethylase inhibiting fungicides. *Fungal Genetics and Biology* 42: 726–735.
- Yan X, Ma W-B, Li Y, Wang H, Que Y-W, Ma Z-H, Talbot NJ, Wang Z-Y. 2011. A sterol 14 α -demethylase is required for conidiation, virulence and for mediating sensitivity to sterol demethylation inhibitors by the rice blast fungus *Magnaporthe oryzae*. *Fungal Genetics and Biology* 48: 144–153.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 The split-marker deletion strategy of *FgCYP51* genes in *Fusarium graminearum* isolate PH-1.

Fig. S2 *In vitro* transcript levels of *ERG* genes.

Fig. S3 Relative quantification of *actin* and β -*tubulin* gene expression *in vitro*.

Fig. S4 Relative quantification of gene expression in infected wheat ears.

Fig. S5 Infection of *Arabidopsis* floral tissue with PH-1 and *FgCYP51* gene deletion mutants.

Fig. S6 Pathogenicity of PH-1 and *FgCYP51* gene deletion mutants on apple sections.

Fig. S7 Infection of tomato fruits with PH-1 and *FgCYP51* gene deletion mutants.

Fig. S8 The proposed sterol biosynthesis pathway in *Fusarium graminearum*. The sterol pathway based on compounds detected in the wild-type PH-1 and *FgCYP51* gene deletion mutants incubated in PDB media for 48 h. The proposed pathway in wild-type PH-1 is shown in solid arrows. The pathway after *FgCYP51B* deletion is shown with dashed arrows in the dashed box. The two additional sterols, 4,4,14-trimethyl ergosta-trienol and 4,4,14-trimethyl ergosta-dienol, in the dashed box were identified as TMS esters with *m/z* 510 and *m/z* 512.

Fig. S9 Alignment of predicted *FgCYP51* amino acid sequences.

Table S1 Primers used in this study

Table S2 Concentrations of fungicides used in azole sensitivity testing

Table S3 The *Fusarium*–*Arabidopsis* disease scoring system

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



About New Phytologist

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <25 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@ornl.gov)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**