



(11) **EP 2 388 332 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
23.11.2011 Bulletin 2011/47

(51) Int Cl.:
C12N 15/82 (2006.01) **C12N 9/02** (2006.01)
A01H 5/00 (2006.01) **C12N 5/10** (2006.01)
C12P 7/64 (2006.01)

(21) Application number: **10187573.0**

(22) Date of filing: **23.11.2006**

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR

- **Syanova, Olga**
Redbourn, St. Albans AL3 7HF (GB)
- **Zank, Thorsten**
68165 Mannheim (DE)

(30) Priority: **24.11.2005 GB 0523915**
30.11.2005 GB 0524420
19.06.2006 GB 0612109

(74) Representative: **Dick, Alexander**
Herzog Fiesser & Partner
Patentanwälte
Bad Kreuznacher Strasse 27-29
68309 Mannheim (DE)

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
06124663.3 / 1 790 731

Remarks:

This application was filed on 01-10-2010 as a divisional application to the application mentioned under INID code 62.

(71) Applicant: **BASF Plant Science GmbH**
67056 Ludwigshafen (DE)

(72) Inventors:
• **Napier, Johnathan A.**
Preston, Hertfordshire SG4 7TR (GB)

(54) **Process for the production of delta5-unsaturated fatty acids in transgenic organisms**

(57) A process for the production of Δ^5 -unsaturated fatty acids in transgenic organisms comprises transforming an organism with nucleic acid encoding a Δ^5 -desaturase.

EP 2 388 332 A1

Description

[0001] The present invention relates to a process for the production of $\Delta 5$ -unsaturated fatty acids in an organism by introducing, into the organism, nucleic acids which encode polypeptides with $\Delta 5$ -desaturase activity. These nucleic acid sequences, if appropriate together with further nucleic acid sequences which encode polypeptides of the biosynthesis of the fatty acid or lipid metabolism, can advantageously be expressed in the organism. The process can be used for the production of non-methylene-interrupted fatty acids (NMIFAs) such as sciadonic acid ($\Delta 5_{cis}$ -11,14-20:3) and juniperonic acid ($\Delta 5_{cis}$ -11,14,17-20:4), which do not occur naturally in most food plants. The invention furthermore relates to a process for the production of oils and/or triacylglycerides with an elevated content of long-chain polyunsaturated fatty acids.

[0002] The invention furthermore relates to the nucleic acid sequences, gene constructs, vectors and organisms comprising the nucleic acid sequences according to the invention, vectors comprising the nucleic acid sequences and/or the gene constructs and to transgenic organisms comprising the abovementioned nucleic acid sequences, gene constructs and/or vectors.

[0003] A further part of the invention relates to oils, lipids and/or fatty acids produced by the process according to the invention and to their use. Moreover, the invention relates to $\Delta 5$ -unsaturated fatty acids and to triglycerides with an elevated content of $\Delta 5$ -unsaturated fatty acids and to their use.

[0004] Non-methylene-interrupted fatty acids (NMIFAs) have been found in algae, slime molds, marine sponges and plants. Seed plants are botanically divided into gymnosperms and angiosperms. It is believed that the angiosperms have evolved from their gymnospermous predecessors in the late Jurassic or early Cretaceous. The seed oils of nearly all gymnosperms contain $\Delta 5$ -NMIFAs, usually in the form of C16-20 monounsaturates (see, for example, Wolff et al, *Lipids* 1999, Oct; 34(10); 1083-1097). Similar $\Delta 5$ -NMIFAs occur in the seed oils of a very few angiosperm species, predominantly in the plant family Ranunculaceae but also found in some other "primitive" plant families, such as Menispermaceae and Limnaceae. In angiosperms, the unusual NMIFAs are invariably found only in the seed oils and do not occur in vegetative tissue; this is in contrast to the gymnosperms, where the presence of these fatty acids in leaves is well-documented.

[0005] Recent studies have shown biomedical benefits of $\Delta 5_{cis}$ -NMIFA-containing oils, such as triglyceride-lowering effect and anti-inflammatory properties (Asset et al, 1999, *Lipids*, 34, 39-44; Berger et al, *Lipids in Health and Disease* 2002, 1:5, and references therein). Oils containing $\Delta 5_{cis}$ -11,14-20:3 sciadonic acid (SA) can also lower plasma cholesterol in experimental animals. Several investigators have shown that the addition of sciadonic acid to animals or cultured cells is effective in reducing the level of 20:4n-6 (arachidonic acid, ARA) in phosphatidylinositol (PI), thereby modifying cellular ARA metabolism (Tanaka et al, 1999, *Biochem. Biophys. Res. Comm.*, 264, 683-688; and Tanaka et al, 2001, *Eur. J. Biochem.*, 268, 4928-4939).

[0006] It would therefore be advantageous to be able to produce $\Delta 5_{cis}$ -NMIFAs such as sciadonic acid and juniperonic acid in other organisms from where they can easily be extracted. One example of this is the production of the fatty acids in the vegetative tissues of angiospermous plants.

[0007] It is known that polyunsaturated fatty acids such as arachidonic acid, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) can be produced in a variety of ways. Thus, EPA or DHA are produced in marine bacteria such as *Vibrio* sp. or *Shewanella* sp. via the polyketide pathway (Yu, R. et al. *Lipids* 35:1061-1064, 2000; Takeyama, H. et al. *Microbiology* 143:2725-2731, 1997).

[0008] An alternative strategy is the alternating activity of desaturases and elongases (Zank, T.K. et al. *Plant Journal* 31:255-268, 2002; Sakuradani, E. et al. *Gene* 238:445-453, 1999). A modification of the above-described pathway by $\Delta 6$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase, $\Delta 5$ -elongase and $\Delta 4$ -desaturase is the Sprecher pathway (Sprecher 2000, *Biochim. Biophys. Acta* 1486:219-231) in mammals. Instead of the $\Delta 4$ -desaturation, a further elongation step is effected here to give C_{24} , followed by a further $\Delta 6$ -desaturation and finally β -oxidation to give the C_{22} chain length. Thus what is known as Sprecher pathway (see figure 1) is, however, not suitable for the production in plants and microorganisms since the regulatory mechanisms are not known.

[0009] Our co-pending UK patent application No. 0523915.7 as well as UK patent application No. 0524420.7 teach the production of polyunsaturated fatty acids in a variety of plants and microorganism by means of genetically modifying these species with a selection of nucleic acid sequences encoding a series of elongase and desaturase enzymes.

[0010] The present inventors have decided to adopt a similar strategy in order to produce $\Delta 5$ -NMIFAs. In order to do this, it was necessary to identify a desaturase capable of introducing a C=C double bond at the $\Delta 5$ -position.

[0011] The biosynthesis of higher plant monounsaturated very long chain fatty acids (VLCFAs) containing a $\Delta 5$ double bond was studied previously by radio labeling of developing *Limnanthes alba* seeds. The seed oil of *Limnanthes* spp is enriched in the $\Delta 5$ -eicosaenoic acid (20:1 $\Delta 5$). Based on in vitro analysis of *Limnanthes* spp. seed extracts, Moreau et al, *Arch Biochem Biophys.* 1981; 209:376-384 suggested that 20:0-CoA is the substrate for the $\Delta 5$ -desaturase. In more recent studies, random sequencing of EST library from *L. douglasii* seeds resulted in the identification of a candidate cDNA for a $\Delta 5$ -desaturase that showed similarity to known acyl-CoA desaturases from animals, yeast and cyanobacteria

EP 2 388 332 A1

(Cahoon et al, 2000, Plant Physiology, 124, 243-251). Co-expression of this desaturase cDNA with an FAE1 (fatty acid elongating activity) homolog from *L. douglasii* in soybean somatic embryos resulted in the accumulation of Δ^5 -monounsaturated 16:1, 18:1 and 20:1 fatty acids, thus confirming the pathway proposed for 20:1 Δ^5 formation in *Limnanthes* spp:

- a) a large flux of palmitic acid (16:0) from the plastid to the endoplasmic reticulum;
- b) microsomal elongation of 16:0, presumably as a CoA ester, to eicosanoic acid (20:0),
- c) Δ^5 -desaturation of 20:0 to form 20:1 Δ^5 (see Cahoon et al, Plant Physiol. 2000 Sep;124(1):243-51 and references therein).

[0012] However, no direct biochemical evidence has been presented to support the assertion that the *Limnanthes* desaturases utilizes acyl-CoA substrates, as opposed to the predominant glycerolipid or ACP-linked desaturation occurring in plants.

[0013] Equally, cDNAs encoding proteins related to the animal and yeast acyl-CoA desaturases (abbreviated to ADS) have been identified in several plant species, though their activity towards acyl-CoA substrates is inferred only from homology (Fukuchi-Mizutani et al, 1998, Plant and Cell Physiology 39: 247-253).


[0014] Two cytoplasmic ADS enzymes from *Arabidopsis*, ADS1 and ADS2, formed Δ^9 -desaturated fatty acids when expressed in yeast (Heilmann et al. 2004, PNAS, 101 (28), 10266-10271), *Arabidopsis* or in *Brassica juncea* (ADS1 only, Yao et al, 2003 Plant Biotechnol 1: 221-229). ADS3, another member of the *Arabidopsis* desaturase (ADS) multigene family, was characterized as a gene, encoding palmitoyl-monogalactosyldiacylglycerol Δ^7 -desaturase. A conifera white spruce (*Picea glauca*) homolog of ADS3 has been characterized as an 18:0 Δ^9 desaturase gene by heterologous expression in yeast (Marillia et al., 2002, FEBS Lett, 526: 49-52).

[0015] Δ^5 cis-C20 NMIFAs occur in seed and leaves in all Coniferae (see, for example, Destailats et al, Lipids, 2001 Nov; 36(11); 1247-1254 and Wolff et al, Lipids. 2001 May; 36(5); 439-451). It was also found that seed oil of several members of the rather primitive angiosperm family Ranunculaceae contains significant amounts of these unusual fatty acids (see Wolff et al. 2002 for review: Lipids. 2002 Jan; 37(1):17-26 and see also Tsevegsuren et al, (1997) Hrc-Journal of High Resolution Chromatography, 20: 237-241).


[0016] *Cimicifuga* spp., *Caltha palustris* L. and *Anemone leveillei/rivularis* are all reported to produce cis Δ^5 desaturated fatty acids, including 16:1 Δ^5 and 18:1 Δ^5 and (predominantly) 20:3 Δ^5 ,11,14 (sciadonic acid). *C. palustris* produces additional Δ^5 desaturation products such as 18:4 Δ^5 ,9,12,15 and 20:4,11,14,17 (juniperonic acid). In *Caltha*, *Cimicifuga* and *A. leveillei* the main Δ^5 cis-C20 NMIFA is sciadonic acid (SA, 20:3, Δ^5 ,11,14). It is found at levels of 19.7% of total fatty acids in seed oils of *C. palustris* and of 9.8% of total fatty acids in *Anemone* seed oils. *C. palustris* produces additional Δ^5 desaturation products such as 18:4 Δ^5 ,9,12,15, 20:1 Δ^5 and 20:4 Δ^5 ,11,14,17 (Smith et al, 1968, Lipids 3: 37-&). We have previously attempted to identify the Δ^5 -desaturase responsible for the synthesis of SA from *Anemone*, and initially considered it likely that this enzyme was a cytochrome b5 fusion desaturase. Although *Anemone* contains several such desaturases, none of them were identified as being involved in SA synthesis (Whitney et al. 2003, Planta, 217(6), 983-992)

[0017] The present inventors therefore hypothesized that the enzyme introducing a double bond at the Δ^5 position to yield SA might be an acyl-CoA desaturase similar to the group of membrane bound fatty acid desaturases recently cloned from *Arabidopsis*, rose and *Limnanthes* and following this reasoning, have obtained nucleotide sequences which encode a Δ^5 -desaturase enzyme. This has made possible the development of a process for the production of Δ^5 -unsaturated NMIFAs in a variety of organisms.

[0018] Therefore, in a first aspect of the present invention, there is provided a process for the production of a compound of formula (I):



in an organism, the process comprising introducing into an organism which comprises a lipid of general formula (III):




at least one nucleic acid sequence comprising :

- a) SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5;
 b) a nucleic acid sequence which hybridizes under stringent conditions with a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.
 c) a nucleic acid sequence which encodes a polypeptide of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6;
 d) a derivative of a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, wherein said polypeptide has $\Delta 5$ -desaturase activity;


and expressing said nucleic acid sequence;
 wherein, in general formulae I and III:

Y = a C₁₀-C₁₈ hydrocarbon chain containing up to four carbon-carbon double bonds;
 R¹ = hydroxyl, coenzyme A (thioester), lysophosphatidylcholine, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysodiphosphatidylglycerol, lysophosphatidylserine, lysophosphatidylinositol, sphingo base or a radical of the formula II



in which

R² = hydrogen, lysophosphatidyl choline, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysodiphosphatidylglycerol, lysophosphatidylserine, lysophosphatidylinositol or saturated or unsaturated C₂-C₂₄-alkylcarbonyl,
 R³ = hydrogen, saturated or unsaturated C₂-C₂₄-alkylcarbonyl, or R² and R³ independently of one another are a radical of the formula Ia:



in which

n = 2, 3, 4, 5, 6, 7 or 9, m = 2, 3, 4, 5 or 6 and p = 0 or 3;
 and wherein an oxygen in the R¹ radical may be replaced by sulphur such that R¹ is bonded to the remainder of the molecule via a thioester linkage.

[0019] In the context of the present invention "hybridizes under stringent conditions" is intended to describe hybridization and washing conditions under which nucleotide sequences with at least 60% homology to one another usually remain hybridized with one another. Conditions are preferably such that sequences with at least approximately 65%, preferably

at least approximately 70% and especially preferably at least 75% or more homology to one another usually remain hybridized to one another. These stringent conditions are known to the skilled worker and described, for example, in Current Protocols in Molecular Biology, John Wiley & Sons, N. Y. (1989), 6.3.1-6.3.6. A preferred nonlimiting example of stringent hybridization conditions is hybridizations in 6 x sodium chloride/sodium citrate (= SSC) at approximately 45°C, followed by one or more washing steps in 0.2 x SSC, 0.1 % SDS at 50 to 65°C. The skilled worker knows that these hybridization conditions differ depending on the type of nucleic acid and, for example when organic solvents are present, regarding temperature and buffer concentration. Under "standard hybridization conditions", for example, the hybridization temperature is, depending on the type of nucleic acid, between 42°C and 58°C in aqueous buffer with a concentration of 0.1 to 5 x SSC (pH 7.2). If organic solvents, for example 50% formamide, are present in the above-mentioned buffer, the temperature under standard conditions is approximately 42°C. The hybridization conditions for DNA:DNA hybrids, for example, are 0.1 x SSC and 20°C to 45°C, preferably 30°C to 45°C. The hybridization conditions for DNA:RNA hybrids are, for example, 0.1 x SSC and 30°C to 55°C, preferably 45°C to 55°C. The abovementioned hybridization conditions are determined by way of example for a nucleic acid with approximately 100 bp (= base pairs) in length and with a G + C content of 50% in the absence of formamide. The skilled worker knows how to determine the required hybridization conditions on the basis of the abovementioned textbooks or textbooks such as Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989; Hames and Higgins (Ed.) 1985, "Nucleic Acids Hybridization: A Practical Approach", IRL Press at Oxford University Press, Oxford; Brown (Ed.) 1991, "Essential Molecular Biology: A Practical Approach", IRL Press at Oxford University Press, Oxford.

[0020] Furthermore, when the present specification refers to isolated nucleic acid molecules of a nucleotide sequence which hybridize with one of the nucleotide sequences shown in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 or with a part thereof under stringent conditions, "a part thereof" is understood as meaning, in accordance with the invention, that at least 25 base pairs (= bp), 50 bp, 75 bp, 100 bp, 125 bp or 150 bp, preferably at least 175 bp, 200 bp, 225 bp, 250 bp, 275 bp or 300 bp, especially preferably 350 bp, 400 bp, 450 bp, 500 bp or more base pairs are used for the hybridization.

[0021] In the context of the present invention "Homologs" of the $\Delta 5$ -desaturase nucleic acid sequences with the sequence SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 means, for example, allelic variants with at least approximately 50 or 60%, preferably at least approximately 60 or 70%, more preferably at least approximately 70 or 80%, 90% or 95% and even more preferably at least approximately 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95 %, 96%, 97%, 98%, 99% or more identity or homology with a nucleotide sequence shown in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5. "Allelic variants" comprise in particular functional variants which can be obtained by deletion, insertion or substitution of nucleotides from/into the sequence, it being intended, however, that the enzyme activity of the resulting proteins which are synthesized is advantageously retained for the insertion of one or more genes. "Homologs" also means bacterial, fungal and plant homologs, truncated sequences, single-stranded DNA or RNA of the coding and noncoding DNA sequence and derivatives such as, for example, promoter variants. The promoters upstream of the nucleotide sequences detailed can be modified by one or more nucleotide exchanges, by insertion(s) and/or deletion (s) without the functionality or activity of the promoters being adversely affected, however. It is furthermore possible that the modification of the promoter sequence enhances their activity or that they are replaced entirely by more active promoters, including those from heterologous organisms.

[0022] In order to determine the percentage of homology (= identity) of two amino acid sequences, the sequences are written one under the other for an optimal comparison (for example, gaps may be introduced into the sequence of a protein or of a nucleic acid in order to generate an optimal alignment with the other protein or the other nucleic acid). Then, the amino acid residue or nucleotides at the corresponding amino acid positions or nucleotide positions are compared. If a position in a sequence is occupied by the same amino acid residue or the same nucleotide as the corresponding position in the other sequence, then the molecules are homologous at this position (i.e. amino acid or nucleic acid "homology" as used in the present context corresponds to amino acid or nucleic acid "identity"). The percentage of homology between the two sequences is a function of the number of positions which the sequences share (i.e. % homology = number of identical positions/total number of positions x 100). The terms homology and identity are therefore to be considered as synonymous.

[0023] The homology was calculated over the entire amino acid or nucleic acid sequence region. The skilled worker has available a series of programs which are based on various algorithms for the comparison of various sequences. Here, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. The program PileUp (J. Mol. Evolution., 25, 351-360, 1987, Higgins et al., CABIOS, 5 1989: 151-153) or the programs Gap and BestFit [Needleman and Wunsch (J. Mol. Biol. 48; 443-453 (1970) and Smith and Waterman (Adv. Appl. Math. 2; 482-489 (1981)), which are part of the GCG software packet [Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711 (1991)], were used for the sequence alignment. The sequence homology values which are indicated above as a percentage were determined over the entire sequence region using the program GAP and the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000. Unless otherwise specified, these settings were always used as standard settings for the sequence alignments.

[0024] In the context of the present invention "Δ5-desaturase activity" is understood as meaning that a protein encoded by a derivative of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 retains an enzymatic activity of at least 10%, preferably 20%, especially preferably 30% and very especially 40% in comparison with the proteins/enzymes encoded by the sequence SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 and can thus catalyse the reaction of a compound of formula III to form a compound of formula I.

[0025] In the context of the present invention, the term "thioester linkage" refers to a compound in which the -O- of an ester group is replaced by a -S-. Thus, in the case where R¹ is lysophosphatidylcholine, the free oxygen of the lysophosphatidyl choline radical may be replaced by sulphur atom such that the linkage to the remainder of the molecule is via a thioester rather than an ester group. The same also applies to other R¹ groups.

[0026] In the context of the present invention, "saturated or unsaturated C₂-C₂₄-alkylcarbonyl chain" refers to a substituted or unsubstituted alkylcarbonyl, alkenylcarbonyl or alkynylcarbonyl group having one or more double and/or triple bonds. Suitable substituents include hydroxyl or epoxy.

[0027] When R² or R³ are C₂-C₂₄-alkylcarbonyl chains, examples of suitable groups include radicals such as ethylcarbonyl, n-propylcarbonyl, n-butylcarbonyl, n-pentylcarbonyl, n-hexylcarbonyl, n-heptylcarbonyl, n-octylcarbonyl, n-nonylcarbonyl, n-decylcarbonyl, n-undecylcarbonyl, n-dodecylcarbonyl, n-tridecylcarbonyl, n-tetradecylcarbonyl, n-pentadecylcarbonyl, n-hexadecylcarbonyl, n-heptadecylcarbonyl, n-octadecylcarbonyl, n-nonadecylcarbonyl, n-eicosylcarbonyl, n-docosanylcarbonyl- or n-tetracosanylcarbonyl, which comprise one or more double bonds.

[0028] Saturated or unsaturated C₁₀-C₂₂-alkylcarbonyl radicals such as n-decylcarbonyl, n-undecylcarbonyl, n-dodecylcarbonyl, n-tridecylcarbonyl, n-tetradecylcarbonyl, n-pentadecylcarbonyl, n-hexadecylcarbonyl, n-heptadecylcarbonyl, n-octadecylcarbonyl, n-nonadecylcarbonyl, n-eicosylcarbonyl, n-docosanylcarbonyl or n-tetracosanylcarbonyl, which comprise one or more double bonds are preferred.


[0029] Especially preferred are saturated and/or unsaturated C₁₀-C₂₂-alkylcarbonyl radicals such as C₁₀-alkylcarbonyl, C₁₁-alkylcarbonyl, C₁₂-alkylcarbonyl, C₁₃-alkylcarbonyl, C₁₄-alkylcarbonyl, C₁₆-alkylcarbonyl, C₁₈-alkylcarbonyl, C₂₀-alkylcarbonyl or C₂₂-alkylcarbonyl radicals which comprise one or more double bonds.

[0030] Even more preferred are saturated or unsaturated C₁₆-C₂₂-alkylcarbonyl radicals such as C₁₆-alkylcarbonyl, C₁₈-alkylcarbonyl, C₂₀-alkylcarbonyl or C₂₂-alkylcarbonyl radicals which comprise one or more double bonds. These advantageous radicals can comprise two, three, four, five or six double bonds. The especially preferred radicals with 20 or 22 carbon atoms in the fatty acid chain comprise up to six double bonds, advantageously three, four, five or six double bonds, especially preferably five or six double bonds. All the abovementioned radicals are derived from the corresponding fatty acids. The substituents R² or R³ in formula II are advantageously and independently of one another saturated or unsaturated C₁₈-C₂₂-alkylcarbonyl, especially advantageously they are, independently of one another, unsaturated C₁₈-, C₂₀- or C₂₂-alkylcarbonyl with at least two double bonds.


[0031] In the compound of general formula (I), it is preferred that the -CH=CH- double bond is in the cis orientation.

[0032] The C₁₀-C₁₈ hydrocarbon chain which forms the group Y may contain up to 4 carbon-carbon double bonds and these will generally be positioned such that the product of formula I is a non-methylene-interrupted fatty acid.

[0033] In the moiety Y, the carbon-carbon double bonds are preferably in the cis orientation. Preferred Y moieties include:



such that the compound of general formula I is sciadonic acid or a derivative thereof; and



such that the compound of general formula (I) is juniperonic acid or a derivative thereof.



[0034] When Y is one of these moieties, the product can be obtained in any organism which naturally produces linoleic (C18:2 n-6) or linolenic (C18:3 n-3) acid by genetically modifying that organism to express a Δ9 elongase enzyme as well as a Δ5-desaturase enzyme. The processes are as follows:

C18:2 n-6 (linoleic acid) is converted by Δ9 elongase to 11,14 C20:2 ; 11,14 C20:2 is converted by Δ5-desaturase to Δ5cis-11,14-20:3 (SA).

C18:3 n-3 (linolenic acid) is converted by Δ9 elongase to 11,14,17 C20:3 ; 11,14,17 C20:3 is converted by Δ5-desaturase to Δ5cis-11,14,17-20:4 (JA).

[0035] Therefore, in a second aspect of the invention, there is provided a process for the production of a compound of general formula (I) as defined above and in which Y is:

EP 2 388 332 A1



5 in an organism which comprises linoleic (C18:2 n-6) or linolenic (C18:3 n-3) acid, the process comprising introducing into the organism:

at least one nucleic acid sequence comprising :

- 10 a) SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5;
b) a nucleic acid sequence which hybridizes under stringent conditions with a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.
c) a nucleic acid sequence which encodes a polypeptide of SEQ ID NO:2, SEQ ID NO: 4 or SEQ ID NO: 6;
15 d) a derivative of a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, wherein said polypeptide has $\Delta 5$ -desaturase activity;

and at least one nucleic acid sequence encoding a polypeptide having $\Delta 9$ elongase activity;
and expressing said nucleic acid sequences.

20 **[0036]** In the process of this aspect of the invention, the nucleic acid sequence encoding a polypeptide having $\Delta 9$ elongase activity may comprise a sequence encoding the C18- $\Delta 9$ elongase from *Isochrysis galbana* (SEQ ID NO: 25) or *Acanthamoeba castellanii* (SEQ ID NO: 24).

25 **[0037]** The processes according to the invention preferably yields total $\Delta 5$ -unsaturated fatty acids in a content of at least 1% by weight, advantageously at least 3% by weight, based on the total fatty acids in the transgenic organisms, preferably in a transgenic plant.

[0038] The fatty acids are advantageously produced in bound form. These $\Delta 5$ -unsaturated fatty acids can, with the aid of the nucleic acids used in the process according to the invention, be positioned at the sn1, sn2 and/or sn3 position of the advantageously produced triglycerides.

30 **[0039]** Since a plurality of reaction steps are performed by the starting compounds linoleic acid (C18:2) and linolenic acid (C18:3) in the process according to the second aspect of invention, the end products of the process such as, for example, sciadonic acid or juniperonic acid are not obtained as absolutely pure products; minor traces of the precursors are always present in the end product. If, for example, both linoleic acid and linolenic acid are present in the starting organism and the starting plant, the end products are present as mixtures.

35 **[0040]** Fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise 6 to 15% of palmitic acid, 1 to 6% of stearic acid, 7-85% of oleic acid, 0.5 to 8% of vaccenic acid, 0.1 to 1% of arachic acid, 7 to 25% of saturated fatty acids, 8 to 85% of monounsaturated fatty acids and 60 to 85% of polyunsaturated fatty acids, in each case based on 100% and on the total fatty acid content of the organisms. Advantageous polyunsaturated fatty acids which are present in the fatty acid esters or fatty acid mixtures are preferably at least 0.1, 0.2, 0.3,
40 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1% of arachidonic acid, based on the total fatty acid content. Moreover, the fatty acid esters or fatty acid mixtures which have been produced by the process of the invention advantageously comprise fatty acids selected from the group of the fatty acids erucic acid (13-docosaenoic acid), sterculic acid (9,10-methyleneoctadec-9-enoic acid), malvalic acid (8,9-methyleneheptadec-8-enoic acid), chaulmoogric acid (cyclo-pentenedodecanoic acid), furan fatty acid (9,12-epoxyoctadeca-9,11-dienoic acid), vernolic acid (9,10-epoxyoctadec-12-enoic acid), tariric acid (6-octadecynoic acid), 6-nonadecynoic acid, santalbic acid (11-octadecen-9-ynoic acid), 6,9-octadecenyonic acid, pyru-
45 lic acid (10-heptadecen-8-ynoic acid), crepenyninic acid (9-octadecen-12-ynoic acid), 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diyonic acid, petroselenic acid (cis-6-octadecenoic acid), 9c,12t-octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid), catalpic acid (9t11 t13c-octadecatrienoic acid), eleostearic acid (9c11 t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), punnic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid), pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or coriolic acid (13-hydroxy-9c,11t-octadecadienoic acid). The abovementioned fatty acids are, as a rule, advantageously only found in traces in the fatty acid esters or fatty acid mixtures produced by the process according to the invention, that is to say that, based on the total fatty acids, they occur to less than 30%, preferably to less than 25%, 24%, 23%, 22% or 21%, especially preferably to less than 20%,
50 15%, 10%, 9%, 8%, 7%, 6% or 5%, very especially preferably to less than 4%, 3%, 2% or 1%. In a further preferred form of the invention, these abovementioned fatty acids occur to less than 0.9%, 0.8%, 0.7%, 0.6% or 0.5%, especially preferably to less than 0.4%, 0.3%, 0.2%, 0.1%, based on the total fatty acids. The fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise less than 0.1%, based on the total fatty

acids, and/or no butyric acid, no cholesterol, no clupanodonic acid (= docosapentaenoic acid, C22:5^{Δ4,8,12,15,21}) and no nisinic acid (tetracosahexaenoic acid, C23:6^{Δ3,8,12,15,18,21}).

[0041] Chemically pure polyunsaturated fatty acids or fatty acid compositions can also be synthesized by the processes described above. To this end, the fatty acids or the fatty acid compositions are isolated from the organisms, such as the microorganisms or the plants or the culture medium in or on which the organisms have been grown, or from the organism and the culture medium, in the known manner, for example via extraction, distillation, crystallization, chromatography or a combination of these methods. These chemically pure fatty acids or fatty acid compositions are advantageous for applications in the food industry sector, the cosmetic sector and especially the pharmacological industry sector.

[0042] Suitable organisms for the production in the process according to the invention are, in principle, any organisms such as microorganisms, nonhuman animals or plants. Plants which are suitable are, in principle, all those plants which are capable of synthesizing fatty acids, such as all dicotyledonous or monocotyledonous plants, algae or mosses. Advantageous plants are selected from the group of the plant families Adolotheciaceae, Anacardiaceae, Asteraceae, Apiaceae, Betulaceae, Boraginaceae, Brassicaceae, Bromeliaceae, Caricaceae, Cannabaceae, Convolvulaceae, Chenopodiaceae, Crypthecodiniaceae, Cucurbitaceae, Ditrichaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Lauraceae, Leguminosae, Linaceae, Euglenaceae, Prasinophyceae or vegetable plants or ornamentals such as Tagetes.

[0043] Examples which may be mentioned are the following plants selected from the group consisting of: Adolotheciaceae such as the genera Physcomitrella, for example the genus and species Physcomitrella patens, Anacardiaceae such as the genera Pistacia, Mangifera, Anacardium, for example the genus and species Pistacia vera [pistachio], Mangifera indica [mango] or Anacardium occidentale [cashew], Asteraceae, such as the genera Calendula, Carthamus, Centaurea, Cichorium, Cynara, Helianthus, Lactuca, Locusta, Tagetes, Valeriana, for example the genus and species Calendula officinalis [common marigold], Carthamus tinctorius [safflower], Centaurea cyanus [cornflower], Cichorium intybus [chicory], Cynara scolymus [artichoke], Helianthus annuus [sunflower], Lactuca sativa, Lactuca crispa, Lactuca esculenta, Lactuca scariola L. ssp. sativa, Lactuca scariola L. var. integrata, Lactuca scariola L. var. integrifolia, Lactuca sativa subsp. romana, Locusta communis, Valeriana locusta [salad vegetables], Tagetes lucida, Tagetes erecta or Tagetes tenuifolia [african or french marigold], Apiaceae, such as the genus Daucus, for example the genus and species Daucus carota [carrot], Betulaceae, such as the genus Corylus, for example the genera and species Corylus avellana or Corylus colurna [hazelnut], Boraginaceae, such as the genus Borago, for example the genus and species Borago officinalis [borage], Brassicaceae, such as the genera Brassica, Camelina, Melanosinapis, Sinapis,

[0044] Arabidopsis, for example the genera and species Brassica napus, Brassica rapa ssp. [oilseed rape], Sinapis arvensis Brassica juncea, Brassica juncea var. juncea, Brassica juncea var. crispifolia, Brassica juncea var. foliosa, Brassica nigra, Brassica sinapioides, Camelina sativa, Melanosinapis communis [mustard], Brassica oleracea [fodder beet] or Arabidopsis thaliana, Bromeliaceae, such as the genera Anana, Bromelia (pineapple), for example the genera and species Anana comosus, Ananas ananas or Bromelia comosa [pineapple], Caricaceae, such as the genus Carica, such as the genus and species Carica papaya [pawpaw], Cannabaceae, such as the genus Cannabis, such as the genus and species Cannabis sativa [hemp], Convolvulaceae, such as the genera Ipomea, Convolvulus, for example the genera and species Ipomea batatas, Ipomea pandurata, Convolvulus batatas, Convolvulus tiliaceus, Ipomea fastigiata, Ipomea tiliacea, Ipomea triloba or Convolvulus panduratus [sweet potato, batate], Chenopodiaceae, such as the genus Beta, such as the genera and species Beta vulgaris, Beta vulgaris var. altissima, Beta vulgaris var. vulgaris, Beta maritima, Beta vulgaris var. perennis, Beta vulgaris var. conditiva or Beta vulgaris var. esculenta [sugarbeet], Crypthecodiniaceae, such as the genus Crypthecodinium, for example the genus and species Crypthecodinium cohnii, Cucurbitaceae, such as the genus Cucurbita, for example the genera and species Cucurbita maxima, Cucurbita mixta, Cucurbita pepo or Cucurbita moschata [pumpkin/squash], Cymbellaceae, such as the genera Amphora, Cymbella, Okedenia, Phaeodactylum, Reimeria, for example the genus and species Phaeodactylum tricornutum, Ditrichaceae, such as the genera Ditrichaceae, Astomiopsis, Ceratodon, Chrysoblastella, Ditrichum, Distichium, Eccremidium, Lophidion, Philibertiella, Pleuridium, Saelania, Trichodon, Skottsbergia, for example the genera and species Ceratodon antarcticus, Ceratodon columbiae, Ceratodon heterophyllus, Ceratodon purpurascens, Ceratodon purpureus, Ceratodon purpureus ssp. convolutus, Ceratodon purpureus ssp. stenocarpus, Ceratodon purpureus var. rotundifolius, Ceratodon ratodon, Ceratodon stenocarpus, Chrysoblastella chilensis, Ditrichum ambiguum, Ditrichum brevisetum, Ditrichum crispatisimum, Ditrichum difficile, Ditrichum falcifolium, Ditrichum flexicaule, Ditrichum giganteum, Ditrichum heteromallum, Ditrichum lineare, Ditrichum lineare, Ditrichum montanum, Ditrichum montanum, Ditrichum pallidum, Ditrichum punctulatum, Ditrichum pusillum, Ditrichum pusillum var. tortile, Ditrichum rhynchostegium, Ditrichum schimperi, Ditrichum tortile, Distichium capillaceum, Distichium hagenii, Distichium inclinatum, Distichium macounii, Eccremidium floridanum, Eccremidium whiteleggei, Lophidion strictus, Pleuridium acuminatum, Pleuridium alternifolium, Pleuridium holdridgei, Pleuridium mexicanum, Pleuridium ravenelii, Pleuridium subulatum, Saelania glaucescens, Trichodon borealis, Trichodon cylindricus or Trichodon cylindricus var. oblongus, Elaeagnaceae, such as the genus Elaeagnus, for example the genus and species Olea europaea [olive], Ericaceae, such as the genus Kalmia, for example the genera and species Kalmia latifolia, Kalmia angustifolia, Kalmia microphylla, Kalmia polifolia, Kalmia occidentalis, Cistus chamaerhodendros

or *Kalmia lucida* [mountain laurel], Euglenaceae, such as the genera *Ascoglena*, *Astasia*, *Colacium*, *Cyclidiopsis*, *Euglena*, *Euglenopsis*, *Hyalaphacus*, *Khawkinea*, *Lepocinclis*, *Phacus*, *Strombomonas*, *Trachelomonas*, for example the genus and species *Euglena gracilis*; Euphorbiaceae, such as the genera *Manihot*, *Janipha*, *Jatropha*, *Ricinus*, for example the genera and species *Manihot utilissima*, *Janipha manihot*, *Jatropha manihot*, *Manihot aipil*, *Manihot dulcis*, *Manihot manihot*, *Manihot melanobasis*, *Manihot esculenta* [cassava] or *Ricinus communis* [castor-oil plant], Fabaceae, such as the genera *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicago*, *Glycine*, *Dolichos*, *Phaseolus*, soybean, for example the genera and species *Pisum sativum*, *Pisum arvense*, *Pisum humile* [pea], *Albizia berteriana*, *Albizia julibrissin*, *Albizia lebbeck*, *Acacia berteriana*, *Acacia littoralis*, *Albizia berteriana*, *Albizzia berteriana*, *Cathormion berteriana*, *Feuillea berteriana*, *Inga fragrans*, *Pithecellobium berterianum*, *Pithecellobium fragrans*, *Pithecolobium berterianum*, *Pseudalbizzia berteriana*, *Acacia julibrissin*, *Acacia nemu*, *Albizia nemu*, *Feuillea julibrissin*, *Mimosa julibrissin*, *Mimosa speciosa*, *Sericanrda julibrissin*, *Acacia lebbeck*, *Acacia macrophylla*, *Albizia lebbeck*, *Feuillea lebbeck*, *Mimosa lebbeck*, *Mimosa speciosa*, *Medicago sativa*, *Medicago falcata*, *Medicago varia* [alfalfa] *Glycine max* *Dolichos soja*, *Glycine gracilis*, *Glycine hispida*, *Phaseolus max*, *Soja hispida* or *Soja max* [soybean], Funariaceae, such as the genera *Aphanorrhagma*, *Entosthodon*, *Funaria*, *Physcomitrella*, *Physcomitrium*, for example the genera and species *Aphanorrhagma serratum*, *Entosthodon attenuatus*, *Entosthodon bolanderi*, *Entosthodon bonplandii*, *Entosthodon californicus*, *Entosthodon drummondii*, *Entosthodon jamesonii*, *Entosthodon leibergii*, *Entosthodon neoscoticus*, *Entosthodon rubrisetus*, *Entosthodon spathulifolius*, *Entosthodon tucsoni*, *Funaria americana*, *Funaria bolanderi*, *Funaria calcarea*, *Funaria californica*, *Funaria calvescens*, *Funaria convoluta*, *Funaria flavicans*, *Funaria groutiana*, *Funaria hygrometrica*, *Funaria hygrometrica* var. *arctica*, *Funaria hygrometrica* var. *calvescens*, *Funaria hygrometrica* var. *convoluta*, *Funaria hygrometrica* var. *muralis*, *Funaria hygrometrica* var. *utahensis*, *Funaria microstoma*, *Funaria microstoma* var. *obtusifolia*, *Funaria muhlenbergii*, *Funaria orcuttii*, *Funaria plano-convexa*, *Funaria polaris*, *Funaria ravenelii*, *Funaria rubriseta*, *Funaria serrata*, *Funaria sonorae*, *Funaria sublimbatus*, *Funaria tucsoni*, *Physcomitrella californica*, *Physcomitrella patens*, *Physcomitrella readeri*, *Physcomitrium australe*, *Physcomitrium californicum*, *Physcomitrium collenchymatum*, *Physcomitrium coloradense*, *Physcomitrium cupuliferum*, *Physcomitrium drummondii*, *Physcomitrium eurytomum*, *Physcomitrium flexifolium*, *Physcomitrium hookeri*, *Physcomitrium hookeri* var. *serratum*, *Physcomitrium immersum*, *Physcomitrium kellermanii*, *Physcomitrium megalocarpum*, *Physcomitrium pyriforme*, *Physcomitrium pyriforme* var. *serratum*, *Physcomitrium rufipes*, *Physcomitrium sandbergii*, *Physcomitrium subsphaericum*, *Physcomitrium washingtoniense*, Geraniaceae, such as the genera *Pelargonium*, *Cocos*, *Oleum*, for example the genera and species *Cocos nucifera*, *Pelargonium grossularioides* or *Oleum cocois* [coconut], Gramineae, such as the genus *Saccharum*, for example the genus and species *Saccharum officinarum*, Juglandaceae, such as the genera *Juglans*, *Wallia*, for example the genera and species *Juglans regia*, *Juglans ailanthifolia*, *Juglans sieboldiana*, *Juglans cinerea*, *Wallia cinerea*, *Juglans bixbyi*, *Juglans californica*, *Juglans hindsii*, *Juglans intermedia*, *Juglans jamaicensis*, *Juglans major*, *Juglans microcarpa*, *Juglans nigra* or *Wallia nigra* [walnut], Lauraceae, such as the genera *Persea*, *Laurus*, for example the genera and species *Laurus nobilis* [bay], *Persea americana*, *Persea gratissima* or *Persea persea* [avocado], Leguminosae, such as the genus *Arachis*, for example the genus and species *Arachis hypogaea* [peanut], Linaceae, such as the genera *Adenolinum*, for example the genera and species *Linum usitatissimum*, *Linum humile*, *Linum austriacum*, *Linum bienne*, *Linum angustifolium*, *Linum catharticum*, *Linum flavum*, *Linum grandiflorum*, *Adenolinum grandiflorum*, *Linum lewisii*, *Linum narbonense*, *Linum perenne*, *Linum perenne* var. *lewisii*, *Linum pratense* or *Linum trigynum* [linseed], Lythraeae, such as the genus *Punica*, for example the genus and species *Punica granatum* [pomegranate], Malvaceae, such as the genus *Gossypium*, for example the genera and species *Gossypium hirsutum*, *Gossypium arboreum*, *Gossypium barbadense*, *Gossypium herbaceum* or *Gossypium thurberi* [cotton], Marchantiaceae, such as the genus *Marchantia*, for example the genera and species *Marchantia berteriana*, *Marchantia foliacea*, *Marchantia macropora*, Musaceae, such as the genus *Musa*, for example the genera and species *Musa nana*, *Musa acuminata*, *Musa paradisiaca*, *Musa* spp. [banana], Onagraceae, such as the genera *Camissonia*, *Oenothera*, for example the genera and species *Oenothera biennis* or *Camissonia brevipes* [evening primrose], Palmae, such as the genus *Elaeis*, for example the genus and species *Elaeis guineensis* [oil palm], Papaveraceae, such as, for example, the genus *Papaver*, for example the genera and species *Papaver orientale*, *Papaver rhoeas*, *Papaver dubium* [poppy], Pedaliaceae, such as the genus *Sesamum*, for example the genus and species *Sesamum indicum* [sesame], Piperaceae, such as the genera *Piper*, *Artanthe*, *Peperomia*, *Steffensia*, for example the genera and species *Piper aduncum*, *Piper amalago*, *Piper angustifolium*, *Piper auritum*, *Piper betel*, *Piper cubeba*, *Piper longum*, *Piper nigrum*, *Piper retrofractum*, *Artanthe adunca*, *Artanthe elongata*, *Peperomia elongata*, *Piper elongatum*, *Steffensia elongata* [cayenne pepper], Poaceae, such as the genera *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea* (maize), *Triticum*, for example the genera and species *Hordeum vulgare*, *Hordeum jubatum*, *Hordeum murinum*, *Hordeum secalinum*, *Hordeum distichon*, *Hordeum aegiceras*, *Hordeum hexastichon*, *Hordeum hexastichum*, *Hordeum irregulare*, *Hordeum sativum*, *Hordeum secalinum* [barley], *Secale cereale* [rye], *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida* [oats], *Sorghum bicolor*, *Sorghum halepense*, *Sorghum saccharatum*, *Sorghum vulgare*, *Andropogon drummondii*, *Holcus bicolor*, *Holcus sorghum*, *Sorghum aethiopicum*, *Sorghum arundinaceum*, *Sorghum caffrorum*, *Sorghum cernuum*, *Sorghum dochna*, *Sorghum drummondii*, *Sorghum durra*, *Sorghum guineense*, *Sorghum lanceolatum*, Sor-

ghum nervosum, Sorghum saccharatum, Sorghum subglabrescens, Sorghum verticilliflorum, Sorghum vulgare, Holcus halepensis, Sorghum miliaceum, Panicum militaceum [millet], Oryza sativa, Oryza latifolia [rice], Zea mays [maize] Triticum aestivum, Triticum durum, Triticum turgidum, Triticum hybernum, Triticum macha, Triticum sativum or Triticum vulgare [wheat], Porphyridiaceae, such as the genera Chrootheca, Flintiella, Petrovanella, Porphyridium, Rhodella, Rhodosorus, Vanhoeffenia, for example the genus and species Porphyridium cruentum, Proteaceae, such as the genus Macadamia, for example the genus and species Macadamia intergrifolia [macadamia], Prasinophyceae, such as the genera Nephroselmis, Prasinococcus, Scherffelia, Tetraselmis, Mantoniella, Ostreococcus, for example the genera and species Nephroselmis olivacea, Prasinococcus capsulatus, Scherffelia dubia, Tetraselmis chui, Tetraselmis suecica, Mantoniella squamata, Ostreococcus tauri, Rubiaceae, such as the genus Coffea, for example the genera and species Coffea spp., Coffea arabica, Coffea canephora or Coffea liberica [coffee], Scrophulariaceae, such as the genus Verbascum, for example the genera and species Verbascum blattaria, Verbascum chaixii, Verbascum densiflorum, Verbascum lagurus, Verbascum longifolium, Verbascum lychnitis, Verbascum nigrum, Verbascum olympicum, Verbascum phlomoides, Verbascum phoenicum, Verbascum pulverulentum or Verbascum thapsus [verbascum], Solanaceae, such as the genera Capsicum, Nicotiana, Solanum, Lycopersicon, for example the genera and species Capsicum annuum, Capsicum annuum var. glabrusculum, Capsicum frutescens [pepper], Capsicum annuum [paprika], Nicotiana tabacum, Nicotiana alata, Nicotiana attenuata, Nicotiana glauca, Nicotiana langsdorffii, Nicotiana obtusifolia, Nicotiana quadri-valvis, Nicotiana repanda, Nicotiana rustica, Nicotiana sylvestris [tobacco], Solanum tuberosum [potato], Solanum melongena [eggplant] Lycopersicon esculentum, Lycopersicon lycopersicum, Lycopersicon pyriforme, Solanum integrifolium or Solanum lycopersicum [tomato], Sterculiaceae, such as the genus Theobroma, for example the genus and species Theobroma cacao [cacao] or Theaceae, such as the genus Camellia, for example the genus and species Camellia sinensis [tea].

[0045] Advantageous microorganisms are, for example, fungi selected from the group of the families Chaetomiaceae, Choanephoraceae, Cryptococcaceae, Cunninghamellaceae, Dematiaceae, Moniliaceae, Mortierellaceae, Mucoraceae, Pythiaceae, Saccharomycetaceae, Saprolegniaceae, Schizosaccharomycetaceae, Sodariaceae or Tuberculariaceae.

[0046] Examples of microorganisms which may be mentioned are those from the groups: Choanephoraceae, such as the genera Blakeslea, Choanephora, for example the genera and species Blakeslea trispora, Choanephora cucurbitarum, Choanephora infundibulifera var. cucurbitarum, Mortierellaceae, such as the genus Mortierella, for example the genera and species Mortierella isabellina, Mortierella polycephala, Mortierella ramanniana, Mortierella vinacea, Mortierella zonata, Pythiaceae, such as the genera Phytium, Phytophthora, for example the genera and species Phytium debaryanum, Phytium intermedium, Phytium irregulare, Phytium megalacanthum, Phytium paroecandrum, Phytium sylvaticum, Phytium ultimum, Phytophthora cactorum, Phytophthora cinnamomi, Phytophthora citricola, Phytophthora citrophthora, Phytophthora cryptogea, Phytophthora drechsleri, Phytophthora erythroseptica, Phytophthora lateralis, Phytophthora megasperma, Phytophthora nicotianae, Phytophthora nicotianae var. parasitica, Phytophthora palmivora, Phytophthora parasitica, Phytophthora syringae, Saccharomycetaceae, such as the genera Hansenula, Pichia, Saccharomyces, Saccharomycodes, Yarrowia, for example the genera and species Hansenula anomala, Hansenula californica, Hansenula canadensis, Hansenula capsulata, Hansenula ciferrii, Hansenula glucozyma, Hansenula henricii, Hansenula holstii, Hansenula minuta, Hansenula nonfermentans, Hansenula philodendri, Hansenula polymorpha, Hansenula saturnus, Hansenula subpelliculosa, Hansenula wickerhamii, Hansenula wingei, Pichia alcoholophila, Pichia angusta, Pichia anomala, Pichia bispora, Pichia burtonii, Pichia canadensis, Pichia capsulata, Pichia carsonii, Pichia cellobiosa, Pichia ciferrii, Pichia farinosa, Pichia fermentans, Pichia finlandica, Pichia glucozyma, Pichia guilliermondii, Pichia haplophila, Pichia henricii, Pichia holstii, Pichia jadinii, Pichia lindnerii, Pichia membranaefaciens, Pichia methanolica, Pichia minuta var. minuta, Pichia minuta var. nonfermentans, Pichia norvegensis, Pichia ohmeri, Pichia pastoris, Pichia philodendri, Pichia pini, Pichia polymorpha, Pichia quercuum, Pichia rhodanensis, Pichia sargentensis, Pichia stipitidis, Pichia strasburgensis, Pichia subpelliculosa, Pichia toletana, Pichia trehalophila, Pichia vini, Pichia xylosa, Saccharomyces acetii, Saccharomyces baillii, Saccharomyces bayanus, Saccharomyces bisporus, Saccharomyces capensis, Saccharomyces carlsbergensis, Saccharomyces cerevisiae, Saccharomyces cerevisiae var. ellipsoideus, Saccharomyces chevalieri, Saccharomyces delbrueckii, Saccharomyces diastaticus, Saccharomyces drosophilarum, Saccharomyces elegans, Saccharomyces ellipsoideus, Saccharomyces fermentati, Saccharomyces florentinus, Saccharomyces fragilis, Saccharomyces heterogenicus, Saccharomyces hienipiensis, Saccharomyces inusitatus, Saccharomyces italicus, Saccharomyces kluyveri, Saccharomyces krusei, Saccharomyces lactis, Saccharomyces marxianus, Saccharomyces microellipsoides, Saccharomyces montanus, Saccharomyces norbensis, Saccharomyces oleaceus, Saccharomyces paradoxus, Saccharomyces pastorianus, Saccharomyces pretoriensis, Saccharomyces rosei, Saccharomyces rouxii, Saccharomyces uvarum, Saccharomycodes ludwigii, Yarrowia lipolytica, Schizosaccharomycetaceae such as the genera Schizosaccharomyces e.g. the species Schizosaccharomyces japonicus var. japonicus, Schizosaccharomyces japonicus var. versatilis, Schizosaccharomyces malidevorans, Schizosaccharomyces octosporus, Schizosaccharomyces pombe var. malidevorans, Schizosaccharomyces pombe var. pombe, Thraustochytriaceae such as the genera Althornia, Aplanochytrium, Japonochytrium, Schizochytrium, Thraustochytrium e.g. the species Schizochytrium aggregatum, Schizochytrium limacinum, Schizochytrium mangrovei, Schizochytrium minutum, Schizochytrium octosporum,

Thraustochytrium aggregatum, *Thraustochytrium amoeboides*, *Thraustochytrium antarcticum*, *Thraustochytrium arundinale*, *Thraustochytrium aureum*, *Thraustochytrium benthicola*, *Thraustochytrium globosum*, *Thraustochytrium indicum*, *Thraustochytrium kerguelense*, *Thraustochytrium kinnei*, *Thraustochytrium motivum*, *Thraustochytrium multirudimentale*, *Thraustochytrium pachydermum*, *Thraustochytrium proliferum*, *Thraustochytrium roseum*, *Thraustochytrium rossii*, *Thraustochytrium striatum* or *Thraustochytrium visurgense*.

[0047] Further advantageous microorganisms are, for example, bacteria selected from the group of the families Bacillaceae, Enterobacteriaceae or Rhizobiaceae.

[0048] Examples which may be mentioned are the following microorganisms selected from the group consisting of: Bacillaceae, such as the genus *Bacillus*, for example the genera and species *Bacillus acidocaldarius*, *Bacillus acidoterrestris*, *Bacillus alcalophilus*, *Bacillus amyloliquefaciens*, *Bacillus amylolyticus*, *Bacillus brevis*, *Bacillus cereus*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus sphaericus* subsp. *fusiformis*, *Bacillus galactophilus*, *Bacillus globisporus*, *Bacillus globisporus* subsp. *marinus*, *Bacillus halophilus*, - *Bacillus lentimorbus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus polymyxa*, *Bacillus psychrosaccharolyticus*, *Bacillus pumilus*, *Bacillus sphaericus*, *Bacillus subtilis* subsp. *spizizenii*, *Bacillus subtilis* subsp. *subtilis* or *Bacillus thuringiensis*; Enterobacteriaceae such as the genera *Citrobacter*, *Edwardsiella*, *Enterobacter*, *Erwinia*, *Escherichia*, *Klebsiella*, *Salmonella* or *Serratia*, for example the genera and species *Citrobacter amalonaticus*, *Citrobacter diversus*, *Citrobacter freundii*, *Citrobacter genomospecies*, *Citrobacter gillenbergii*, *Citrobacter intermedium*, *Citrobacter koseri*, *Citrobacter murliniae*, *Citrobacter* sp., *Edwardsiella hoshiniae*, *Edwardsiella ictaluri*, *Edwardsiella tarda*, *Erwinia alni*, *Erwinia amylovora*, *Erwinia ananatis*, *Erwinia aphidicola*, *Erwinia billingiae*, *Erwinia cacticida*, *Erwinia cancerogena*, *Erwinia carnegieana*, *Erwinia carotovora* subsp. *atroseptica*, *Erwinia carotovora* subsp. *betavasculorum*, *Erwinia carotovora* subsp. *odorifera*, *Erwinia carotovora* subsp. *wasabiae*, *Erwinia chrysanthemi*, *Erwinia cypripedii*, *Erwinia dissolvens*, *Erwinia herbicola*, *Erwinia mallotivora*, *Erwinia milletiae*, *Erwinia nigrifluens*, *Erwinia nimipressuralis*, *Erwinia persicina*, *Erwinia psidii*, *Erwinia pyrifoliae*, *Erwinia quercina*, *Erwinia rhamnontici*, *Erwinia rubrifaciens*, *Erwinia salicis*, *Erwinia stewartii*, *Erwinia tracheiphila*, *Erwinia uredovora*, *Escherichia adecarboxylata*, *Escherichia anindolica*, *Escherichia aurescens*, *Escherichia blattae*, *Escherichia coli*, *Escherichia coli* var. *communior*, *Escherichia coli*-mutabile, *Escherichia fergusonii*, *Escherichia hermannii*, *Escherichia* sp., *Escherichia vulneris*, *Klebsiella aerogenes*, *Klebsiella edwardsii* subsp. *atlantae*, *Klebsiella ornithinolytica*, *Klebsiella oxytoca*, *Klebsiella planticola*, *Klebsiella pneumoniae*, *Klebsiella pneumoniae* subsp. *pneumoniae*, *Klebsiella* sp., *Klebsiella terrigena*, *Klebsiella trevisanii*, *Salmonella abony*, *Salmonella arizonae*, *Salmonella bongori*, *Salmonella choleraesuis* subsp. *arizonae*, *Salmonella choleraesuis* subsp. *bongori*, *Salmonella choleraesuis* subsp. *choleraesuis*, *Salmonella choleraesuis* subsp. *diarizonae*, *Salmonella choleraesuis* subsp. *houtenae*, *Salmonella choleraesuis* subsp. *indica*, *Salmonella choleraesuis* subsp. *salamae*, *Salmonella daressalaam*, *Salmonella enterica* subsp. *houtenae*, *Salmonella enterica* subsp. *salamae*, *Salmonella enteritidis*, *Salmonella gallinarum*, *Salmonella heidelberg*, *Salmonella panama*, *Salmonella senftenberg*, *Salmonella typhimurium*, *Serratia entomophila*, *Serratia ficaria*, *Serratia fonticola*, *Serratia grimesii*, *Serratia liquefaciens*, *Serratia marcescens*, *Serratia marcescens* subsp. *marcescens*, *Serratia marinorubra*, *Serratia odorifera*, *Serratia plymuthensis*, *Serratia plymuthica*, *Serratia proteamaculans*, *Serratia proteamaculans* subsp. *quinovora*, *Serratia quinivorans* or *Serratia rubidaea*; Rhizobiaceae, such as the genera *Agrobacterium*, *Carbophilus*, *Chelatobacter*, *Ensifer*, *Rhizobium*, *Sinorhizobium*, for example the genera and species *Agrobacterium atlanticum*, *Agrobacterium ferrugineum*, *Agrobacterium gelatinovorum*, *Agrobacterium larrymoorei*, *Agrobacterium meteor*, *Agrobacterium radiobacter*, *Agrobacterium rhizogenes*, *Agrobacterium rubi*, *Agrobacterium stellulatum*, *Agrobacterium tumefaciens*, *Agrobacterium vitis*, *Carbophilus carboxidus*, *Chelatobacter heintzii*, *Ensifer adhaerens*, *Ensifer arboris*, *Ensifer fredii*, *Ensifer kostiensis*, *Ensifer kummerowiae*, *Ensifer medicae*, *Ensifer meliloti*, *Ensifer sahelii*, *Ensifer terangae*, *Ensifer xinjiangensis*, *Rhizobium ciceri*, *Rhizobium etli*, *Rhizobium fredii*, *Rhizobium galegae*, *Rhizobium gallicum*, *Rhizobium giardinii*, *Rhizobium hainanense*, *Rhizobium huakuii*, *Rhizobium huautlense*, *Rhizobium indigoferae*, *Rhizobium japonicum*, *Rhizobium leguminosarum*, *Rhizobium loessense*, *Rhizobium loti*, *Rhizobium lupini*, *Rhizobium mediterraneum*, *Rhizobium meliloti*, *Rhizobium mongolense*, *Rhizobium phaseoli*, *Rhizobium radiobacter*, *Rhizobium rhizogenes*, *Rhizobium rubi*, *Rhizobium sullae*, *Rhizobium tianshanense*, *Rhizobium trifolii*, *Rhizobium tropici*, *Rhizobium undicola*, *Rhizobium vitis*, *Sinorhizobium adhaerens*, *Sinorhizobium arboris*, *Sinorhizobium fredii*, *Sinorhizobium kostiensis*, *Sinorhizobium kummerowiae*, *Sinorhizobium medicae*, *Sinorhizobium meliloti*, *Sinorhizobium morelense*, *Sinorhizobium sahelii* or *Sinorhizobium xinjiangense*.

[0049] Further examples of advantageous microorganisms for the process according to the invention are protists or diatoms selected from the group of the families Dinophyceae, Turaniellidae or Oxytrichidae, such as the genera and species: *Cryptocodinium cohnii*, *Phaeodactylum tricornutum*, *Stylonychia mytilus*, *Stylonychia pustulata*, *Stylonychia putrina*, *Stylonychia notophora*, *Stylonychia* sp., *Colpidium campylum* or *Colpidium* sp. Those which are advantageously applied in the process according to the invention are transgenic organisms such as fungi, such as *Mortierella* or *Thraustochytrium*, yeasts such as *Saccharomyces* or *Schizosaccharomyces*, mosses such as *Physcomitrella* or *Ceratodon*, nonhuman animals such as *Caenorhabditis*, algae such as *Nephroselmis*, *Pseudoscofieldia*, *Prasinococcus*, *Scherffelia*, *Tetraselmis*, *Mantoniella*, *Ostreococcus*, *Cryptocodinium* or *Phaeodactylum* or plants such as dicotyledonous or monocotyledonous plants. Organisms which are especially advantageously used in the process according to the invention

are organisms which belong to the oil-producing organisms, that is to say which are used for the production of oil, such as fungi, such as *Mortierella* or *Thraustochytrium*, algae such as *Nephroselmis*, *Pseudoscourfielda*, *Prasinococcus*, *Scherffelia*, *Tetraselmis*, *Mantoniella*, *Ostreococcus*, *Cryptocodinium*, *Phaeodactylum*, or plants, in particular plants, preferably oilseed or oil crop plants which comprise large amounts of lipid compounds, such as peanut, oilseed rape, canola, sunflower, safflower (*Carthamus tinctoria*), poppy, mustard, hemp, castor-oil plant, olive, sesame, *Calendula*, *Punica*, evening primrose, verbascum, thistle, wild roses, hazelnut, almond, macadamia, avocado, bay, pumpkin/squash, linseed, soybean, pistachios, borage, trees (oil palm, coconut or walnut) or arable crops such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, *Tagetes*, Solanaceae plants such as potato, tobacco, eggplant and tomato, *Vicia* species, pea, alfalfa or bushy plants (coffee, cacao, tea), *Salix* species, and perennial grasses and fodder crops. Preferred plants according to the invention are oil crop plants such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, *Calendula*, *Punica*, evening primrose, pumpkin/squash, linseed, soybean, borage, trees (oil palm, coconut). Especially preferred for the second aspect of the invention are plants which are high in C18:2- and/or C18:3-fatty acids, such as sunflower, safflower, tobacco, verbascum, sesame, cotton, pumpkin/squash, poppy, evening primrose, walnut, linseed, hemp or thistle. Very especially preferred plants are plants such as safflower, sunflower, poppy, evening primrose, walnut, linseed or hemp.

[0050] It is also be advantageous for the above-described method according to the invention additionally to introduce, into the organism, further nucleic acids which encode enzymes of the fatty acid or lipid metabolism, in addition to the nucleic acids introduced in the processes of the first and second aspects of the invention.

[0051] Nucleic acids used in the process according to the invention are advantageously derived from plants such as algae, for example algae of the family of the *Prasinophyceae* such as the genera *Heteromastix*, *Mammella*, *Mantoniella*, *Micromonas*, *Nephroselmis*, *Ostreococcus*, *Prasinocladus*, *Prasinococcus*, *Pseudoscourfielda*, *Pycnococcus*, *Pyramimonas*, *Scherffelia* or *Tetraselmis* such as the genera and species *Heteromastix longifillis*, *Mamiella gilva*, *Mantoniella squamata*, *Micromonas pusilla*, *Nephroselmis olivacea*, *Nephroselmis pyriformis*, *Nephroselmis rotunda*, *Ostreococcus tauri*, *Ostreococcus* sp. *Prasinocladus ascus*, *Prasinocladus lubricus*, *Pycnococcus provasolii*, *Pyramimonas amyliifera*, *Pyramimonas disomata*, *Pyramimonas obovata*, *Pyramimonas orientalis*, *Pyramimonas parkeae*, *Pyramimonas spinifera*, *Pyramimonas* sp., *Tetraselmis apiculata*, *Tetraselmis carteriaformis*, *Tetraselmis chui*, *Tetraselmis convolutae*, *Tetraselmis desikacharyi*, *Tetraselmis gracilis*, *Tetraselmis hazeni*, *Tetraselmis impellucida*, *Tetraselmis inconspicua*, *Tetraselmis levis*, *Tetraselmis maculata*, *Tetraselmis marina*, *Tetraselmis striata*, *Tetraselmis subcordiformis*, *Tetraselmis suecica*, *Tetraselmis tetrabrachia*, *Tetraselmis tetrathele*, *Tetraselmis verrucosa*, *Tetraselmis verrucosa* fo. *rubens* or *Tetraselmis* sp. or from algae of the family *Euglenaceae* such as the genera *Ascoglena*, *Astasia*, *Colacium*, *Cyclidiopsis*, *Euglena*, *Euglenopsis*, *Hyalophacus*, *Khawkinea*, *Lepocinclis*, *Phacus*, *Strombomonas* or *Trachelomonas*, such as the genera and species *Euglena acus*, *Euglena geniculata*, *Euglena gracilis*, *Euglena mixocylindracea*, *Euglena rostrifera*, *Euglena viridis*, *Colacium stentorium*, *Trachelomonas cylindrica* or *Trachelomonas volvocina*. The nucleic acids used are advantageously derived from algae of the genera *Euglena*, *Mantoniella* or *Ostreococcus*.

[0052] Further advantageous plants are algae such as *Isochrysis* or *Cryptocodinium*, algae/diatoms such as *Thalassiosira* or *Phaeodactylum*, mosses such as *Physcomitrella* or *Ceratodon*, or higher plants such as the *Primulaceae* such as *Aleuritia*, *Calendula stellata*, *Osteospermum spinescens* or *Osteospermum hyoseroides*, microorganisms such as fungi, such as *Aspergillus*, *Thraustochytrium*, *Phytophthora*, *Entomophthora*, *Mucor* or *Mortierella*, bacteria such as *Shewanella*, yeasts or animals such as nematodes such as *Caenorhabditis*, insects, frogs, abalone, or fish. The isolated nucleic acid sequences according to the invention are advantageously derived from an animal of the order of the vertebrates. Preferably, the nucleic acid sequences are derived from the classes of the Vertebrata; Euteleostomi, Actinopterygii; Neopterygii; Teleostei; Euteleostei, Protacanthopterygii, Salmoniformes; Salmonidae or *Oncorhynchus* or Vertebrata, Amphibia, Anura, Pipidae, *Xenopus* or Evertebrata such as Protochordata, Tunicata, Holothuroidea, Cionidae such as *Amaroucium constellatum*, *Botryllus schlosseri*, *Ciona intestinalis*, *Molgula citrina*, *Molgula manhattensis*, *Perophora viridis* or *Styela partita*. The nucleic acids are especially advantageously derived from fungi, animals, or from plants such as algae or mosses, preferably from the order of the Salmoniformes, such as the family of the Salmonidae, such as the genus *Salmo*, for example from the genera and species *Oncorhynchus mykiss*, *Trutta trutta* or *Salmo trutta fario*, from algae, such as the genera *Mantoniella* or *Ostreococcus*, or from the diatoms such as the genera *Thalassiosira* or *Phaeodactylum* or from algae such as *Cryptocodinium*.

[0053] In a preferred embodiment, the process furthermore comprises the step of obtaining a cell or an intact organism which comprises the nucleic acid sequences used in the process, where the cell and/or the organism is transformed with a nucleic acid sequence according to the invention which encodes the $\Delta 5$ -desaturase, a gene construct or a vector as described above, alone or in combination with further nucleic acid sequences which encode proteins of the fatty acid or lipid metabolism. In a further preferred embodiment, this process furthermore comprises the step of obtaining the oils, lipids or free fatty acids from the organism or from the culture. The culture can, for example, take the form of a fermentation culture, for example in the case of the cultivation of microorganisms, such as, for example, *Mortierella*, *Thalassiosira*, *Mantoniella*, *Ostreococcus*, *Saccharomyces* or *Thraustochytrium*, or a greenhouse- or field-grown culture of a plant. The cell or the organism produced thus is advantageously a cell of an oil-producing organism, such as an oil crop, such

as, for example, peanut, oilseed rape, canola, linseed, hemp, peanut, soybean, safflower, hemp, sunflowers or borage.

[0054] In the case of plant cells, plant tissue or plant organs, "growing" is understood as meaning, for example, the cultivation on or in a nutrient medium, or of the intact plant on or in a substrate, for example in a hydroponic culture, potting compost or on arable land.

[0055] For the purposes of the invention, "transgenic" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette (= gene construct) or a vector comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, expression cassettes or vectors according to the invention, all those constructions brought about by recombinant methods in which either

- a) the nucleic acid sequence according to the invention, or
- b) a genetic control sequence which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or
- c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. The natural genetic environment is understood as meaning the natural genomic or chromosomal locus in the original organism or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, most preferably at least 5000 bp. A naturally occurring expression cassette - for example the naturally occurring combination of the natural promoter of the nucleic acid sequences with the corresponding $\Delta 5$ -desaturase gene - becomes a transgenic expression cassette when this expression cassette is modified by non-natural, synthetic ("artificial") methods such as, for example, mutagenic treatment. Suitable methods are described, for example, in US 5,565,350 or WO 00/15815.

[0056] A transgenic organism or transgenic plant for the purposes of the invention is therefore understood as meaning, as above, that the nucleic acids used in the process are not at their natural locus in the genome of an organism, it being possible for the nucleic acids to be expressed homologously or heterologously. However, as mentioned, transgenic also means that, while the nucleic acids according to the invention are at their natural position in the genome of an organism, the sequence has been modified with regard to the natural sequence, and/or that the regulatory sequences of the natural sequences have been modified. Transgenic is preferably understood as meaning the expression of the nucleic acids according to the invention at an unnatural locus in the genome, i.e. homologous or, preferably, heterologous expression of the nucleic acids takes place. Preferred transgenic organisms are fungi such as *Mortierella* or *Phytophthora*, mosses such as *Physcomitrella*, algae such as *Mantoniella*, *Euglena*, *Cryptothecodinium* or *Ostreococcus*, diatoms such as *Thalassiosira* or *Phaeodactylum*, or plants such as the oil crops.

[0057] Organisms or host organisms for the nucleic acids, the expression cassette or the vector used in the process according to the invention are, in principle, advantageously all organisms which are capable of synthesizing fatty acids, specifically unsaturated fatty acids, and/or which are suitable for the expression of recombinant genes. Examples which may be mentioned are plants such as *Arabidopsis*, *Asteraceae* such as *Calendula* or crop plants such as soybean, peanut, castor-oil plant, sunflower, maize, cotton, flax, oilseed rape, coconut, oil palm, safflower (*Carthamus tinctorius*) or cacao bean, microorganisms, such as fungi, for example the genus *Mortierella*, *Thraustochytrium*, *Saprolegnia*, *Phytophthora* or *Pythium*, bacteria, such as the genus *Escherichia* or *Shewanella*, yeasts, such as the genus *Saccharomyces*, cyanobacteria, ciliates, algae such as *Mantoniella*, *Euglena*, *Thalassiosira* or *Ostreococcus*, or protozoans such as dinoflagellates, such as *Cryptothecodinium*. Preferred organisms are those which are naturally capable of synthesizing substantial amounts of oil, such as fungi, such as *Mortierella alpina*, *Pythium insidiosum*, *Phytophthora infestans*, or plants such as soybean, oilseed rape, coconut, oil palm, safflower, flax, hemp, castor-oil plant, *Calendula*, peanut, cacao bean or sunflower, or yeasts such as *Saccharomyces cerevisiae* with soybean, flax, oilseed rape, safflower, sunflower, *Calendula*, *Mortierella* or *Saccharomyces cerevisiae* being especially preferred. In principle, host organisms are, in addition to the abovementioned transgenic organisms, also transgenic animals, advantageously nonhuman animals, for example *C. elegans*, *Ciona intestinalis* or *Xenopus laevis*.

[0058] Further utilizable host cells are detailed in: Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990).

[0059] Expression strains which can be used, for example those with a lower protease activity, are described in: Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128.

[0060] These include plant cells and certain tissues, organs and parts of plants in all their phenotypic forms such as anthers, fibers, root hairs, stalks, embryos, calli, cotyledons, petioles, harvested material, plant tissue, reproductive tissue and cell cultures which are derived from the actual transgenic plant and/or can be used for bringing about the

transgenic plant.

[0061] Transgenic plants which comprise the polyunsaturated fatty acids synthesized in the process according to the invention can advantageously be marketed directly without there being any need for the oils, lipids or fatty acids synthesized to be isolated. Plants for the process according to the invention are listed as meaning intact plants and all plant parts, plant organs or plant parts such as leaf, stem, seeds, root, tubers, anthers, fibers, root hairs, stalks, embryos, calli, cotyledons, petioles, harvested material, plant tissue, reproductive tissue and cell cultures which are derived from the actual transgenic plant and/or can be used for bringing about the transgenic plant. In this context, the seed comprises all parts of the seed such as the seed coats, epidermal cells, seed cells, endosperm or embryonic tissue. However, the compounds produced in the process according to the invention can also be isolated from the organisms, advantageously plants, in the form of their oils, fats, lipids and/or free fatty acids. Polyunsaturated fatty acids produced by this process can be obtained by harvesting the organisms, either from the crop in which they grow, or from the field. This can be done via pressing or extraction of the plant parts, preferably the plant seeds. In this context, the oils, fats, lipids and/or free fatty acids can be obtained by what is known as cold-beating or cold-pressing without applying heat. To allow for greater ease of disruption of the plant parts, specifically the seeds, they are previously comminuted, steamed or roasted. The seeds which have been pretreated in this manner can subsequently be pressed or extracted with solvents such as warm hexane. The solvent is subsequently removed. In the case of microorganisms, the latter are, after harvesting, for example extracted directly without further processing steps or else, after disruption, extracted via various methods with which the skilled worker is familiar. In this manner, more than 96% of the compounds produced in the process can be isolated. Thereafter, the resulting products are processed further, i.e. refined. In this process, substances such as the plant mucilages and suspended matter are first removed. What is known as desliming can be effected enzymatically or, for example, chemico-physically by addition of acid such as phosphoric acid. Thereafter, the free fatty acids are removed by treatment with a base, for example sodium hydroxide solution. The resulting product is washed thoroughly with water to remove the alkali remaining in the product and then dried. To remove the pigment remaining in the product, the products are subjected to bleaching, for example using filler's earth or active charcoal. At the end, the product is deodorized, for example using steam.

[0062] The fatty acids produced by the processes of the present invention can be isolated from the organism in the form of an oil, a lipid or a free fatty acid. Suitable organisms are, for example, those mentioned above. Preferred organisms are transgenic plants. One embodiment of the invention is therefore oils, lipids or fatty acids of formula I or fractions thereof which have been produced by the above-described process, especially preferably oil, lipid or a fatty acid composition comprising a compound of formula I and being derived from transgenic plants.

[0063] As described above, these oils, lipids or fatty acids advantageously comprise 6 to 15% of palmitic acid, 1 to 6% of stearic acid, 7-85% of oleic acid, 0.5 to 8% of vaccenic acid, 0.1 to 1% of arachic acid, 7 to 25% of saturated fatty acids, 8 to 85% of monounsaturated fatty acids and 60 to 85% of polyunsaturated fatty acids, in each case based on 100% and on the total fatty acid content of the organisms. Advantageous polyunsaturated fatty acids which are present in the fatty acid esters or fatty acid mixtures are preferably at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1% of arachidonic acid, based on the total fatty acid content. Moreover, the fatty acid esters or fatty acid mixtures which have been produced by the process of the invention advantageously comprise fatty acids selected from the group of the fatty acids erucic acid (13-docosaenoic acid), sterculic acid (9,10-methyleneoctadec-9-enoic acid), malvalic acid (8,9-methyleneheptadec-8-enoic acid), chaulmoogric acid (cyclo-pentenedodecanoic acid), furan fatty acid (9,12-epoxyoctadec-9,11-dienoic acid), vernolic acid (9,10-epoxyoctadec-12-enoic acid), tariric acid (6-octadecynoic acid), 6-nonadecynoic acid, santalbic acid (11-octadecen-9-ynoic acid), 6,9-octadecenynoic acid, pyrulic acid (10-heptadecen-8-ynoic acid), crepenyninic acid (9-octadecen-12-ynoic acid), 13,14-dihydrooropheic acid, octadecen-13-ene-9,11-diynoic acid, petroselenic acid (cis-6-octadecenoic acid), 9c,12t-octadecadienoic acid, calendulic acid (8t10t12c-octadecatrienoic acid), catalpic acid (9t11t13c-octadecatrienoic acid), eleostearic acid (9c11t13t-octadecatrienoic acid), jacaric acid (8c10t12c-octadecatrienoic acid), punnic acid (9c11t13c-octadecatrienoic acid), parinaric acid (9c11t13t15c-octadecatetraenoic acid), pinolenic acid (all-cis-5,9,12-octadecatrienoic acid), laballenic acid (5,6-octadecadienallenic acid), ricinoleic acid (12-hydroxyoleic acid) and/or coriolic acid (13-hydroxy-9c, 11 t-octadecadienoic acid). The abovementioned fatty acids are, as a rule, advantageously only found in traces in the fatty acid esters or fatty acid mixtures produced by the process according to the invention, that is to say that, based on the total fatty acids, they occur to less than 30%, preferably to less than 25%, 24%, 23%, 22% or 21%, especially preferably to less than 20%, 15%, 10%, 9%, 8%, 7%, 6% or 5%, very especially preferably to less than 4%, 3%, 2% or 1%. In a further preferred form of the invention, these abovementioned fatty acids occur to less than 0.9%, 0.8%, 0.7%, 0.6% or 0.5%, especially preferably to less than 0.4%, 0.3%, 0.2%, 0.1%, based on the total fatty acids. The fatty acid esters or fatty acid mixtures produced by the process according to the invention advantageously comprise less than 0.1%, based on the total fatty acids, and/or no butyric acid, no cholesterol, no clupanodonic acid (= docosapentaenoic acid, C22:5^{Δ4,8,12,15,21}) and no nisinic acid (tetracosahexaenoic acid, C23:6^{Δ3,8,12,15,18,21}).

[0064] A further embodiment according to the invention is the use of the oil, lipid, the fatty acids and/or the fatty acid composition in feedstuffs, foodstuffs, cosmetics or pharmaceuticals. The oils, lipids, fatty acids or fatty acid mixtures

according to the invention can be used in the manner with which the skilled worker is familiar for mixing with other oils, lipids, fatty acids or fatty acid mixtures of animal origin, such as, for example, fish oils. These oils, lipids, fatty acids or fatty acid mixtures, which are composed of vegetable and animal constituents, may also be used for the preparation of feedstuffs, foodstuffs, cosmetics or pharmacologicals.

5 **[0065]** The term "oil", "lipid" or "fat" is understood as meaning a fatty acid mixture comprising unsaturated, saturated, preferably esterified, fatty acid(s). The oil, lipid or fat is preferably high in polyunsaturated free or, advantageously, esterified fatty acid(s), in particular linoleic acid, γ -linolenic acid, dihomo- γ -linolenic acid, arachidonic acid, α -linolenic acid, stearidonic acid, eicosatetraenoic acid, eicosapentaenoic acid, docosapentaenoic acid or docosahexaenoic acid.

10 **[0066]** The amount of unsaturated esterified fatty acids preferably amounts to approximately 30%, a content of 50% is more preferred, a content of 60%, 70%, 80% or more is even more preferred. For the analysis, the fatty acid content can, for example, be determined by gas chromatography after converting the fatty acids into the methyl esters by transesterification. The oil, lipid or fat can comprise various other saturated or unsaturated fatty acids, for example calendulic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid and the like. The content of the various fatty acids in the oil or fat can vary, in particular depending on the starting organism.

15 **[0067]** The $\Delta 5$ -unsaturated fatty acids which are produced in the process are, as described above, for example sphingolipids, phosphoglycerides, lipids, glycolipids, phospholipids, monoacylglycerol, diacylglycerol, triacylglycerol or other fatty acid esters.

20 **[0068]** Starting from the polyunsaturated fatty acids with advantageously at least five or six double bonds, which acids have been prepared in the process according to the invention, the polyunsaturated fatty acids which are present can be liberated for example via treatment with alkali, for example aqueous KOH or NaOH, or acid hydrolysis, advantageously in the presence of an alcohol such as methanol or ethanol, or via enzymatic cleavage, and isolated via, for example, phase separation and subsequent acidification via, for example, H_2SO_4 . The fatty acids can also be liberated directly without the above-described processing step.

25 **[0069]** After their introduction into an organism, advantageously a plant cell or plant, the nucleic acids used in the process can either be present on a separate plasmid or, advantageously, integrated into the genome of the host cell. In the case of integration into the genome, integration can be random or else be effected by recombination such that the native gene is replaced by the copy introduced, whereby the production of the desired compound by the cell is modulated, or by the use of a gene in trans, so that the gene is linked operably with a functional expression unit which comprises at least one sequence which ensures the expression of a gene and at least one sequence which ensures the polyadenylation of a functionally transcribed gene. The nucleic acids are advantageously introduced into the organisms via
30 multiexpression cassettes or constructs for multiparallel expression, advantageously into the plants for the multiparallel seed-specific expression of genes.

35 **[0070]** If microorganisms such as yeasts, such as *Saccharomyces* or *Schizosaccharomyces*, fungi such as *Mortierella*, *Aspergillus*, *Phytophthora*, *Entomophthora*, *Mucor* or *Thraustochytrium*, algae such as *Isochrysis*, *Mantoniella*, *Euglena*, *Ostreococcus*, *Phaeodactylum* or *Cryptocodinium* are used as organisms in the process according to the invention, these organisms are advantageously grown in fermentation cultures.

40 **[0071]** If microorganisms are used as organisms in the process according to the invention, they are grown or cultured in the manner with which the skilled worker is familiar, depending on the host organism. As a rule, microorganisms are grown in a liquid medium comprising a carbon source, usually in the form of sugars, a nitrogen source, usually in the form of organic nitrogen sources such as yeast extract or salts such as ammonium sulfate, trace elements such as salts of iron, manganese and magnesium and, if appropriate, vitamins, at temperatures of between 0°C and 100°C, preferably between 10°C and 60°C, while passing in oxygen. The pH of the liquid medium can either be kept constant, that is to say regulated during the culturing period, or not. The cultures can be grown batchwise, semi-batchwise or continuously. Nutrients can be provided at the beginning of the fermentation or fed in semicontinuously or continuously. The polyunsaturated fatty acids produced can be isolated from the organisms as described above by processes known to the skilled worker, for example by extraction, distillation, crystallization, if appropriate precipitation with salt, and/or chromatography. To this end, the organisms can advantageously be disrupted beforehand.

45 **[0072]** If the host organisms are microorganisms, the process according to the invention is advantageously carried out at a temperature of between 0°C and 95°C, preferably between 10°C and 85°C, especially preferably between 15°C and 75°C, very especially preferably between 15°C and 45°C.

50 **[0073]** In this process, the pH value is advantageously kept between pH 4 and 12, preferably between pH 6 and 9, especially preferably between pH 7 and 8.

55 **[0074]** The process according to the invention can be operated batchwise, semibatchwise or continuously. An overview over known cultivation methods can be found in the textbook by Chmiel (Bioprozeßtechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess technology 1. Introduction to Bioprocess technology] (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and peripheral equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

[0075] The culture medium to be used must suitably meet the requirements of the strains in question. Descriptions of

culture media for various microorganisms can be found in the textbook "Manual of Methods für General Bacteriology" of the American Society for Bacteriology (Washington D. C., USA, 1981).

[0076] As described above, these media which can be employed in accordance with the invention usually comprise one or more carbon sources, nitrogen sources, inorganic salts, vitamins and/or trace elements.

[0077] Preferred carbon sources are sugars, such as mono-, di- or polysaccharides. Examples of very good carbon sources are glucose, fructose, mannose, galactose, ribose, sorbose, ribulose, lactose, maltose, sucrose, raffinose, starch or cellulose. Sugars can also be added to the media via complex compounds such as molasses or other byproducts from sugar raffination. The addition of mixtures of a variety of carbon sources may also be advantageous. Other possible carbon sources are oils and fats such as, for example, soya oil, sunflower oil, peanut oil and/or coconut fat, fatty acids such as, for example, palmitic acid, stearic acid and/or linoleic acid, alcohols and/or polyalcohols such as, for example, glycerol, methanol and/or ethanol, and/or organic acids such as, for example, acetic acid and/or lactic acid.

[0078] Nitrogen sources are usually organic or inorganic nitrogen compounds or materials comprising these compounds. Examples of nitrogen sources comprise ammonia in liquid or gaseous form or ammonium salts such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate or ammonium nitrate, nitrates, urea, amino acids or complex nitrogen sources such as cornsteep liquor, soya meal, soya protein, yeast extract, meat extract and others. The nitrogen sources can be used individually or as a mixture.

[0079] Inorganic salt compounds which may be present in the media comprise the chloride, phosphorus and sulfate salts of calcium, magnesium, sodium, cobalt, molybdenum, potassium, manganese, zinc, copper and iron.

[0080] Inorganic sulfur-containing compounds such as, for example, sulfates, sulfites, dithionites, tetrathionates, thio-sulfates, sulfides, or else organic sulfur compounds such as mercaptans and thiols may be used as sources of sulfur for the production of sulfur-containing fine chemicals, in particular of methionine.

[0081] Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts may be used as sources of phosphorus. Chelating agents may be added to the medium in order to keep the metal ions in solution. Particularly suitable chelating agents include dihydroxyphenols such as catechol or protocatechuate and organic acids such as citric acid.

[0082] The fermentation media used according to the invention for culturing microorganisms usually also comprise other growth factors such as vitamins or growth promoters, which include, for example, biotin, riboflavin, thiamine, folic acid, nicotinic acid, panthothenate and pyridoxine. Growth factors and salts are frequently derived from complex media components such as yeast extract, molasses, cornsteep liquor and the like. It is moreover possible to add suitable precursors to the culture medium. The exact composition of the media compounds heavily depends on the particular experiment and is decided upon individually for each specific case. Information on the optimization of media can be found in the textbook "Applied Microbiol. Physiology, A Practical Approach" (Editors P.M. Rhodes, P.F. Stanbury, IRL Press (1997) pp. 53-73, ISBN 0 19 963577 3). Growth media can also be obtained from commercial suppliers, for example Standard 1 (Merck) or BHI (brain heart infusion, DIFCO) and the like.

[0083] All media components are sterilized, either by heat (20 min at 1.5 bar and 121°C) or by filter sterilization. The components may be sterilized either together or, if required, separately. All media components may be present at the start of the cultivation or added continuously or batchwise, as desired.

[0084] The culture temperature is normally between 15°C and 45°C, preferably at from 25°C to 40°C, and may be kept constant or may be altered during the experiment. The pH of the medium should be in the range from 5 to 8.5, preferably around 7.0. The pH for cultivation can be controlled during cultivation by adding basic compounds such as sodium hydroxide, potassium hydroxide, ammonia and aqueous ammonia or acidic compounds such as phosphoric acid or sulfuric acid. Foaming can be controlled by employing antifoams such as, for example, fatty acid polyglycol esters. To maintain the stability of plasmids it is possible to add to the medium suitable substances having a selective effect, for example antibiotics. Aerobic conditions are maintained by introducing oxygen or oxygen-containing gas mixtures such as, for example, ambient air into the culture. The temperature of the culture is normally 20° to 40°C and preferably 25°C to 40°C. The culture is continued until formation of the desired product is at a maximum. This aim is normally achieved within 10 to 160 hours.

[0085] The fermentation broths obtained in this way, in particular those containing polyunsaturated fatty acids, usually contain a dry mass of from 7.5 to 25% by weight. The fermentation broth can then be processed further. The biomass may, according to requirement, be removed completely or partially from the fermentation broth by separation methods such as, for example, centrifugation, filtration, decanting or a combination of these methods or be left completely in said broth. It is advantageous to process the biomass after its separation.

[0086] However, the fermentation broth can also be thickened or concentrated without separating the cells, using known methods such as, for example, with the aid of a rotary evaporator, thin-film evaporator, falling-film evaporator, by reverse osmosis or by nanofiltration. Finally, this concentrated fermentation broth can be processed to obtain the fatty acids present therein.

[0087] The fatty acids obtained in the process are also suitable as starting material for the chemical synthesis of further products of interest. For example, they can be used in combination with one another or alone for the preparation of

pharmaceuticals, foodstuffs, animal feeds or cosmetics.

[0088] All of the nucleic acid sequences used in the process according to the invention are advantageously derived from a eukaryotic organism such as a plant, a microorganism or an animal. The nucleic acid sequences are preferably derived from the order Salmoniformes, algae such as Mantonella, Cryptocodinium, Euglena or Ostreococcus, fungi

such as the genus Phytophthora or from diatoms such as the genera Thalassiosira or Phaeodactylum.

[0089] The invention furthermore relates to isolated nucleic acid sequences encoding polypeptides with $\Delta 5$ -desaturase activity.

[0090] Therefore, in a further aspect of the invention there is provided an isolated nucleic acid sequence which encodes a polypeptide with $\Delta 5$ -desaturase activity and which is selected from the group consisting of:

a) SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5;

b) nucleic acid sequences which hybridize under stringent conditions with a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5.

c) an isolated nucleic acid sequence which encodes a polypeptide with $\Delta 5$ -desaturase activity, wherein the polypeptide is selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6;

d) A derivative of a nucleic acid sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6, wherein said polypeptide has $\Delta 5$ -desaturase activity.

[0091] In still another aspect of the invention there is provided an amino acid sequence which is encoded by a nucleic acid sequence in (a), (b), (c) or (d) above.

[0092] Advantageously, the proteins encoded by these nucleic acid molecules have at least approximately 50%, preferably at least approximately 60% and more preferably at least approximately 70%, 80% or 90% and most preferably at least approximately 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity with the amino acid sequences shown in SEQ ID NO: 2, SEQ ID NO: 4 or SEQ ID NO: 6.

[0093] The nucleic acid sequences used in the process are advantageously introduced into an expression cassette which makes possible the expression of the nucleic acids in organisms such as microorganisms or plants.

[0094] Therefore, in another aspect of the invention there is provided a gene construct comprising a nucleic acid sequence which encodes a polypeptide with $\Delta 5$ -desaturase activity as set out above, operably linked with one or more regulatory sequences.

[0095] In the expression cassette, the nucleic acid sequence which encode $\Delta 5$ -desaturase, is linked operably with one or more regulatory sequences, advantageously for enhancing gene expression. These regulatory sequences are intended to make possible the specific expression of the genes and proteins. Depending on the host organism, this may mean, for example, that the gene is expressed and/or overexpressed only after induction has taken place, or else that it is expressed and/or overexpressed immediately. For example, these regulatory sequences take the form of sequences to which inducers or repressors bind, thus controlling the expression of the nucleic acid. In addition to these novel regulatory sequences, or instead of these sequences, the natural regulatory elements of these sequences may still be present before the actual structural genes and, if appropriate, may have been genetically modified in such a way that their natural regulation is eliminated and the expression of the genes is enhanced. However, the expression cassette (= expression construct = gene construct) can also be simpler in construction, that is to say no additional regulatory signals have been inserted before the nucleic acid sequence or its derivatives, and the natural promoter together with its regulation was not removed. Instead, the natural regulatory sequence has been mutated in such a way that regulation no longer takes place and/or gene expression is enhanced. These modified promoters can also be positioned on their own before the natural gene in the form of part-sequences (= promoter with parts of the nucleic acid sequences used in accordance with the invention) in order to enhance the activity. Moreover, the gene construct may advantageously also comprise one or more what are known as enhancer sequences in operable linkage with the promoter, which make possible an enhanced expression of the nucleic acid sequence. Additional advantageous sequences, such as further regulatory elements or terminator sequences, may also be inserted at the 3' end of the DNA sequences. The $\Delta 12$ -desaturase, $\omega 3$ -desaturase, $\Delta 4$ -desaturase, $\Delta 5$ -desaturase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 5$ -elongase, $\Delta 6$ -elongase and/or $\Delta 9$ -elongase genes may be present in one or more copies of the expression cassette (= gene construct). Preferably, only one copy of the genes is present in each expression cassette. This gene construct or the gene constructs can be expressed together in the host organism. In this context, the gene construct(s) can be inserted in one or more vectors and be present in the cell in free form, or else be inserted in the genome. It is advantageous for the insertion of further genes in the genome when the genes to be expressed are present together in one gene construct.

[0096] In this context, the regulatory sequences or factors can, as described above, preferably have a positive effect on the gene expression of the genes introduced, thus enhancing it. Thus, an enhancement of the regulatory elements, advantageously at the transcriptional level, may take place by using strong transcription signals such as promoters and/or enhancers. In addition, however, enhanced translation is also possible, for example by improving the stability of

the mRNA.

[0097] The regulatory sequences include, in particular, plant sequences such as promoter and terminator sequences. The constructs can advantageously be stably propagated in microorganisms, in particular in *E. coli* and *Agrobacterium tumefaciens*, under selective conditions and make possible the transfer of heterologous DNA into plants or microorganisms.

[0098] Useful regulatory sequences are present, for example, in promoters such as the *cos*, *tac*, *trp*, *tet*, *trp-tet*, *lpp*, *lac*, *lpp-lac*, *lacIq*, T7, T5, T3, *gal*, *trc*, *ara*, SP6, λ -PR or λ -PL promoter and are advantageously employed in Gram-negative bacteria. Further advantageous regulatory sequences are, for example, present in the Gram-positive promoters *amy* and *SPO2*, in the yeast or fungal promoters *ADC1*, *MF α* , *AC*, *P-60*, *CYC1*, *GAPDH*, *TEF*, *rp28*, *ADH* or in the plant promoters *CaMV/35S* [Franck et al., *Cell* 21 (1980) 285-294], *PRP1* [Ward et al., *Plant. Mol. Biol.* 22 (1993)], *SSU*, *OCS*, *lib4*, *usp*, *STLS1*, *B33*, *nos* or in the ubiquitin or phaseolin promoter. Advantageous in this context are also inducible promoters, such as the promoters described in EP-A-0 388 186 (benzenesulfonamide-inducible), *Plant J.* 2, 1992: 397-404 (Gatz et al., tetracycline-inducible), EP-A-0 335 528 (abscissic acid-inducible) or WO 93/21334 (ethanol- or cyclohexenol-inducible) promoters. Further suitable plant promoters are the cytosolic FBPase promoter or the ST-LSI promoter of potato (Stockhaus et al., *EMBO J.* 8, 1989, 2445), the glycine max phosphoribosylpyrophosphate amidotransferase promoter (Genbank Accession No. U87999) or the node-specific promoter described in EP-A-0 249 676.

[0099] Especially advantageous promoters are promoters which make possible the expression in tissues which are involved in the biosynthesis of fatty acids. Very especially advantageous are seed-specific promoters, such as the USP promoter as described, but also other promoters such as the *LeB4*, *DC3*, phaseolin or napin promoter. Further especially advantageous promoters are seed-specific promoters which can be used for monocotyledonous or dicotyledonous plants and which are described in US 5,608,152 (oilseed rape napin promoter), WO 98/45461 (*Arabidopsis oleosin* promoter), **[0100]** US 5,504,200 (*Phaseolus vulgaris* phaseolin promoter), WO 91/13980 (*Brassica Bce4* promoter), by Bäumlein et al., *Plant J.*, 2, 1992:233-239 (*LeB4* promoter from a legume), these promoters being suitable for dicots. Examples of promoters which are suitable for monocots are the barley *lpt-2* or *lpt-1* promoter (WO 95/15389 and WO 95/23230), the barley hordein promoter and other suitable promoters described in WO 99/16890.

[0101] In principle, it is possible to use all natural promoters together with their regulatory sequences, such as those mentioned above. It is also possible and advantageous to use synthetic promoters, either in addition or alone, in particular when they mediate seed-specific expression, such as those described in WO 99/16890.

[0102] In order to achieve a particularly high $\Delta 5$ -unsaturated fatty acid content, especially in transgenic plants, the $\Delta 5$ -desaturase genes should advantageously be expressed in oil crops in a seed-specific manner. To this end, seed-specific promoters can be used, or those promoters which are active in the embryo and/or in the endosperm. In principle, seed-specific promoters can be isolated both from dicotyledonous and from monocotyledonous plants. Preferred promoters are listed hereinbelow: USP (= unknown seed protein) and vicilin (*Vicia faba*) [Bäumlein et al., *Mol. Gen. Genet.*, 1991, 225(3)], napin (oilseed rape) [US 5,608,152], acyl carrier protein (oilseed rape) [US 5,315,001 and WO 92/18634], oleosin (*Arabidopsis thaliana*) [WO 98/45461 and WO 93/20216], phaseolin (*Phaseolus vulgaris*) [US 5,504,200], *Bce4* [WO 91/13980], legumines B4 (*LegB4* promoter) [Bäumlein et al., *Plant J.*, 2,2, 1992], *Lpt2* and *Lpt1* (barley) [WO 95/15389 and WO95/23230], seed-specific promoters from rice, maize and wheat [WO 99/16890], *Amy32b*, *Amy 6-6* and *aleurain* [US 5,677,474], *Bce4* (oilseed rape) [US 5,530,149], glycinin (soybean) [EP 571 741], phosphoenol pyruvate carboxylase (soybean) [JP 06/62870], *ADR12-2* (soybean) [WO 98/08962], isocitrate lyase (oilseed rape) [US 5,689,040] or α -amylase (barley) [EP 781 849].

[0103] Plant gene expression can also be facilitated via a chemically inducible promoter (see review in Gatz 1997, *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 48:89-108). Chemically inducible promoters are particularly suitable when it is desired that gene expression should take place in a time-specific manner. Examples of such promoters are a salicylic-acid-inducible promoter (WO 95/19443), a tetracycline-inducible promoter (Gatz et al. (1992) *Plant J.* 2, 397-404) and an ethanol-inducible promoter.

[0104] To ensure the stable integration of the biosynthesis genes into the transgenic plant over a plurality of generations, each of the nucleic acids which encode $\Delta 2$ -desaturase, $\omega 3$ -desaturase, $\Delta 9$ -elongase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase, $\Delta 5$ -elongase and/or $\Delta 4$ -desaturase and which are used in the process should be expressed under the control of a separate promoter, preferably a promoter which differs from the other promoters, since repeating sequence motifs can lead to instability of the T-DNA, or to recombination events. In this context, the expression cassette is advantageously constructed in such a way that a promoter is followed by a suitable cleavage site, advantageously in a polylinker, for insertion of the nucleic acid to be expressed and, if appropriate, a terminator sequence is positioned behind the polylinker. This sequence is repeated several times, preferably three, four or five times, so that up to five genes can be combined in one construct and introduced into the transgenic plant in order to be expressed. Advantageously, the sequence is repeated up to three times. To express the nucleic acid sequences, the latter are inserted behind the promoter via a suitable cleavage site, for example in the polylinker. Advantageously, each nucleic acid sequence has its own promoter and, if appropriate, its own terminator sequence. Such advantageous constructs are disclosed, for example, in DE 101 02 337 or DE 101 02 338. However, it is also possible to insert a plurality of nucleic

acid sequences behind a promoter and, if appropriate, before a terminator sequence. Here, the insertion site, or the sequence, of the inserted nucleic acids in the expression cassette is not of critical importance, that is to say a nucleic acid sequence can be inserted at the first or last position in the cassette without its expression being substantially influenced thereby. Advantageously, different promoters such as, for example, the USP, LegB4 or DC3 promoter, and different terminator sequences can be used in the expression cassette. However, it is also possible to use only one type of promoter in the cassette. This, however, may lead to undesired recombination events.

[0105] As described above, the transcription of the genes which have been introduced should advantageously be terminated by suitable terminator sequences at the 3' end of the biosynthesis genes which have been introduced (behind the stop codon). An example of a sequence which can be used in this context is the OCS 1 terminator sequence. As is the case with the promoters, different terminator sequences should be used for each gene.

[0106] Advantageously, in particular for the process of the second aspect of the invention, the gene construct additionally comprises a nucleic acid sequence encoding a polypeptide having $\Delta 9$ -elongase activity, for example a sequence encoding the C18- $\Delta 9$ elongase from *Isochrysis galbana* (SEQ ID NO: 25) or *Acanthamoeba castellanii* (SEQ ID NO: 24)

[0107] The gene construct of the present invention may also comprise biosynthesis genes of the fatty acid or lipid metabolism selected from the group acyl-CoA dehydrogenase(s), acyl-ACP [= acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyltransferase(s), acyl-CoA:lysophospholipid acyltransferase(s), fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acylcoenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allenoxide synthases, hydroperoxide lyases or fatty acid elongase(s) and desaturase(s) such as $\Delta 4$ -desaturase, $\Delta 5$ -desaturase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 9$ -desaturase, $\Delta 12$ -desaturase or $\Delta 6$ -elongase.

[0108] These additional nucleic acids or genes can be cloned into the expression cassettes, which are then used for transforming plants with the aid of vectors such as *Agrobacterium*.

[0109] Here, the regulatory sequences or factors can, as described above, preferably have a positive effect on, and thus enhance, the expression genes which have been introduced. Thus, enhancement of the regulatory elements can advantageously take place at the transcriptional level by using strong transcription signals such as promoters and/or enhancers. However, an enhanced translation is also possible, for example by improving the stability of the mRNA. In principle, the expression cassettes can be used directly for introduction into the plants or else be introduced into a vector. Therefore, in yet another aspect of the invention, there is provided a vector comprising a nucleic acid or a gene construct in any of the aspects of the invention described above.

[0110] In one embodiment, the vector may be a cloning vector.

[0111] The nucleic acid sequences of the invention may be introduced alone, or preferably, in combination with an expression cassette (nucleic acid construct) into an organism. To introduce the nucleic acids, the latter are advantageously amplified and ligated in the known manner. Preferably, a procedure following the protocol for Pfu DNA polymerase or a Pfu/Taq DNA polymerase mixture is followed. The primers are selected taking into consideration the sequence to be amplified. The primers should advantageously be chosen in such a way that the amplificate comprises the entire codogenic sequence from the start codon to the stop codon. After the amplification, the amplificate is expediently analyzed. For example, a gel-electrophoretic separation can be carried out, which is followed by a quantitative and a qualitative analysis. Thereafter, the amplificate can be purified following a standard protocol (for example Qiagen). An aliquot of the purified amplificate is then available for the subsequent cloning step. Suitable cloning vectors are generally known to the skilled worker. These include, in particular, vectors which are capable of replication in microbial systems, that is to say mainly vectors which ensure efficient cloning in yeasts or fungi and which make possible the stable transformation of plants. Those which must be mentioned in particular are various binary and cointegrated vector systems which are suitable for the T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they comprise at least the vir genes required for the *Agrobacterium*-mediated transformation and the T-DNA-delimiting sequences (T-DNA border). These vector systems advantageously also comprise further cis-regulatory regions such as promoters and terminator sequences and/or selection markers, by means of which suitably transformed organisms can be identified. While in the case of cointegrated vector systems vir genes and T-DNA sequences are arranged on the same vector, binary systems are based on at least two vectors, one of which bears vir genes, but no T-DNA, while a second one bears T-DNA, but no vir gene. Owing to this fact, the last-mentioned vectors are relatively small, easy to manipulate and to replicate both in *E. coli* and in *Agrobacterium*. These binary vectors include vectors from the series pBIB-HYG, pPZP, pBecks, pGreen. In accordance with the invention, Bin19, pBI101, pBinAR, pGPTV and pCAMBIA are used by preference. An overview of the binary vectors and their use is found in Hellens et al, Trends in Plant Science (2000) 5, 446-451. In order to prepare the vectors, the vectors can first be linearized with restriction endonuclease(s) and then modified enzymatically in a suitable manner. Thereafter, the vector is purified, and an aliquot is employed for the cloning step. In the cloning step, the enzymatically cleaved and, if appropriate, purified amplificate is cloned with vector fragments which have been prepared in a similar manner, using ligase. In this context, a particular nucleic acid construct, or vector or plasmid construct, can have one or else more than one codogenic gene segment. The codogenic gene segments in these constructs are preferably linked operably with regulatory sequences. The regulatory sequences include, in par-

ticular, plant sequences such as the above-described promoters and terminator sequences. The constructs can advantageously be stably propagated in microorganisms, in particular in *E. coli* and *Agrobacterium tumefaciens*, under selective conditions and make possible the transfer of heterologous DNA into plants or microorganisms.

[0112] The nucleic acids used in the process, the inventive nucleic acids and nucleic acid constructs, can be introduced into organisms such as microorganisms or advantageously plants, advantageously using cloning vectors, and thus be used in the transformation of plants such as those which are published and cited in: *Plant Molecular Biology and Biotechnology* (CRC Press, Boca Raton, Florida), Chapter 6/7, p. 71-119 (1993); F.F. White, *Vectors for Gene Transfer in Higher Plants*; in: *Transgenic Plants, Vol. 1, Engineering and Utilization*, Ed.: Kung and R. Wu, Academic Press, 1993, 15-38; B. Jenes et al., *Techniques for Gene Transfer*, in: *Transgenic Plants, Vol. 1, Engineering and Utilization*, Ed.: Kung and R. Wu, Academic Press (1993), 128-143; Potrykus, *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42 (1991), 205-225. Thus, the nucleic acids, the inventive nucleic acids and nucleic acid constructs, and/or vectors used in the process can be used for the recombinant modification of a broad spectrum of organisms, advantageously plants, so that the latter become better and/or more efficient producers of $\Delta 5$ -unsaturated fatty acids.

[0113] A series of mechanisms exist by which a modification of the $\Delta 5$ -desaturase protein is possible, so that the yield, production and/or production efficiency of the advantageous $\Delta 5$ -unsaturated fatty acids in a plant, preferably in an oil crop plant or a microorganism, can be influenced directly owing to this modified protein. The number or activity of the proteins or genes can be increased, so that greater amounts of the gene products and, ultimately, greater amounts of the compounds of the general formula I are produced. A de novo synthesis in an organism which has lacked the activity and ability to biosynthesize the compounds prior to introduction of the corresponding gene(s) is also possible. This applies analogously to the combination with further desaturases or elongases or further enzymes of the fatty acid and lipid metabolism. The use of various divergent sequences, i.e. sequences which differ at the DNA sequence level, may also be advantageous in this context, or else the use of promoters for gene expression which make possible a different gene expression in the course of time, for example as a function of the degree of maturity of a seed or an oil-storing tissue.

[0114] Owing to the introduction of a $\Delta 5$ -desaturase gene into an organism, alone or in combination with other genes in a cell, it is not only possible to increase biosynthesis flux towards the end product, but also to increase, or to create de novo the corresponding triacylglycerol composition. Likewise, the number or activity of other genes which are involved in the import of nutrients which are required for the biosynthesis of one or more fatty acids, oils, polar and/or neutral lipids, can be increased, so that the concentration of these precursors, cofactors or intermediates within the cells or within the storage compartment is increased, whereby the ability of the cells to produce $\Delta 5$ -unsaturated fatty acids as described below is enhanced further. By optimizing the activity or increasing the number of one or more $\Delta 5$ -desaturase genes which are involved in the biosynthesis of these compounds, or by destroying the activity of one or more genes which are involved in the degradation of these compounds, an enhanced yield, production and/or efficiency of production of fatty acid and lipid molecules in organisms, advantageously in plants, is made possible.

[0115] Nucleic acids which can advantageously be used in the process are derived from bacteria, fungi, diatoms, animals such as *Caenorhabditis* or *Oncorhynchus* or plants such as algae or mosses, such as the genera *Shewanella*, *Physcomitrella*, *Thraustochytrium*, *Fusarium*, *Phytophthora*, *Ceratodon*, *Mantoniella*, *Ostreococcus*, *Isochrysis*, *Aleurita*, *Muscarioides*, *Mortierella*, *Borago*, *Phaeodactylum*, *Cryptocodinium*, specifically from the genera and species *Oncorhynchus mykiss*, *Xenopus laevis*, *Ciona intestinalis*, *Thalassiosira pseudonona*, *Mantoniella squamata*, *Ostreococcus* sp., *Ostreococcus tauri*, *Euglena gracilis*, *Physcomitrella patens*, *Phytophthora infestans*, *Fusarium gramineum*, *Cryptocodinium cohnii*, *Ceratodon purpureus*, *Isochrysis galbana*, *Acanthamoeba castellanii*, *Aleurita farinosa*, *Thraustochytrium* sp., *Muscarioides vialii*, *Mortierella alpina*, *Borago officinalis*, *Phaeodactylum tricornutum*, *Caenorhabditis elegans* or especially advantageously from *Oncorhynchus mykiss*, *Euglena gracilis*, *Thalassiosira pseudonona* or *Cryptocodinium cohnii*.

[0116] In an alternative embodiment, the vector may be an expression vector designed to transform an organism in which the nucleic acid is to be expressed and the compound of formula I is synthesized.

[0117] These advantageous vectors, preferably expression vectors, comprise the nucleic acids which encode the $\Delta 5$ -desaturase and which is used in the process of the first or second aspects

[0118] As used in the present context, the term "vector" refers to a nucleic acid molecule which is capable of transporting another nucleic acid to which it is bound. One type of vector is a "plasmid", a circular double-stranded DNA loop into which additional DNA segments can be ligated. A further type of vector is a viral vector, it being possible for additional DNA segments to be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they have been introduced (for example bacterial vectors with bacterial replication origin). Other vectors are advantageously integrated into the genome of a host cell when they are introduced into the host cell, and thus replicate together with the host genome. Moreover, certain vectors can govern the expression of genes with which they are in operable linkage. These vectors are referred to in the present context as "expression vectors". Usually, expression vectors which are suitable for DNA recombination techniques take the form of plasmids.

[0119] In the present description, where the term "plasmid" is used, it should be understood that plasmids can be substituted for other types of expression vector, such as viral vectors, which exert similar functions. Furthermore, the

term "vector" is also intended to comprise other vectors with which the skilled worker is familiar, such as phages, viruses such as SV40, CMV, TMV, transposons, IS elements, phasmids, phagemids, cosmids, linear or circular DNA.

[0120] The recombinant expression vectors advantageously used in the process comprise the nucleic acids described below or the above-described gene construct in a form which is suitable for expressing the nucleic acids used in a host cell, which means that the recombinant expression vectors comprise one or more regulatory sequences, selected on the basis of the host cells used for the expression, which regulatory sequence(s) is/are linked operably with the nucleic acid sequence to be expressed. In a recombinant expression vector, "linked operably" means that the nucleotide sequence of interest is bound to the regulatory sequence(s) in such a way that the expression of the nucleotide sequence is possible and they are bound to each other in such a way that both sequences carry out the predicted function which is ascribed to the sequence (for example in an in-vitro transcription/translation system, or in a host cell if the vector is introduced into the host cell). The term "regulatory sequence" is intended to comprise promoters, enhancers and other expression control elements (for example polyadenylation signals). These regulatory sequences are described, for example, in Goeddel: Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990), or see: Gruber and Crosby, in: Methods in Plant Molecular Biology and Biotechnology, CRC Press, Boca Raton, Florida, Ed.: Glick and Thompson, Chapter 7, 89-108, including the references cited therein. Regulatory sequences comprise those which govern the constitutive expression of a nucleotide sequence in many types of host cell and those which govern the direct expression of the nucleotide sequence only in specific host cells under specific conditions. The skilled worker knows that the design of the expression vector can depend on factors such as the choice of host cell to be transformed, the desired expression level of the protein and the like. The recombinant expression vectors used can be designed for the expression of $\Delta 5$ -desaturase in prokaryotic or eukaryotic cells. This is advantageous since intermediate steps of the vector construction are frequently carried out in microorganisms for the sake of simplicity. For example, the $\Delta 5$ -desaturase gene can be expressed in bacterial cells, insect cells (using Baculovirus expression vectors), yeast and other fungal cells (see Romanos, M.A., et al. (1992) "Foreign gene expression in yeast: a review", Yeast 8:423-488; van den Hondel, C.A.M.J.J., et al. (1991) "Heterologous gene expression in filamentous fungi", in: More Gene Manipulations in Fungi, J.W. Bennet & L.L. Lasure, Ed., pp. 396-428: Academic Press: San Diego; and van den Hondel, C.A.M.J.J., & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of Fungi, Peberdy, J.F., et al., Ed., pp. 1-28, Cambridge University Press: Cambridge), algae (Falciatore et al., 1999, Marine Biotechnology.1, 3:239-251), ciliates of the types: Holotrichia, Peritrichia, Spirotrichia, Suctorina, Tetrahymena, Paramecium, Colpidium, Glaucoma, Platyophrya, Potomacus, Desaturaseudocohnilembus, Euplotes, Engelmaniella and Stylonychia, in particular of the genus Stylonychia lemnae, using vectors in a transformation method as described in WO 98/01572 and, preferably, in cells of multi-celled plants (see Schmidt, R. and Willmitzer, L. (1988) "High efficiency Agrobacterium tumefaciens-mediated transformation of Arabidopsis thaliana leaf and cotyledon explants" Plant Cell Rep.:583-586; Plant Molecular Biology and Biotechnology, C Press, Boca Raton, Florida, Chapter 6/7, pp.71-119 (1993); F.F. White, B. Jenens et al., Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press (1993), 128-43; Potrykus, Annu. Rev. Plant Physiol. Plant Molec. Biol. 42 (1991), 205-225 (and references cited therein)). Suitable host cells are furthermore discussed in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). As an alternative, the recombinant expression vector can be transcribed and translated in vitro, for example using T7-promoter regulatory sequences and T7-polymerase.

[0121] In most cases, the expression of proteins in prokaryotes involves the use of vectors comprising constitutive or inducible promoters which govern the expression of fusion or nonfusion proteins. Typical fusion expression vectors are, inter alia, pGEX (Pharmacia Biotech Inc; Smith, D.B., and Johnson, K.S. (1988) Gene 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ), where glutathione S-transferase (GST), maltose-E binding protein and protein A, respectively, is fused with the recombinant target protein.

[0122] Examples of suitable inducible nonfusion E. coli expression vectors are, inter alia, pTrc (Amann et al. (1988) Gene 69:301-315) and pET 11 d (Studier et al., Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, California (1990) 60-89). The target gene expression from the pTrc vector is based on the transcription from a hybrid trp-lac fusion promoter by the host RNA polymerase. The target gene expression from the vector pET 11 d is based on the transcription of a T7-gn10-lac fusion promoter, which is mediated by a viral RNA polymerase (T7 gn1), which is coexpressed. This viral polymerase is provided by the host strains BL21 (DE3) or HMS174 (DE3) from a resident λ -prophage which harbors a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

[0123] Other vectors which are suitable for prokaryotic organisms are known to the skilled worker, these vectors are, for example in E. coli pLG338, pACYC184, the pBR series such as pBR322, the pUC series such as pUC18 or pUC19, the M113mp series, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III113-B1, λ gt11 or pBdCl, in Streptomyces pIJ101, pIJ364, pIJ702 or pIJ361, in Bacillus pUB110, pC194 or pBD214, in Corynebacterium pSA77 or pAJ667.

[0124] In a further embodiment, the expression vector is a yeast expression vector. Examples for vectors for expression in the yeast *S. cerevisiae* comprise pYeDesaturasec1 (Baldari et al. (1987) Embo J. 6:229-234), pMFa (Kurjan and

Herskowitz (1982) Cell 30:933-943), pJRY88 (Schultz et al. (1987) Gene 54:113-123) and pYES2 (Invitrogen Corporation, San Diego, CA). Vectors and processes for the construction of vectors which are suitable for use in other fungi, such as the filamentous fungi, comprise those which are described in detail in: van den Hondel, C.A.M.J.J., & Punt, P.J. (1991) "Gene transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of fungi, J.F. Peberdy et al., Ed., pp. 1-28, Cambridge University Press: Cambridge, or in: More Gene Manipulations in Fungi [J.W. Bennet & L.L. Lasure, Ed., pp. 396-428: Academic Press: San Diego]. Further suitable yeast vectors are, for example, pAG-1, YEp6, YEp13 or pEMBLye23.

[0125] As an alternative, $\Delta 5$ -desaturase can be expressed in insect cells using Baculovirus vectors. Baculovirus expression vectors which are available for the expression of proteins in cultured insect cells (for example Sf9 cells) comprise the pAc series (Smith et al. (1983) Mol. Cell Biol. 3:2156-2165) and the pVL series (Lucklow and Summers (1989) Virology 170:31-39).

[0126] The abovementioned vectors are only a small overview over suitable vectors which are possible. Further plasmids are known to the skilled worker and are described, for example, in: Cloning Vectors (Ed. Pouwels, P.H., et al., Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). For further suitable expression systems for prokaryotic and eukaryotic cells, see the Chapters 16 and 17 in Sambrook, J., Fritsch, E.F., and Maniatis, T., Molecular Cloning: A Laboratory Manual, 2. edition, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

[0127] In a further embodiment of the process, the $\Delta 5$ -desaturase can be expressed in single-celled plant cells (such as algae), see Falcatore et al., 1999, Marine Biotechnology 1 (3):239-251 and references cited therein, and in plant cells from higher plants (for example spermatophytes such as arable crops). Examples of plant expression vectors comprise those which are described in detail in: Becker, D., Kemper, E., Schell, J., and Masterson, R. (1992) "New plant binary vectors with selectable markers located proximal to the left border", Plant Mol. Biol. 20:1195-1197; and Bevan, M.W. (1984) "Binary Agrobacterium vectors for plant transformation", Nucl. Acids Res. 12:8711-8721; Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, p. 15-38.

[0128] A plant expression cassette preferably comprises regulatory sequences which are capable of governing the expression of genes in plant cells and which are linked operably so that each sequence can fulfill its function, such as transcriptional termination, for example polyadenylation signals. Preferred polyadenylation signals are those which are derived from *Agrobacterium tumefaciens* T-DNA, such as gene 3 of the Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3 (1984) 835 et seq.), which is known as octopine synthase, or functional equivalents thereof, but all other terminator sequences which are functionally active in plants are also suitable.

[0129] Since plant gene expression is very often not limited to the transcriptional level, a plant expression cassette preferably comprises other sequences which are linked operably, such as translation enhancers, for example the overdrive sequence, which enhances the tobacco mosaic virus 5' - untranslated leader sequence, which increases the protein/RNA ratio (Gallie et al., 1987, Nucl. Acids Research 15:8693-8711).

[0130] As described above, plant gene expression must be linked operably with a suitable promoter which triggers gene expression with the correct timing or in a cell- or tissue-specific manner. Utilizable promoters are constitutive promoters (Benfey et al., EMBO J. 8 (1989) 2195-2202), such as those which are derived from plant viruses, such as 35S CaMV (Franck et al., Cell 21 (1980) 285-294), 19S CaMV (see also US 5352605 and WO 84/02913), or plant promoters, such as the promoter of the Rubisco subunit, which is described in US 4,962,028.

[0131] Other preferred sequences for use in operable linkage in plant gene expression cassettes are targeting sequences, which are required for steering the gene product into its corresponding cell compartment (see a review in Kermode, Crit. Rev. Plant Sci. 15, 4 (1996) 285-423 and references cited therein), for example into the vacuole, into the nucleus, all types of plastids, such as amyloplasts, chloroplasts, chromoplasts, the extracellular space, the mitochondria, the endoplasmic reticulum, elaioplasts, peroxisomes and other compartments of plant cells.

[0132] As described above, plant gene expression can also be achieved via a chemically inducible promoter (see review in Gatz 1997, Annu. Rev. Plant Physiol. Plant Mol. Biol., 48:89-108). Chemically inducible promoters are particularly suitable when it is desired that the gene expression takes place in a time-specific manner. Examples of such promoters are a salicylic-acid-inducible promoter (WO 95/19443), a tetracyclin-inducible promoter (Gatz et al. (1992) Plant J. 2, 397-404) and an ethanol-inducible promoter.

[0133] Promoters which respond to biotic or abiotic stress conditions are also suitable, for example the pathogen-induced PRP1 gene promoter (Ward et al., Plant Mol. Biol. 22 (1993) 361-366), the heat-inducible tomato hsp80 promoter (US 5,187,267), the chill-inducible potato alpha-amylase promoter (WO 96/12814) or the wound-inducible pinII promoter (EP-A-0 375 091).

[0134] Especially preferred are those promoters which bring about the gene expression in tissues and organs in which the biosynthesis of fatty acids, lipids and oils takes place, in seed cells, such as cells of the endosperm and of the developing embryo. Suitable promoters are the oilseed rape napin promoter (US 5,608,152), the *Vicia faba* USP promoter (Baeumlein et al., Mol Gen Genet, 1991, 225 (3):459-67), the *Arabidopsis* oleosin promoter (WO 98/45461), the Phaseolus

vulgaris phaseolin promoter (US 5,504,200), the Brassica Bce4 promoter (WO 91/13980) or the legumine B4 promoter (LeB4; Baeumlein et al., 1992, Plant Journal, 2 (2):233-9), and promoters which bring about the seed-specific expression in monocotyledonous plants such as maize, barley, wheat, rye, rice and the like. Suitable noteworthy promoters are the barley lpt2 or lpt1 gene promoter (WO 95/15389 and WO 95/23230) or the promoters from the barley hordein gene, the rice glutelin gene, the rice oryzin gene, the rice prolamine gene, the wheat gliadine gene, the wheat glutelin gene, the maize zeine gene, the oat glutelin gene, the sorghum kasirin gene or the rye secalin gene, which are described in WO 99/16890.

[0135] For the process of the second aspect, it may be desired to bring about the multiparallel expression of the $\Delta 5$ -desaturase and the $\Delta 9$ -elongase used in the process. Such expression cassettes can be introduced via the simultaneous transformation of a plurality of individual expression constructs or, preferably, by combining a plurality of expression cassettes on one construct. Also, a plurality of vectors can be transformed with in each case a plurality of expression cassettes and then transferred into the host cell.

[0136] Other promoters which are likewise especially suitable are those which bring about a plastid-specific expression, since plastids constitute the compartment in which the precursors and some end products of lipid biosynthesis are synthesized. Suitable promoters, such as the viral RNA polymerase promoter, are described in WO 95/16783 and WO 97/06250, and the clpP promoter from Arabidopsis, described in WO 99/46394.

[0137] Vector DNA can be introduced into prokaryotic and eukaryotic cells via conventional transformation or transfection techniques. The terms "transformation" and "transfection", conjugation and transduction, as used in the present context, are intended to comprise a multiplicity of methods known in the prior art for the introduction of foreign nucleic acid (for example DNA) into a host cell, including calcium phosphate or calcium chloride coprecipitation, DEAE-dextran-mediated transfection, lipofection, natural competence, chemically mediated transfer, electroporation or particle bombardment. Suitable methods for the transformation or transfection of host cells, including plant cells, can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual., 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) and other laboratory textbooks such as Methods in Molecular Biology, 1995, Vol. 44, Agrobacterium protocols, Ed.: Gartland and Davey, Humana Press, Totowa, New Jersey.

[0138] In a further aspect of the invention there is provided a transgenic non human organism comprising at least one nucleic acid, gene construct or vector according to a previous aspect of the invention.

[0139] The transgenic nonhuman organism may be a microorganism, a nonhuman animal or a plant.

[0140] Host cells which are suitable in principle for taking up the nucleic acid according to the invention, the gene product according to the invention or the vector according to the invention are all prokaryotic or eukaryotic organisms. The host organisms which are advantageously used are microorganisms such as fungi or yeasts, or plant cells, preferably plants or parts thereof. Fungi, yeasts or plants are preferably used, especially plants, for example plants such as oil crops, which are high in lipid compounds, such as oilseed rape, evening primrose, hemp, thistle, peanut, canola, linseed, soybean, safflower, sunflower, borage, or plants such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, Tagetes, Solanacea plants such as potato, tobacco, eggplant and tomato, Vicia species, pea, alfalfa, bushy plants (coffee, cacao, tea), Salix species, trees (oil palm, coconut), and perennial grasses and fodder crops. Especially preferred plants according to the invention are oil crops such as soybean, peanut, oilseed rape, canola, linseed, hemp, evening primrose, sunflower, safflower, trees (oil palm, coconut).

[0141] In an advantageous embodiment, the term "nucleic acid (molecule)" as used in the present context additionally comprises the untranslated sequence at the 3' and at the 5' end of the coding gene region: at least 500, preferably 200, especially preferably 100 nucleotides of the sequence upstream of the 5' end of the coding region and at least 100, preferably 50, especially preferably 20 nucleotides of the sequence downstream of the 3' end of the coding gene region. An "isolated" nucleic acid molecule is separate from other nucleic acid molecules which are present in the natural source of the nucleic acid. An "isolated" nucleic acid preferably has no sequences which naturally flank the nucleic acid in the genomic DNA of the organism from which the nucleic acid is derived (for example sequences which are located at the 5' and 3' ends of the nucleic acid). In various embodiments, the isolated $\Delta 12$ -desaturase, $\omega 3$ -desaturase, $\Delta 9$ -elongase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 6$ -elongase, $\Delta 5$ -desaturase, $\Delta 5$ -elongase or $\Delta 4$ -desaturase molecule can comprise for example fewer than approximately 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in the genomic DNA of the cell from which the nucleic acid is derived.

[0142] The nucleic acid molecules of the present invention, for example a nucleic acid molecule with a nucleotide sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 or of a part thereof can be isolated using molecular-biological standard techniques and the sequence information provided herein. Also, for example a homologous sequence or homologous, conserved sequence regions can be identified at the DNA or amino acid level with the aid of comparative algorithms. They can be used as hybridization probe and standard hybridization techniques (such as, for example, those described in Sambrook et al., Molecular Cloning: A Laboratory Manual. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) for isolating further nucleic acid sequences which can be used in the process.

[0143] Moreover, a nucleic acid molecule comprising a complete sequence of SEQ ID NO: 1, SEQ ID NO: 3 or SEQ

ID NO: 5 or a part thereof can be isolated by polymerase chain reaction, where oligonucleotide primers which are used on the basis of this sequence or parts thereof (for example a nucleic acid molecule comprising the complete sequence or part thereof can be isolated by polymerase chain reaction using oligonucleotide primers which have been generated based on this same sequence). For example, mRNA can be isolated from cells (for example by means of the guanidinium thiocyanate extraction method of Chirgwin et al. (1979) *Biochemistry* 18:5294-5299) and cDNA by means of reverse transcriptase (for example Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD, or AMV reverse transcriptase, available from Seikagaku America, Inc., St.Petersburg, FL).

[0144] Synthetic oligonucleotide primers for the amplification by means of polymerase chain reaction can be generated based on one of the sequences shown in SEQ ID NO: 1, SEQ ID NO: 3 or SEQ ID NO: 5 or with the aid of the amino acid sequences detailed in SEQ ID NO: 2, SEQ ID NO: 4 and SEQ ID NO: 6.

[0145] A nucleic acid according to the invention can be amplified by standard PCR amplification techniques using cDNA or, alternatively, genomic DNA as template and suitable oligonucleotide primers. The nucleic acid amplified thus can be cloned into a suitable vector and characterized by means of DNA sequence analysis. Oligonucleotides which correspond to a desaturase nucleotide sequence can be generated by standard synthetic methods, for example using an automatic DNA synthesizer.

[0146] The abovementioned nucleic acids and protein molecules with $\Delta 5$ -desaturase activity are used in the process according to the invention for the modulation of the production of $\Delta 5$ -unsaturated fatty acids in transgenic organisms, advantageously in plants, such as maize, wheat, rye, oats, triticale, rice, barley, soybean, peanut, cotton, *Linum* species such as linseed or flax, Brassica species such as oilseed rape, canola and turnip rape, pepper, sunflower, borage, evening primrose and Tagetes, Solanaceae plants such as potato, tobacco, eggplant and tomato, Vicia species, pea, cassava, alfalfa, bushy plants (coffee, cacao, tea), Salix species, trees (oil palm, coconut) and perennial grasses and fodder crops, either directly (for example when the overexpression or optimization of a fatty acid biosynthesis protein has a direct effect on the yield, production and/or production efficiency of the fatty acid from modified organisms) and/or can have an indirect effect which nevertheless leads to an enhanced yield, production and/or production efficiency of the $\Delta 5$ -unsaturated fatty acids or a reduction of undesired compounds (for example when the modulation of the metabolism of lipids and fatty acids, cofactors and enzymes lead to modifications of the yield, production and/or production efficiency or the composition of the desired compounds within the cells, which, in turn, can affect the production of one or more fatty acids).

[0147] The combination of various precursor molecules and biosynthesis enzymes leads to the production of various fatty acid molecules, which has a decisive effect on lipid composition, since $\Delta 5$ -unsaturated fatty acids are not only incorporated into triacylglycerol but also into membrane lipids.

[0148] Brassicaceae, Boraginaceae, Primulaceae, or Linaceae are particularly suitable for the production of PUFAs, for example stearidonic acid, eicosapentaenoic acid and docosahexaenoic acid. Linseed (*Linum usitatissimum*) is especially advantageously suitable for the production of PUFAs with the nucleic acid sequences according to the invention, advantageously, as described, in combination with further desaturases and elongases.

[0149] Lipid synthesis can be divided into two sections: the synthesis of fatty acids and their binding to sn-glycerol-3-phosphate, and the addition or modification of a polar head group. Usual lipids which are used in membranes comprise phospholipids, glycolipids, sphingolipids and phosphoglycerides. Fatty acid synthesis starts with the conversion of acetyl-CoA into malonyl-CoA by acetyl-CoA carboxylase or into acetyl-ACP by acetyl transacylase. After condensation reaction, these two product molecules together form acetoacetyl-ACP, which is converted via a series of condensation, reduction and dehydration reactions so that a saturated fatty acid molecule with the desired chain length is obtained. The production of the unsaturated fatty acids from these molecules is catalyzed by specific desaturases, either aerobically by means of molecular oxygen or anaerobically (regarding the fatty acid synthesis in microorganisms, see F.C. Neidhardt et al. (1996) *E. coli* and *Salmonella*. ASM Press: Washington, D.C., pp. 612-636 and references cited therein; Lengeler et al. (Ed.) (1999) *Biology of Prokaryotes*. Thieme: Stuttgart, New York, and the references therein, and Magnuson, K., et al. (1993) *Microbiological Reviews* 57:522-542 and the references therein). To undergo the further elongation steps, the resulting phospholipid-bound fatty acids must be returned to the fatty acid CoA ester pool. This is made possible by acyl-CoA:lysophospholipid acyltransferases. Moreover, these enzymes are capable of transferring the elongated fatty acids from the CoA esters back to the phospholipids. If appropriate, this reaction sequence can be followed repeatedly.

[0150] Examples of precursors for the biosynthesis of sciadonic acid and juniperonic acid are linoleic acid and linolenic acid. The C18-carbon fatty acids must be elongated to C20 in order to obtain the required fatty acids and this can be achieved with a $\Delta 9$ -elongase. With the aid of the $\Delta 5$ -desaturases of the present invention, sciadonic acid and juniperonic acid can be produced and subsequently employed in various applications regarding foodstuffs, feedstuffs, cosmetics or pharmaceuticals.

[0151] The synthesized $\Delta 5$ -unsaturated fatty acids are obtained in the process according to the invention in the form of the free fatty acid or in the form of their esters, for example in the form of their glycerides.

[0152] The term "glyceride" is understood as meaning glycerol esterified with one, two or three carboxyl radicals (mono-, di- or triglyceride). "Glyceride" is also understood as meaning a mixture of various glycerides. The glyceride or

glyceride mixture may comprise further additions, for example free fatty acids, antioxidants, proteins, carbohydrates, vitamins and/or other substances.

[0153] For the purposes of the invention, a "glyceride" is furthermore understood as meaning glycerol derivatives. In addition to the above-described fatty acid glycerides, these also include glycerophospholipids and glyceroglycolipids. Preferred examples which may be mentioned in this context are the glycerophospholipids such as lecithin (phosphatidylcholine), cardiolipin, phosphatidylglycerol, phosphatidylserine and alkylacylglycerophospholipids.

[0154] Furthermore, fatty acids must subsequently be translocated to various modification sites and incorporated into the triacylglycerol storage lipid. A further important step in lipid synthesis is the transfer of fatty acids to the polar head groups, for example by glycerol fatty acid acyltransferase (see Frentzen, 1998, *Lipid*, 100(4-5):161-166). Publications on plant fatty acid biosynthesis and on the desaturation, the lipid metabolism and the membrane transport of lipidic compounds, on beta-oxidation, fatty acid modification and cofactors, triacylglycerol storage and triacylglycerol assembly, including the references therein, see the following papers: Kinney, 1997, *Genetic Engineering*, Ed.: JK Setlow, 19: 149-166; Ohlrogge and Browse, 1995, *Plant Cell* 7:957-970; Shanklin and Cahoon, 1998, *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 49:611-641; Voelker, 1996, *Genetic Engineering*, Ed.: JK Setlow, 18:111-13; Gerhardt, 1992, *Prog. Lipid R.* 31:397-417; Gühnemann-Schäfer & Kindl, 1995, *Biochim. Biophys Acta* 1256:181-186; Kunau et al., 1995, *Prog. Lipid Res.* 34:267-342; Szymne et al., 1993, in: *Biochemistry and Molecular Biology of Membrane and Storage Lipids of Plants*, Ed.: Murata and Somerville, Rockville, American Society of Plant Physiologists, 150-158, Murphy & Ross 1998, *Plant Journal*. 13(1):1-16.

[0155] Phospholipids for the purposes of the invention are understood as meaning phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine, phosphatidylglycerol and/or phosphatidylinositol, advantageously phosphatidylcholine. The terms production or productivity are known in the art and comprise the concentration of the fermentation product (compounds of the formula I) which is formed within a specific period of time and in a specific fermentation volume (for example kg of product per hour per liter). It also comprises the productivity within a plant cell or a plant, that is to say the content of the desired fatty acids produced in the process relative to the content of all fatty acids in this cell or plant. The term production efficiency comprises the time required for obtaining a specific production quantity (for example the time required by the cell to establish a certain throughput rate of a fine chemical). The term yield or product/carbon yield is known in the art and comprises the efficiency of the conversion of the carbon source into the product (i.e. the fine chemical). This is usually expressed for example as kg of product per kg of carbon source. By increasing the yield or production of the compound, the amount of the molecules obtained of this compound, or of the suitable molecules of this compound obtained in a specific culture quantity over a specified period of time is increased. The terms biosynthesis or biosynthetic pathway are known in the art and comprise the synthesis of a compound, preferably an organic compound, by a cell from intermediates, for example in a multi-step and strongly regulated process. The terms catabolism or catabolic pathway are known in the art and comprise the cleavage of a compound, preferably of an organic compound, by a cell to give catabolites (in more general terms, smaller or less complex molecules), for example in a multi-step and strongly regulated process. The term metabolism is known in the art and comprises the totality of the biochemical reactions which take place in an organism. The metabolism of a certain compound (for example the metabolism of a fatty acid) thus comprises the totality of the biosynthetic pathways, modification pathways and catabolic pathways of this compound in the cell which relate to this compound.

[0156] Owing to their homology to the $\Delta 5$ -desaturase, nucleic acids disclosed here, nucleic acid molecules which are advantageous for the process according to the invention can be isolated following standard hybridization techniques under stringent hybridization conditions, using the sequences or part thereof as hybridization probe. In this context it is possible, for example, to use isolated nucleic acid molecules which are least 15 nucleotides in length and which hybridize under stringent conditions with the nucleic acid molecules which comprise a nucleotide sequence of

[0157] The invention will now be described in greater detail with reference to the following Examples and to the drawings in which:

Figure 1 is a comparison of the deduced amino acid sequences of *A. leveillei* and *C. palustris* desaturases AL10, AL21 and Cal2 with related sequences.

Figure 2 is a GC-MS analysis of fatty acid methyl esters derived from leaf tissue of wildtype *Arabidopsis* (A) or transgenic lines, expressing *A. leveillei* acyl-CoA desaturase AL21 (B). The presence of additional peaks is indicated with arrows. The figure shows that AL21 inserts $\Delta 5$ -double bonds into a range of substrates

Figure 3 shows mass spectral identification of $\Delta 5$ -fatty acids from transgenic *Arabidopsis* expressing *A. leveillei* acyl-CoA desaturase AL21: a) 18:1 $\Delta 5$; b) authentic compound; c) 16:1 $\Delta 5$; d) 18:3 $\Delta 5,9,12$; e). authentic fatty acid derivative from a *Pinus* sp. Diagnostic ions are starred.

Figure 4 is a comparison of Acyl-CoA profiles of transgenic *Arabidopsis* expressing *A. leveillei* acyl-CoA desaturase AL21. Acyl-CoA pools of *Arabidopsis* leaves from wild-type plants (A) and AL21 transgenic plants (B) were analysed by HPLC. The presence of additional peaks is indicated with stars. The internal standard is 17:0. The figure shows that desaturation products (*) are present in the acyl-CoA pool.

Figure 5 is a series of GC profiles showing the expression of *A. leveillei* acyl-CoA desaturases in transgenic *Arabidopsis*. Total fatty acid methyl esters derived from leaf tissue of wildtype *Arabidopsis* (a), $\Delta 9$ elongase line CA1-9 (b), double-transgenics CA1-9+ AL21 (C) or CA1-9 + AL10. C20 elongated products are marked with a star, $\Delta 5$ -unsaturated fatty acids are indicated with solid arrows. The figure demonstrates that co-expression with $\Delta 9$ elongase reveals the function of A10 $\Delta 5$ -desaturase.

Figure 6 shows MS identification of 20:35, 11, 14 and 20:45, 11, 14, 17 from transgenic *Arabidopsis* plants co-expressing the *A. leveillei* acyl-CoA desaturases A10 and A121 and IgASE1.

Figure 7 is a series of Acyl-CoA profiles of transgenic *Arabidopsis* co-expressing *A. leveillei* acyl-CoA desaturase AL21 and IgASE1. Acyl-CoA pools of *Arabidopsis* leaves from $\Delta 9$ elongase line CA1-9 (A) and double-transgenic CA1-9 + AL10 plants (B) were analysed by HPLC. The presence of additional peaks is indicated with stars. The internal standard is 17:0. $\Delta 5$ -unsaturated fatty acids are indicated with solid arrows. The figure shows that C20 NMIFAs are present in the acyl-CoA pool of transgenic plants co-expressing A10 and IgELO9.

Figure 8 Functional expression in yeast in the presence of exogenous substrates.

FIGURE 9a is a GC profiles of an *A. thaliana* leaf fatty acid methyl esters, extracted from a double transgenic plant expressing the *A. leveillei* $\Delta 5$ -desaturase A121 and *I. galbana* $\Delta 9$ -elongase.

FIGURE 9b is a GC profiles of an *A. thaliana* leaf fatty acid methyl esters, extracted from a single transgenic plant expressing the *I. galbana* $\Delta 9$ -elongase.

FIGURE 10 shows an acyl CoA profile of an *Arabidopsis* single transgenic plant expressing IgASE1. The accumulation of 20:2 and 20:3 is observed, representing the elongation of LA and ALA, and the failure of these elongation products to accumulate in glycerolipids.

FIGURE 11 shows an acyl CoA profile of a double transgenic plant expressing IgASE1 and *A. leveillei* $\Delta 5$ -desaturase A121.

[0158] For reasons of clarity:

A110 shall be identical to A1Coa10

A121 shall be identical to A1Coa21

Cal2 shall be identical to CalCoa2

Example 1

Materials and Methods.

Plant material.

[0159] Seeds were obtained from Chiltern Seeds, Cumbria, UK.

Nucleic acid manipulation.

[0160] Total RNA was isolated from developing seeds of *Anemone leveillei* and *Caltha palustris* using the RNeasy kit (Qiagen).

PCR-based cloning.

[0161] Two degenerate primers were designed to conserved amino acid sequences corresponding to the histidine boxes and identified in previously characterized acyl-CoA desaturases: forward primer Des2F:

5'- TGGGTI(A/T)(G/C)IA(C/T)ICA(T/C)(C/A)GITA(T/C)CA(T/C)-

[0162] (SEQ ID NO 7) designed to encode histidine box II (WVSTHRYHHQF) and reverse primer DesR:

5'- GC(A/G)TG(A/G)TG(A/G)TT(A/G)TT(A/G)TGCCAICC(T/C)TCICC,

[0163] (SEQ ID NO: 8) designed to encode the complement of histidine box III (GEGWHNNHHA), where I substitutes inosine.

[0164] These primers were used for PCR amplification with cDNAs transcribed from total RNA isolated from developing seeds of *Anemone leveillei* and *Caltha palustris* using previously described protocol (Sayanova et al, PNAS, 1997; 94: 4211-4216). The PCR fragments of the expected length (420 bp) were cloned into TOPO TA vector (Invitrogen) and

EP 2 388 332 A1

sequenced. Data bank searches and alignments with these new sequences showed similarities to acyl-CoA desaturases for two different fragments from *A. leveillei* and one from *C. palustris*. The derived sequence data were used to design primers for the amplification of 5'- and 3' ends of putative desaturases using a SMART RACE cDNA Amplification kit (BD-Clontech).

5 **[0165]** The resulting 5'- and 3'- RACE products were used to amplify full-length copies of the two putative desaturases from cDNAs of *Anemone leveillei* designated AI10 (SEQ ID NO: 5) and AI21 (SEQ ID NO: 3) respectively and one putative desaturases from cDNAs of *Caltha palustris* designated Cal2 (SEQ ID NO: 1). Gene -specific primers were designed to the 5'- and 3' -ends of the coding regions of the corresponding desaturase sequences with restriction sites to facilitate cloning into plant vectors. The following pairs of forward/reverse (F/R) primers were used (restriction sites are indicated in bold).

10 **[0166]** Cal2:

SalF: GGTTCGACATGGCTCTAATTGCAACAACCCCCAAA (SEQ ID NO: 9);
ClaF: GATCGATATGGCTCTAATTGCAACAACCCCCAAA (SEQ ID NO: 10);
15 SacR: GAGCTCTTACTCGCTGAAACACATCC (SEQ ID NO: 11);
ClaR: GATCGATTTACTCGCTGAAACACATCC (SEQ ID NO: 12);

[0167] AI10

20 XbaF: GGTCTAGAATGGATCTCACATCAATGG (SEQ ID NO: 13);
NotF: GGCGGCCGCATGGATCTCACATCAATGG (SEQ ID NO: 14);
ClaF: GATCGATATGGATCTCACATCAATGG (SEQ ID NO: 15);
SalR: GGTCTGACTCAATTTTTGAAAGACATCTTACGCTTG (SEQ ID NO: 16);
25 NotR: GGCGGCCGCCTCAATTTTTGAAAGACATCTTACGCTTG (SEQ ID NO: 17).

[0168] AI21

30 SmaF: GGCCCGGGATGGAAGTCCCAGCGAT (SEQ ID NO: 18);
NotF: GGCGGCCGCATGGAACTCCCAGCGAT (SEQ ID NO: 19);
ClaF: GATCGATATGGAAGTCCCAGCGAT (SEQ ID NO: 20);
SalR: GGTCTGACTTACTTGTCTGTTTACACAGAAC (SEQ ID NO: 21);
NotR: GGCGGCCGCCTTACTTGTCTGTTTACACAGAAC (SEQ ID NO: 22);
ClaR: GATCGATTTACTTGTCTGTTTACACAGAAC (SEQ ID NO: 23).

35 Functional expression in yeast.

[0169] All coding regions corresponding to *A. leveillei* desaturases were inserted as Cla1/Cla1 fragments into the yeast pYES2.1 TOPO TA expression vector. Coding region corresponding to *Isochrysis galbana* C18-Δ9-elongase, ASE1, was inserted as a Kpn1-BamH1 fragment into the pYES3 expression vector. Forward primers were designed to contain a G at position -3 and +4 to improve translation initiation in eukaryotic cells. ORFs encoding putative desaturation activities were introduced in *S. cerevisiae* strain W303-1A by a lithium acetate method as previously described (Sayanova et al., 2003, FEBS Letters, 542,100-104). Cultures were grown at 22°C in the presence of 2% (v/v) raffinose and expression of the transgenes was induced by the addition of galactose to 2% (w/v) in the presence of 0.5 mM of the corresponding fatty acid and 1% (w/v) tergitol -Nonidet P-40 (Sigma) as described. Yeast transformants containing pYES2-derived constructs were grown on synthetic minimal medium minus uracil; pYES3-derived constructs were grown on minimal medium minus tryptophan. Co-transformed yeast was grown on minimal medium minus uracil and tryptophan.

Plant transformation constructs.

50 **[0170]** All coding regions used were placed in CaMV 35S promoter- nos terminator expression cassettes. The coding regions of AI10 and AI21 were inserted as XbaI/SalI-and SmaI/SalI fragments respectively into the binary vector pBIN19-35S, kindly provided by Dr. P. Buchner.

[0171] The coding region of CalCoa2 was inserted as a SalI/SacI fragment into the vector pJD330 and excised with BamHI/XbaI for insertion into pBin19. For seed specific expression both coding regions of AlCoa10 and AlCoa21 were inserted as NotI fragments into expression vector pKMS2 (donated by Dr. E. Cahoon).

Plant transformation.

[0172] Binary plasmids were transferred to *Agrobacterium tumefaciens* strain GV3101 by electroporation and kanamycin-resistant colonies were selected. Wild type *Arabidopsis thaliana* ecotype Columbia plants and transgenic *Arabidopsis* plants harbouring the $\Delta 9$ -specific elongase from *Isochrysis galbana*, IgASE1 (Qi et al. 2004 *Nature Biotechnology* 22: 739-745), were transformed by the floral dipping method (Clough and Bent, 1998, *Plant Journal* 16: 735-743). Kanamycin-resistant plants for each construct were selected and transplanted to soil and analysed.

Fatty acid analysis.

[0173] Fatty acids were extracted and methylated as described (Sayanova et al., 2003, *FEBS Letters*, 542,100-104). Methyl ester derivatives of total fatty acids extracted from leaves were analysed by GC and GC-MS. For the determination of the double bond positions fatty acids methyl esters were converted to the 4,4-dimethyloxazoline (DMOX) derivatives (Fay and Richli, 1991, *Journal of Chromatography* 541: 89-98). The derivatives were submitted to gas chromatography-mass spectrometry (GC-MS), with a Hewlett Packard 5890 Series II plus gas chromatograph attached to an HP model 5989 MS engine. The latter was used in the electron impact mode at 70 eV with a source temperature of 250°C. The GC was fitted on-column injection, and was equipped with a capillary column of fused silica coated with Supelcowax 10TM (0.25 mm x 25 m, 0.25 μ m film; Supelco UK, Poole, UK). After holding the temperature at 80°C for 3 min, the column was temperature-programmed at 20°C/min to 180°C, then at 2°C/min to 280°C, where it was held for 15 min. Helium was the carrier gas at a constant flow-rate of 1mL/min.

Acyl-CoA analysis.

[0174] For acyl-CoA analysis the method developed by Larson and Graham for plant tissues was used (Larson and Graham, 2001, *Plant Journal* 25: 115-125).

Results

Isolation of putative acyl-CoA desaturases from Ranunculaceae species.

[0175] The seed oils of *A. leveillei* and *C. palustris* contain significant amounts of sciadonic acid. Therefore, the developing seed of these two species were used for RNA extraction and further analysis.

[0176] A PCR based approach was used to identify cDNAs for $\Delta 5$ cis-desaturase. Degenerate primers (SEQ ID NO: 7 and SEQ ID NO: 8) designed to the histidine boxes identified in previously isolated putative acyl-CoA desaturases from rose and *Arabidopsis* were used.

[0177] The resulting 420 bp fragments were sequenced yielding two different types of sequences from *A. leveillei* and one type of sequence from *C. palustris*. All of them showed a high level of identity to putative acyl-CoA desaturase-like polypeptides from higher plants such as *Arabidopsis*; they also show some similarity to the partial sequence described by Cahoon et al (2000, *Plant Physiology* 124: 243-251) for the *Limnanthes* $\Delta 5$ -desaturase. These partial sequences were used to design 5' and 3'-RACE primers to amplify both ends of three putative acyl-CoA desaturases. Total RNA isolated from *A. leveillei* and *C. palustris* seeds was used as a template in all the RACE reactions. The sequence data acquired from 3' and 5' RACE were used to design primers to the 5' and 3'- ends of the coding regions of each of the three genes (SEQ ID NOS: 9-23). The primers also contained restriction sites to facilitate the cloning of these genes into the binary vectors for plant expression. Full-length copies of these cDNA clones were amplified directly from the corresponding cDNAs. These primers successfully amplified two full-sized cDNA clones from *A. leveillei*, designated AI10 (SEQ ID NO: 3) and AI21 (SEQ ID NO: 5) and two cDNA clones from *C. palustris*, designated Cal1 and Cal2 (SEQ ID NO: 1). The cDNAs encode polypeptides of 378 (Cal1), 376 (Cal2, SEQ ID NO: 2), 312 (AI10, SEQ ID NO: 4) and 321 (AI21, SEQ ID NO: 6) amino acids. Polypeptides corresponding to cDNA clones Cal1 and Cal2 were almost identical except Cal1 contained an insertion of two amino acids at the position 60 (Leu Asp) (sequence not included). For further analysis we used cDNA clone Cal2. All four polypeptides were found to share up to 25% amino acid sequence identity with $\Delta 9$ -acyl CoA desaturases from rat and human, but were most related to acyl-CoA desaturase like polypeptides from *L. douglasii*, rose and *Arabidopsis* (Fig. 1).

[0178] Expression of *A. leveillei* and *C. palustris* putative Acyl-CoA Desaturases in *A. thaliana*. The coding regions for the three polypeptides, Cal2, AI10 and AI21, were introduced into plant expression vectors under the control of the CAMV 35S promoter and resulting constructs were used to transform *A. thaliana* plants. GC profiles of total fatty acids extracted from leaves of wild-type and transgenic plants transformed with AI10 and AI21 constructs show the presence of several mono- and polyunsaturated fatty acids that were not detected in wild type *Arabidopsis* (Fig. 2). These novel fatty acids were identified by GC-MS analysis of a DMOX derivatives of their methyl esters as previously described (Fay and Richli,

1991, Journal of Chromatography 541: 89-98). This confirmed their identities as $\Delta 5$ isomers of 16:1, 18:1, 18:3 and 18:4 fatty acids (Fig. 3). A double bond in position 5 in a DMOX derivative is usually characterized by a fingerprint of ions at $m/z = 153, 166, \text{ and } 180$. Thus, the double bond position of 18:1 was confirmed from the spectrum shown in Fig 3a which can be compared with that of the authentic compound (Fig 3b). 16:1 $\Delta 5$ was essentially the same in the key region of the spectrum, differing only in the area of molecular ion (Fig. 3c). Fatty acid 18:3 $\Delta 5,9,12$ was identified from comparisons of authentic spectra shown in Fig. 3d,e. The abundant ion at $m/z = 180$ is highly characteristic of a 5,9-double bond system. MS analysis confirmed the presence of double bonds in positions 9 and 12 (located by gaps of 12 amu between $m/z = 194 \text{ and } 206$ and $234 \text{ and } 246$, respectively) and in the position 5, indicated by the especially abundant key diagnostic ion at $m/z = 180$ representing cleavage at the centre of the bis-methylene-interrupted double bond system.

[0179] The most abundant of these fatty acids was 16:1 $\Delta 5$ which accounted for 1.8% (w/w) of the total fatty acids. The $\Delta 5$ isomers of 18:1, 18:3 and 18:4 each participated to less than 1% of the total fatty acids in the transgenic Arabidopsis. However, no $\Delta 5$ -unsaturated fatty acids were found in transgenic Arabidopsis plants expressing Al10. Plants expressing Cal2 produced very small amounts of 16:0 (<1%).

[0180] Analysis of the acyl-CoA pools of leaf tissues of Arabidopsis transformed with both constructs, Al10 and Al21 was carried out (Fig. 4). This revealed the presence of the $\Delta 5$ -desaturated fatty acids corresponding to 16:1, 18:1 18:3 and 18:4 in transgenic plants, expressing Al21. Especially high were levels of 16:1 $\Delta 5$ in transgenic plants expressing AL21. In the case of the acyl-CoA profiles for the leaves of the plants expressing AL10, the presence of the $\Delta 5$ -desaturated fatty acids was not detected.

Co-expression of *A. leveillei* and *C. palustris* Acyl-CoA Desaturases in IgASE1 transgenic *A. thaliana*.

[0181] Since the synthesis of SA requires the presence of 20:2 n-6 as a substrate, we used Arabidopsis lines engineered to accumulate this fatty acid. Kanamycin resistance was used as a selectable marker in transformation of Liberty herbicide resistant IgASE1-line (which expresses the C18- $\Delta 9$ -elongase from *Isochrysis galbana* (SEQ ID NO: 25) and therefore synthesises 20:2, n-6 and 20:3, n-3) with constructs containing *A. leveillei* and *C. palustris* acyl-CoA desaturases coding regions under the control of the CaMV 35S promoter. 40 independent double transgenic lines (i.e. BASTA and Kan resistant) for each of the constructs, AL10 and Al21, were obtained by *A. tumefaciens* - mediated transformation. Two additional peaks appeared in the GC profiles of the leaf fatty acids of the double transgenic lines harbouring both *A. leveillei* putative acyl-CoA desaturase activities, Al10 and Al21, compared to its single transgenic parent (Fig. 5). These were identified by GC-MS analysis as sciadonic acid (C20:3 $\Delta 5_{cis}, 11, 14$) and juniperonic acid (JA, C20:4 $\Delta 5_{cis}, 11, 14, 17$), respectively, on the basis of the retention times of known standards and m/z (Fig. 6). GC- MS analysis of the leaf fatty acids of double transgenic Arabidopsis plants expressing AL21 also confirmed the presence of additional peak corresponding to 16:1 $\Delta 5$.

55 50 45 40 35 30 25 20 15 10 5

Table Ia. Total fatty acid composition of leaves from transgenic *A. thaliana* plants

Fatty acids	16:0	16:1 ⁵	16:1 ⁹	16:2	16:3	18:0	18:1	18:2	18:3	20:2 ^{11,14}	20:3 ^{5,11,14}	20:3 ^{11,14,17}	20:4 ^{5,11,14,17}
Elo9	14.9±0.8	-	3.8±0.3	2.0±0.1	19.4±0.1	0.9±0.1	1.6±0.4	8.0±1.0	47±2.5	3.3±1.5	-	3.8±2.0	-
AL21+ Elo9	11.1±0.3	0.6±0.1	2.7±0.4	2.5±0.2	22.9±0.8	0.5±0.2	1.7±0.3	5.2±1.5	42±1.1	0.74±0.3	0.5±0.1	2.9±1.0	1.3±0.1
AL10+ Elo9	13.4±0.5	-	3.3±0.2	3.4±0.1	18.8±1.5	0.6±0.4	2.2±0.2	5.3±0.9	43±1.1	1.9±0.4	1.5±0.2	3.1±1.1	1.4±0.3
The values are mol% ±SD (n=4).													

[0182] SA and JA accounted for 0.5% and 1.3 % of total fatty acids in plants expressing AL21 (Table Ia) and represented a conversion of 40% and 31% of their respective substrates. In plants expressing AL10 SA and JA accounted for 1.5% and 1.1% respectively and represented a conversion of 44% and 31% of their respective substrates. Single transgenic plants expressing $\Delta 9$ -elongase activity accumulated C20:2 and C20:3 fatty acids to 3.3 % and 3.8 % of total fatty acids.

[0183] SA and JA accounted for 1% and 0.8% of total fatty acids (Table Ib) and represented a conversion of 53% and 44% of their respective substrates. Single transgenic plants expressing $\Delta 9$ -elongase activity accumulated C20:2 and C20:3 fatty acids to 4.5 % and 3.0 % of total fatty acids, similar to those observed previously.

Table Ib. Total fatty acid composition of leaves from single and double transgenic *A. thaliana* plant

Fatty acids	16:0	16:1	16:2	16:3	18:0	18:1	18:2	18:3	20:2	SA	20:3	JA
AI1028	14.0	4.0	0.7 4	13.8	1.6	3.6	12.6 5	45. 7	0.8	0.92	1.0	0.8
IgASE1	11.0	2.8	0.	13.5. 0	0.8	3.5	8.2	30. 5	5.5	-	4.4	4.4 -

Acyl-CoA profiling of double- and single- transgenic A. thaliana plants.
Recent studies of PUFAs biosynthesis in transgenic plants suggest that the microsomal elongase uses acyl CoA as a substrate.

Table II. Fatty acid profile of the Acyl-CoA pool of a single and double transgenic *A.thaliana*. plants.

FA	16:0	18:0/22:3	18:1	18:2	18:2/JA	18:3	20:0	20:1	20:2	20:3	20:3/SA
IgASE1	16.3	1.8	1.1	2.4	-	6.0	2.3	1.0	23.6	17.3	-
AI1028	14.2	3.4	1.7	-	12.4	4.4	2.3	1.3	27.0	27.0 -	32.1

[0184] Analysis of the acyl-CoA pool of the *Ischrysis galbana* $\Delta 9$ (IsoD9) elongase-containing transgenic plants (IgASE1) revealed the accumulation of high levels of acyl-CoAs representing the elongation products of linoleic acid (LA) and linolenic acid (ALA) (i.e. 20:2 n-6 and 20:3 n-3). These latter products represent potential substrates for the Anemone desaturase, especially if the enzyme utilises acyl-CoA substrates. It should also be noted that the accumulation of 20:2-CoA and 20:3-CoA in the IgASE1 lines is likely to represent poor transfer of these esters from the CoA pool into glycerolipids, and may imply they are poor substrates for endogenous *Arabidopsis* acyltransferases. When the acyl-CoA pool for lines co-expressing the IgASE1 elongase and Anemone desaturase were examined, considerable conversion of 20:2 and 20:3 to SA and JA was observed (see Fig 7 for details). As has previously been observed with analysis of PUFA acyl-CoAs, separation of regioisomers is potentially difficult. However, SA accumulates as an overlapping peak with the 20:3 n-3 peak (i.e. the elongation product of ALA). In the case of JA, this elutes close to 18:2, and we reason that the JA peak is likely to be the larger of the two (based on retention time and no alteration to endogenous 18:2 fatty acid levels in these lines).

Functional characterization of A. leveillei AL 10 and AL21 desaturases in yeast.

[0185] The two full-length cDNAs corresponding to the two *A. leveillei* AI10 and AI21 ORFs were cloned in pYES2 vector and expressed in yeast under the control of galactose inducible GAL1 promoter in the presence of either 20:2 or 20:3 exogenous substrates. GC analysis of fatty acid methyl esters (FAMES) revealed that transformed yeast cells were able to produce additional fatty acids that were not detected in controls. These fatty acids were identified by GC-MS-analysis as $\Delta 5$ cis-20:3 -and 20:4 NMIP-fatty acids. Each new putative $\Delta 5$ -unsaturated fatty acid accounted for <1% of the total fatty acids of transgenic yeast (Fig. 8). To obtain higher levels of $\Delta 5$ -unsaturated fatty acids in transgenic yeast we co-expressed both *A. leveillei* $\Delta 5$ desaturases with the C18-09 PUFA-specific elongase IgASE1 from *Ischrysis galbana* in the presence of two different substrates, 18:2, n-6 or 18:3, n-3. The results indicated that combination of elongation and desaturation activities has not generated better amounts of Sciadonic and Juniperonic acids in yeast (results not shown).

[0186] We therefore conclude that we have functionally identified the $\Delta 5$ -desaturase responsible for the NMI-desaturation and the synthesis of sciadonic and juniperonic acid. Moreover, we show that this enzyme is an acyl-CoA desaturase, and therefore represents the first cloned and characterised example of such an activity from higher plants. It is very likely that it would be possible to increase the accumulation of SA and JA in transgenic plants by optimising their exchange out of the acyl-CoA pool, by the co-expression of a suitable acyltransferases.

Example 2**Generation of transgenic plants**

5 a) *Generation of transgenic oilseed rape plants (modified method of Moloney et al., 1992, Plant Cell Reports, 8:238-242)*

10 **[0187]** Binary vectors in *Agrobacterium tumefaciens* C58C1:pGV2260 or *Escherichia coli* (Deblaere et al, 1984, Nucl. Acids. Res. 13, 4777-4788) can be used for generating transgenic oilseed rape plants. To transform oilseed rape plants (Var. Drakkar, NPZ Nordeutsche Pflanzenzucht, Hohenlieth, Germany), a 1:50 dilution of an overnight culture of a positively transformed agrobacterial colony in Murashige-Skoog medium (Murashige and Skoog 1962 Physiol. Plant. 15, 473) supplemented with 3% sucrose (3MS medium) is used. Petiols or hypocotyls of freshly germinated sterile oilseed rape plants (in each case approx. 1 cm²) are incubated with a 1:50 agrobacterial dilution for 5-10 minutes in a Petri dish. This is followed by 3 days of coincubation in the dark at 25°C on 3MS medium supplemented with 0.8% Bacto agar. The cultures are then grown for 3 days at 16 hours light/8 hours dark and the cultivation is continued in a weekly rhythm on MS medium supplemented with 500 mg/l Claforan (cefotaxim sodium), 50 mg/l kanamycin, 20 µM benzylaminopurine (BAP), now supplemented with 1.6 g/l of glucose. Growing shoots are transferred to MS medium supplemented with 2% sucrose, 250 mg/l Claforan and 0.8% Bacto agar. If no roots develop after three weeks, 2-indolebutyric acid was added to the medium as growth hormone for rooting. Regenerated shoots were obtained on 2MS medium supplemented with kanamycin and Claforan; after rooting, they were transferred to compost and, after growing on for two weeks in a controlled-environment cabinet or in the greenhouse, allowed to flower, and mature seeds were harvested and analyzed by lipid analysis for elongase expression, such as $\Delta 5$ -elongase or $\Delta 6$ -elongase activity or $\omega 3$ -desaturase activity. In this manner, lines with elevated contents of polyunsaturated C₂₀- and C₂₂-fatty acids can be identified.

25 b) *Generation of transgenic linseed plants*

30 **[0188]** Transgenic linseed plants can be generated for example by the method of Bell et al., 1999, In Vitro Cell. Dev. Biol.-Plant. 35(6):456-465 by means of particle bombardment. In general, linseed was transformed by an agrobacteria-mediated transformation, for example by the method of Mlynarova et al. (1994), Plant Cell Report 13: 282-285.

35

40

45

50

55

EP 2 388 332 A1

SEQUENCE LISTING

5 <110> BASF

<120> Process for the Production of Delta-5 Unsaturated Fatty Acids in
 10 Transgenic Organisms

<130> DY-04

15 <160> 25

<170> Patent In version 3.1

20 <210> 1

<211> 1131

25 <212> DNA

<213> *Caltha palustris*

30 <220>

<221> CDS

<222> (1)..(1131)

35 <223>

<400> 1

40 atg gct cta att gca aca acc ccc aaa gcc aca ttc tat ccc tct tct 48
 Met Ala Leu Ile Ala Thr Thr Pro Lys Ala Thr Phe Tyr Pro Ser Ser
 1 5 10 15

cca tac ctt cgt cca act agg cgt gga aac ccc aaa tcc att tcc act 96
 Pro Tyr Leu Arg Pro Thr Arg Arg Gly Asn Pro Lys Ser Ile Ser Thr
 20 25 30

45 tat ttc aaa gtt ttt tct tca att tca ttt att tcc aag aca aat tcc 144
 Tyr Phe Lys Val Phe Ser Ser Ile Ser Phe Ile Ser Lys Thr Asn Ser
 35 40 45

50 agt acc att tgt ttt ccc aat act tgc cta act ttg gaa agg aga aac 192
 Ser Thr Ile Cys Phe Pro Asn Thr Cys Leu Thr Leu Glu Arg Arg Asn
 50 55 60

aag aga tgt aca aga tcc ata tct tgt tcg gcg gcc aca act gga gaa 240
 Lys Arg Cys Thr Arg Ser Ile Ser Cys Ser Ala Ala Thr Thr Gly Glu
 65 70 75 80

55 gac tcg ggg aag atc ttt tta tct gat gtg caa gtg ata cga cga cca 288
 Asp Ser Gly Lys Ile Phe Leu Ser Asp Val Gln Val Ile Arg Arg Pro

EP 2 388 332 A1

	85					90					95						
5	cag G n	gtt Val	t at Tyr	t gg Tr p 100	gga G y	aga Arg	aaa Lys	t gg Tr p	aac Asn 105	aca Thr	gta Val	gat Asp	att Ile	gct Ala 110	act Thr	ggc G y	336
	t ca Ser	gta Val	gtc Val 115	t tg Leu	ggg G y	atg Met	cac His	att Ile 120	ctg Leu	t cc Ser	t tg Leu	ttt Phe	gcc Ala 125	cct Pro	ttt Phe	aca Thr	384
10	t tc Phe	agt Ser 130	t gg Tr p	t ct Ser	gca Ala	ctt Leu	agt Ser 135	atg Met	gca Ala	t tc Phe	gtg Val	ctt Leu 140	t at Tyr	gtg Val	gta Val	aca Thr	432
15	ggg G y 145	ctg Leu	ttt Phe	ggg G y	att Ile	act Thr 150	ctg Leu	t ca Ser	t at Tyr	cat His	aga Arg 155	aat Asn	ctt Leu	gct Ala	cat His	aaa Lys 160	480
	gca Ala	t tc Phe	aag Lys	ctt Leu	cct Pro 165	aaa Lys	t gg Tr p	ctt Leu	gaa G u	t ac Tyr 170	t ta Leu	ttt Phe	gca Ala	t at Tyr	t gc Cys 175	ggg G y	528
20	gca Ala	cag G n	gca Ala	t tg Leu 180	cag G n	ggg G y	aat Asn	cca Pro	att Ile 185	gat Asp	t gg Tr p	gtt Val	agt Ser	act Thr 190	cac His	agg Arg	576
25	t ac Tyr	cac His	cac His 195	cag G n	t tc Phe	t gt Cys	gat Asp	t cc Ser 200	gaa G u	aga Arg	gac Asp	cct Pro	cat His 205	agt Ser	cct Pro	ctt Leu	624
	gaa G u	gga G y 210	t tc Phe	t gg Tr p	t tc Phe	agt Ser	cat His 215	ata Ile	agt Ser	t gg Tr p	t tc Phe	ttt Phe 220	gac Asp	acc Thr	aaa Lys	t ct Ser	672
30	ctt Leu 225	act Thr	gag G u	aag Lys	t gt Cys	gga G y 230	gtg Val	ccc Pro	aac Asn	aat Asn	gtg Val 235	t ct Ser	gat Asp	t ta Leu	gag G u	aag Lys 240	720
35	caa G n	cca Pro	t tc Phe	t ac Tyr	aaa Lys 245	ttt Phe	atg Met	gaa G u	aag Lys	aca Thr 250	t at Tyr	cct Pro	ctt Leu	cat His 255	cca Pro	att Ile	768
	gca Ala	ctt Leu	gca Ala	gct Ala 260	gtg Val	cta Leu	t at Tyr	gca Ala	cta Leu 265	gga G y	gga G y	ttt Phe	ccc Pro	ttt Phe 270	ctt Leu	gta Val	816
40	t gg Tr p	agc Ser	atg Met 275	ggc G y	gta Val	agg Arg	gtt Val	gta Val 280	t gg Tr p	gtg Val	t at Tyr	cat His	att Ile 285	aca Thr	t gg Tr p	ctc Leu	864
45	gta Val	aat Asn 290	t ca Ser	gct Ala	t gt Cys	cat His	gta Val 295	t gg Tr p	gga G y	agc Ser	cag G n	t cg Ser 300	t gg Tr p	aat Asn	acg Thr	ggc G y	912
	gac Asp 305	t ta Leu	t ct Ser	agg Arg	aac Asn 310	aac Asn	t gg Tr p	t gg Tr p	gtg Val	gct Ala	t tg Leu 315	ctt Leu	gca Ala	ttt Phe	gga G y	gag G u 320	960
50	ggt G y	t gg Tr p	cat His	aac Asn	aat Asn 325	cac His	cat His	gca Ala	ttt Phe	gaa G u 330	ttt Phe	t ca Ser	gct Ala	cgg Arg	cat His 335	gga G y	1008
	t tg Leu	gaa G u	t gg Tr p	t gg Tr p	caa G n	ctt Leu	gac Asp	t tg Leu	aca Thr 345	t gg Tr p	t at Tyr	act Thr	gta Val	cga Arg 350	att Ile	ctt Leu	1056
55	caa G n	gca Ala	ctt Leu	gga G y	t tg Leu	gcg Ala	aca Thr	gat Asp	gtg Val	aaa Lys	gtt Val	cca Pro	t cc Ser	gag G u	gta Val	cag G n	1104

EP 2 388 332 A1

355 360 365
 aag caa agg atg tgt ttc agc gag taa 1131
 Lys Gln Arg Met Cys Phe Ser Glu
 370 375
 5
 <210> 2
 <211> 376
 10
 <212> PRT
 <213> Caltha palustris
 15
 <400> 2
 Met Ala Leu Ile Ala Thr Thr Pro Lys Ala Thr Phe Tyr Pro Ser Ser
 1 5 10 15
 20
 Pro Tyr Leu Arg Pro Thr Arg Arg Gly Asn Pro Lys Ser Ile Ser Thr
 20 25 30
 Tyr Phe Lys Val Phe Ser Ser Ile Ser Phe Ile Ser Lys Thr Asn Ser
 25 30 35 40 45
 Ser Thr Ile Cys Phe Pro Asn Thr Cys Leu Thr Leu Glu Arg Arg Asn
 50 55 60
 30
 Lys Arg Cys Thr Arg Ser Ile Ser Cys Ser Ala Ala Thr Thr Gly Glu
 65 70 75 80
 Asp Ser Gly Lys Ile Phe Leu Ser Asp Val Gln Val Ile Arg Arg Pro
 85 90 95
 35
 Gln Val Tyr Trp Gly Arg Lys Trp Asn Thr Val Asp Ile Ala Thr Gly
 100 105 110
 40
 Ser Val Val Leu Gly Met His Ile Leu Ser Leu Phe Ala Pro Phe Thr
 115 120 125
 Phe Ser Trp Ser Ala Leu Ser Met Ala Phe Val Leu Tyr Val Val Thr
 130 135 140
 45
 Gly Leu Phe Gly Ile Thr Leu Ser Tyr His Arg Asn Leu Ala His Lys
 145 150 155 160
 50
 Ala Phe Lys Leu Pro Lys Trp Leu Glu Tyr Leu Phe Ala Tyr Cys Gly
 165 170 175
 Ala Gln Ala Leu Gln Gly Asn Pro Ile Asp Trp Val Ser Thr His Arg
 180 185 190
 55
 Tyr His His Gln Phe Cys Asp Ser Glu Arg Asp Pro His Ser Pro Leu

EP 2 388 332 A1

	195		200		205												
5	G u	G y	Phe	Trp	Phe	Ser	His	Ile	Ser	Trp	Phe	Phe	Asp	Thr	Lys	Ser	
		210					215					220					
	Leu	Thr	G u	Lys	Cys	G y	Val	Pro	Asn	Asn	Val	Ser	Asp	Leu	G u	Lys	
	225					230					235					240	
10	G n	Pro	Phe	Tyr	Lys	Phe	Met	G u	Lys	Thr	Tyr	Pro	Leu	His	Pro	Ile	
					245					250					255		
15	Al a	Leu	Al a	Al a	Val	Leu	Tyr	Al a	Leu	G y	G y	Phe	Pro	Phe	Leu	Val	
				260					265					270			
	Trp	Ser	Met	G y	Val	Arg	Val	Val	Trp	Val	Tyr	His	Ile	Thr	Trp	Leu	
			275					280					285				
20	Val	Asn	Ser	Al a	Cys	His	Val	Trp	G y	Ser	G n	Ser	Trp	Asn	Thr	G y	
		290					295					300					
25	Asp	Leu	Ser	Arg	Asn	Asn	Trp	Trp	Val	Al a	Leu	Leu	Al a	Phe	G y	G u	
	305				310						315					320	
	G y	Trp	His	Asn	Asn	His	His	Al a	Phe	G u	Phe	Ser	Al a	Arg	His	G y	
				325						330					335		
30	Leu	G u	Trp	Trp	G n	Leu	Asp	Leu	Thr	Trp	Tyr	Thr	Val	Arg	Ile	Leu	
				340					345					350			
35	G n	Al a	Leu	G y	Leu	Al a	Thr	Asp	Val	Lys	Val	Pro	Ser	G u	Val	G n	
			355					360					365				
	Lys	G n	Arg	Met	Cys	Phe	Ser	G u									
		370					375										
40	<210>	3															
	<211>	966															
	<212>	DNA															
45	<213>	Anemone l evei l l ei															
	<220>																
50	<221>	CDS															
	<222>	(1) . . (966)															
	<223>																
55	<400>	3															

EP 2 388 332 A1

	atg	gaa	ctc	cca	gcg	atg	gcg	ttg	caa	agt	gaa	cta	aag	caa	gac	aaa	48
	Met	Gu	Leu	Pro	Ala	Met	Ala	Leu	Gln	Ser	Gu	Leu	Lys	Gln	Asp	Lys	
	1				5					10					15		
5	ttt	cca	tcc	tcc	gat	gtg	ccg	gtg	aaa	gac	aag	acg	aag	aag	ata	act	96
	Phe	Pro	Ser	Ser	Asp	Val	Pro	Val	Lys	Asp	Lys	Thr	Lys	Lys	Ile	Thr	
				20					25					30			
10	tgg	gtt	agg	gta	tgg	aac	tcg	acc	gat	atc	ttc	cac	ata	att	ctc	gta	144
	Trp	Val	Arg	Val	Trp	Asn	Ser	Thr	Asp	Ile	Phe	His	Ile	Ile	Leu	Val	
			35					40					45				
15	ggc	ggt	ctg	cat	gtt	ttg	tgc	ttg	tct	gca	cct	ttc	acc	ttc	agt	tgg	192
	Gly	Gly	Leu	His	Val	Leu	Cys	Leu	Ser	Ala	Pro	Phe	Thr	Phe	Ser	Trp	
		50					55					60					
20	agt	gca	ttt	tgg	ttg	tcc	tta	acg	cta	tac	gcc	gtc	tgt	gga	gtc	ttc	240
	Ser	Ala	Phe	Trp	Leu	Ser	Leu	Thr	Leu	Tyr	Ala	Val	Cys	Gly	Val	Phe	
						70				75					80		
25	ggg	act	act	ctg	tca	tac	cac	agg	aac	ctt	act	cat	aga	agt	ttc	aag	288
	Gly	Thr	Thr	Leu	Ser	Tyr	His	Arg	Asn	Leu	Thr	His	Arg	Ser	Phe	Lys	
					85					90					95		
30	ctt	cca	aag	tat	ctc	gag	tac	ttc	ttc	gcc	tat	gtt	gga	ctt	cat	gct	336
	Leu	Pro	Lys	Tyr	Leu	Gu	Tyr	Phe	Phe	Ala	Tyr	Val	Gly	Leu	His	Ala	
				100					105					110			
35	tta	cag	ggg	gat	cct	gta	tgg	tgg	gtg	agc	aca	cac	agg	tac	cat	cac	384
	Leu	Gln	Gly	Asp	Pro	Val	Trp	Trp	Val	Ser	Thr	His	Arg	Tyr	His	His	
			115				120						125				
40	aag	ttc	acc	gac	aca	tac	ctg	gat	cca	cac	agt	cca	att	gaa	gga	ttt	432
	Lys	Phe	Thr	Asp	Thr	Tyr	Leu	Asp	Pro	His	Ser	Pro	Ile	Gu	Gly	Phe	
			130				135					140					
45	tgg	ttc	tct	cac	ata	ttc	tgg	ctg	ttc	gac	tct	aaa	tat	ata	ctc	gaa	480
	Trp	Phe	Ser	His	Ile	Phe	Trp	Leu	Phe	Asp	Ser	Lys	Tyr	Ile	Leu	Gu	
					150						155				160		
50	gag	tgc	gga	cgg	tac	gaa	aat	gcc	ggc	gat	cta	ctg	aaa	caa	agc	tat	528
	Gu	Cys	Gly	Arg	Tyr	Gu	Asn	Ala	Gly	Asp	Leu	Leu	Lys	Gln	Ser	Tyr	
					165				170						175		
55	tac	agg	ttt	cta	gaa	agg	act	ttt	gtg	ttt	cac	gtt	tac	cta	caa	gca	576
	Tyr	Arg	Phe	Leu	Gu	Arg	Thr	Phe	Val	Phe	His	Val	Tyr	Leu	Gln	Ala	
				180					185					190			
60	gct	ttg	ctg	tat	atg	ttc	gga	gga	ttt	cct	ttc	atc	gtc	tgg	gga	atg	624
	Ala	Leu	Leu	Tyr	Met	Phe	Gly	Gly	Phe	Pro	Phe	Ile	Val	Trp	Gly	Met	
				195				200					205				
65	gca	gta	aga	act	gta	tgg	gga	ttc	cat	gcc	tct	tgg	cta	gtg	aat	tca	672
	Ala	Val	Arg	Thr	Val	Trp	Gly	Phe	His	Ala	Ser	Trp	Leu	Val	Asn	Ser	
			210				215					220					
70	gta	tgc	cac	aga	tat	gga	cac	caa	gca	tgg	gac	aca	ggg	gac	ttg	tca	720
	Val	Cys	His	Arg	Tyr	Gly	His	Gln	Ala	Trp	Asp	Thr	Gly	Asp	Leu	Ser	
					225			230			235				240		
75	aca	aac	aac	tgg	ttc	ata	gcg	atg	ctt	aca	tcg	ggt	gag	ggc	tgg	cat	768
	Thr	Asn	Asn	Trp	Phe	Ile	Ala	Met	Leu	Thr	Ser	Gly	Gu	Gly	Trp	His	
					245				250					255			
80	aac	aac	cat	cac	gct	ttt	gag	tat	tca	gct	cg	cat	gga	ata	gag	tgg	816
	Asn	Asn	His	His	Ala	Phe	Gu	Tyr	Ser	Ala	Arg	His	Gly	Ile	Gu	Trp	
				260					265					270			

EP 2 388 332 A1

	tgg	caa	ata	gat	acg	aca	tgg	tat	gta	ata	aag	cta	ctc	gaa	tat	ctc	864
	Trp	G n	I l e	Asp	Thr	Thr	Trp	Tyr	Val	I l e	Lys	Leu	Leu	G u	Tyr	Leu	
			275					280					285				
5	gga	ttg	gca	acg	gat	gtg	aag	gtg	cct	tca	gag	gtt	cac	aag	cga	aag	912
	G y	Leu	Al a	Thr	Asp	Val	Lys	Val	Pro	Ser	G u	Val	Hi s	Lys	Arg	Lys	
		290					295					300					
10	atg	tct	ttc	aaa	aat	tgc	gtt	cag	gat	aaa	cag	ttc	tgt	gtg	aac	gac	960
	Met	Ser	Phe	Lys	Asn	Cys	Val	G n	Asp	Lys	G n	Phe	Cys	Val	Asn	Asp	
	305					310					315					320	
	aag	taa															966
	Lys																
15	<210>	4															
	<211>	321															
	<212>	PRT															
20	<213>	Anemone	I	l	e	i	I	l	e	i							
	<400>	4															
25	Met	G u	Leu	Pro	Al a	Met	Al a	Leu	G n	Ser	G u	Leu	Lys	G n	Asp	Lys	
	1				5					10					15		
30	Phe	Pro	Ser	Ser	Asp	Val	Pro	Val	Lys	Asp	Lys	Thr	Lys	Lys	I l e	Thr	
				20					25					30			
	Trp	Val	Arg	Val	Trp	Asn	Ser	Thr	Asp	I l e	Phe	Hi s	I l e	I l e	Leu	Val	
			35					40					45				
35	G y	G y	Leu	Hi s	Val	Leu	Cys	Leu	Ser	Al a	Pro	Phe	Thr	Phe	Ser	Trp	
		50					55					60					
40	Ser	Al a	Phe	Trp	Leu	Ser	Leu	Thr	Leu	Tyr	Al a	Val	Cys	G y	Val	Phe	
	65					70					75					80	
	G y	Thr	Thr	Leu	Ser	Tyr	Hi s	Arg	Asn	Leu	Thr	Hi s	Arg	Ser	Phe	Lys	
					85					90					95		
45	Leu	Pro	Lys	Tyr	Leu	G u	Tyr	Phe	Phe	Al a	Tyr	Val	G y	Leu	Hi s	Al a	
				100					105					110			
50	Leu	G n	G y	Asp	Pro	Val	Trp	Trp	Val	Ser	Thr	Hi s	Arg	Tyr	Hi s	Hi s	
			115					120					125				
	Lys	Phe	Thr	Asp	Thr	Tyr	Leu	Asp	Pro	Hi s	Ser	Pro	I l e	G u	G y	Phe	
		130					135					140					
55	Trp	Phe	Ser	Hi s	I l e	Phe	Trp	Leu	Phe	Asp	Ser	Lys	Tyr	I l e	Leu	G u	
	145					150					155					160	

EP 2 388 332 A1

5 Glu Cys Gly Arg Tyr 165 Glu Asn Ala Gly Asp 170 Leu Leu Lys Gln Ser Tyr
 Tyr Arg Phe 180 Leu Glu Arg Thr Phe Val 185 Phe His Val Tyr Leu Gln Ala
 10 Ala Leu 195 Leu Tyr Met Phe Gly Gly 200 Phe Pro Phe Ile Val 205 Trp Gly Met
 Ala Val 210 Arg Thr Val Trp Gly 215 Phe His Ala Ser Trp 220 Leu Val Asn Ser
 15 Val 225 Cys His Arg Tyr Gly 230 His Gln Ala Trp Asp 235 Thr Gly Asp Leu Ser 240
 Thr Asn Asn Trp Phe 245 Ile Ala Met Leu Thr 250 Ser Gly Glu Gly Trp His 255
 20 Asn Asn His 260 His Ala Phe Glu Tyr Ser 265 Ala Arg His Gly Ile Glu Trp 270
 25 Trp Gln Ile 275 Asp Thr Thr Trp Tyr 280 Val Ile Lys Leu Leu 285 Glu Tyr Leu
 Gly Leu 290 Ala Thr Asp Val Lys 295 Val Pro Ser Glu Val 300 His Lys Arg Lys
 30 Met Ser Phe Lys Asn Cys 310 Val Gln Asp Lys Gln 315 Phe Cys Val Asn Asp 320

35 Lys

40 <210> 5

<211> 939

<212> DNA

<213> Anemone levei llei

45 <220>

<221> CDS

50 <222> (1)..(939)

<223>

55 <400> 5
 atg gat ctc aca tca atg gca atg caa gaa aca acc gct gct gca gag
 Met Asp Leu Thr Ser Met Ala Met Gln Gu Thr Thr Ala Ala Ala Glu

48

EP 2 388 332 A1

	1		5		10		15										
5	gaa G u	gac A s p	aga A r g	ctt L e u 20	cca P r o	tgc C y s	tca S e r	gaa G u	ggt V a l 25	cca P r o	gtg V a l	aag L y s	gag G u	aag L y s 30	aca T h r	aag L y s	96
	aag L y s	atc I l e	act T h r 35	tgg T r p	ggt V a l	agg A r g	aaa L y s	tgg T r p 40	aac A s n	tcc S e r	acc T h r	gac A s p	gtc V a l 45	ttc P h e	cac H i s	att I l e	144
10	ctg L e u	ttt P h e 50	att I l e	ggt G y	ggt G y	ttg L e u	cac H i s 55	ggt V a l	ttg L e u	tgc C y s	ttg L e u	ttt P h e 60	gca A l a	cct P r o	tcc S e r	acc T h r	192
15	ttc P h e 65	agc S e r	tgg T r p	aaa L y s	tct S e r	ttt P h e 70	tgg T r p	gtg V a l	tgc C y s	ttc P h e	gcg A l a 75	cta L e u	tat T y r	gcc A l a	atc I l e	tgt C y s 80	240
	gga G y	gtg V a l	ttc P h e	ggg G y	act T h r 85	act T h r	ttg L e u	tcc S e r	ttc P h e	cac H i s 90	agg A r g	aac A s n	ctt L e u	aca T h r	cac H i s 95	aga A r g	288
20	agt S e r	ttc P h e	aag L y s	ctt L e u 100	cca P r o	aag L y s	tat T y r	ctc L e u	gag G u 105	tac T y r	ttc P h e	ttc P h e	gct A l a	tat T y r 110	gtc V a l	gga G y	336
25	ctt L e u	cat H i s	gct A l a 115	cta L e u	cag G n	ggt G y	gat A s p	cca P r o 120	gtg V a l	tgg T r p	tgg T r p	gtg V a l	agc S e r 125	aca T h r	cat H i s	aga A r g	384
	tac T y r	cat H i s 130	cac H i s	aag L y s	tac T y r	act T h r	gac A s p 135	aca T h r	tac T y r	ctg L e u	gat A s p	cca P r o 140	cac H i s	agt S e r	cca P r o	atc I l e	432
30	gaa G u 145	gga G y	ttt P h e	tgg T r p	ttc P h e	tgc C y s 150	cac H i s	ata I l e	ttc P h e	tgg T r p	ctt L e u 155	ttc P h e	gac A s p	tcc S e r	aaa L y s	tat T y r 160	480
35	ata I l e	atc I l e	gaa G u	aag L y s	tgc C y s 165	gga G y	agg A r g	tat T y r	gaa G u	aat A s n 170	gca A l a	ggg G y	gat A s p	ctt L e u	atg M e t 175	aaa L y s	528
	cag G n	agc S e r	tat T y r	tac T y r 180	agg A r g	ttt P h e	ctg L e u	gag G u	agg A r g 185	act T h r	ttc P h e	gtg V a l	tat T y r 190	cat H i s	ggt V a l	tac T y r	576
40	tta L e u	caa G n	gca A l a 195	gct A l a	ttg L e u	ttg L e u	tat T y r	ctg L e u 200	ttc P h e	gga G y	ggg G y	ttt P h e	ccg P r o 205	ttc P h e	atc I l e	ggt V a l	624
45	tgg T r p	gga G y	atg M e t 210	gca A l a	gta V a l	aga A r g	act T h r 215	ata I l e	ttg L e u	gga G y	ttc P h e	cat H i s 220	ctc L e u	tct S e r	tgg T r p	cta L e u	672
	gtg V a l 225	aat A s n	tca S e r	ggt V a l	tgc C y s	cat H i s 230	aga A r g	tgg T r p	gga G y	aac A s n	cga A r g 235	cca P r o	tgg T r p	aac A s n	acc T h r	ggg G y 240	720
50	gac A s p	ttg L e u	tca S e r	aca T h r	aac A s n 245	aat A s n	tgg T r p	ttc P h e	att I l e	gct A l a 250	atg M e t	ctt L e u	aca T h r	tcg S e r	ggt G y 255	gag G u	768
	ggt G y	tgg T r p	cat H i s	aac A s n 260	aac A s n	cat H i s	cat H i s	gcg A l a	ttt P h e 265	gag G u	tat T y r	tca S e r	gct A l a	cgg A r g 270	cat H i s	gga G y	816
55	ata I l e	gag G u	tgg T r p	tgg T r p	caa G n	ata I l e	gac A s p	acg T h r	acg T h r	tgg T r p	tac T y r	ata I l e	ata I l e	aag L y s	cta L e u	ctc L e u	864

EP 2 388 332 A1

	275	280	285	
5	gag t at ct c gga ttg gca act gat at c aag gtg ccc t ca gaa at t cac			912
	Gu Tyr Leu Gly Leu Ala Thr	Asp Ile Lys Val Pro Ser	Glu Ile His	
	290	295	300	
	aag cgt aag atg tct ttc aaa aat tga			939
	Lys Arg Lys Met Ser Phe Lys Asn			
	305	310		
10	<210> 6			
	<211> 312			
	<212> PRT			
15	<213> Anemone levei llei			
	<400> 6			
20	Met Asp Leu Thr Ser Met Ala Met Gn Gu Thr Thr Ala Ala Ala Gu			
	1 5 10 15			
25	Gu Asp Arg Leu Pro Cys Ser Gu Val Pro Val Lys Gu Lys Thr Lys			
	20 25 30			
	Lys Ile Thr Trp Val Arg Lys Trp Asn Ser Thr Asp Val Phe His Ile			
	35 40 45			
30	Leu Phe Ile Gly Gly Leu His Val Leu Cys Leu Phe Ala Pro Ser Thr			
	50 55 60			
35	Phe Ser Trp Lys Ser Phe Trp Val Cys Phe Ala Leu Tyr Ala Ile Cys			
	65 70 75 80			
	Gly Val Phe Gly Thr Thr Leu Ser Phe His Arg Asn Leu Thr His Arg			
	85 90 95			
40	Ser Phe Lys Leu Pro Lys Tyr Leu Gu Tyr Phe Phe Ala Tyr Val Gly			
	100 105 110			
45	Leu His Ala Leu Gn Gly Asp Pro Val Trp Trp Val Ser Thr His Arg			
	115 120 125			
	Tyr His His Lys Tyr Thr Asp Thr Tyr Leu Asp Pro His Ser Pro Ile			
	130 135 140			
50	Gu Gly Phe Trp Phe Cys His Ile Phe Trp Leu Phe Asp Ser Lys Tyr			
	145 150 155 160			
	Ile Ile Gu Lys Cys Gly Arg Tyr Gu Asn Ala Gly Asp Leu Met Lys			
	165 170 175			
55	Gn Ser Tyr Tyr Arg Phe Leu Gu Arg Thr Phe Val Tyr His Val Tyr			

EP 2 388 332 A1

	180	185	190													
5	Leu	G n	Al a	Al a	Leu	Leu	Tyr	Leu	Phe	G y	G y	Phe	Pro	Phe	I l e	Val
			195					200					205			
	Trp	G y	M et	Al a	Val	Arg	Thr	I l e	Leu	G y	Phe	Hi s	Leu	Ser	Trp	Leu
		210					215					220				
10	Val	Asn	Ser	Val	Cys	Hi s	Arg	Trp	G y	Asn	Arg	Pro	Trp	Asn	Thr	G y
	225					230					235					240
15	Asp	Leu	Ser	Thr	Asn	Asn	Trp	Phe	I l e	Al a	M et	Leu	Thr	Ser	G y	G u
					245					250					255	
	G y	Trp	Hi s	Asn	Asn	Hi s	Hi s	Al a	Phe	G u	Tyr	Ser	Al a	Arg	Hi s	G y
				260					265					270		
20	I l e	G u	Trp	Trp	G n	I l e	Asp	Thr	Thr	Trp	Tyr	I l e	I l e	Lys	Leu	Leu
			275					280						285		
25	G u	Tyr	Leu	G y	Leu	Al a	Thr	Asp	I l e	Lys	Val	Pro	Ser	G u	I l e	Hi s
		290					295					300				
	Lys	Arg	Lys	M et	Ser	Phe	Lys	Asn								
	305					310										
30	<210>	7														
	<211>	24														
	<212>	DNA														
35	<213>	Artificial Sequence														
	<220>															
40	<223>	Forward primer designed to encode conserved amino acid sequence c														
		orresponding to histidine box II in previously characterized acyl														
		-CoA desaturases														
	<220>															
45	<221>	misc_feature														
	<222>	(6)..(6)														
	<223>	inosine														
50	<220>															
	<221>	misc_feature														
55	<222>	(9)..(9)														
	<223>	inosine														

5 <220>
 <221> m i s c _ f e a t u r e
 <222> (12) . . (12)
 <223> n i s i n o s i n e
 10

<220>
 <221> m i s c _ f e a t u r e
 15 <222> (18) . . (18)
 <223> n i s i n o s i n e

20 <400> 7
 t g g g t n w s n a y n c a y m g n t a y c a y 24

<210> 8
 25 <211> 29
 <212> DNA
 <213> A r t i f i c i a l S e q u e n c e

30 <220>
 <223> R e v e r s e p r i m e r d e s i g n e d t o e n c o d e t h e c o m p l e m e n t o f h i s t i d i n e b o x
 35 I I I i n p r e v i o u s l y c h a r a c t e r i z e d a c y l - C o A d e s a t u r a s e s

<220>
 <221> m i s c _ f e a t u r e
 <222> (21) . . (21)
 40 <223> n i s i n o s i n e

<220>
 45 <221> m i s c _ f e a t u r e
 <222> (27) . . (27)
 <223> n i s i n o s i n e

50 <400> 8
 g c r t g r t g r t t r t t r t g c c a n c c y t c n c c 29

55 <210> 9

EP 2 388 332 A1

<211> 34
 <212> DNA
 5 <213> Cal t ha pal ustr i s

 <400> 9
 10 ggt cgacat g gct ct aat t g caacaacccc caaa 34

 <210> 10
 <211> 34
 15 <212> DNA
 <213> Cal t ha pal ustr i s

 <400> 10
 20 gat cgat at g gct ct aat t g caacaacccc caaa 34

 <210> 11
 25 <211> 26
 <212> DNA
 <213> Cal t ha pal ustr i s

 30 <400> 11
 gagct ct t ac t cgct gaaac acat cc 26

 <210> 12
 35 <211> 27
 <212> DNA
 <213> Cal t ha pal ustr i s

 40 <400> 12
 gat cgat t t a ct cgct gaaa cacat cc 27

 <210> 13
 <211> 27
 <212> DNA
 50 <213> Anemone l evei l l ei

 <400> 13
 55 ggt ct agaat ggat ct caca t caat gg 27

EP 2 388 332 A1

<210> 14
 <211> 28
 5 <212> DNA
 <213> Anemone levei l ei

 10 <400> 14
 ggcggccgca tggat ct cac at caat gg 28

 <210> 15
 15 <211> 26
 <212> DNA
 <213> Anemone levei l ei

 20 <400> 15
 gat cgat at g gat ct cacat caat gg 26

 <210> 16
 25 <211> 35
 <212> DNA
 30 <213> Anemone levei l ei

 <400> 16
 ggt cgact ca atttttgaaa gacat ctt ac gcttg 35
 35 <210> 17
 <211> 37
 <212> DNA
 40 <213> Anemone levei l ei

 <400> 17
 45 ggcggccgct caat ttttga aagacat ctt acgcttg 37

 <210> 18
 <211> 25
 50 <212> DNA
 <213> Anemone levei l ei

 <400> 18
 55 ggcccgggat ggaact ccca gcgat 25

EP 2 388 332 A1

5 <210> 19
<211> 24
<212> DNA
<213> Anemone levei llei
10
<400> 19
ggcggccgca gttcaccgac atac 24
15 <210> 20
<211> 24
<212> DNA
20 <213> Anemone levei llei
<400> 20
gatcgat atg gaactcccag cgat 24
25 <210> 21
<211> 29
<212> DNA
30 <213> Anemone levei llei
<400> 21
35 ggtcgactta cttgtcgttc acacagaac 29
<210> 22
<211> 31
40 <212> DNA
<213> Anemone levei llei
<400> 22
45 ggcggccgct tacttgcgt tcacacagaa c 31
<210> 23
50 <211> 29
<212> DNA
<213> Anemone levei llei
55

EP 2 388 332 A1

<400> 23
gat c g a t t t a c t t g t c g t t c a c a c a g a a c 29

5 <210> 24
<211> 891
<212> DNA

10 <213> A. cast e l l a n i i e l o n g a s e

<400> 24
15 at g g c g g c t g c g a c g g c g a c g a c g g c a a c g a c g g c g g t g a t g g a g c a a g t g c c c a t t a c g 60
g a g g c c a t c t t c c g g c c g g a c c t c t g g g t c g g a c g g g a c c a g t g g g a g g c g a a t g c c g t g 120
a g c t t c g t a t g g a g g t a c t g g t g g t t c t t c c t g g t g a t g g g c g t g g c a t a c c t g c c c a t c 180
a t c t t c g g c c t c a a g t a c t g g a t g a a g g a t c g t c c g g c c t t c a a c c t c c g t c g g c c g c t c 240
20 a t c t t g t g g a a t a t c t t c a t g g c g a c g t t c t c g a c c g c c g g c t t c c t g t c g a t c g t c t a c 300
c c c c t c a t c g a g a a c t g g g t c t a c c c c g g c g g c g g c c t c a c c c c g c a t g a g t t c a t c t g c 360
t c g g c c a g c t a c t c c t a c a a g t t t g g t g a t t g c g c c a t c t g g g t g t t c c t c t t c a a c a t g 420
25 t c g a a g a t c c t c g a g t t c g t c g a c a c c a t c t t c a t c g t c c c a g g a a g a c c a c c t c g g c 480
t t c c t c c a c t a c t a c c a c c a c a t c a t c a c c t a c t c c t t c t g c c t c t a c g c c g g c c a g t a c 540
a t g c a c c a c t a c a a c t g t g g c g g c t a t t t c t t c t g c c t c a t g a a c t t c t t c g t c c a c g g c 600
30 a t c a t g t a c t t c t a c t a c g c t c t c c g c t c c a t g g g c t t c c g t c c c t c c t t c g a t a t t g g c 660
a t c a c c t t c c t c c a g a t t t t g c a a a t g g t g c t c g g c g t g g c c a t c a t c a c c a t c t c c g c c 720
g g c t g c g a g a a g g t g g a c c c c a t c g g a a c g a c c t t c g g c t a c t t t a t t t a t t t c t c g t t c 780
35 t t c g t c c t c t t c t g c a a g t t c t t c t a c t a c c g c t a c a t c g c c a c g c c c g c c a a g a a g c c c 840
g a g g c c g c c g c c a a g t c g c c a g c c a c c a a g c c c a a g a g g a a g c a c g a c t a a 891

<210> 25
40 <211> 777
<212> DNA


<213> I s o c h r y s i s g a l b a n a

45 <400> 25
at g g c c c t c g c a a a c g a c g c g g g a g a g c g c a t c t g g g c g g c t g t g a c c g a c c c g g a a a t c 60
50 c t c a t t g g c a c c t t c t c g t a c t t g c t a c t c a a a c c g c t g c t c c g c a a t t c c g g g c t g g t g 120
g a t g a g a a g a g g g c g c a t a c a g g a c g t c c a t g a t c t g g t a c a a c g t t c t g c t g g c g c t c 180
t t c t c t g c g c t g a g c t t c t a c g t g a c g g c g a c c g c c c t c g g c t g g g a c t a t g g t a c g g g c 240
55 g c g t g g c t g c g c a g g c a a a c c g g c g a c a c a c c g c a g c c g c t c t t c c a g t g c c c g t c c c c g 300
g t t t g g g a c t c g a a g c t c t t c a c a t g g a c c g c c a a g g c a t t c t a t t a c t c c a a g t a c g t g 360


gagt acct cg acacggcct g gct gagggc t cct t t ct cc aggcct t cca ccaact t t ggc 420
 5 gcgccgt ggg at gt gt acct cggcat t cgg ct gcacaacg agggcgt at g gat ct t cat g 480
 t t t t t caact cgt t cat t ca caccat cat g t acacct act acggcct cac cgccgccggg 540
 t at aagt t ca aggccaagcc gct cat cacc gcgat gcaga t ct gccagt t cgt gggcggc 600
 10 t t cct gt t gg t ct gggact a cat caacgt c ccct gct t ca act cggacaa agggaagt t g 660
 t t cagct ggg ct t t caact a t gcat acgt c ggct cggc t t ct t gct ct t ct gccact t t 720
 t t ct accagg acaact t ggc aacgaagaaa t cggccaagg cgggcaagca gct ct ag 777

Claims

1. A process for the production of a compound of formula (I):



in an organism, the process comprising introducing into an organism which comprises a lipid of general formula (III):




at least one nucleic acid sequence comprising :

- a) SEQ ID NO: 3;
- b) a nucleic acid sequence which hybridizes under stringent conditions with a nucleic acid sequence of SEQ ID NO: 3.
- c) a nucleic acid sequence which encodes a polypeptide of SEQ ID NO: 4; or
- d) a derivative of a nucleic acid sequence of SEQ ID NO: 3 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 4, wherein said polypeptide has $\Delta 5$ -desaturase activity;

and expressing said nucleic acid sequence;
 wherein, in general formulae (I) and (III):


Y = a C₁₀-C₁₈ hydrocarbon chain containing up to four carbon-carbon double bonds;
 R¹ = hydroxyl, coenzyme A (thioester), lysophosphatidylcholine, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysodiphosphatidylglycerol, lysophosphatidylserine, lysophosphatidylinositol, sphingo base or a radical of the formula II:



in which

R² = hydrogen, lysophosphatidyl choline, lysophosphatidylethanolamine, lysophosphatidylglycerol, lysodi-phosphatidylglycerol, lysophosphatidylserine, lysophosphatidylinositol or saturated or unsaturated C₂-C₂₄-alkylcarbonyl,

R³ = hydrogen, saturated or unsaturated C₂-C₂₄-alkylcarbonyl, or R² and R³ independently of one another are a radical of the formula Ia:



in which

n = 2, 3, 4, 5, 6, 7 or 9, m = 2, 3, 4, 5 or 6 and p = 0 or 3;


and wherein an oxygen in the R¹ radical may be replaced by sulphur such that R¹ is bonded to the remainder of the molecule via a thioester linkage.

2. A process as claimed in claim 1 wherein, in the compound of formula (I), the -CH=CH- double bond is in the cis orientation.
3. A process as claimed in claim 1 or claim 2 wherein, in the moiety Y, the carbon-carbon double bonds are in the cis orientation.
4. A process as claimed in claim 1 or claim 2, wherein in the compound of general formula (I) Y is:


-(CH₂)₄-CH=CH-CH₂-CH=CH-(CH₂)₄-CH₃, such that the compound of general formula (I) is sciadonic acid or a derivative thereof;

-(CH₂)₄-CH=CH-CH₂-CH=CH-CH₂-CH=CH-CH₂-CH₃, such that the compound of general formula (I) is juniperonic acid or a derivative thereof.

5. A process for the production of a compound of general formula (I) as defined in claim 1 and in which Y is:



or



in an organism which comprises linoleic (C18:2 n-6) or linolenic (C18:3 n-3) acid, the process comprising introducing into the organism:

at least one nucleic acid sequence comprising :

a) SEQ ID NO: 3;

b) a nucleic acid sequence which hybridizes under stringent conditions with a nucleic acid sequence of

SEQ ID NO: 3.

c) a nucleic acid sequence which encodes a polypeptide of SEQ ID NO: 4; or

d) a derivative of a nucleic acid sequence of SEQ ID NO: 3 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 4 wherein said polypeptide has $\Delta 5$ -desaturase activity; and

5

at least one nucleic acid sequence encoding a polypeptide having $\Delta 9$ elongase activity; and expressing said nucleic acid sequences;

10

6. A process as claimed in claim 5, wherein the nucleic acid sequence encoding a polypeptide having $\Delta 9$ elongase activity comprises a sequence encoding the C18- $\Delta 9$ elongase from *Isochrysis galbana* (SEQ ID NO: 25) or *Acanthamoeba castellanii* (SEQ ID NO: 24)

15

7. The process according to any of claims 1 to 6, wherein the substituents R2 or R3 independently of one another are saturated or unsaturated C18-C22-alkylcarbonyl.

20

8. The process according to any of claims 1 to 7, wherein the substituents R2 or R3 independently of one another are unsaturated C18-, C20- or C22-alkylcarbonyl with at least two double bonds.

25

9. The process according to any of claims 1 to 8, wherein the transgenic organism is a transgenic microorganism or a transgenic plant.

30

10. The process according to any of claims 1 to 9, wherein the transgenic organism is an oil-producing plant, a vegetable plant or an ornamental.

11. The process according to any of claims 1 to 10, wherein the transgenic organism is a transgenic plant selected from the group of the plant families: Adelotheceaceae, Anacardiaceae, Asteraceae, Apiaceae, Betulaceae, Boraginaceae, Brassicaceae, Bromeliaceae, Caricaceae, Cannabaceae, Convolvulaceae, Chenopodiaceae, Cryptocodiniaceae, Cucurbitaceae, Ditrichaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Lauraceae, Leguminosae, Linaceae or Prasinophyceae.

35

12. The process according to any of claims 1 to 11, wherein the compounds of the formula I are isolated from the organism in the form of their oils, lipids or free fatty acids.

13. The process according to any of claims 1 to 12, wherein the compounds of the formula I are isolated in a concentration of at least 1% by weight based on the total lipid content of the transgenic organism.

40

14. The oil, lipid or fatty acid, or a fraction thereof, produced by the process according to any of claims 1 to 13.

15. An oil, lipid or fatty acid composition which comprises an oil, lipid or fatty acid produced by a process as claimed in any one of claims 1 to 14 and which is derived from transgenic plants.

45

16. A process for the production of oils, lipids or fatty acid compositions by mixing oils, lipids or fatty acids according to claim 15 or oil, lipid or fatty acid compositions according to claim 15 with animal oils, lipids or fats.

17. An isolated nucleic acid sequence which encodes a polypeptide with $\Delta 5$ -desaturase activity and which is selected from the group consisting of SEQ ID NO: 3.

50

18. A derivative of a nucleic acid sequence of SEQ ID NO:3 which encodes a polypeptide with at least 40% identity at the amino acid level with SEQ ID NO: 4, wherein said polypeptide has $\Delta 5$ -desaturase activity.

19. An amino acid sequence which is encoded by a nucleic acid sequence according to claim 17 or claim 18.

55

20. A nucleic acid construct comprising a nucleic acid sequence according to claim 17 or claim 18, operably linked with one or more regulatory sequences.

21. A gene construct according to claim 20, further comprising a nucleic acid sequence encoding a polypeptide having $\Delta 9$ -elongase activity.

EP 2 388 332 A1

22. A gene construct according to claim 21, wherein the nucleic acid sequence encoding a polypeptide having $\Delta 9$ elongase activity comprises a sequence encoding the C18- $\Delta 9$ elongase from *Isochrysis galbana* (SEQ ID NO:25) or *Acanthamoeba castellanii* (SEQ ID No:24).
- 5 23. A gene construct according to claim 21 or claim 22, wherein the nucleic acid construct comprises additional biosynthesis genes of the fatty acid or lipid metabolism selected from the group acyl-CoA dehydrogenase(s), acyl-ACP [= acyl carrier protein] desaturase(s), acyl-ACP thioesterase(s), fatty acid acyltransferase(s), acyl-CoA:lysophospholipid acyltransferase(s), fatty acid synthase(s), fatty acid hydroxylase(s), acetyl-coenzyme A carboxylase(s), acyl-coenzyme A oxidase(s), fatty acid desaturase(s), fatty acid acetylenases, lipoxygenases, triacylglycerol lipases, allenoxide synthases, hydroperoxide lyases or fatty acid elongase(s).
- 10 24. The gene construct according to claim 22 or claim 23, wherein the nucleic acid construct comprises additional biosynthesis genes of the fatty acid or lipid metabolism selected from the group $\Delta 4$ -desaturase, $\Delta 5$ -desaturase, $\Delta 6$ -desaturase, $\Delta 8$ -desaturase, $\Delta 9$ -desaturase, $\Delta 12$ -desaturase or $\Delta 6$ -elongase.
- 15 25. A vector comprising a nucleic acid according to claim 17 or claim 18 or a gene construct according to any one of claims 20 to 24.
- 20 26. A transgenic nonhuman organism comprising at least one nucleic acid according to claim 17 or claim 18, a gene construct according to any one of claims 19 to 23 or a vector according to claim 24.
- 25 27. The transgenic nonhuman organism according to claim 26, which organism is a microorganism, a nonhuman animal or a plant.
- 30 28. The transgenic nonhuman organism according to claim 26 or claim 27, which organism is a plant.
- 35
- 40
- 45
- 50
- 55

```

alcoa21 : ~~~~~*~~~~~20~~~~~*~~~~~40~~~~~*~~~~~60~~~~~*~~~~~80~~~~~*
alcoa11 : ~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*
calcoa2 : ~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*
ADS1 : ~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*
pined9 : MSALVLSLAFSTPARGGESREAEKALRRRLNASSCNASKNQAQLMCERRILSMCYGIRRRNSAGLWGKATRVQV--SVAVEEEGGRILLS : 90
limd5 : ~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~*~~~~~* : 65

alcoa21 : DVPKDKTKITVV-RVNNSTDIHFHILVGGLEHVLCDSAPBETRSMSAFMLSLTLY-AVCEVFETTLTSLYHRNDTHRSEKLPKYLEYFFAYVGL : 110
alcoa11 : EWPVKEKTKITVV-RKNNSTDVHFHIFIGGLHVLCDFAPSTESMSKSFVCFALY-AICEVFETTLTSLYHRNDTHRSEKLPKYLEYFFAYVGL : 113
calcoa2 : DVQVIRR-PQVYMG-RKNNSTVDTATGVSVLGMVHILSLEAPETESMSALSMAFVLY-VVTGLFETTLTSLYHRNLAHKAFLPKKMLEYLEAYCGA : 177
ADS1 : KAENGRK-KRAMME-RKNNSTVDTATGVSVLGMVHILSLEAPETESMSALSMAFVLY-VVTGLFETTLTSLYHRNLAHKAFLPKKMLEYLEAYCGA : 106
pined9 : DVVKKKRPPELLLELRMSLGDIGAAGTVFGMHADCLLAPPETMKAAGVFAVLY-VVTGLFETTLTSLYHRNLAHKAFLPKKMLEYLEAYCGV : 181
limd5 : EVVLERKAKGF-RRKNNPRDIQNAVTLILVLEHAAAMABEYESMDAFWISFILLGFASEVLEITTCFHECTHGGCKPKLVLEYEFAAYCGS : 155

alcoa21 : HALQGDPEVMMVSTHRYHHKFTDTYLDPHSEIEGEMPSHIFMLEDSKYIIEE-CGRYENAGDLLKQSYRLELRTEVPHVYVLAALLLYMFSGE : 201
alcoa11 : HALQGDPEVMMVSTHRYHHKFTDTYLDPHSEIEGEMPSHIFMLEDSKYIIEE-CGRYENAGDLKQSYRLELRTEVPHVYVLAALLLYMFSGE : 204
calcoa2 : QALQGNFIDMMVSTHRYHHQFCDSERDPHSPEEGEMPSHISMFEDTKSLTEK-CGVPNNVSDLEKQPFYKEMKRYPLHPHIALAAVLYIALSGE : 268
ADS1 : IALQGDPEVMMVSTHRYHHQFTDSDRDPHSPEEGEMPSHISMFEDTKSLTEK-CGRRTNVEDDLKRWYKQRYKQRYVLYHLLTFGDLIYFSGE : 197
pined9 : QAVQGDPELDMMVSTHRYHHHEYCDVSKDPHSPEEGEMPSHISMSMEDEKTMFDR-VGTRNNVSDLEMQPFYRFRIRDYIIHPHIAAGLILYALSGP : 272
limd5 : IALQGDPEVMMVSNHRYHHQFVDTERDVHSFTQSEMECHICGVLDKDLFVEKRGERRNNVNDLKKQAFKRELRQRYMYHQLALIALLYVYSGE : 247

alcoa21 : PEIVMGAVRTVWGFHASMLVNSVCHRYEHOAMDGDLSLNNMFIAMLTSEGGMNHHAFEYSARHGLEMMQIDPTMYVTKLLEYLGLATD : 293
alcoa11 : PEIVMGAVRTILGFHLSSMLVNSVCHRMEVNRPNPCDLSLNNMFIAMLTSEGGMNHHAFEYSARHGLEMMQIDPTMYVTKLLEYLGLATD : 296
calcoa2 : PEIVMGAVRIVWVYHITWLVNSACHVMSQSNNFTGLSRNNMVALLAFSEGGMNHHAFEFSSARQGLEMMQIDPTMYVTKLLEYLGLATD : 360
ADS1 : SELTMGGIGVAMEHHTVCLINSLCHVMGSRMKNDPTSRNNMWSVFSFEGSMNHHAFESSARQGLEMMQIDPTMYVTKLLEYLGLATD : 289
pined9 : PEIVMGAVRIVWVYHITWLVNSAAHVMEYQANNFTGLSRNNMVALALAFSEGGMNHHAFQYSARHGLEMMQIDPTMYVTKLLEYLGLATD : 364
limd5 : PEIVMGAVRIVWVYHITWLVNSAAHVMEYQANNFTGLSRNNMVALALAFSEGGMNHHAFQYSARHGLEMMQIDPTMYVTKLLEYLGLATD : 339

```

Figure 1

	*	380	*	
alcoa21	:	MKVPSEVHKRKM	SKNCVQDKQFCVNDK*	: 321
alcoa11	:	IKVFSIHKRKM	SFKN*	: 312
calcoa2	:	MKVPSEVQKQRM	CFSE*	: 376
ADS1	:	MKLFSESQRRRM	AMVR	: 305
pined9	:	MRAPLEQHKTRM	SSKPGL	: 382
limd5	:	MKLPTEAQKQKLLKAKSA		: 356

Figure 1 cont.

Figure 2




Figure 3






Figure 3 (contd.)





Figure 4




Figure 5




Figure 6




Figure 7




Figure 7 (contd.)





Figure 8






Figure 9a




Figure 9b

Figure 10





Figure 11





EUROPEAN SEARCH REPORT

Application Number
EP 10 18 7573

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	US 2004/224413 A1 (CAHOON EDGAR B [US] ET AL) 11 November 2004 (2004-11-11) * the whole document *	1-3,5, 7-12,14, 15, 18-20, 25-28	INV. C12N15/82 C12N9/02 A01H5/00 C12N5/10 C12P7/64
X	MARILLIA E-F ET AL: "Expression of meadowfoam Des5 and FAE1 genes in yeast and in transgenic soybean somatic embryos, and their roles in fatty acid modification", PLANT PHYSIOLOGY AND BIOCHEMISTRY, GAUTHIER-VILLARS, PARIS, FR, vol. 40, no. 10, 2002, pages 821-828, XP002288917, ISSN: 0981-9428 * abstract, p. 822-825 *	14-16	
X	QI B ET AL: "Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants", NATURE BIOTECHNOLOGY, NATURE PUBLISHING GROUP, NEW YORK, NY, US, vol. 22, no. 6, June 2004 (2004-06), pages 739-745, XP002348313, ISSN: 1087-0156 * abstract, p. 739-743, fig. 1,3,4, table 1 *	14-16	TECHNICAL FIELDS SEARCHED (IPC) C12N C12P
The present search report has been drawn up for all claims			
3	Place of search Munich	Date of completion of the search 8 August 2011	Examiner Puonti-Kaerlas, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.02 (P04C01)



EUROPEAN SEARCH REPORT

 Application Number
 EP 10 18 7573

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WU G ET AL: "STEPWISE ENGINEERING TO PRODUCE HIGH YIELDS OF VERY LONG-CHAIN POLYUNSATURATED FATTY ACIDS IN PLANTS", NATURE BIOTECHNOLOGY, NATURE PUBLISHING GROUP, NEW YORK, NY, US, vol. 23, no. 8, August 2005 (2005-08), pages 1013-1017, XP007900180, ISSN: 1087-0156 * p. 1013-1016, fig. 1 table 1 * -----	14-16	TECHNICAL FIELDS SEARCHED (IPC)
X	WOLFF ROBERT L ET AL: "General characteristics of Pinus spp. seed fatty acid compositions, and importance of DELTA5-olefinic acids in the taxonomy and phylogeny of the genus", LIPIDS, vol. 35, no. 1, January 2000 (2000-01), pages 1-22, XP009079462, ISSN: 0024-4201 * the whole document * -----	14-16	
X	WOLFF R L ET AL: "ARACHIDONIC, EICOSAPENTAENOIC, AND BIOSYNTHETICALLY RELATED FATTY ACIDS IN THE SEED LIPIDS FROM A PRIMITIVE GYMNOSPERM, AGATHIS ROBUSTA", LIPIDS, CHAMPAIGN, IL, US, vol. 34, no. 10, 1999, pages 1083-1097, XP001015626, ISSN: 0024-4201 * the whole document * ----- -/--	14-16	
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 8 August 2011	Examiner Puonti-Kaerlas, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

 3
 EPO FORM 1503 03.82 (P04C01)



EUROPEAN SEARCH REPORT

 Application Number
 EP 10 18 7573

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
A	I. HEILMANN: "Identification of the Arabidopsis Palmitoyl-Monogalactosyldiacylglycerol 7-Desaturase Gene FAD5, and Effects of Plastidial Retargeting of Arabidopsis Desaturases on the fad5 Mutant Phenotype", PLANT PHYSIOLOGY, vol. 136, no. 4, 1 December 2004 (2004-12-01), pages 4237-4245, XP55004199, ISSN: 0032-0889, DOI: 10.1104/pp.104.052951 -----	1-28	TECHNICAL FIELDS SEARCHED (IPC)
A	MARILLIA E-F ET AL: "A desaturase-like protein from white spruce is a DELTA<9> desaturase", FEBS LETTERS, ELSEVIER, AMSTERDAM, NL, vol. 526, no. 1-3, 28 August 2002 (2002-08-28), pages 49-52, XP004379222, ISSN: 0014-5793, DOI: 10.1016/S0014-5793(02)03112-5 -----	1-28	
A	WO 2005/012316 A (BASF PLANT SCIENCE GMBH [DE]; ZANK THORSTEN [DE]; BAUER JOERG [DE]; CI) 10 February 2005 (2005-02-10) -----	1-28	
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 8 August 2011	Examiner Puonti-Kaerlas, J
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

3

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 10 18 7573

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-08-2011

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2004224413 A1	11-11-2004	US 2009138992 A1 US 6838594 B1	28-05-2009 04-01-2005
-----	-----	-----	-----
WO 2005012316 A	10-02-2005	BR PI0413073 A CA 2533613 A1 CN 1860211 A EP 1654344 A2 EP 2166067 A2 EP 2166068 A2 EP 2166069 A2 EP 2166089 A2 EP 2172536 A2 EP 2169052 A2 EP 2169053 A2 EP 2166090 A2 EP 2166070 A2 EP 2166071 A2 JP 2007500504 A JP 2011087578 A US 2008155705 A1	03-10-2006 10-02-2005 08-11-2006 10-05-2006 24-03-2010 24-03-2010 24-03-2010 24-03-2010 07-04-2010 31-03-2010 31-03-2010 24-03-2010 24-03-2010 24-03-2010 18-01-2007 06-05-2011 26-06-2008
-----	-----	-----	-----

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- GB 0523915 A [0009]
- GB 0524420 A [0009]
- US 5565350 A [0055]
- WO 0015815 A [0055]
- EP 0388186 A [0098]
- EP 0335528 A [0098]
- WO 9321334 A [0098]
- EP 0249676 A [0098]
- US 5608152 A [0099] [0102] [0134]
- WO 9845461 A [0099] [0102] [0134]
- US 5504200 A [0100] [0102] [0134]
- WO 9113980 A [0100] [0102] [0134]
- WO 9515389 A [0100] [0102] [0134]
- WO 9523230 A [0100] [0102] [0134]
- WO 9916890 A [0100] [0101] [0102] [0134]
- US 5315001 A [0102]
- WO 9218634 A [0102]
- WO 9320216 A [0102]
- US 5677474 A [0102]
- US 5530149 A [0102]
- EP 571741 A [0102]
- JP 6062870 A [0102]
- WO 9808962 A [0102]
- US 5689040 A [0102]
- EP 781849 A [0102]
- WO 9519443 A [0103] [0132]
- DE 10102337 [0104]
- DE 10102338 [0104]
- WO 9801572 A [0120]
- US 5352605 A [0130]
- WO 8402913 A [0130]
- US 4962028 A [0130]
- US 5187267 A [0133]
- WO 9612814 A [0133]
- EP 0375091 A [0133]
- WO 9516783 A [0136]
- WO 9706250 A [0136]
- WO 9946394 A [0136]

Non-patent literature cited in the description

- WOLFF et al. *Lipids*, October 1999, vol. 34 (10), 1083-1097 [0004]
- ASSET et al. *Lipids*, 1999, vol. 34, 39-44 [0005]
- BERGER et al. *Lipids in Health and Disease*, 2002, vol. 1, 5 [0005]
- TANAKA et al. *Biochem. Biophys. Res. Comm.*, 1999, vol. 264, 683-688 [0005]
- TANAKA et al. *Eur. J. Biochem.*, 2001, vol. 268, 4928-4939 [0005]
- YU, R. et al. *Lipids*, 2000, vol. 35, 1061-1064 [0007]
- TAKEYAMA, H. et al. *Microbiology*, 1997, vol. 143, 2725-2731 [0007]
- ZANK, T.K. et al. *Plant Journal*, 2002, vol. 31, 255-268 [0008]
- SAKURADANI, E. et al. *Gene*, 1999, vol. 238, 445-453 [0008]
- SPRECHER. *Biochim. Biophys. Acta*, 2000, vol. 1486, 219-231 [0008]
- MOREAU et al. *Arch Biochem Biophys.*, 1981, vol. 209, 376-384 [0011]
- CAHOON et al. *Plant Physiology*, 2000, vol. 124, 243-251 [0011] [0177]
- CAHOON et al. *Plant Physiol.*, September 2000, vol. 124 (1), 243-51 [0011]
- FUKUCHI-MIZUTANI et al. *Plant and Cell Physiology*, vol. 39, 247-253 [0013]
- HEILMANN et al. *PNAS*, 2004, vol. 101 (28), 10266-10271 [0014]
- YAO et al. *Plant Biotechnol*, 2003, vol. 1, 221-229 [0014]
- MARILLIA et al. *FEBS Lett*, 2002, vol. 526, 49-52 [0014]
- DESTAILLATS et al. *Lipids*, November 2001, vol. 36 (11), 1247-1254 [0015]
- WOLFF et al. *Lipids*, May 2001, vol. 36 (5), 439-451 [0015]
- WOLFF et al. *Lipids*, January 2002, vol. 37 (1), 17-26 [0015]
- TSEVEGSUREN et al. *Hrc-Journal of High Resolution Chromatography*, 1997, vol. 20, 237-241 [0015]
- SMITH et al. *Lipids*, 1968, vol. 3, 37 [0016]
- WHITNEY et al. *Planta*, 2003, vol. 217 (6), 983-992 [0016]
- Current Protocols in Molecular Biology. John Wiley & Sons, 1989, 6.3.1-6.3.6 [0019]
- SAMBROOK et al. *Molecular Cloning*. Cold Spring Harbor Laboratory, 1989 [0019]
- Nucleic Acids Hybridization: A Practical Approach. IRL Press at Oxford University Press, 1985 [0019]
- Essential Molecular Biology: A Practical Approach. IRL Press at Oxford University Press, 1991 [0019]
- J. Mol. Evolution. 1987, vol. 25, 351-360 [0023]

- **HIGGINS et al.** *CABIOS*, 1989, vol. 5, 151-153 [0023]
- **NEEDLEMAN ; WUNSCH.** *J. Mol. Biol.* 1970, vol. 48, 443-453 [0023]
- **SMITH ; WATERMAN.** *Adv. Appl. Math.*, 1981, vol. 2, 482-489 [0023]
- **GOEDDEL.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185 [0058] [0120]
- **GOTTESMAN, S.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185, 119-128 [0059]
- **CHMIEL.** *Bioprozeßtechnik 1. Einführung in die Bioverfahrenstechnik.* Gustav Fischer Verlag, 1991 [0074]
- **STORHAS.** *Bioreaktoren und periphere Einrichtungen.* Vieweg Verlag, 1994 [0074]
- *Manual of Methods für General Bacteriology.* American Society for Bacteriology, 1981 [0075]
- *Applied Microbiol. Physiology, A Practical Approach.* IRL Press, 1997, 53-73 [0082]
- **FRANCK et al.** *Cell*, 1980, vol. 21, 285-294 [0098] [0130]
- **WARD et al.** *Plant. Mol. Biol.*, 1993, vol. 22 [0098]
- **GATZ.** *Plant J.*, 1992, vol. 2, 397-404 [0098]
- **STOCKHAUS et al.** *EMBO J.*, 1989, vol. 8, 2445 [0098]
- **BAEUMLEIN et al.** *Plant J.*, 1992, vol. 2 (2), 233-239 [0100]
- **BÄUMLEIN et al.** *Mol. Gen Genet.*, 1991, vol. 225 (3 [0102]
- **BÄUMLEIN et al.** *Plant J.*, 1992, 2, 2 [0102]
- **GATZ.** *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 1997, vol. 48, 89-108 [0103] [0132]
- **GATZ et al.** *Plant J.*, 1992, vol. 2, 397-404 [0103] [0132]
- **HELLENS et al.** *Trends in Plant Science*, 2000, vol. 5, 446-451 [0111]
- *Plant Molecular Biology and Biotechnology.* CRC Press, 1993, 71-119 [0112]
- *Vectors for Gene Transfer in Higher Plants.* **F.F. WHITE.** *Transgenic Plants, Vol. 1, Engineering and Utilization.* Academic Press, 1993, vol. 1, 15-38 [0112]
- *Techniques for Gene Transfer.* **B. JENES et al.** *Transgenic Plants, Vol. 1, Engineering and Utilization.* Academic Press, 1993, vol. 1, 128-143 [0112]
- **POTRYKUS.** *Annu. Rev. Plant Physiol. Plant Molec. Biol.*, 1991, vol. 42, 205-225 [0112] [0120]
- **GRUBER ; CROSBY.** *Methods in Plant Molecular Biology and Biotechnology.* CRC Press, vol. 7, 89-108 [0120]
- **ROMANOS, M.A. et al.** *Foreign gene expression in yeast: a review.* *Yeast*, vol. 8, 423-488 [0120]
- *Heterologous gene expression in filamentous fungi.* **VAN DEN HONDEL, C.A.M.J.J. et al.** *More Gene Manipulations in Fungi.* Academic Press, 1991, 396-428 [0120]
- *Gene transfer systems and vector development for filamentous fungi.* **VAN DEN HONDEL, C.A.M.J.J. ; PUNT, P.J. et al.** *Applied Molecular Genetics of Fungi.* Cambridge University Press, 1991, 1-28 [0120]
- **FALCIATORE et al.** *Marine Biotechnology*, 1999, vol. 1 (3), 239-251 [0120] [0127]
- **SCHMIDT, R. ; WILLMITZER, L.** *High efficiency Agrobacterium tumefaciens-mediated transformation of Arabidopsis thaliana leaf and cotyledon explants.* *Plant Cell Rep.*, 1988, 583-586 [0120]
- *Plant Molecular Biology and Biotechnology.* C Press, 1993, 71-119 [0120]
- *Techniques for Gene Transfer.* **F.F. WHITE ; B. JENES et al.** *Transgenic Plants, Vol. 1, Engineering and Utilization.* Academic Press, 1993, 128-43 [0120]
- **JOHNSON, K.S.** *Gene*, 1988, vol. 67, 31-40 [0121]
- **AMANN et al.** *Gene*, 1988, vol. 69, 301-315 [0122]
- **STUDIER et al.** *Gene Expression Technology: Methods in Enzymology.* Academic Press, 1990, vol. 185, 60-89 [0122]
- **BALDARI et al.** *Embo J.*, 1987, vol. 6, 229-234 [0124]
- **KURJAN ; HERSKOWITZ.** *Cell*, 1982, vol. 30, 933-943 [0124]
- **SCHULTZ et al.** *Gene*, 1987, vol. 54, 113-123 [0124]
- *Gene transfer systems and vector development for filamentous fungi.* **VAN DEN HONDEL, C.A.M.J.J. ; PUNT, P.J. et al.** *Applied Molecular Genetics of fungi.* Cambridge University Press, 1991, 1-28 [0124]
- *More Gene Manipulations in Fungi.* Academic Press, 396-428 [0124]
- **SMITH et al.** *Mol. Cell Biol.*, 1983, vol. 3, 2156-2165 [0125]
- **LUCKLOW ; SUMMERS.** *Virology*, 1989, vol. 170, 31-39 [0125]
- *Cloning Vectors.* Elsevier, 1985 [0126]
- **SAMBROOK, J. ; FRITSCH, E.F. ; MANIATIS, T.** *Molecular Cloning: A Laboratory Manual.* Cold Spring Harbor Laboratory, 1989 [0126]
- **BECKER, D. ; KEMPER, E. ; SCHELL, J. ; MASTERSON, R.** *New plant binary vectors with selectable markers located proximal to the left border.* *Plant Mol. Biol.*, 1992, vol. 20, 1195-1197 [0127]
- **BEVAN, M.W.** *Binary Agrobacterium vectors for plant transformation.* *Nucl. Acids Res.*, 1984, vol. 12, 8711-8721 [0127]
- *Vectors for Gene Transfer in Higher Plants.* *Transgenic Plants, Vol. 1, Engineering and Utilization.* Academic Press, 1993, vol. 1, 15-38 [0127]
- **GIELEN et al.** *EMBO J.*, 1984, vol. 3, 835 [0128]
- **GALLIE et al.** *Nucl. Acids Research*, 1987, vol. 15, 8693-8711 [0129]
- **BENFEY et al.** *EMBO J.*, 1989, vol. 8, 2195-2202 [0130]
- **KERMODE.** *Crit. Rev. Plant Sci.*, 1996, vol. 15 (4), 285-423 [0131]

EP 2 388 332 A1

- **WARD et al.** *Plant. Mol. Biol.*, 1993, vol. 22, 361-366 [0133]
- **BAEUMLEIN et al.** *Mol Gen Genet*, 1991, vol. 225 (3), 459-67 [0134]
- **BAEUMLEIN et al.** *Plant Journal*, 1992, vol. 2 (2), 233-9 [0134]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1989 [0137]
- *Agrobacterium protocols. Methods in Molecular Biology*. Humana Press, 1995, vol. 44 [0137]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1989 [0142]
- **CHIRGWIN et al.** *Biochemistry*, 1979, vol. 18, 5294-5299 [0143]
- **F.C. NEIDHARDT et al.** *E. coli and Salmonella*. ASM Press, 1996, 612-636 [0149]
- *Biology of Prokaryotes*. Thieme, 1999 [0149]
- **MAGNUSON, K. et al.** *Microbiological Reviews*. 1993, vol. 57, 522-542 [0149]
- **FRENTZEN.** *Lipid*, 1998, vol. 100 (4-5), 161-166 [0154]
- **KINNEY.** *Genetic Engineering*. 1997, vol. 19, 149-166 [0154]
- **OHLROGGE ; BROWSE.** *Plant Cell*, 1995, vol. 7, 957-970 [0154]
- **SHANKLIN ; CAHOON.** *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 1998, vol. 49, 611-641 [0154]
- **VOELKER.** *Genetic Engineering*. 1996, vol. 18, 111-13 [0154]
- **GERHARDT.** *Prog. Lipid R.* 1992, vol. 31, 397-417 [0154]
- **GÜHNEMANN-SCHÄFER ; KINDL.** *Biochim. Biophys Acta*, 1995, vol. 1256, 181-186 [0154]
- **KUNAU et al.** *Prog. Lipid Res.*, 1995, vol. 34, 267-342 [0154]
- **STYMNE et al.** *Biochemistry and Molecular Biology of Membrane and Storage Lipids of Plants*. American Society of Plant Physiologists, 1993, 150-158 [0154]
- **MURPHY ; ROSS.** *Plant Journal*, 1998, vol. 13 (1), 1-16 [0154]
- **SAYANOVA et al.** *PNAS*, 1997, vol. 94, 4211-4216 [0164]
- **SAYANOVA et al.** *FEBS Letters*, 2003, vol. 542, 100-104 [0169] [0173]
- **QI et al.** *Nature Biotechnology*, 2004, vol. 22, 739-745 [0172]
- **CLOUGH ; BENT.** *Plant Journal*, 1998, vol. 16, 735-743 [0172]
- **FAY ; RICHLI.** *Journal of Chromatography*, 1991, vol. 541, 89-98 [0173] [0178]
- **LARSON ; GRAHAM.** *Plant Journal*, 2001, vol. 25, 115-125 [0174]
- **DEBLAERE et al.** *Nucl. Acids. Res.*, 1984, vol. 13, 4777-4788 [0187]
- **VAR. DRAKKAR.** *NPZ Nordeutsche Pflanzenzucht* [0187]
- **MURASHIGE ; SKOOG.** *Physiol. Plant.*, 1962, vol. 15, 473 [0187]
- **BELL et al.** *In Vitro Cell. Dev. Biol.-Plant.*, 1999, vol. 35 (6), 456-465 [0188]
- **MLYNAROVA et al.** *Plant Cell Report*, 1994, vol. 13, 282-285 [0188]