



- (51) **International Patent Classification:**  
C12N1/00 (2006.01)
- (21) **International Application Number:**  
PCT/GB2013/052553
- (22) **International Filing Date:**  
1 October 2013 (01.10.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
1217524.6 1 October 2012 (01.10.2012) GB
- (71) **Applicant:** ROTHAMSTED RESEARCH LIMITED [GB/GB]; West Common, Harpenden, Hertfordshire, AL5 2JQ (GB).
- (72) **Inventors:** NAPIER, Johnathan A.; Department of Biological Chemistry, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ (GB). SAYANOVA, Olga; Department of Biological Chemistry, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ (GB). HAMILTON, Mary; Department of Biological Chemistry, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ (GB). VAEZI, Royah; Department of Biological Chemistry, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ (GB).
- (74) **Agent:** MARKS & CLERK LLP; 62-68 Hills Road, Cambridge, Cambridgeshire, CB2 1LA (GB).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**  
— with international search report (Art. 21(3))

(54) **Title:** RECOMBINANT ORGANISMS




FIGURE 1 20°C 60 μmol photons m<sup>-2</sup> s<sup>-1</sup>

(57) **Abstract:** The invention relates to genetically modified organisms with enhanced production of omega-3 long chain polyunsaturated fatty acids.

## Recombinant organisms

### Field of the invention

5 The invention relates to transgenic organisms, in particular transgenic microalgae, with enhanced production of omega-3 long chain polyunsaturated fatty acids, related methods and uses.

### Introduction

10

Long chain polyunsaturated fatty acids (LC-PUFAs) have a carbon backbone of at least 20 carbons in length and contain multiple double-bond desaturations. Long chain polyunsaturated fatty acids can be grouped into either an omega-3 (TO-3) or omega-6 (TO-6) category based on the position of the first double bond from the methyl, or TO,  
15 fatty acid terminus.

15

It is now well established that omega-3 LC-PUFAs, especially eicosapentaenoic acid (EPA; 20:5  $\Delta$ 5,8,11,14,17) and docosahexaenoic acid (DHA; 22:6  $\Delta$ 4,7, 10,13,16,19) are essential constituents of human nutrition and have key roles in growth and  
20 development of infants and children and in maintaining health through their effects on immune system (Voigt et al., 2000; Calder, 2003). There is growing evidence from clinical studies that the presence of omega-3 LC-PUFAs in the human diet has therapeutic effect in conditions such as cardiovascular diseases, obesity, metabolic syndrome and eczema (Navarro et al., 2000; Nugent, 2004; Das, 2002).

25

Although marine fish is the main dietary source of EPA and DHA, the depletion of fish stocks and pollution of the marine environment indicate an urgent need for an alternative and sustainable source of LC-PUFAs. Marine microorganisms are the primary producers of LC-PUFAs in the aquatic food chain and EPA- and DHA-rich  
30 microalgae have been demonstrated to be a promising alternative source to fish oils for human consumption. Thus, commercial cultivation of *Cryptocodinium cohnii* and *Schizochytrium* sp. have been successfully developed for DHA production and some marine microorganisms have demonstrated potential for the industrial production of EPA (*Nannochloropsis* species, *Phaeodactylum* species, *Nitzshia* spp.) (Harwood and  
35 Guschina, 2009). However, commercial production of highly valuable products like

omega-3 LC-PUFAs is expensive to maintain and represents a substantial technological challenge.

5 One of the approaches to increase the levels of LC-PUFAs is to use acyl-CoA dependent desaturases (Venegas-Caleron et al., 2010). In recent years, considerable focus has been placed on engineering higher plants for the production of very long chain polyunsaturated fatty acids (VLC-PUFAs) in their seed oils. Recently, the advantages of using an acyl-CoA-dependent  $\Delta 6$ -desaturase from *Ostreococcus tauri* (OtD6) to synthesize LC-PUFAs in transgenic Arabidopsis and Camelina plants have  
10 been demonstrated (Sayanova O., et al, 2012, Ruiz-Lopez N., et al., 2012). These studies indicate that the first step in the LC-PUFA pathway, the  $\Delta 6$ -desaturation, is rate-limiting.

15 As an alternative way of producing LC-PUFAs, there is increasing interest in the metabolic engineering of microalgae and genetic modification of algal strains represents a promising strategy to produce sustainable omega-3 oils. Effective recombinant engineering of microalgae to produce increased levels of LC-PUFAs for commercial production would address a global need and microalgae manipulated in this way would be useful as food additives and animal feed, including aquaculture, to  
20 meet global demand.

*Phaeodactylum tricorutum* is an unicellular diatom which accumulates up to 30% EPA and only traces of DHA and is considered a good source for the industrial production of EPA (Molina Grima et al., 1996). The first labelling experiments with [14C]acetate  
25 suggested that *P. tricorutum* synthesized EPA de novo by elongation and aerobic desaturation of fatty acids (Moreno et al., 1979). In pulse-chase experiments Arao and Yamada have demonstrated that EPA can be synthesized by 4 different routes and that the preferred route involved intermediates of both omega-6 and omega-3 pathways (Arao and Yamada, 1994). The majority of the EPA was found in galactolipids as  
30 opposed to neutral lipids such as triacylglycerol (Arao et al., 1987; Yongmanitchai and Ward, 1993). Recently, the genes encoding the  $\Delta 5$ - and  $\Delta 6$ -desaturases involved in EPA biosynthesis in *P. tricorutum* have been cloned and characterized (Domergue et al., 2002). It was shown that both desaturases were microsomal enzymes contributing equally to both pathways and they supported the preferred route acting simultaneously  
35 in omega-6 and omega-3 pathways. This suggests that  $\Delta 6$ - and  $\Delta 5$ - desaturation and

5  $\Delta 6$ - elongation involved in biosynthesis of EPA in *P. tricornutum* take place in the endoplasmic reticulum (ER) and newly synthesized EPA is imported after into the plastids. The presence of only minor amounts of all the intermediates of EPA biosynthetic pathway indicates that *P. tricornutum* have developed highly efficient mechanism towards the accumulation of EPA as a single end-product (Arao and Yamada, 1994). In several microalgae DHA can be synthesized by the elongation of EPA to docosapentaenoic acid (DPA; 22:5 $\Delta 7,10, 13, 16, 19$ ) by a specific  $\Delta \delta$ -elongase, with DPA then converted to DHA by a A4-desaturase.

10 The present invention is aimed at mitigating the shortcomings in the production of LC-PUFAs in various organisms, in particular in algae.

### Summary of the invention

15 The invention generally relates to transgenic organisms, in particular transgenic microalgae, with enhanced production of LC-PUFAs, in particular omega-3 LC-PUFAs such as DHA and/or EPA. The transgenic organisms, in particular transgenic microalgae, express one or more heterologous nucleic acid encoding for a polypeptide involved in the LC-PUFAs biosynthesis pathway. The invention also relates to methods  
20 for making transgenic organisms, in particular transgenic microalgae, uses of the transgenic organisms, in particular transgenic microalgae, and methods for increasing the production of LC-PUFAs, in particular omega-3 LC-PUFAs, more particular DHA and/or EPA in an organism, in particular microalgae. The invention also relates to isolated nucleic acids and their uses in methods for the enhanced production of LC-  
25 PUFAs, in particular omega-3 LC-PUFAs, in transgenic organisms.

The inventors have shown that microalgae can be manipulated using recombinant methods to produce an increased amount of LC-PUFAs, in particular EPA and DHA using heterologous gene expression. The inventors have surprisingly demonstrated  
30 that heterologous expression of  $\Delta \delta$ -elongase from *Ostreococcus tauri* alone results in increased accumulation of DHA in *P. tricornutum* with DHA levels in transgenic strains reaching up to 13% of total fatty acids. The inventors have also shown that overexpression of OtD6 in *P. tricornutum* has a positive effect on EPA levels. These findings provide evidence for the efficacy of expressing heterologous genes and  
35 enhancing the LC-PUFAs biosynthetic pathway through metabolic engineering in

transgenic microalgae. Furthermore, other organisms that make EPA/DHA, including animals and plants, can be manipulated in the same way by overexpression of  $\Delta 5$ -elongase from *Ostreococcus tauri*.

- 5 Accordingly, in one aspect, the invention relates to a transgenic microalgae with increased production of one or more omega-3 LC-PUFA. In one embodiment, the omega-3 LC-PUFA is selected from DHA and/or EPA. In another aspect, the invention relates to the use of a transgenic microalgae in producing omega-3 LC-PUFAs. In another aspect, the invention relates to a method for producing transgenic microalgae with increased omega-3 LC-PUFAs content. In another aspect, the invention relates to a method for increasing production of one of more omega-3 LC-PUFA in microalgae comprising
- 10
- a) introducing and expressing in a microalgae a heterologous nucleic acid,
  - b) cultivating said microalgae and
  - 15 c) obtaining said one of more omega-3 LC-PUFA from the transgenic microalgae.

In another aspect, the invention relates to a method for increasing production of DHA in microalgae. In another aspect, the invention relates to a method for increasing production of EPA in microalgae.

20

The invention also relates to an oil isolated from a microalgae described herein or a composition comprising a transgenic microalgae described or product therefrom herein and uses thereof.

25

In another aspect, the invention relates to a method for making a feedstuff comprising

- a) cultivating a transgenic microalgae described herein and
- b) obtaining said one of more omega-3 LC-PUFA from the transgenic microalgae.

30 In another aspect, the invention relates to an isolated nucleic acids comprising SEQ ID No. 7 or 9 encoding a  $\Delta 6$ -desaturase (Ost809A6) comprising SEQ ID No. 8 or 10, a functional variant thereof or a  $\Delta 6$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 10 and uses thereof. The invention also relates to

35 an isolated nucleic acid comprising SEQ ID No. 15 or 17 encoding a A4-desaturase

(Ost809A4) comprising SEQ ID No. 16 or 18, a functional variant thereof or a  $\Delta 4$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 18 and uses thereof. In another aspect, the invention relates to an isolated nucleic acid comprising SEQ ID No. 19 encoding A6-elongase (FcEL06) comprising SEQ ID No. 20, a functional variant thereof or a A6-elongase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 and an isolated nucleic acid comprising SEQ ID No. 21 encoding  $\Delta\delta$ -desaturase comprising SEQ ID No. 22, a functional variant thereof or a  $\Delta\delta$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22 and uses thereof.

In another aspect, the invention relates to the use of an isolated nucleic described herein in increasing the production of omega-3 LC-PUFAs, in particular DHA and/or EPA, in microalgae or higher plants.

Further, the invention relates to a transgenic organism, preferably a microalgae, with increased DHA levels expressing a heterologous  $\Delta\delta$ -elongase.

20

### Figures

The invention is further described in the following non-limiting figures.

**Fig. 1.** EPA content in WT and transgenic *P. tricornutum* expressing *O.tauri*  $\Delta 6$  desaturase under different growth conditions at two different growth stages: a) 20°C 60  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ; b) 20°C 25  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ; c) 18°C 25  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$

**Fig. 2a.** Total fatty acid composition of WT and transgenic *P. tricornutum* cells expressing OtElo5 during the exponential (E) and stationary (S) phases. Cultures were grown at 20°C under constant illumination 60  $\mu\text{mol photons. m}^{-2} \text{s}^{-1}$  with agitation. Each value represents the mean  $\pm$  SD of 3 separate experiments.

b. EPA< DPA and DHA content in WT and transgenic *P. tricornutum* expressing OtElo5. Cultures were grown at 20°C 60  $\mu\text{mol m}^{-2}\text{s}^{-1}$  under constant agitation at 70 rpm. Each measurement is the average of 3 biological replicates.

**Fig. 3.** The acyl-CoA profiles of WT (A) and transgenic *P. tricornutum* expressing the *Ostreococcus* Elo5 (B). The accumulation of LC-PUFA acyl-CoAs in B is boxed with a dotted line. The internal standard (istd) is 17:00 acyl-CoA.

35

**Fig. 4.** EPA and DHA content in the total FA extracts of WT and transgenic OtElo5 *P. tricornutum* cells.

**Fig. 5A.** The distribution of TAG species from WT and transgenic *P. tricornutum* at stationary phase of growth.

5 **Fig. 5B.** The distribution of TAG species from WT and transgenic *P. tricornutum* at different stages of growth.

**Fig. 6.** The distribution of DHA in TAG species from WT and transgenic *P. tricornutum* expressing OtElo5 at different stages of the growth cycle: A-DHA in specific TAGs; B-% of TAG containing DHA.

10 **Fig. 7.** Omega-3 PUFA biosynthetic pathway (schematic representation).

**Fig. 8.** Expression of Ost809A6-desaturase in transgenic yeast in the presence of the exogenous substrate 18:3n-3 (ALA). (BPX72 column). Note the conversion of ALA to the higher unsaturated form (SDA - arrowed). No conversion occurs with yeast strains containing the empty vector (pYES2 - C), and only when the expression of the Ost809 desaturase is induced by the addition of galactose (Gal +; B)

15 **Fig. 9.** Functional characterization of Ost809A6 in yeast (BPX72 column). Yeast cells supplemented with LA and ALA. Expression of *Ostreococcus 809*  $\Delta$ 6 in yeast, supplemented with both 18:2 (LA) and 18:3 (ALA). Note the specific conversion of ALA, but not LA, to a higher unsaturated. No conversion occurs with yeast strains containing the empty vector (pYES2 - C), and only when the expression of the Ost809 desaturase is induced by the addition of galactose (Gal +; B)

20 **Fig. 10.** FAMES profile of transgenic yeast expressing Ost809A4 desaturase in the presence of DPA (C22:5n-3). *Expression of Ostreococcus 809*  $\Delta$ 4 in yeast cells supplemented with exogenous 22:5 (DPA). Note the conversion of 22:5n-3 to the higher unsaturated form (22:6n-3; DHA - arrowed). No conversion occurs with yeast strains containing the empty vector (pYES2 - C), and only when the expression of the Ost809 D4 desaturase is induced by the addition of galactose (Gal +; B). NB. These C22 PUFAs are best resolved on a HP1 GC column - in this case, the (poly)unsaturated fatty acids eluted earlier than less saturated forms - this is the inverse compared to BPX72 column used above

25 **Fig. 11.** FAMES profile of transgenic yeast expressing FcElo6 (BPX72 column). Yeast were supplemented with 18:3n-6 (GLA). *Expression of Fragilariopsis cylindrus* Elo6 in yeast cells supplemented with exogenous 18:3 (GLA). Note the conversion of 18:3 ALA to the elongated form 20:3n-3 (arrowed). No conversion occurs with yeast strains

containing the empty vector (pYES2 - C), and only when the expression of the *Fragilariopsis* Elo6 is induced by the addition of galactose (Gal +; B).

**Fig. 12.** Phylogenetic tree showing relationship between n-3 specific Ost809A6 desaturase and other  $\Delta 6$ -desaturases.

5 **Fig. 13.** Expression of FcElo6 resulted in increase of DHA levels up to 14-17%. GC-MS analysis of total FA profiles from Pt cells expressing FcElo6.

**Fig. 14.** Schematic representation of vector system pPTOS2.

**Figure 15.** Co-expression of two heterologous omega-3 LC-PUFA biosynthetic activities in *P. tricornutum*. Fatty acid composition of Pt\_WT, pPhOS2.1 (expressing OtElo5) and pPhOS2.2 (expressing OtD6Pt and OtElo5) cells during the S phase of growth at 16°C and 20°C. Values are the average of three experiments (+/- standard error).

10

**Fig. 16.** Fatty acid composition of pPhOS\_Ppglut (expressing OtElo5 and Ppglucose transporter) cells during the S phase of growth at 20°C, 100  $\mu\text{mol l}^{-1}$  m<sup>-2</sup>s<sup>-1</sup> under constant agitation at 70 rpm. N=1 .

15

**Fig. 17.** Fatty acid composition of pPhOS\_Hsglut (expressing OtElo5 and human glucose transporter) cells during the S phase of growth at 20°C, 100  $\mu\text{mol l}^{-1}$  m<sup>-2</sup>s<sup>-1</sup> under constant agitation at 70 rpm. N=1 .

**Fig. 18.** Growth of Wt and pPhOS\_Ppglut Pt cells in the dark.

20

### Detailed description

The present invention will now be further described. In the following passages, different aspects of the invention are defined in more detail. Each aspect so defined may be combined with any other aspect or aspects unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous.

25

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of microbiology, tissue culture, molecular biology, chemistry, biochemistry and recombinant DNA technology, which are within the skill of the art. Such techniques are explained fully in the literature.

30

The invention relates to the genetic manipulation of the fatty acid biosynthetic pathway in microalgae. In particular, the invention relates to methods for increasing the

35



production of LC-PUFAs, in particular omega-3 LC-PUFAs, for example one of more omega-3 LC-PUFA in an organism, in particular in microalgae.

5 Polyunsaturated fatty acids can be classified into two major families, depending on the position (n) of the first double bond nearest the methyl end of the fatty acid carbon chain. Thus, the omega-6 fatty acids (TO-6) have the first unsaturated double bond six carbon atoms from the omega (methyl) end of the molecule and additionally have a total or two or more double bonds, with each subsequent unsaturation occurring 3 additional carbon atoms toward the carboxyl end of the molecule. In contrast, the  
10 omega-3 fatty acids (w-3) have the first unsaturated double bond three carbon atoms away from the omega end of the molecule and additionally have a total of three or more double bonds with each subsequent unsaturation occurring 3 additional carbon atoms towards the carboxyl end of the molecule.

15 Table I summarizes the common names of omega-3 fatty acids and the abbreviations that will be used throughout the specification:

Table I

Common Name	Abbreviation	Shorthand notation
oleic acid	OA	18:1 <sup>Δ9</sup>
Linoleic acid	LA	18:2 <sup>Δ9,12</sup>
γ-Linolenic acid	GLA	18:3 <sup>Δ6,9,12</sup>
di-homo γ-linolenic acid	DGLA	20:3 <sup>Δ8,11,14</sup>
Arachidonic acid	ARA	20:4 <sup>Δ5,8,11,14</sup>
α- linolenic acid	ALA	18:3 <sup>Δ9,12,15</sup>
stearidonic acid	SDA	18:4 <sup>Δ6,9,12,15</sup>
eicosatetraenoic acid	ETA	20:4 <sup>Δ8,11,14,17</sup>
eicosapentaenoic acid	EPA	20:5 <sup>Δ5,8,11,14,17</sup>
docosapentaenoic acid	DPA	22:5 <sup>Δ7,10,13,16,19</sup>
docosahexaenoic acid	DHA	22:6 <sup>Δ4,7,10,13,16,19</sup>

20 There are a number of enzymes that are involved in the omega-3 PUFA biosynthetic pathway as shown in figure 7. These include desaturases and elongases.

A variety of genes involved in oil production have been identified through genetic means in different organisms and the DNA sequences of some of these genes are publicly available. Non-limiting examples are shown below:

5	Accession No.	Description
	AY131238	<i>Argania spinosa</i> $\Delta 6$ -desaturase
	Y055118	<i>Echium pitardii</i> var. <i>pitardii</i> $\Delta 6$ -desaturase
	AY055117	<i>Echium gentianoides</i> $\Delta 6$ -desaturase
	AF296076	<i>Mucor rouxii</i> $\Delta 6$ -desaturase
10	AF007561	<i>Borago officinalis</i> $\Delta 6$ -desaturase
	L11421	<i>Synechocystis</i> sp $\Delta 6$ -desaturase
	NM_031344	<i>Rattus norvegicus</i> $\Delta 6$ fatty acid desaturase
	AF465283,	<i>Mortierella alpine</i> $\Delta 6$ fatty acid desaturase
	AF465282	<i>Mortierella isabellina</i> $\Delta 6$ fatty acid desaturase
15	AF419296	<i>Pythium irregulare</i> $\Delta 6$ fatty acid desaturase
	AB052086	<i>Mucor circinelloides</i> D6d mRNA for $\Delta 6$ fatty acid desaturase
	AJ250735	<i>Ceratodon purpureus</i> mRNA for $\Delta 6$ fatty acid desaturase
	AF126799	<i>Homo sapiens</i> $\Delta 6$ fatty acid desaturase
20	AF126798	<i>Mus musculus</i> $\Delta 6$ fatty acid desaturase
	AF199596,	<i>Homo sapiens</i> $\Delta 5$ desaturase
	AF320509	<i>Rattus norvegicus</i> liver $\Delta 5$ desaturase
	AB072976	<i>Mus musculus</i> D5D mRNA for $\Delta 5$ desaturase
	AF489588	<i>Thraustochytrium</i> sp. ATCC21685 $\Delta 5$ desaturase
25	AJ510244	<i>Phytophthora megasperma</i> mRNA for $\Delta 5$ fatty acid desaturase
	AF419297	<i>Pythium irregulare</i> $\Delta 5$ fatty acid desaturase
	AF07879	<i>Caenorhabditis elegans</i> $\Delta 5$ fatty acid desaturase
	AF067654	<i>Mortierella alpina</i> $\Delta 5$ fatty acid desaturase
30	AB022097	<i>Dictyostelium discoideum</i> mRNA for $\Delta 5$ fatty acid desaturase
	AF489589.1	<i>Thraustochytrium</i> sp. ATcc21685 $\Delta 4$ fatty acid desaturase
	AY332747	<i>Pavlova lutheri</i> $\Delta 4$ fatty acid desaturase (desl) mRNA
35	AAG36933	<i>Emericella nidulans</i> oleate $\Delta 12$ desaturase

	AF1 10509,	<i>Mortierella alpina</i> $\Delta$ 12 fatty acid desaturase mRNA
	AAL13300	<i>Mortierella alpina</i> $\Delta$ 12 fatty acid desaturase mRNA
	AF417244	<i>Mortierella alpina</i> ATCC 16266 $\Delta$ 12 fatty acid desaturase
	AF161219	<i>Mucor rouxii</i> $\Delta$ 12 desaturase mRNA
5	X86736 S	<i>Pirulinea platensis</i> $\Delta$ 12 desaturase
	AF240777	<i>Caenorhabditis elegans</i> $\Delta$ 12 desaturase
	AB007640	<i>Chlamydomonas reinhardtii</i> $\Delta$ 12 desaturase
	AB075526	<i>Chlorella vulgaris</i> $\Delta$ 12 desaturase
	AP002063	<i>Arabidopsis thaliana</i> microsomal $\Delta$ 12 desaturase
10	NP_441622,	<i>Synechocystis</i> sp. PCC6803 $\Delta$ 15 desaturase
	AAL36934	<i>Perilla frutescens</i> $\Delta$ 15 desaturase

All references to sequence IDs herein are specifically incorporated by reference.

15 Additionally, the patent literature provides many additional DNA sequences of genes (and/or details concerning several of the genes above and their methods of isolation) involved in polyunsaturated fatty acid production (see, for example: U.S. Pat. No. 5,968,809 ( $\Delta\delta$ -desaturases); U.S. Pat. No. 5,972,664 and U.S. Pat. No. 6,075,183 ( $\Delta$ 5 desaturases); WO 91/13972 and U.S. Pat. No. 5,057,419 (A9-desaturases); WO  
20 93/1 1245 (A15-desaturases); WO 94/1 1516. U.S. Pat. No. 5,443,974 and WO 03/099216 (A12-desaturases); U.S. 2003/0196217 A1 (A17-desaturase); WO 02/090493 (A4-desaturases); and WO 00/12720 and U.S. 2002/01 39974A1 (elongases)).

25 The term "desaturases" as used herein refers to a polypeptide component of a multi-enzyme complex that can desaturate, i.e. introduce a double bond in one or more fatty acids to produce a mono- or polyunsaturated fatty acid or precursor of interest. Some desaturases have activity on two or more substrates. It may be desirable to empirically determine the specificity of a fatty acid desaturase by transforming a suitable host with  
30 the gene for the fatty acid desaturase and determining its effect on the fatty acid profile of the host. Nucleic acids that encode for desaturases are isolated from various organisms can be used according to the various aspects of the invention and examples are described herein, including *Ostreococcus* sp.

Desaturases include omega-3-desaturase,  $\Delta 6$ -desaturase,  $\Delta \delta$ -desaturase,  $\Delta 12$ -desaturase, A19-desaturase, A17-desaturase and A4-desaturase.

5 The term "elongase" as used herein refers to a polypeptide that can elongate a fatty acid carbon chain to produce an acid two carbons longer than the fatty acid substrate that the elongase acts upon. Nucleic acids that encode for elongases isolated from various organisms can be used according to the various aspects of the invention and examples are described herein, including *Ostreococcus sp.*

10 Examples of reactions catalyzed by elongase systems are the conversion of GLA to DGLA, SDA to ETA, ARA to DTA and EPA to DPA. In general, the substrate selectivity of elongases is somewhat broad but segregated by both chain length and the degree and type of unsaturation.

15 For example, a C14/16 elongase will utilize a C14 substrate (e.g., myristic acid), a C16/18 elongase will utilize a C16 substrate (e.g., palmitate), a C18/20 elongase will utilize a C18 substrate (e.g., GLA, SDA, LA, ALA) and a C20/22 elongase (also referred to as a  $\Delta \delta$ -elongase) will utilize a C20 substrate (e.g., ARA, EPA).

20 Since some elongases have broad specificity, a single enzyme may be capable of catalyzing several elongase reactions (e.g., thereby acting as both a C16/18 elongase and C18/20 elongase). It may be desirable to empirically determine the specificity of a fatty acid elongase by transforming a suitable host with the gene for the fatty acid elongase and determining its effect on the fatty acid profile of the host.

25

Elongases include  $\Delta 6$ -,  $\Delta 5$ - and A9-elongases.  $\Delta \delta$ -elongase is not generally viewed as rate limiting in the production of DHA and it is generally assumed that the first step in the LC-PUFA pathway, the D6-saturation, is rate-limiting.

30 Embodiments of the invention relating to the production of omega-3 LC-PUFAs in transgenic microalgae are described below. A skilled person would understand that these embodiments are not limited to transgenic microalgae, but can be applied to other organisms to produce omega-3 LC-PUFAs. The organism may be an animal, for example a mammal. In one embodiment, humans are specifically excluded. In another  
35 embodiment, the organism is a plant, for example a crop plant.

In a first aspect, the invention relates to a transgenic microalgae with increased production of omega-3 LC-PUFAs, for example one or more omega-3 LC-PUFA or total omega-3 LC-PUFA content. According to the various aspects of the invention, the omega-3 LC-PUFAs may be selected from SDA, ETA, EPA, DPA or DHA. In one embodiment, the omega-3 LC-PUFAs is DHA. In another embodiment, the omega-3 fatty acid is EPA.

According to the various aspects of the invention described herein, the increase in the production of DHA or EPA is measured as an individual content of different omega-3 LC-PUFAs in total fatty acids (TFA). In other words, the increase is measured as a percentage of the total fatty acid content. Preferably, the increase is at least 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15% or more compared to a control microalgae (mol %).

15

In one embodiment, the omega-3 LC-PUFAs is DHA. In the transgenic microalgae of the invention, the DHA content is increased by at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15% or more compared to a control microalgae. In one embodiment, the omega-3 LC-PUFAs is DHA. In the transgenic microalgae of the invention, the DHA content is at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9 or at least 10 fold higher than in a control microalgae. Preferably, the total DHA content is at least 10% of the total fatty acid content (mol %).

20

In another embodiment, the omega-3 LC-PUFAs is EPA. In the transgenic microalgae according to the various aspects of the invention, the EPA content is increased by at least 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%. Preferably, the total EPA content is at least 20% of the total fatty acid content (mol %).

25

According to the various aspects of the invention, the total fatty acid content, LC-PUFAs content, omega-3 LC-PUFAs content or the content of individual fatty acids such as DHA is increased compared to a control microalgae. A control microalgae as used herein is a microalgae which has not been modified according to the methods of the invention. Accordingly, the control microalgae has not been genetically modified to express a nucleic acid as described herein to alter LC-PUFA content. In one embodiment, the control microalgae is a wild type microalgae. In another embodiment,

30  
35

the control microalgae is a microalgae that does not carry a transgene according to the methods described herein, but expresses a different transgene. The control microalgae is typically of the same algae species.

5 The term "total fatty acids content" herein refers to the sum of all cellular fatty acids that can be derivitized to fatty acid methyl esters by the base transesterification method in a given sample (known as the art, for example as described in Sayanova et al., (1997); Sayanova et al., (2003) FEBS Lett. 2003 May 8;542(1-3):100-4).

10 According to the various aspects of the invention, the increase is measured in the stationary phase.

According to the various aspects of the invention, the term microalgae encompasses all microalgae which have the capacity to make LC-PUFAs. The algae may be a  
15 heterotrophic or autotrophic algae.

A skilled person would know that the term "microalgae" includes unicellular, photosynthetic microorganisms from several distinct biological groups, comprising, for example, eukaryotic chlorophyta, rhodophyta, heterokont, haptophyta divisions of algae  
20 and prokaryotic cyanobacteria.

EPA has been found in a wide variety of marine microalgae including in the classes *Bacillariophyceae* (diatoms), *Chlorophyceae*, *Chrysophyceae*, *Cryptophyceae*, *Eustigmatophyceae* and *Prasinophyceae* (see Table II). Accordingly, according to the  
25 various aspects of the invention, the microalgae may be selected from these orders, classes or species.

According to the various aspects of the invention, the microalgae may be selected from a microalgae listed in Table II.

30

Table II: Proportions of PUFAs in marine microalgae

\**Emiliania huxleyi* the now accepted name for *Coccolithus huxleyi*

35

Omega-3 LC-PUFAs (% of Total Fatty acids)				
Mircoalgae (Order/class/sp.)	sp.	EPA	DHA	References
Chlorophyta (green algae)				
<u>Chlorophyceae</u>				
	<i>Chlorella minutissima</i>	45.0	-	Seto et al., (1984)
<u>Prasinophyceae</u>				
	<i>Ostreococcus tauri</i>	2.0	12.0	Wagner M. et al., (2010)
	<i>Ostreococcus lucimarinus</i>	2.1	3.8	Ahmann et al., (2011)
	<i>Hetermastrix rotundra</i>	28	7	Yongmanitchai and Ward, (1989)
Haptophyta				
<u>Pavlovophyceae</u>				
	<i>Pavlova lutheri</i>	11.6	9.1	Tonon et al., (2002)
<u>Prymnesiophyceae</u>				
	<i>Isochrysis galbana</i>	22.6	8.4	Molina Grima et al., (1995)
	<i>Emilinaia huxleyi</i> *	17	-	Yongmanitchai and Ward, (1989)
Cryptophyceae				
<u>Cryptomonadaceae</u>				
	<i>Cryptomonas maculate</i>	17	-	Yongmanitchai and Ward, (1989)
	<i>Chromonas sp.</i>	12	6.6	Renaud et al., (1999)
	<i>Cryptomonas sp.</i>	16	10	Yongmanitchai and Ward, (1989)
	<i>Rhodomonas sp.</i>	8.7	4.6	Renaud et al., (1999)
Heterokont				
<u>Bacillariophyceae (diatoms)</u>				

<i>Asterionella japonica</i>	20	-	Yongmanitchai and Ward, (1989)
<i>Amphora coffeaformis</i>	1.39	0.39	Renaud et al., (1999)
<i>Biddulphia sinensis</i>	24.0	1.0	Yongmanitchai and Ward, (1989)
<i>Chaetoceros sp.</i>	16.7	0.8	Renaud et al., (1999)
<i>Cylindrotheca fusiformis</i>	18.8	-	Tan and Johns, (1996)
<i>Fragilaria pinnata</i>	6.8	1.0	Renaud et al., (1999)
<i>Nitzschia angularis</i>	21	-	Kyle et al., (1992)
<i>Navicula incerta</i>	25.2	-	Tan and Johns, (1996)
<i>Navicula pelliculosa</i>	9.4	-	Tan and Johns, (1996)
<i>Navicula saprophila</i>	16.0	-	Kitano et al., (1997)
<i>Nitzschia closterium</i>	15.2	-	Renaud et al., (1994)
<i>Nitzschia frustulum</i>	23.1	-	Renaud et al., (1994)
<i>Nitzschia laevis</i>	19.1	-	Wen and Chen, (2001)
<i>Phaeodactylum tricornutum</i>	34.5	-	Yongmanitchai and Ward, (1991)
<i>Skeletonema costatum</i>	29.2	3.4	Blanchemain and Grizeau, (1999)
<i>Thalassiosira pseudonana</i>	12.2	-	Tonon et al., (2002)
<u>Chrysophyceae (golden algae)</u>			
<i>Monochrysis lutheri</i>	19	-	Yongmanitchai and Ward, (1989); Kyle, (1992)
<i>Pseudopedinella sp.</i>	27	-	Yongmanitchai and Ward, (1989)
<i>Crisosphaera carterae</i>	20	-	Yongmanitchai and Ward, (1989)
<i>C.elongate</i>	28	-	Yongmanitchai and Ward, (1989)
<u>Eustigmatophyceae</u>			
<i>Nannochloropsis salina</i>	15	-	Yongmanitchai and Ward, (1989)
<i>Nannochloropsis sp.</i>	35	-	Sukenik, (1991)
<i>Nannochloris sp.</i>	27	-	Yongmanitchai and Ward, (1989)
<i>Monodus subterraneus</i>	32.9	-	Quiang et al., (1997)

In one embodiment, autotrophic microalgae which are as the primary producers of PUFAs are preferred. For example, the microalgae may be selected from *Phaeodactylum*, *Nannochloropsis*, *Thraustochytrium* or *Schizochytrium*. Other genera



include *Spirulina*, *Dunaliella*, *Chlorella*, *Thalassiosira*, *Isochrysis*, *Porphyridium*, *Nannochloropsis*, *Pavlova*, *Chaetoceros*, *Cryptothecodinium*, *Fragilariopsis* and *Nitzshia*.

5 For example, the microalgae may be selected from *Chaetoceros calcitrans*, *Isochrysis galbana*, *Pavlova lutheri*, *Pseudoisochrysis paradoxa*, *Tetraselmis suecica* and *Skeletonema costatum*, *Nannochloropsis oculata*, *Thalassiosira pseudonana*, *Pavlova lutheria*, *Porphyridium irregular*, *Cryptothecodinium cohnii*, *Porphyridium purpureum* and *Porphyridium omentum*.

10 In one embodiment, the microalgae is a diatom. Diatoms are brown algae found throughout marine and freshwater ecosystems that are responsible for around 20% of global primary productivity. A defining feature of diatoms is their ornately patterned silicified cell wall (known as frustule), which display species-specific nanoscale-structures.

15 The diatom may be a centric diatoms or a pennate diatom. In one embodiment, the diatom belongs to the order of Naviculales. In one embodiment, the diatom is *P. tricorutum* or *Thalassiosira pseudonana*. In a preferred embodiment, the diatom is *P. tricorutum*. In another embodiment, the diatom is *Fragilariopsis sp.* for example *Fragilariopsis cylindrus*.

20 A skilled person would understand that the aspects of the invention are not limited to *P. tricorutum*. Indeed, a skilled person would understand that the invention can be applied to any microalgae that has the capacity to synthesise EPA and/or DHA.

The transgenic microalgae according to the various aspects of the invention expresses one or more heterologous transgenes which encode for one or more nucleic acid  
25 involved in the biosynthesis of LC-PUFAs. "Heterologous" with respect to sequence means a sequence that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. The heterologous transgene is preferably derived or isolated from a microalgae. In one embodiment, the heterologous transgene is derived  
30 or isolated from *Prasinophyceae*, for example *Ostreococcus sp.* Sequences of heterologous transgenes may be modified to be codon optimised for expression in the target organism. Thus, the invention relates to transgenic organisms obtained through recombinant methods.

For example, the heterologous transgene may encode for one or more of a  $\Delta 15$ -desaturase, a  $\Delta 6$ -desaturase, a  $\Delta \delta$ -desaturase, a  $\Delta 4$ -desaturase, a  $\Delta 12$ -desaturase, a  $\Delta \delta$ -elongase,  $\Delta 6$ -elongase or combinations thereof.

5

In one embodiment, the transgenic microalgae expresses a heterologous nucleic acid encoding a  $\Delta \delta$ -elongase. Thus, in one aspect, the invention relates to a transgenic microalgae expressing a nucleic acid encoding a  $\Delta \delta$ -elongase. For example, the transgenic microalgae expresses a nucleic acid encoding a  $\Delta \delta$ -elongase, but does not  
10 express any other transgene encoding for a polypeptide involved in the regulation of the LC-PUFAs biosynthetic pathway. In other embodiments, the transgenic microalgae expresses a nucleic acid encoding a  $\Delta \delta$ -elongase and one or more additional heterologous transgene involved in the regulation of the LC-PUFAs biosynthetic pathway, for example a  $\Delta 6$ -desaturase such as OtD6 as shown in example 4. Thus,  
15 embodiments where nucleic acids encoding a  $\Delta \delta$ -elongase and a  $\Delta 6$ -desaturase are co-expressed are specifically part of the invention.  $\Delta \delta$ -elongases and  $\Delta 6$ -desaturases are as defined herein.

In one embodiment, the transgenic microalgae described herein co-expresses a  
20 heterologous nucleic acid which is not involved in the regulation of the LC-PUFAs biosynthetic pathway, for example a glucose transporter gene as shown in example  $\delta$  together with a heterologous nucleic acid involved in the regulation of the LC-PUFAs biosynthetic pathway such as OiEI $\delta$ . As shown in the example, a vector can be used allowing co-expression of two heterologous nucleic acids involved in the regulation of  
25 different traits - one for omega-3s, and one which allows the alga to be grown in the dark, by the expression of a glucose transporter. If the cells are then provided with an exogenous carbon source such as glucose, they can grow in the dark. Thus, in one embodiment, an exogenous carbon source such as glucose is provided when culturing algae expressing a gene involved in the regulation of the LC-PUFAs biosynthetic  
30 pathway such as OiEI $\delta$  and a glucose reporter. Examples of nucleic acids that can be used according to the invention encoding a glucose reporter are shown in SEQ ID No. 23 and SEQ ID No. 25. Respective peptides are shown in SEQ ID No. 24 and SEQ ID No. 26.

As used herein, the words "nucleic acid", "nucleic acid sequence", "nucleotide", or "polynucleotide" are intended to include DNA molecules (e.g. cDNA or genomic DNA), RNA molecules (e.g., mRNA), natural occurring, mutated, synthetic DNA or RNA molecules, and analogs of the DNA or RNA generated using nucleotide analogs. It can be single-stranded or double-stranded. Such nucleic acids or polynucleotides include, but are not limited to, coding sequences of structural genes, anti-sense sequences, and non-coding regulatory sequences that do not encode mRNAs or protein products. These terms also encompass a gene. The term "gene" or "gene sequence" is used broadly to refer to a DNA nucleic acid associated with a biological function. Thus, genes may include introns and exons as in genomic sequence, or may comprise only a coding sequence as in cDNAs, and/or may include cDNAs in combination with regulatory sequences. In one embodiment of the various aspects of the invention, cDNA sequences synthetic (deduced) open reading frames, analogous to cDNA are preferred.

For the purposes of the invention, "transgenic", "transgene" or "recombinant" means with regard to, for example, a nucleic acid sequence, an expression cassette, gene construct, a vector or an autonomous replicating element such as an artificial chromosome comprising the nucleic acid sequence or an organism transformed with the nucleic acid sequences, expression cassettes or vectors according to the invention, all those constructions brought about by recombinant methods in which either

(a) the nucleic acid sequences encoding proteins useful in the methods of the invention, or

(b) genetic control sequence(s) which is operably linked with the nucleic acid sequence according to the invention, for example a promoter, or

(c) a) and b)

are not located in their natural genetic environment or have been modified by recombinant methods, such as mutagenesis, it being possible for the modification to take the form of, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. The natural genetic environment is understood as meaning the natural genomic or chromosomal locus in the original microalgae or the presence in a genomic library.

A transgenic microalgae for the purposes of the invention is thus understood as meaning a microalgae which comprises within its nuclear and or plastidial genome a

heterologous polynucleotide. The heterologous polynucleotide is preferably stably integrated within the genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant DNA construct.

5

In the context of the present invention, a  $\Delta\delta$ -elongase catalyzes the conversion of EPA to DPA. Thus, any nucleic acid that encodes a  $\Delta\delta$ -elongase that catalyzes the conversion of EPA to DPA may be used according to the various aspects of the invention as a transgene. In one embodiment, the  $\Delta\delta$ -elongase used in the present invention is derived or isolated from *Ostreococcus*, preferably *Ostreococcus tauri*. Preferably, the  $\Delta\delta$ -elongase is  $\Delta\delta$  derived or isolated from *Ostreococcus tauri*. In one embodiment, the transgenic microalgae according to the invention expresses a nucleic acid comprising SEQ ID No. 1, a functional variant thereof or a sequence that encodes for a  $\Delta\delta$ -elongase wherein said elongase has at least 60%, at least  $\delta\delta\%$ , at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 2. In a preferred embodiment, the microalgae is *P. tricornutum* and the nucleic acid encodes a  $\Delta\delta$ -elongase comprising or consisting of SEQ ID No. 2.

10  
15  
20

A functional variant as used according to the aspects of the invention is a biologically active variant. For example, a biologically active variant of SEQ ID No. 1 is a nucleic acid sequence, which, when expressed in a microalgae such as *P. tricornutum*, increases production of DHA. The term variant includes sequences which have been altered for codon optimisation for expression in the target organism for example for expression in *P. tricornutum*.

25

Thus, it is understood, as those skilled in the art will appreciate, that the aspects of the invention, which use certain polynucleotides including the methods and uses, encompasses more than the sequence specified, but also include alterations in the peptide that do not affect the biological function. For example, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not affect the functional properties of the encoded polypeptide, are well known in the art. For example, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue,

30  
35

such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

10

In one embodiment, the said nucleic acid according to the various aspects of the invention is operably linked to a regulatory sequence.

15

20

25

30

The terms "regulatory element" is used interchangeably herein with "control sequence" and "promoter" and all terms are to be taken in a broad context to refer to regulatory nucleic acid sequences capable of effecting expression of the sequences to which they are ligated. The term "promoter" typically refers to a nucleic acid control sequence located upstream from the transcriptional start of a gene and which is involved in recognising and binding of RNA polymerase and other proteins, thereby directing transcription of an operably linked nucleic acid. Encompassed by the aforementioned terms are transcriptional regulatory sequences derived from a classical eukaryotic genomic gene (including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence) and additional regulatory elements (i.e. upstream activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. Also included within the term is a transcriptional regulatory sequence of a classical prokaryotic gene, in which case it may include a -35 box sequence and/or -10 box transcriptional regulatory sequences. The term "regulatory element" also encompasses a synthetic fusion molecule or derivative that confers, activates or enhances expression of a nucleic acid molecule in a cell, tissue or organ.

Suitable promoters are identified in the examples. For example, if the microalgae is *P. tricornutum*, the promoter may be the *P. tricornutum* promoter fcpA. However, a skilled person would understand that other promoters can also be used. For example, suitable

promoters may also be selected from inducible promoters which respond to specific environmental or chemical stimuli.

5 The term "operably linked" as used herein refers to a functional linkage between the promoter sequence and the gene of interest, such that the promoter sequence is able to initiate transcription of the gene of interest.

10 The transgene may be part of a vector which, in addition to one or more regulatory sequences also comprises selection markers. These are known in the art. Transformation of microalgae may be carried out by standard procedures known in the art, for example by particle bombardment or electroporation.

15 The transgenic microalgae expressing a nucleic acid encoding a  $\Delta\delta$ -elongase is characterised by an increase in DHA and DPA compared to a control microalgae. In particular, the increase, as measured as a percentage of the total fatty acid content is at least 2, at least 3, at least 4, at least 5, at least 6, at least, at least 8, at least 9 or at least 10 fold higher than in a control microalgae. Specifically, the DHA content is at least 2, at least 3, at least 4, at least 5, at least 6, at least, at least 8, at least 9 or at least 10 fold higher than in a control microalgae. Preferably, the total DHA content is at least 10% of the total LC-PUFAs content (%mol). In one embodiment, the transgenic microalgae expressing a nucleic acid encoding a  $\Delta\delta$ -elongase does not express a second transgene encoding for another polypeptide involved in the regulation of the LC-PUFAs pathway, preferably in the regulation of the omega-3 LC-PUFAs pathway.

25 In one embodiment of the various aspects of the invention, the transgenic microalgae expressing a heterologous nucleic acid encoding a  $\Delta\delta$ -elongase may further express one or more additional heterologous nucleic acid encoding for one or more polypeptide involved in the regulation of the LC-PUFAs pathway, preferably in the regulation of the omega-3 LC-PUFAs pathway. In other words, the transgenic microalgae comprises one or more further transgene encoding for one or more polypeptide involved in the regulation of the LC-PUFAs pathway. The polypeptide is preferably selected from any desaturase or elongase involved in the omega-3 PUFA biosynthetic pathway as shown in figure 7. Any combination of desaturase and elongase may also be used. Thus, the nucleic acid may encode for one or more of a  $\Delta^5$ -desaturase, a  $\Delta^6$ -desaturase, a  $\Delta^5$ -

desaturase, a A4-desaturase, a  $\Delta 6$ -desaturase, a  $\Delta \delta$ -elongase,  $\Delta 6$ -elongase or combinations thereof.

5 In one embodiment, the nucleic acid encodes a  $\Delta 6$ -desaturase. In the context of the present invention, a  $\Delta 6$ -desaturase catalyzes the conversion of ALA to SDA and also LA to GLA. A6-Desaturases are described in WO 93/06712, US 5,614, 393, US 5614393, WO 96/21022, WO 02/1557 and WO 99/271 11 and their application to production in transgenic organisms is also described, e.g. in WO 98/46763, WO 98/46764 and WO 98/46765. In one embodiment, the  $\Delta \delta$ -desaturase used in the present invention is derived or isolated from *Ostreococcus*, preferably OtD6 from *Ostreococcus tauri* (Domergue et al (2005), AY746357). In one embodiment, the nucleic acid comprises SEQ ID No. 3 or 5 and encodes a 6A-desaturase comprising or consisting of SEQ ID No. 4 or 6, a functional variant thereof or a polypeptide that encodes for a 6A-desaturase that has at least 50%, at least 55%, at least 60%, at least 10 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 4 or 6.

20 In another embodiment, the A6-desaturase is from the microalgae *Ostreococcus* RCC 809. Preferably, the nucleic acid comprises SEQ ID No. 7 or 9 and encodes a  $6\Delta$ -desaturase from the microalgae *Ostreococcus* RCC 809 comprising or consisting of SEQ ID No. 8 or 10, a functional variant thereof or a sequence that encodes for a  $6\Delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, 25 at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 8 or 10.

30 In another embodiment, the nucleic acid encodes for a A4-desaturase. According to the various aspects of the invention, a A4-desaturase may be derived or isolated from *E. huxleyi*. Thus, in one embodiment, the nucleic acid comprises SEQ ID No. 11 encoding a A4-desaturase comprising or consisting of SEQ ID No. 12, a functional variant thereof or a A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, 35 at least 98% or at least 99% homology to SEQ ID No. 12.

In another embodiment, the A4-desaturase is derived or isolated from *T. pseudonana*. Thus, in one embodiment, the nucleic acid comprises SEQ ID No. 13 encoding a  $\Delta$ 4-desaturase comprising or consisting of SEQ ID No. 14, a functional variant thereof or a  
5 A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 14.

10 In another embodiment, the A4-desaturase is derived or isolated from *Ostreococcus* RCC809. In one embodiment, the nucleic acid comprises SEQ ID No. 15 or 17 encoding a A4-desaturase comprising or consisting of SEQ ID No. 16 or 18, a functional variant thereof or a A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least  
15 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 16 or 18.

In another embodiment, a A6-elongase is from *Fragilariopsis cylindrus*. In one embodiment, the nucleic acid comprises SEQ ID No 19 encoding a A6-elongase  
20 comprising or consisting of SEQ ID No. 20, a functional variant thereof or a  $\Delta$ 6-elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 20.

25 In another embodiment, a  $\Delta\delta$ -desaturase is from *Fragilariopsis cylindrus*. In one embodiment, the nucleic acid comprises SEQ ID No 21 encoding a  $\Delta\delta$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta$ 6-elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%,  
30 at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 22.



In another aspect, the transgenic microalgae of the invention expresses a heterologous nucleic acid encoding a  $\Delta 6$ -desaturase, a  $\Delta \delta$ -desaturase, a A4-desaturase,  $\Delta 6$ -elongase or combinations thereof. These enzymes are defined herein.

5 In one aspect, a transgenic microalgae of the invention expresses a heterologous nucleic acid encoding a  $\Delta 6$ -desaturase. Thus, in another aspect, the invention also relates to transgenic microalgae expressing a heterologous nucleic acid encoding a  $\Delta 6$ -desaturase. For example, the transgenic microalgae expresses a nucleic acid encoding a  $\Delta 6$ -desaturase, but does not express any other transgene involved in the  
10 regulation of the LC-PUFAs biosynthetic pathway. In other embodiments, the transgenic microalgae expresses a  $\Delta 6$ -desaturase and additional transgenes involved in the regulation of the LC-PUFAs biosynthetic pathway, for example a  $\Delta \delta$ -elongase such as OtElo5 as shown in the examples.

15 In one embodiment, the microalgae is *P. triconutum*. In one embodiment, the nucleic acid comprising or consisting of SEQ ID No. 3 or 5 encodes a  $\Delta 6$ -desaturase or a sequence that encodes for a  $\Delta 6$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%,  
20 at least 97%, at least 98%, at least 99% homology to SEQ ID No. 4 or 6. In a preferred embodiment, the microalgae is *P. triconutum* and the nucleic acid encodes a  $\Delta 6$ -desaturase comprising or consisting of SEQ ID No. 4 or 6.

The transgenic microalgae expressing a nucleic acid encoding a A6-desaturase is  
25 characterised in that the total fatty acids content, specifically the omega 3 LC-PUFA content, is altered compared to a control microalgae. In particular, the omega-3 LC-PUFA content is increased by at least 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15% or more. Specifically, the EPA content is increased by at least 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15% compared to a  
30 control microalgae. Preferably, the total EPA content is at least 20% of the total LC-PUFAs content (mol %). Moreover, the DHA content in the transgenic algae is also increased by at least 0.5%.

In one embodiment, the various aspects of the invention exclude embodiments that  
35 relate to the production of biofuels.

In another aspect, the invention relates to a method for producing transgenic microalgae with increased omega-3 LC-PUFA content comprising introducing and expressing in a microalgae a heterologous nucleic acid which encodes for a polypeptide involved in the LC-PUFAs biosynthetic pathway. The omega-3 fatty acid may be selected from ALA, SDA, ETA, EPA, DPA or DHA. In one embodiment, the omega-3 LC-PUFAs is DHA. In another embodiment, the omega-3 fatty acid is EPA. The nucleic acid may encode  $\Delta 6$ -desaturase,  $\Delta \delta$ -desaturase, A4-desaturase,  $\Delta 5$ -elongase,  $\Delta 6$ -elongase or combinations thereof.

10

In one embodiment, the method relates to producing transgenic microalgae with increased DHA levels said method comprising transforming a microalgae with a heterologous nucleic acid encoding a  $\Delta \delta$ -elongase. According to this embodiment, the method may further comprise transforming said microalgae with one or more additional heterologous nucleic acid that regulates the production of omega-3 fatty acids, for example transforming with a nucleic acid encoding a  $\Delta 6$ -desaturase. In another embodiment, no additional nucleic acid that regulates the production of omega-3 fatty acids is introduced into said microalgae and expressed as heterologous nucleic acids.

15

In another embodiment, the invention relates to a method for producing transgenic microalgae with increased EPA levels said method comprising transforming a microalgae with a nucleic acid encoding a  $\Delta 6$ -desaturase. According to this embodiment, the method may further comprise transforming said microalgae with one or more additional nucleic acid that regulates the production of omega-3 LC-PUFAs. In another embodiment, no additional nucleic acid that regulates the production of omega-3 fatty acids is introduced into said microalgae.

20

25

In one embodiment, the method comprises transforming said microalgae with one or more additional nucleic acid that does not regulates the production of omega-3 LC-PUFAs, for example a glucose transporter gene.

30

Microalgae obtained or obtainable by those methods are also within the scope of the invention.

In another aspect, the invention relates to a method for increasing production of one of more omega-3 LC-PUFA in microalgae comprising

- a) cultivating a transgenic microalgae described herein and
- b) obtaining said one of more omega-3 LC-PUFA from the transgenic  
5 microalgae.

Specifically, the invention relates to a method for increasing the production of one or more omega-3 LC-PUFAs in microalgae comprising:

- a) introducing and expressing in a microalgae a heterologous nucleic acid which  
10 encodes for a polypeptide involved in the LC-PUFAs biosynthetic pathway,
- b) cultivating a transgenic microalgae expressing said heterologous nucleic acid and
- c) obtaining one or more omega -3 fatty acid from the transgenic microalgae.

The transgenic microalgae is as described herein and is cultivated under conditions  
15 which allow for the production of one or more omega-3 LC-PUFAs. The nucleic acid may encode a A15-desaturase, a A6-desaturase, a  $\Delta\delta$ -desaturase, a A4-desaturase, a A12-desaturase,  $\Delta\delta$ -elongase, A6-elongase or combinations thereof as described herein.

20 In one embodiment, the method relates to increasing DHA production in microalgae comprising

- a) introducing and expressing in a microalgae a heterologous nucleic acid encoding a  $\Delta\delta$ -elongase,
- b) cultivating a transgenic microalgae expressing said heterologous nucleic acid and
- c) obtaining DHA from the transgenic microalgae.  
25

The microalgae as described herein. The  $\Delta\delta$ -elongase is as described herein. In one embodiment, the microalgae does not include and express a second heterologous nucleic acid encoding an enzyme involved in the regulation of the synthesis of omega-3  
30 LC-PUFAs. In another embodiment, the microalgae includes and expresses a second heterologous nucleic acid encoding a polypeptide involved in the regulation of the synthesis of omega-3 LC-PUFAs. In another embodiment, the microalgae includes and expresses a second heterologous nucleic acid encoding a polypeptide not involved in the regulation of the synthesis of omega-3 LC-PUFAs, for example a glucose

transporter. The transgenic microalgae is cultivated under conditions which allow for the production of DHA.

5 In one embodiment, the method relates to increasing DHA production in microalgae comprising

- a) introducing and expressing in *P. triconutum* a heterologous nucleic acid encoding a  $\Delta\delta$ -elongase,
- b) cultivating *P. triconutum* expressing said heterologous nucleic acid and
- c) obtaining said DHA from *P. triconutum*.

10

The microalgae as described herein. The  $\Delta\delta$ -elongase is as described herein. In one embodiment, the microalgae does not include and express a second heterologous nucleic acid encoding an enzyme involved in the regulation of the synthesis of omega-3 LC-PUFAs. In another embodiment, the microalgae includes and expresses a second  
15 heterologous nucleic acid encoding an enzyme involved in the regulation of the synthesis of omega-3 LC-PUFAs. In another embodiment, the microalgae includes and expresses a second heterologous nucleic acid encoding a polypeptide not involved in the regulation of the synthesis of omega-3 LC-PUFAs, for example a glucose transporter.

20

*P. triconutum* is cultivated under conditions which allow for the production of DHA. These conditions will be apparent to the skilled person. For example, preferred culture conditions for *P. triconutum* are about 20°C under constant illumination in about 60-80  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . In one embodiment, the method comprises transforming said  
25 microalgae with one or more additional nucleic acid that does not regulates the production of omega-3 LC-PUFAs, for example a glucose transporter gene and supplying an exogenous carbon source. The algae can be grown in the dark.

30

In another embodiment, the method relates to increasing EPA in microalgae comprising:

- a) introducing and expressing in a microalgae a heterologous nucleic acid encoding a  $6\Delta$ -desaturase,
- b) cultivating the transgenic microalgae and
- c) obtaining said EPA from the transgenic microalgae.

35

The microalgae as described herein The  $6\Delta$ -desaturase is as described herein. The microalgae is cultivated under conditions which allow for the production of EPA.

5 In one embodiment, the method relates to increasing EPA production in microalgae comprising

- a) introducing and expressing in *P. triconutum* a heterologous nucleic acid encoding a  $6\Delta$ -desaturase,
- b) cultivating *P. triconutum* and
- c) obtaining said EPA from *P. triconutum*.

10

The microalgae as described herein The  $\Delta 6$ -desaturase is as described herein. *P. triconutum* is cultivated under conditions which allow for the production of EPA.

15

These conditions will be apparent to the skilled person. For example, preferred culture conditions for *P. triconutum* are about 20°C under constant illumination in about 0- $80\mu\text{mol photons m}^{-2}\text{s}^{-1}$  or preferably about 18°C under constant illumination in about 25 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . In one embodiment, the method comprises transforming said microalgae with one or more additional nucleic acid that does not regulate the production of omega-3 LC-PUFAs, for example a glucose transporter gene and

20 supplying an exogenous carbon source. The algae can be grown in the dark.

20

In another aspect, the invention relates to a method for the manufacture of an oil, lipid or fatty acid composition comprising

25

- a) cultivating a transgenic microalgae as described herein under conditions which allow for the production one or more omega-3 LC-PUFAs and
- b) obtaining said one or more omega-3 LC-PUFAs from the transgenic microalgae.

In preferred embodiment, the omega-3 LC-PUFAs is DHA or EPA.

30

In another aspect, the invention relates to an omega-3 LC-PUFAs or oil isolated from a transgenic microalgae as described herein.

35

The fatty acids produced by the processes of the present invention can be isolated from the microalgae in the form of an oil, a lipid or a free fatty acid. One embodiment of the invention is therefore oils, lipids or fatty acids or fractions thereof which have been

produced by the methods of the invention, especially preferably oil, lipid or a fatty acid composition comprising EPA or DHA and being derived from the transgenic microalgae.

5 The term "oil", or "lipid" is understood as meaning a fatty acid mixture comprising unsaturated, preferably esterified, fatty acid(s). The oil or lipid is preferably high in omega-3 polyunsaturated or, advantageously, esterified fatty acid(s). In a particularly preferred embodiment the oil or lipid has a high ALA, ETA, EPA, DPA and/or DHA content, preferably a high EPA and/or DHA content.

10

For the analysis, the fatty acid content can, for example, be determined by gas chromatography after converting the fatty acids into the methyl esters by transesterification of the lipids such as triacylglycerides and/or phospholipids.

15

The omega-3 polyunsaturated acids produced in the method of the present invention, for example EPA and DHA, may be in the form of fatty acid derivatives, for example sphingolipids, phosphoglycerides, lipids, glycolipids, phospholipids, monoacylglycerol, diacylglycerol, triacylglycerol or other fatty acid esters.

20

The omega-3 and other polyunsaturated fatty acids which are present can be liberated for example via treatment with alkali, for example aqueous KOH or NaOH, or acid hydrolysis, advantageously in the presence of an alcohol such as methanol or ethanol, or via enzymatic cleavage, and isolated via, for example, phase separation and subsequent acidification via, for example  $H_2SO_4$ . The fatty acids can also be liberated

25

directly without the above-described processing step.

30

If further purification is necessary, standard methods can be employed. Such methods may include extraction, treatment with urea, fractional crystallization, HPLC, fractional distillation, silica gel chromatography, high-speed centrifugation or distillation, or combinations of these techniques. Protection of reactive groups, such as the acid or alkenyl groups, may be done at any step through known techniques (e.g. alkylation, iodination, use of butylated hydroxytoluene (BHT)). Methods used include methylation of the fatty acids to produce methyl esters. Similarly, protecting groups may be removed at any step. Desirably, purification of fractions containing, for example, ALA, STA, ETA,

EPA, DPA and DHA may be accomplished by treatment with urea and/or fractional distillation.

5 Large scale purification methods of fatty acids from algae are known in the art. For example, a microalgae strain is cultivated to increase cell density using photobioreactors, open ponds, race ways or hybrid systems. Algal cells are separated from culture media by filtration, flocculation or centrifugation, followed by drying to improve extraction. Lipid extraction is then commonly performed using a non-water miscible organic solvent. Larger scale extraction is typically carried out with hexane as  
10 a solvent. Subsequently, unsaturated fatty acids are separated from the total lipids by fractional (molecular) distillation or winterization, whereby oil temperature is reduced to precipitate the more saturated lipids. Further processing to improve the quality, shelf-life and quantity of PUFA oil can include filtration, bleaching, deodorization, polishing and antioxidant addition. These methods are all known to a person skilled in the art.

15

In another aspect, the invention also relates to the use of the transgenic organism, preferably microalgae, as described herein in the production of fatty acids, preferably a omega-3 fatty acids. The invention encompasses the use of a transgenic organism, preferably microalgae, as described herein or of the oil, lipid, the fatty acids obtained  
20 from a transgenic organism, preferably microalgae, as described herein in feedstuffs, foodstuffs, cosmetics, nutraceutical or pharmaceuticals. The invention encompasses the use of a transgenic organism, preferably microalgae as described herein, in producing feedstuffs, foodstuffs, cosmetics, nutraceutical or pharmaceuticals. In another aspect, the invention also relates to the use of the transgenic microalgae, as described herein  
25 as a feedstuff for animals, preferably fish.

In another aspect, the invention also relates to a composition comprising the transgenic microalgae as described herein or a fatty acid, preferably a omega-3 fatty acid, oil, or lipid obtained from said microalgae. In a preferred embodiment, the composition  
30 comprises the transgenic microalgae as described herein or a product obtained or obtainable therefrom., such as an oil. In one embodiment, the composition may be a pharmaceutical composition, a cosmetic, a foodstuff, including food supplements, or feedstuff for animals. In particular, the invention relates to a foodstuff comprising the transgenic microalgae as described herein or fatty acid, preferably a omega-3 fatty  
35 acid, oil, or lipid obtained from said algae. This can be in the form of a dietary

supplement, including fish oils. The invention also relates to an animal feed, especially for aquaculture, comprising the transgenic microalgae as described herein or fatty acid, preferably a omega-3 fatty acid, oil, or lipid obtained from said algae.

- 5 In another aspect, the invention relates to a composition comprising the transgenic microalgae as described herein, a fatty acid, preferably a omega-3 fatty acid, oil, or lipid obtained from said microalgae for use in medicine. In particular, the composition may be used to lower both blood pressure and heart rate in hypertensive individuals reducing the risk of sudden death, reduce inflammation, and to reduce the long-term
- 10 risk of atherosclerosis and ischemic heart disease. The composition may also be used to treat eczema or metabolic syndrome. Also, a DHA rich diet is associated with increased cognitive abilities and depression and has a positive effect on arthritis and type II diabetes (Horrocks et al, 1999). Thus, the invention also relates to a composition comprising the transgenic microalgae as described herein or fatty acid, preferably a
- 15 omega-3 fatty acid, oil, or lipid obtained from said microalgae for use in the treatment or prevention of cardiovascular conditions, including atherosclerosis, thrombosis, high blood pressure, myocardial infarction and atherosclerosis, inflammatory conditions, depression, cognitive decline, arthritis, and type II diabetes. Also encompassed in the scope of the invention are methods of treating or preventing cardiovascular and
- 20 inflammatory conditions, depression, cognitive decline, arthritis and type II diabetes administering a composition comprising a therapeutic amount of the transgenic microalgae as described herein, a fatty acid, preferably a omega-3 fatty acid, oil, or lipid obtained from said microalgae to a patient in need thereof. The invention also relates to the use of a composition comprising the transgenic microalgae as described
- 25 herein in the manufacture of a medicament for treating cardiovascular conditions, including atherosclerosis, thrombosis, high blood pressure, myocardial infarction and atherosclerosis, inflammatory conditions, depression, cognitive decline, arthritis, and type II diabetes.
- 30 In preferred embodiments, the composition may comprise or be obtained from a transgenic microalgae expressing a nucleic acid encoding a  $\Delta 6$ -desaturase and/or a transgenic microalgae expressing a nucleic acid encoding a  $\Delta \delta$ -elongase as described herein.



The inventors have shown that microalgae can be manipulated using recombinant methods to produce an increased amount of LC-PUFAs, in particular EPA and DHA using heterologous gene expression. The inventors have surprisingly demonstrated that heterologous expression of  $\Delta\delta$ -elongase from *Ostreococcus tauri* alone results in increased accumulation of DHA in *P. tricornutum* with DHA levels in transgenic strains reaching up to 13% of total fatty acids. A skilled person would understand that the invention is not restricted to algae and can indeed be applied to any organism that makes EPA/DHA. Thus, the invention also relates to a transgenic organism with increased DHA levels expressing a heterologous  $\Delta\delta$ -elongase, preferably a  $\Delta 5$ -elongase from *Ostreococcus tauri*. In one embodiment, no other transgenes are expressed in the transgenic organism. In another embodiment, further transgenes may be expressed as described herein. Furthermore, the invention also relates to methods for increasing the production of DHA in a transgenic organism. This is achieved by expressing a heterologous  $\Delta\delta$ -elongase, preferably a  $\Delta\delta$ -elongase from *Ostreococcus tauris* in said organism. Details of said methods are described herein.

The organism may be an animal, for example a mammal. In one embodiment, humans are specifically excluded. In another embodiment, the organism is a plant, for example a monocot or dicot plant, for example crop plant. Crop plants include but are not limited to maize, rice, wheat, oilseed rape/canola, sorghum, soybean, sunflower, alfalfa, potato, tomato, tobacco, grape, barley, pea, bean, field bean, lettuce, cotton, sugar cane, sugar beet, broccoli or other vegetable brassicas or poplar.

In another aspect, the invention relates to isolated nucleic acids encoding for novel forms of the desaturases and elongases which may be useful in the heterologous reconstitution of the omega-3 long chain polyunsaturated fatty acid biosynthetic pathway in algae and higher plants. Specifically, the invention relates to isolated nucleic acids encoding  $\Delta 6$ -desaturase (Ost809A6), A4-desaturase (Ost809A4) and  $\Delta 6$ -elongase (FcEL06) and their corresponding polypeptides.

In one embodiment, the invention relates to an isolated nucleic acids comprising SEQ ID No. 7 or 9 encoding  $\Delta 6$ -desaturase (Ost809A6) comprising or consisting of SEQ ID No. 8 or 10, a functional variant thereof or a  $\Delta 6$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%,

at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 10. The sequence may also be codon optimised for expression the target organism.

5 In one embodiment, the invention relates to an isolated nucleic acid comprising SEQ ID No. 15 or 17 encoding a A4-desaturase (Ost809A4) comprising or consisting of SEQ ID No. 16 or 18, a functional variant thereof or a A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 10 18. The sequence may also be codon optimised for expression the target organism.

In one embodiment, the invention relates to an isolated nucleic acid comprising SEQ ID No. 19 encoding A6-elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant thereof or a A6-elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20. The sequence may also be codon optimised for expression the target organism.

20 In one embodiment, the invention relates to an isolated nucleic acid comprising SEQ ID No. 21 encoding a  $\Delta\delta$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta\delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22. The sequence may also be codon optimised for expression the target organism.

The invention also relates to a vector comprising one or more of the isolated nucleic acids as specified above. The vector may further comprise a regulatory sequence.

30

The invention also relates to a transgenic microalgae with increased production of omega-3 LC-PUFAs wherein said microalgae expresses a nucleic acid comprises SEQ ID No. 7, 9, 15, 17, 19 or 21 or a sequence that encodes for a peptide that has at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%

35

homology to SEQ ID No. 8, 10, 16, 18, 20 or 22. Compositions comprising the transgenic microalgae, oil or lipids isolated therefrom and uses of as described herein in medicine or the formulation of a medicament, methods of treatment or feedstuff, foodstuff, pharmaceuticals or nutraceutical are also within the scope of the invention.

5

Without wishing to be bound by theory, the inventors believe that the activities of these nucleotides will prove useful in the heterologous reconstitution of the omega-3 long chain polyunsaturated fatty acid biosynthetic pathway in algae and plants. For example, the superior substrate-preference of the Ost809A6 enzyme distinguishes it from other *Ostreococcus* D6-desaturases, and can be used to maximise the flux of substrate through the n-3 pathway. Similarly, the Ost809A4 activity will prove useful in the specific conversion of DPA to DHA in transgenic photosynthetic organisms, whilst the FcEL06 activity provides a means by which GLA can be elongated to 20:3n-6.

15

In another embodiment, the invention therefore relates to the use of an isolated nucleic acid selected from a nucleic acid comprising or consisting of SEQ ID No. 7 or 9 encoding  $\Delta 6$ -desaturase (Ost809A6) comprising or consisting of SEQ ID No. 8 or 10, a functional variant thereof or a  $\Delta 6$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 10, a nucleic acid comprising or consisting of SEQ ID No. 16 or 18, a functional variant thereof or a A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 18, a nucleic acid comprising or consisting of SEQ ID No. 19 encoding A6-elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant thereof or a  $\Delta 6$ -elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding a  $\Delta \delta$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta \delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least

35

96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22 in the production of a transgenic organism with increased omega-3 fatty acid content. In particular, the invention relates to the use of isolated nucleic acids encoding a  $\Delta 6$ -desaturase (Ost809A6) to maximise the flux of substrate through the n-3 pathway and produce enhanced levels of EPA and/or DHA. In another embodiment, the invention relates to the use of an isolated nucleic acid encoding a A4-desaturase (Ost809A4) to convert DPA to DHA. In another embodiment, the invention relates to the use of an isolated nucleic acid encoding a  $\Delta 6$ -elongase to elongate GLA to 20:3.

10 In another embodiment, the invention relates to the use of an isolated nucleic acid selected from a nucleic acid comprising or consisting of SEQ ID No. 19 encoding  $\Delta 6$ -elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant thereof or a  $\Delta 6$ -elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding  $\Delta \delta$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta \delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22 in increasing DHA content. As shown in the examples and figure 13, DHA is increased by at least 10%, for example 14-17%.

25 In another embodiment, the invention relates to a method for producing a transgenic organism with increased of omega-3 LC-PUFAs production, in particular DHA and/or EPA, comprising transforming an organism with an isolated nucleic acid comprising or consisting of SEQ ID No. 7 or 9 encoding A6-desaturase (Ost809A6) comprising or consisting of SEQ ID No. 8 or 10, a functional variant thereof or a A6-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 10, a nucleic acid comprising or consisting of SEQ ID No. 16 or 18, a functional variant thereof or a A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least

85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 18, a nucleic acid comprising or consisting of SEQ ID No. 19 encoding  $\Delta 6$ -elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant thereof or a  $\Delta 6$ -elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding a  $\Delta \delta$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta \delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22.

In one embodiment, the invention relates to a method for producing a transgenic organism with increased of DHA production, comprising transforming an organism with an isolated nucleic acid nucleic acid selected from a nucleic acid comprising or consisting of SEQ ID No. 19 encoding  $\Delta 6$ -elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant thereof or a  $\Delta 6$ -elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding a  $\Delta 5$ -desaturase comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta \delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22 in increasing DHA content. As shown in the examples and Figures 13, DHA is increased by at least 10%, for example 14-17%.

In another embodiment, the invention relates to a method for increasing the production of omega-3 fatty acid transforming an organism with an isolated nucleic acid comprising or consisting of SEQ ID No. 7 or 9 encoding  $\Delta 6$ -desaturase (Ost809A6) comprising or consisting of SEQ ID No. 8 or 10, a functional variant thereof or a  $\Delta 6$ -

desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 10, a nucleic acid comprising or consisting of SEQ ID No. 16 or 18, a functional variant thereof or a  
5 A4-desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 18, a nucleic acid comprising or consisting of SEQ ID No. 19 encoding A6-elongase (FcEL06) comprising or consisting of SEQ ID  
10 No. 20, a functional variant thereof or a A6-elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding a  $\Delta\delta$ -desaturase  
15 comprising or consisting of SEQ ID No. 22, a functional variant thereof or a  $\Delta 5$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22.

20

In one embodiment, the invention relates to a method for increasing the production of omega-3 fatty acid transforming an organism with an isolated nucleic acid nucleic acid selected from a nucleic acid comprising or consisting of SEQ ID No. 19 encoding  $\Delta 6$ -elongase (FcEL06) comprising or consisting of SEQ ID No. 20, a functional variant  
25 thereof or a A6-elongase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20 or a nucleic acid comprising or consisting of SEQ ID No. 21 encoding a  $\Delta\delta$ -desaturase comprising or consisting of  
30 SEQ ID No. 22, a functional variant thereof or a  $\Delta\delta$ -desaturase that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22 in increasing DHA content. As shown in the examples and Figures 13, DHA is  
35 increased by at least 10%, for example 14-17%.

In one embodiment of the methods, method may further comprise transforming said microalgae with one or more additional nucleic acid that regulates the production of omega-3 fatty acids. In another embodiment, no additional nucleic acid that regulates the production of omega-3 fatty acids are introduced into said microalgae. Other heterologous nucleic acids, for example encoding a glucose transporter may be included.

In another aspect, invention relates to a host cell transformed with a vector comprising one or more of the isolated nucleic acids defined herein, specifically an isolated nucleic acid comprising SEQ ID No. 1, 3, 5, 7, 9, 15, 17, 19 or 21. In one embodiment, the host cell is transformed with a vector comprising one of the isolated nucleic acids defined herein and no other heterologous transgenes involved in the regulation of the LC-PUFAs biosynthetic pathway are expressed in said organism.

The host cell may be an algae or a higher plant cell. For example, the host cell is a microalgae. In one embodiment, the host cell is a diatom. The host cell may also comprise one or more additional transgene. For example, the host cell may be a transgenic microalgae described herein expressing a nucleic acid encoding for a  $\Delta 5$ -elongase.

The transgenic organism according to the methods described above may a microalgae or a higher plant. Preferably, the transgenic organism according to the methods described is a microalgae. The term microalgae is defined elsewhere herein and includes a diatom. In one embodiment, the microalgae is *P. tricornutum*. The term higher plant includes monocot and dicot plants. In one embodiment, the plant is a crop plant as described herein.

All references cited in this disclosure are herewith incorporated by reference with respect to their entire disclosure content and the disclosure content specifically mentioned in this application.

"and/or" where used herein is to be taken as specific disclosure of each of the multiple specified features or components with or without the other at each combination unless otherwise dictated. For example "A, B and/or C" is to be taken as specific disclosure of

each of (i) A, (ii) B, (iii) C, (iv) A and B, (v) B and C or (vi) A and B and C, just as if each is set out individually herein.

5 Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the invention and apply equally to all aspects and embodiments which are described.

The invention is further described in the following non-limiting examples.

## 10 Examples

### Example 1 Generation of transgenic algae over-expressing A6-desaturases and Generation of transgenic algae over-expressing $\Delta$ 5-elongase

#### 15 Materials and Methods

##### *Strains and growth conditions*

*P. tricornutum* UTEX 646 was grown in ESAW medium (Harrison et al., 1980) at 18°C and 20°C with moderate shaking under white fluorescent lights in constant illumination (30  $\mu\text{mol}$  and 60  $\mu\text{mol}$  photons  $\text{m}^{-2}\text{s}^{-1}$ ). Analysis of the wild-type and transgenic algae  
20 have been performed during exponential and stationary growth phases.

##### *Plasmid design and cloning*

The coding sequences for  $\Delta$ 6 - desaturase from *Ostreococcus tauri*, OtD6 (Domergue et al., 2005) and *O.tauri*  $\Delta$ 5- elongase OtElo5 (Meyer et al., 2004 ) were inserted as  
25 *Kpn*-*Xba* and *EcoRV*-*SacI* fragments, respectively, into pPha-T1 vector (Zaslavskaja et al., 2000), kindly provided by Dr. P.G.Kroth, (Universitat Konstanz, Germany). The coding region of OtD6 was used as a template to chemically synthesize (Genscript Corporation, NJ) codon-optimized nucleotide sequence OtD6PT for expression in  
30 *P.tricornutum*. This codon-optimized  $\Delta$ 6 - desaturase sequence was cloned into pPha-T1 vector, using *EcoRV*-*SacI* sites. The coding sequences for  $\Delta$ 6 - desaturase from *P. tricornutum*, PtD6 (Domergue et al., 2002) was inserted as *Bam**HI* -*Xba**I* fragment into pPha-T1 vector (Zaslavskaja et al., 2000).

##### *Biolistic transformation*



Biolistic transformation of *P. tricornutum* was performed according to previously described (Zaslavskaja et al., 2000; Kroth 2007). Bombarded cells were transferred onto ESAW agar plates containing 75 µg/ml zeocin. The zeocin plates were placed in 24 h light under fluorescent lights ( $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) and incubated at 20°C for 3 weeks.

5 Selected zeocin-resistant colonies were transferred to fresh zeocin plates and 2 ml ESAW+ zeocin cultures before being transferred to liquid medium minus antibiotic for lipid analysis.

#### *Fatty acid analysis*

10 Algae or yeast cells were harvested by centrifugation. Fatty acids were extracted and methylated as described (Garces and Mancha, 1993) with minor modifications. A 15ml aliquot of algal culture was harvested; following methylation the heptane fraction was concentrated and re-suspended in 40 µl solvent prior to injection of 1 µl on to the GC column. Methyl ester derivatives of total fatty acids extracted were analysed by GC  
15 using an Agilent DB-225 column and identified using known standards.

#### *Acyl-CoA profiling*

Algal cells were harvested by centrifugation, frozen in liquid nitrogen and extracted after Larson and Graham (2001), for reverse-phase LC with either quantitative analysis  
20 of fluorescent acyl-etheno-CoA derivatives or with electrospray ionization tandem mass spectrometry (multi reaction monitoring) in positive ion mode. For the analysis of etheno-CoA derivatives HPLC (Agilent 1200 LC system; Phenomenex LUNA 150 · 2 mm C18(2) column) was performed using the methodology and gradient conditions described previously (Larson and Graham 2001); whilst LC-MS/MS +MRM analysis  
25 followed the methods described by Haynes et al. 2008 (Agilent 1200 LC system; Gemini C18 column, 2 mm inner diameter, 150 mm with 5 mm particles). For the purpose of identification and calibration, standard acyl-CoA esters with acyl chain lengths from C14 to C20 were purchased from Sigma as free acids or lithium salts.

#### 30 *Lipid Profiling*

The molecular species of TAGs and PLs were analysed by electrospray ionisation triple quadrupole mass spectrometry (API 4000 QTRAP; Applied Biosystems). The molecular species of polar lipid were defined by the presence of a head-group fragment and the mass/charge of the intact lipid ion formed by ESI (Welti et al., 2002; Devaiah  
35 et al., 2006 with modifications described by Xiao et al. 2010). Such tandem ESI-MS/MS

precursor and product ion scanning, based on head group fragment, do not determine the individual fatty acyl species. Instead, polar lipids are identified at the level of class, total acyl carbons, and total number of acyl carbon-carbon double bonds. Polar lipids were quantified in comparison with a series of polar lipid internal standards.

5 Triacylglycerols (TAGs) measured after Krank et al. (2007) were defined by the presence of one acyl fragment and the mass/charge of the ion formed from the intact lipid (neutral loss profiling). This allows identification of one TAG acyl species and the total acyl carbons and total number of acyl double bonds in the other two chains. The procedure does not allow identification of the other two fatty acids individually nor the

10 positions (sn-1, sn-2, or sn-3) that individual acyl chains occupy on the glycerol. TAGs were quantified in a manner similar to the polar lipids, including background subtraction, smoothing, integration, isotope deconvolution and comparison of sample peaks with those of the internal standard (using LipidView, Applied Biosystems). However, whereas polar lipids within a class exhibit similar mass spectral response

15 factors, the mass spectral responses of various TAG species are variable, owing to differential ionization of individual molecular TAG species. In the data shown herein, no response corrections were applied to the data. The data were normalized to the internal standards tri15:0 and tri19:0

## 20 Results

### *Generation of transgenic algae over-expressing A6-desaturases.*

The native coding OtD6 and codon-optimized for expression in *P. tricornutum* nucleotide sequences for *O.tauri*  $\Delta 6$  - desaturase were cloned into pPha- T1 vector, generating expression cassettes OtD6N and OtD6Pt respectively, and the resulted

25 constructs were used to transform *P. tricornutum*.

### *Expression of OtD6N construct*

13 zeocin resistant colonies were obtained by transformation with OtD6N and selected for further screening. Selected colonies were transferred into liquid medium and

30 several positive transformants containing OtD6N were identified. We have studied the effects of temperature and light on the production of EPA and total fatty acids in Wt and transgenic *P.tricornutum*. Cultures were grown at different temperatures (18°C and 20°C) under constant illumination in different light intensity (25  $\mu\text{mol}$  and 60  $\mu\text{mol}$  photons  $\text{m}^{-2}\text{s}^{-1}$ ). GC-MS analyses have been performed during the exponential (E) and

35 stationary (S) phases of cell growth. Fatty acid profiling of WT and mutants showed

that palmitoleic acid (16:1 $\Delta^9$ ), EPA (20:5 n-3), palmitic acid (16:0) and myristic acid (14:0) were the major FAs detected in algal cells grown in both stages. Similarly to the results obtained by Tonon et al. (Tonon 2002) from the studies of *P. tricornutum* (CCAP 1052/1A) cell cultures grown at 18°C with 240  $\mu\text{E m}^{-2}\text{s}^{-1}$ , there was decrease in the amount of EPA and DHA as the cells of *P. tricornutum* UTEXS 646 used in our study shifted from exponential to stationary phase. Fatty acid analysis revealed that in cells transformed with Otd6N and grown at 20°C in light intensity 25  $\mu\text{E m}^{-2}\text{s}^{-1}$  and 60  $\mu\text{E m}^{-2}\text{s}^{-1}$  photons  $\text{m}^{-2}\text{s}^{-1}$  EPA and DHA decreased upon transition to stationary phase. However, the levels of EPA and DHA in Otd6N cells grown at 20°C, 60  $\mu\text{E m}^{-2}\text{s}^{-1}$  in stationary phase were higher than those of WT *P. tricornutum* (21.2% of EPA and 1.8% of DHA in Otd6N compared to 18.5% of EPA and 1.3% of DHA in WT (Table III, Fig. 1). In contrast, we found that in transgenic Otd6N cells grown at 18°C, 25  $\mu\text{E m}^{-2}\text{s}^{-1}$  levels of EPA and DHA increased in stationary phase compared to exponential phase and are significantly higher than in WT samples (30.2% of EPA and 1.8% of DHA in Otd6N compared to 16.5% of EPA and 0.9% of DHA in WT). Fatty acids profiles from Wt and Otd6N transgenic *P. tricornutum* showed no differences in  $\Delta^6$  - unsaturated fatty acids (GLA and SDA) composition, which were barely present.

#### *Expression of OtD6PT construct*

4 zeocin resistant colonies obtained by transformation with OtD6PT were selected to inoculate cultures for further screening and GC-MS analysis. The same trend towards decreasing levels of EPA and DHA in the stationary phase was observed for transgenic Otd6Pt cells grown at different light intensity and temperatures (Table III, Fig.1). Recombinant cells expressed higher levels of EPA (20.8% in the stationary phase at 20°C, 60  $\mu\text{E m}^{-2}\text{s}^{-1}$  and 22.2% at 18°C, 25  $\mu\text{E m}^{-2}\text{s}^{-1}$  compared to 18.5% and 16.8% in WT respectively). In addition to detection of higher levels of EPA we also observed an increase in DHA levels with minor variation between the two phases of growth (Table III, Fig.1).

#### *Generation of transgenic algae over-expressing OtElo5*

3 zeocin resistant clones obtained by transformation with OtElo5 were identified in an initial screen and used to inoculate cultures for further screening and GC-MS analysis. Cultures were grown at 20°C under constant illumination in 60  $\mu\text{E m}^{-2}\text{s}^{-1}$  photons  $\text{m}^{-2}\text{s}^{-1}$ . FAMES analysis of *P. tricornutum* transformed with OtElo5 have been performed during the exponential (E) and stationary (S) phases of cell growth and revealed the presence

of DPA in the range of 2.8-4.7% in transgenic clones which was not detected in WT cells (Table IV, Fig 2a). Levels of EPA in transformed clones were decreased to an average of 17.7% compared to 35.9% in WT in the exponential phase of growth and to 8.2% in clones over-expressing the Elo5 gene compared to 18.5% in WT during the stationary phase of growth. A substantial increase in DHA was observed in all 3 transgenic clones averaging 7.4% in exponential phase and 10.4% in stationary phase compared to 2.0% and 1.3% respectively in WT. DHA accumulation has been increased upon transition to stationary phase.

#### 10 *Determination of acyl-CoA pool composition*

To better understand the processes of acyl desaturation in diatoms the composition of the acyl-CoA pool was determined for the wild-type (WT) and transgenic *P. tricornutum*, expressing OtElo5-elongase (Fig.3). The study of acyl-CoA profile of WT *P. tricornutum* in the stationary phase of growth revealed that palmitic, palmitoleic, stearic, oleic and EPA-CoA were the most abundant, thus demonstrating the direct relationship between the levels of native fatty acids in the acyl-CoA pool vs the total fatty acids. EPA-CoA represented 5.7% of the acyl-CoA pool, indicating that this level of EPA-CoA could potentially act as an intermediate in the synthesis of DHA through elongation to 22:5n-3 and desaturation to 22:6n-3. Only traces (<1 .0) of 22:4 n-6, 22:5 n-3 (DPA) and DHA were detected in the CoA pool of WT *P. tricornutum*. As can be seen in Figure 3, similar analysis of transgenic *P. tricornutum* demonstrated a significant increase in the levels of 22:4 n-6, 22:5 n-3 (EPA) and DHA accompanying by the decrease in EPA levels. As shown in Figure 4, detailed analysis of the composition of the acyl-CoA pool through different stages of cell growth revealed that EPA and DHA were accumulating progressively from exponential to stationary phase displaying maximum levels of 5.2% and 6.3% in stationary phase.

#### *Profiling of TAG molecular species*

In this study we identified and compared the molecular species of TAGs formed by WT and OtElo5 transgenic *P. tricornutum* and investigated changes in TAG synthesis in response to transition from exponential to stationary phase. Cultures were grown at 20°C under constant illumination in 60  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  and analysed using ESI-MS. The mass spectrum obtained from direct infusion ESI-MS of algal lipid extracts shows that a majority of the molecular ions are observed between 750 and 950 mass/charge (m/z). We detected 26 individual TAG species in WT *P. tricornutum*. The

oil extracts of WT were predominantly composed of TAGs 46:1, 48:2 48:1, 48:2, and 48:3 and 50:3, having palmitic (18:0), palmitoleic (16:1), and myristic (14:0) acid substituents. TAG 48:1 (16:0/16:0/16:1) and 48:2 (16:0/16:1/16:1) constitute the main TAG molecular species that is expressed throughout the time course analysis of *P. tricornutum* cells (Figs 5a and 5B). An increase in the diversity of TAG molecular species (with as much as 29 individual TAGs) was detected from cells expressing OtElo5 -elongase. Specifically, new TAG species, 54:8, 54:9 and 56:8 were observed and transgenic cells show significantly higher levels of 54:7. DHA was incorporated in TAGs 52:7, 54:7, 54:8, 54:9 and 56:8. The time course (Fig. 6) also revealed that TAGs 54:7 and 56:8 appear to have more DHA incorporated into TAGs as the cells shift from the exponential growth phase to the stationary phase. TAGs molecular species 52:7, 54:8 and 54:9 demonstrated more or less constant DHA proportions when cultures were shifted from exponential to stationary phase. Levels of TAGs containing DHA averaged 12.5% in exponential stage and 10.5% in the stationary phase.

15

Table III. Fatty acid composition (molar %) of WT and transgenic *P. tricornutum* expressing *O. tauri*  $\Delta 6$  desaturase under different growth conditions at two growth stage, where E is the exponential and S is the stationary growth phases. Each measurement is the average of three biological replicates.

20

Cell strain	20°C 00μηιοΙ photons		20°C 25μηιοΙ photons		18°C 25μηιοΙ photons	
	E	S	E	S	E	S
Otd6N 14:0	6.3±1.1	5.6±1.6	11.5±0.7	7.6±1.5	13.0±1.1	10.9±1.0
16:0	16.0±0.5	21.0±1.3	12.8±0.9	16.8±1.6	15.3±0.8	16.6±1.1
16:1	28.3±1.7	36.5±1.6	32.8±0.2	30.3±1.9	35.1 ±2.1	34.4±2.5
16:3	2.5±0.2	0.9±0.2	4.0±0.6	0.9±0.1	3.6±0.0	2.7±0.2
18:0	0.5±0.0	0.7±0.0	0.3±0.0	0.4±0.0	ND	ND
18:1	6.2±1.4	8.6±1.5	18.1 ±0.0	24.9±0.3	2.1±0.2	2.5±0.2
18:2n-6	1.5±0.1	0.6±0.0	ND	ND	1.4±0.2	1.4±0.2
18:3 n-6	0.7±0.3	1.3±0.3	ND	ND	ND	ND
18:4 n-3	0.8±0.1	0.8±0.1	ND	0.4±0.0	1.0±0.4	1.0±0.4
20:5 n-3	32.2±3.6	21.2±1.9	20.6±1.1	17.8±2.6	27.1 ±2.7	30.2±3.2
22:6 n-3	2.3±0.2	1.8±0.3	1.4±0.1	1.0±0.1	1.4±0.4	1.8±0.3
Others	6.89±0.6	4.3±0.6	12.2±1.8	6.0±0.2	5.7±0.4	6.2±0.6

Otd6Pt	14:0	7.0±1.4	4.9±1.0	5.6±0.2	4.9±0.2	12.8±0.1	7.4±0.4
	16:0	16.3±1.3	20.2±1.5	9.5±0.3	16.8±0.7	17.0±0.9	20.4±0.2
	16:1	27.1 ±4.0	38.6±3.6	24.5±0.2	33.4±7.9	28.3±1.2	35.8±2.6
	16:3	2.5±0.2	1.1±0.3	4.0±0.6	1.4±0.1	2.9±0.0	5.2±1.1
	18:0	0.5±0.1	0.6±0.1	0.3±0.0	0.4±0.0	ND	ND
	18:1	7.8±0.2	8.7±0.4	26.9±5.4	24.9±0.3	6.0±0.9	8.5±0.9
	18:2 n-6	1.1±0.2	1.1±0.1	ND	ND	1.2±0.0	1.2±0.0
	18:3 n-6	1.2±0.2	0.8±0.0	0.2±0.0	0.2±0.0	ND	ND
	18:4 n-3	1.1±0.1	1.2±0.1	0.6±0.1	0.6±0.0	1.5±0.0	1.5±0.0
	20:5 n-3	33.2±1.4	20.8±3.5	27.0±4.0	16.6±2.0	25.8±0.1	22.2±1.3
	22:6 n-3	1.7±0.3	1.5±0.4	1.3±0.1	1.2±0.6	1.1±0.0	1.3±0.2
	Others	9.2±0.6	4.3±0.9	12.3±1.8	5.5±3.6	7.3±0.3	3.1±0.3
	WT	14:0	7.7±0.5	4.8±0.1	5.1 ±0.2	4.8±0.5	10.9±0.5
16:0		16.5±0.4	22.2±0.6	11.0±2.0	16.6±3.2	19.7±0.4	21.1±1.3
16:1		28.4±0.6	41.8±0.5	22.3±1.1	32.2±4.1	35.8±0.6	42.1 ±2.5
16:3		2.4±0.3	1.0±0.1	2.6±0.6	0.6±0.1	2.4±0.3	1.4±0.0
18:0		0.4±0.0	0.5±0.0	0.3±0.1	0.3±0.1	ND	ND
18:1		3.8±0.8	7.3±0.2	28.9±1.4	25.7±4.9	6.1±0.3	8.2±0.1
18:2n-6		1.4±0.1	0.6±0.0	ND	ND	1.1±0.1	0.8±0.1
18:3n-6		0.7±0.0	0.6±0.0	ND	ND	ND	ND
18:4 n-3		0.8±0.0	1.0±0.0	0.6±0.0	0.4±0.1	1.0±0.7	0.6±0.8
20:5n-3		35.9±1.6	18.5±0.4	27.6±2.3	17.1 ±2.5	22.2±0.7	16.8±2.8
22:6n-3		2.0±0.3	1.3±0.0	1.8±0.1	1.3±0.3	0.8±0.1	0.9±0.2
Others		6.8±0.3	2.4±0.3	10.0±0.9	5.1±0.8	4.9±0.5	2.9±0.3

Table IV. Fatty acid composition (molar %) of WT and transgenic *P. tricornutum* expressing Ot Elo5 during exponential (E) and stationary (S) phases. Cultures were grown at 20°C 60  $\mu\text{mol m}^{-2}\text{s}^{-1}$  under constant agitation at 70 rpm. Each measurement is the average of 3 biological replicates.

5

Fatty acids	WT		OtElo5	
	E	S	E	S
14:0	7.7±0.5	4.8±0.5	8.4±1.2	5.3±1.6
16:0	16.5±0.5	22.1±0.6	16.8±0.6	17.4±1.3
16:1	28.4±0.6	41.8±0.5	32.9±0.4	42.5±1.6
16:3	2.4±0.3	1.0±0.0	3.6±0.6	1.7±0.6
18:0	0.4±0.0	0.5±0.0	0.6±0.0	0.5±0.0
18:1	3.8±0.8	7.3±0.2	6.8±1.1	6.8±1.5
18:2 n-6	1.4±0.1	0.6±0.0	0.6±0.0	0.3±0.0
18:3n-6	0.7±0.0	0.6±0.0	0.2±0.0	0.2±0.2
18:4 n-3	0.8±0.0	1.0±0.0	1.6±0.0	2.0±0.1
20:5 n-3	35.9±1.6	18.5±0.4	17.7±2.4	8.2±2.0
22:5 n-3	ND	ND	3.3±0.5	3.4±1.2
22:6 n-3	2.0±0.3	1.3±0.1	7.4±1.2	10.4±0.3
24:0	5.2±0.2	2.1±0.0	5.2±0.4	3.1±0.4
Others	1.8±0.3	0.3±0.3	4.1±0.4	2.4±0.6

### Discussion

Many marine microbes produce high levels of EPA and DHA but only few species have the ability to partition these fatty acids into storage lipids in the form of triacylglycerols (TAGs). The majority of algal species accumulate saturated and mono-unsaturated fatty acids in TAGs (Harwood, 1998; Roessler, 1990b). Partitioning of LC-PUFAs into TAGs have been observed in *Parietochloris incise* (Bigogno et al., 2002), the freshwater red microalga *Porphyridium omentum* (Cohen et al., 2000), and marine microalgae *Nannochloropsis oculata*, *Phaeodactylum tricornutum*, *Thalassiosira pseudonana* and *Pavlova lutheri*, (Tonon et al., 2002). Thus these species are good candidates for further studies, in order to understand the processes responsible for the incorporation of LC-PUFAs into storage oils in microalgae.

At present it is generally accepted that oleaginous algae produce small quantities of TAG under optimal growth conditions (Hu et al. 2008). Among major factors affecting triacylglycerol accumulation and fatty acid composition in microalgae are temperature and light intensity. Generally, it is considered that fatty acid unsaturation increases with temperature decrease and low light favours the formation of PUFAs. For example, in *P.tricornutum* UTEXS 640 optimal culture temperature for EPA production was 21.5 to 23°C (Yongmanitchai W. and Ward O., 1991). A temperature shift strategy has been

25

employed to enhance the overall n-3 PUFAs (including EPA) production because the optimal temperature for microalgal growth is often higher than that for n-3 PUFAs formation (Jiang and Chen, 2000). Such a phenomenon has been observed in many different algal species including *P. omentum* (Springer et al., 1994), *Nannochloropsis* sp. (Sukenik, 1991) and *P. irregular* (Stinson et al., 1991). However, Ohta et al. (1993) observed that the optimal temperature for growth of *P. purpureum* also yields a biomass with the highest EPA content. These results suggest that the effect of temperature on cell growth and n-3 PUFA production should be carefully studied for individual microalgal species.

Profiling of TAG species in *P. triornutum* has been previously reported (Yongmanitchai and Ward 1993; Yu et al., 2009). We observed the same predominant fatty acids (i.e., 14:0, 16:0, 16:1, 16:3, and 20:5) incorporated in TAGs as described in these earlier studies. Yongmanitchai and Ward 1993 identified only 18 TAG molecular species via reverse-phase HPLC analysis. Due to the high resolution and sensitivity of ESI-MS, Yu et al., 2009 were able to detect twofold more species in algal oil extracts (14 of the 18 species they detected by HPLC, at comparable percentage composition. However, TAGs 48:7, 48:9, 48:12, and 54:10 were not detected which could be explained by the difference in the *P. triornutum* strains and culture conditions.

## Example 2

### Identification and characterization of new activities for PUFAs biosynthesis in algae and plants

#### 2.1 Identification of a $\Delta 6$ - desaturase from the microalga *Ostreococcus* RCC809

Genome of green alga *Ostreococcus* RCC809 was analysed with BLAST using already known N-terminal cytochrome b5-fusion desaturases as query. This analysis revealed the presence of several genes coding for putative PUFA desaturases. The deduced open reading frames were used as templates to chemically synthesise (Genscript Corporation, NJ) codon-optimised nucleotide sequences for expression in diatoms.

#### *Functional characterization of putative *Ostreococcus* RCC809 $\Delta 6$ -desaturase in yeast.*

The codon-optimised open reading frame of the putative  $\Delta 6$  - desaturase (SEQ ID No.s 7 to 10, hereafter designated Ost809A6) was inserted as *KpnI-SacI* fragment behind the galactose -inducible GAL1 promoter of the yeast expression vector pYES2 (Invitrogen, NJ). Ost809A6



The *S. cerevisiae* strain W303-1A was transformed with plasmid DNA using a lithium acetate method. Cultures were grown at 22°C in the presence of 2% (v/v) raffinose for 48 h, and expression of the transgene was induced by addition of galactose to 2% in the presence of 0.5 mM of linoleic acid (LA, 18:2n-6) and 1% (w/v) tertgitol NP-40 (Sigma) as described (Sayanova et al., 2001).

The predicted function of the candidate desaturase Ost809A6 (predicted to encode a C18  $\Delta$ 6-desaturase of 461 amino acids) was investigated by expression studies in *S. cerevisiae* in the presence of a range of potential fatty acid substrates. Total fatty acid methyl esters from yeast cells were then analysed by GC-FID and the identity of novel peaks confirmed by GC-MS and co-migration with authentic standards. As shown in Fig. 8, expression of a synthetic ORF encoding Ost809A6, confirmed the enzymatic capability to convert exogenously supplied substrate ( $\alpha$ -Linolenic acid, ALA; C18:A9,12,15) to the A6-desaturated product SDA (18:4, n-3). In the absence of galactose, the exogenous substrate ALA is not converted to SDA. Thus, on the basis of these results, Ost809A6 was confirmed as a D6-desaturase. The substrate selectivity of Ost809A6 was determined by exogenously supplying equal quantities of LA and ALA in the growth media. As it is shown in Figure 9, Ost809A6 only recognised the n-3 fatty acid ALA as a substrate, whereas the n-6 substrate was not desaturated. This is distinct from a  $\Delta$ 6-desaturase identified from *Ostreococcus tauri* (Domergue et al, 2005), which showed activity towards both LA and ALA as substrates. Thus Ost809A6 is superior and distinct for the exclusive production of  $\Delta$ 6-desaturated n-3 fatty acids.

Yeast cultures were supplemented with different potential FA substrates (listed in Table V) but desaturation activity of O809d6 was detected only in the presence of ALA.

### 2.2 Identification of putative $\Delta$ 4 - desaturase from O809

The genome sequence of *Ostreococcus* RCC809 [http://genome.jgi-psf.org/OstRCC809\\_2/OstRCC809\\_2.home.html](http://genome.jgi-psf.org/OstRCC809_2/OstRCC809_2.home.html) was searched with previously functionally characterised sequences of A4-desaturases and the presence of an apparent candidate (JGI protein ID # 40461) for a A4-desaturase was detected. The deduced open reading frame was used as a template to chemically synthesise (Genscript Corporation, NJ) codon-optimised nucleotide sequences for expression in diatom *P. tricornutum* (SEQ ID No.s 15 to 18).

*Functional characterization of putative  $\Delta 4$  - desaturase from O809 in yeast.*

The codon-optimised for expression in *P. tricornutum* open reading frame of the putative  $\Delta 4$  - desaturase was inserted as *KpnI-SacI* fragment behind the galactose -  
 5 inducible GAL1 promoter of the yeast expression vector pYES2 (Invitrogen, NJ).

As can be seen in Fig 10, galactose-dependent expression of the Ost809 protein 40461 resulted in the  $\Delta 4$ -desaturation of DPA to DHA, confirming the function of this ORF as a C22  $\Delta 4$ -desaturase and on this basis we designated this gene as Ost809A4.

10 Note that in the absence of the inducer (galactose), no DHA is detected, nor in the absence of the Ost809A4 ORF.

2.3 Identification of a A6-elongase from *Fragilariopsis cylindrus*

The publically available genome sequence of the marine diatom *Fragilariopsis cylindrus* (<http://genome.jgi-psf.org/Fracy1/Fracy1.home.html>) was analysed with  
 15 BLAST using already known  $\Delta 6$ -elongase sequences (such as the  $\Delta 6$ -elongase from *C.elegans* - Beaudoin et al, 2000) as query and a candidate open reading frame (designated Frag #177742) was used as a template to chemically synthesise (Genscript Corporation, NJ) codon-optimised nucleotide sequence for expression in  
 20 *T.pseudonana*.

*Functional characterization of Fc A6-elongase in transgenic yeast*

Heterologous expression of Frag #177742 in *S. cerevisiae* was carried out exactly as described above, with the codon-optimised ORF cloned into the yeast expression  
 25 vector pYES2. Galactose-mediated induction of this construct was used to confirm that this ORF functioned as a  $\Delta 6$ -elongase, specifically elongating C18  $\Delta 6$ -unsaturated substrates such as GLA to a C20 form. As can be seen in figure 11, elongation of GLA to 20:3 only occurs in the presence of galactose and the ORF Frag #177742. On the basis of these results, this was redesignated FcEL06.

30 Table V. List of Substrates Tested:

Ost809 D6

18:2, ALA, GLA, 18:2 & 18:3, 20:4n-6 (ARA), 20:2, ERA, ETA, 22:5n-6 (DPA)

FcElo6

35 18:2, GLA, GLA & SDA

Ost809A4DPA

(Substrates underlined are those which worked)

Table VI. Fatty acid composition of yeast cells expressing Ost809A6, FcElo6 or Ost809A4 and substrate specificities of each of these

5

Fatty Acid Composition (molar %)										
Construct										
FA	O809 $\Delta 6$ Gal -	O809 $\Delta 6$ Gal +	O809 $\Delta 6$ Gal -	O809 $\Delta 6$ Gal +	Fc Elo6 Gal -	Fc Elo6 Gal +	O809 d4 Gal -	O809 d4 Gal +	pYes2 BPX72	pYes2 HP1
<b>16:0</b>	26.2	26.0	24.8	22.4	25.2	23.2	22.8	20.4	26.1	22.2
<b>16:1</b>	25.6	28.8	26.3	27.9	23.7	26.3	49.2	51.0	29.2	51.5
<b>18:0</b>	ND	ND	ND	ND	ND	ND	4.2	4.4	ND	3.9
<b>18:1</b>	15.2	16.3	13.6	15.4	ND	ND	20.2	21.6	17.5	19.7
<b>18:2</b>	5.8	6.8	ND	ND	ND	ND	ND	ND	ND	ND
<b>GLA</b>	ND	ND	ND	ND	38.7	22.8	ND	ND	ND	ND
<b>ALA</b>	25.6	11.9	32.9	15.7	ND	ND	ND	ND	27.2	ND
<b>SDA</b>	1.6	10.3	2.3	18.5	ND	ND	ND	ND	ND	ND
<b>DHGLA</b>	ND	ND	ND	ND	ND	14.1	ND	ND	ND	ND
<b>DPA</b>	ND	ND	ND	ND	ND	ND	2.9	2.3	ND	2.7
<b>DHA</b>	ND	ND	ND	ND	ND	ND	ND	0.4	ND	ND

Table VII. Substrate specificity

10

Substrate Specificity		
Construct	Substrate	%
<b>Ost809<math>\Delta 6</math></b>	18:2	0.0
<b>Ost809<math>\Delta 6</math></b>	18:3 ALA	54.1
<b>FcElo6</b>	18:3 GLA	38.1
<b>Ost809<math>\Delta 4</math></b>	22:5 DPA	13.5

15

On the basis of the identification of novel forms of the  $\Delta 6$ -desaturase (Ost809A6),  $\Delta 4$ -desaturase (Ost809A4) and the  $\Delta 6$ -elongase (FcEL06), it is very likely that these activities will prove useful in the heterologous reconstitution of the omega-3 long chain polyunsaturated fatty acid biosynthetic pathway in algae and plants. For example, the superior substrate-preference of the Ost809A6 enzyme distinguishes it from other

Ostreococcus  $\Delta$  6-desaturases, and can be used to maximise the flux of substrate through the n-3 pathway. Similarly, the Ost809A4 activity will prove useful in the specific conversion of DPA to DHA in transgenic photosynthetic organisms, whilst the FcEL06 activity provides a means by which GLA can be elongated to 20:3n-3.

5

### Example 3

#### Expression of single omega-3 LC-PUFA biosynthetic genes in *Pheodactylum tricornutum* can increase the endogenous accumulation of DHA

#### 10 Materials and methods

##### *Strains and growth conditions*

*P. tricornutum* UTEX 646 was grown in ESAW medium (Harrison et al., 1980) at 20°C with moderate shaking under white fluorescent lights in constant illumination (100  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ). Analysis of the wild-type and transgenic algae have been performed during stationary growth phase.

15

##### *Plasmid design and cloning*

The coding sequence for A6-elongase FcElo6 (protein ID 177742) was used as a template to chemically synthesize (Genscript Corporation, NJ) a codon-optimized nucleotide sequence for expression in *T. pseudonana*. The codon-optimized sequence was inserted as *EcoRV-SacI* fragments, respectively, into pPha-T1 vector (Kroth, 2007; Zaslavskaja et al., 2000).

20

#### Results

25 Expression of FcElo6 resulted in increase of DHA levels up to 14-17% (Figure 13).

### Example 4

#### Co-expression of two genes

##### Material and methods

##### 30 *Design of double-gene vector pPhOS2 and transformation cassettes*

The *EcoRI* –*HindIII* fragment of of pPha -T1 vector containing MCS was replaced by the synthetic sequence comprising of *fcpA* terminator and *fcpA* promoter flanked by 3 multiple cloning sites (MCSs) with unique restriction sites (Figure 14). The coding sequences for *O. tauri*  $\Delta\delta$ -elongase OtElo5 was inserted as *KpnI-SacI* fragment into

35

position 1 of pPhOS vector generating pPhOS2.1.1 construct. The codon optimized for expression in *P.tricornutum* coding sequences for *O. tauri*  $\Delta$ 6-desaturase OtD6Pt was inserted as *Bam*HI- *Xba*I fragment into position 2 of pPhOS2.1.1 generating pPhOS2.2.1 construct.

5

## Results and Discussion

### *Multigene expression in transgenic P. tricornutum*

To facilitate the expression of multiple heterologous genes in *P. tricornutum*, a new vector (designated pPhOS2- Figure 14) was constructed. This vector is based on previously described pPha-T1 vector (Zaslavskaja et al., 2000) and contains two multiple cloning sites (MCS) with unique restriction sites for inserting genes of interest. Each of these MCS is flanked by the promoter and terminator regions of the *FcpA* gene (Zaslavskaja et al., 2000) to promote the co-expression of two inserted genes. The coding sequence for *O. tauri*  $\Delta$ 6-elongase OtElo5 was inserted into position 1 of pPhOS2 vector and the resulting construct pPhOS2.1.1 was used to transform *P. tricornutum*. Cultures were grown at 20°C and 16°C under constant illumination (60  $\mu$ mol photons  $m^{-2}s^{-1}$ ). Multiple (5) independent zeocin-resistant colonies were obtained and used to inoculate cultures for further GC-MS analysis. The mean levels of DHA in analysed pPhOS2.1.1 strains was 9.0% (Table VIII; Figure 1), similar to levels previously observed with OtElo5 expression in pPha-T1, confirming the functionality of this modified vector. The codon-optimized coding sequences for *O. tauri*  $\Delta$ 6-desaturase OtD6Pt was subsequently inserted into position 2 of construct pPhOS2.1.1, generating the two-gene (plus the selectable marker gene *ble*) pPhOS2.2.1 vector. This expression plasmid was introduced into *P. tricornutum* via biolistics and multiple independent zeocin-resistant colonies were obtained and used to inoculate cultures for further screening. Cultures were grown at 16 and 20°C under constant illumination (60  $\mu$ mol photons  $m^{-2}s^{-1}$ ). FAMES analysis of transgenic strains expressing either single or double gene constructs revealed a further increase in DHA levels in transgenic strains co-expressing both OtElo5 and OtD6Pt, indicating the here-demonstrated potential for iterative metabolic engineering in *P. tricornutum* for high value lipid traits (Figure 15, Table VIII).

30

**Table VIII.** Fatty acid composition (Mol %) of wild-type (Pt\_WT) and transgenic *P. tricornutum* expressing pPhOS2.1 and pPhOS2.2 at 16°C and 20°C. Each measurement is the average of 3 biological replicates ( $\pm$  Standard Error).

5

Fatty Acids	Pt_WT		pPhOS2.1		pPhOS2.2	
	16°C	20°C	16°C	20°C	16°C	20°C
	5.3 $\pm$ 0.2	4.8 $\pm$ 0.1	5.1 $\pm$ 0.2	5.3 $\pm$ 0.3	6.7 $\pm$ 0.2	6.3 $\pm$ 0.1
	22.3 $\pm$ 1.0	22.1 $\pm$ 0.4	19.2 $\pm$ 0.4	18.9 $\pm$ 1.4	17.7 $\pm$ 0.5	18.4 $\pm$ 0.3
14:0	39.2 $\pm$ 1.6	41.8 $\pm$ 0.3	39.0 $\pm$ 0.6	40.1 $\pm$ 1.7	43.6 $\pm$ 1.0	40.6 $\pm$ 0.5
16:0	0.8 $\pm$ 0.4	1.0 $\pm$ 0.1	1.2 $\pm$ 0.1	1.8 $\pm$ 0.4	nd	2.0 $\pm$ 0.1
16:1	0.5 $\pm$ 0.0	0.5 $\pm$ 0.1	0.6 $\pm$ 0.1	0.3 $\pm$ 0.1	0.5 $\pm$ 0.0	0.3 $\pm$ 0.1
16:3	6.8 $\pm$ 0.0	4.3 $\pm$ 0.1	2.6 $\pm$ 0.1	2.2 $\pm$ 0.4	1.2 $\pm$ 0.6	0.6 $\pm$ 0.4
18:0	2.2 $\pm$ 0.1	2.8 $\pm$ 0.1	2.1 $\pm$ 0.2	4.2 $\pm$ 0.3	2.7 $\pm$ 0.1	3.7 $\pm$ 1.0
18:1 n-9	1.0 $\pm$ 0.1	1.0 $\pm$ 0.1	1.7 $\pm$ 0.1	1.1 $\pm$ 0.1	1.6 $\pm$ 0.0	1.1 $\pm$ 0.1
18:1 n-11	20.3 $\pm$ 1.9	18.5 $\pm$ 0.1	10.4 $\pm$ 0.3	9.8 $\pm$ 1.0	10.0 $\pm$ 0.4	8.2 $\pm$ 0.1
	nd	nd	3.4 $\pm$ 0.4	1.9 $\pm$ 0.3	5.5 $\pm$ 0.1	2.2 $\pm$ 0.3
18:4 n-7	1.5 $\pm$ 0.2	1.3 $\pm$ 0.1	9.0 $\pm$ 0.3	9.4 $\pm$ 1.0	10.3 $\pm$ 0.4	11.4 $\pm$ 0.2
20:5 n-3	2.9 $\pm$ 0.4	2.4 $\pm$ 0.1	3.2 $\pm$ 0.1	2.3 $\pm$ 0.2	3.3 $\pm$ 0.1	2.2 $\pm$ 0.8
22:5 n-3	2.0 $\pm$ 0.5	1.9 $\pm$ 0.1	1.1 $\pm$ 0.1	2.9 $\pm$ 0.5	2.9 $\pm$ 0.3	3.2 $\pm$ 0.2
22:6 n-3						
24:0						
Others						

## Example 5

### Auxorophic growth

#### Material and methods

#### 10 Design of double-gene vector pPhOS2 and transformation cassettes

The EcoRI –HindIII fragment of of pPha -T1 vector containing MCS was replaced by the synthetic sequence comprising of *fcpA* terminator and *fcpA* promoter flanked by 3 multiple cloning sites (MCSs) with unique restriction sites (Fig. 16). The coding sequences for *O. tauri*  $\Delta\delta$ -elongase OtElo5 was inserted as *KpnI-SacI* fragment into position 1 of pPhOS vector generating pPhOS2.1 .1 construct. The codon optimized for expression in *P.tricornutum* coding sequences for glucose transporters from

15

*Physcomitrella patens* (designated Ppglutl), and human erythrocytes (designated Hsglutl), were inserted as *Bam*HI-*Xba*I fragments into position 2 of pPhOS2.1.1 generating pPhOS\_Ppglut and pPhOS\_HSglut constructs. The resulting constructs were used to transform *P. tricornutum* via biolistics.

5

## Results

Multiple (>10) independent zeocin-resistant colonies were obtained by transformations with these two expression cassettes and used to inoculate cultures for further GC-MS analysis. Transgenic *P. tricornutum* strains expressing pPhOS\_Ppglut and pPhOS\_HSglut constructs accumulating DPA and elevated levels of DHA were selected for further analysis. (Fig. 16 and Fig. 17). The transformants were transferred to solid medium containing 0.5% of glucose, placed in complete darkness and monitored for growth (Fig. 18).

15

## REFERENCES

- Ahmann, K., Heilmann, M., Feussner, I., 2011. Identification of a D4-desaturase from the microalga *Ostreococcus lucimarinus*. *Eur. J. Lipid Sci. Technol* 113,7, 832-840.
- Arao, T., Kawaguchi, A., Yamada, M., 1987. Positional distribution of fatty acids in lipids of the marine diatom, *Phaeodactylum tricornutum*. *Phytochemistry*26, 2573-2576.
- Arao, T., Yamada, M., 1994. Biosynthesis of polyunsaturated fatty acids in the marine diatom, *Phaeodactylum tricornutum*. *Phytochemistry*35,1 177-181 .
- Bigogno, C , Khozin-Goldberg, I., Boussiba, S., Vonshak, A., Cohen, Z., 2002. Lipid and fatty acid composition of the green oleaginous alga *Parietochloris incisa*, the richest plant source of arachidonic acid. *Phytochemistry*60, 497-503.
- Blanchemain, A., Grizeau, D., 1999. Increased production of eicosapentaenoic acid by *Skeletonema costatum* cells after decantation at low temperature. *Biotechnol. Tech* 13. 497-501 .
- Calder, P.C., 2003. N-3 polyunsaturated fatty acids and inflammation: from molecular biology to the clinic. *Lipids* 38, 343-352.
- Cohen, Z., Khozin-Goldberg, I., Adlrestein, D., Bigogno, C , 2000. The role of triacylglycerols as a reservoir of polyunsaturated fatty acids for the rapid production of chloroplastic lipids in certain microalgae. *Biochem. Soc. Trans.* 28, 740-743.
- Das, U., N., 2002. The lipids that matter from infant nutrition to insulin resistance. *Prostaglandins Leukot Essent Fatty Acids* 67, 1-12.

30

- Deviah, S.P., Roth M.R., Baughman E., Li M., Tamura P., Jeannotte R., Welti R., Wang X., 2006. Quantitative profiling of polar glycerolipid species from organs of wild-type *Arabidopsis* and a PHOSPHOLIPASE Da1 knockout mutant. *Phytochemistry* 67, 1907-1924.
- 5 Domergue F., Lerchl J., Zaehring U., Heinz E., 2002. Cloning and functional characterization of *Phaeodactylum tricornutum* front-end desaturases involved in eicosapentaenoic acid biosynthesis. *Eur J Biochem* 269, 4105-4113.
- Domergue F, Abbadi A, Zaehring U, Moreau H, Heinz E, 2005. In vivo characterization of the first acyl-CoA A6-desaturase from a member of the plant kingdom, the  
10 microalgae *Ostreococcus tauri*. *Biochem J* 389, 483-490.
- Garces, M., Mancha, R., 1993. One-Step Lipid Extraction and Fatty Acid Methyl Esters Preparation from Fresh Plant Tissues. *Analytical Biochemistry* 211, 139-143.
- Harwood, J.L., Guschina I.A., 2009. The versatility of algae and their lipid metabolism. *Biochimie*. 91, 679-684.
- 15 Harrison, P.J., Waters R.E., Taylor F.J.R., 1980. A broad spectrum artificial medium for coastal and open ocean phytoplankton. *J. Phycol.* 16, 28-35.
- Haynes, C.A., Allegood, J.C., Sims, K., Wang, E.W., Cameron Sullards, M., Merrill, A.H., 2008. Quantitation of fatty acyl-coenzyme As in mammalian cells by liquid chromatography-electrospray ionization tandem mass spectrometry, *J. Lipid Res.* 49, 1113-1125.
- 20 Harwood, J.L., 1998. Membrane lipids in algae. In *Lipids in Photosynthesis: Structure, Function and Genetics* (Siegenthaler, P.A. and Murata, N., eds). Dordrecht, The Netherlands: Kluwer Academic Publishers 53-64.
- Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., Darzins, A.,  
25 2008. Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *Plant J.* 54, 621-639.
- Horrocks, L.A., Yeo, Y.K., 1999. Health benefits of DHA. *Pharmacological Research* 40, 211-225.
- Jiang, Y., Chen F., 2000. Effects of temperature and temperature shift on docosahexaenoic acid production by the marine microalga *Cryptohecodinium cohnii*. *J. Am. Oil. Chem. Soc* 77 613-617.
- 30 Kitano, M., Matsukawa R., Karube, I., 1997. Changes in eicosapentaenoic acid content of *Navicula saprophiila*, *Rhodomonas salina* and *Nitzschia* sp. under mixotrophic conditions. *J. Appl. Phycol* 9, 559-563.



- Krank J., Murphy R. C , Barkley R. M., Duchoslav, E., McAnoy, A., 2007. Qualitative analysis and quantitative assessment of changes in neutral glycerol lipid molecular species within cells. *Methods in Enzymology* 432, 1-20.
- Kroth, P., 2007. Genetic transformation: a tool for study protein targeting in diatoms. *Methods in Molecular Biology* (Clifton, NJ) 390, 257.
- 5 Kyle, D.J., Sicotte, V.J., Singer, J. and Reeb, S.E., 1992. Bioproduction of docosahexaenoic acid (DHA) by microalgae. In *Industrial Applications of Single Cell Oils* (Kyle, D.J. and Ratledge, C , eds). Champaign, IL: American Oil Chemists' Society. 287-300.
- 10 Larson, T.R. and Graham, I.A., 2001 . A novel technique for the sensitive quantification of acyl CoA esters from plant tissues. *Plant J.* 25, 115-125.
- Meyer A., Kirsch H, Domergue F, Abbadi A, Sperling P, Bauer J, Cirpus P, Zank TK, Moreau H, Roscoe TJ, Zahringer U, Heinz E., 2004. Novel fatty acid elongases and their use for the reconstitution of docosahexaenoic acid biosynthesis. *Journal of Lipid*
- 15 *Research* 45, 1899-1909.
- Molina Grima, E., Sanchez Perez, J.A., Garcia Sanchez, J.L, Garcia Camacho, F., Lopez Alonso, D., 1992. EPA from *!sochrysis galbana*. Growth conditions and productivity. *Process Biochem* 27, 299-305.
- Molina Grima, E., Robles Medina, A., Gimenez Gimenez, A., Ibanez Gonzalez, M.J.
- 20 1996. Gram-scale purification of eicosapentaenoic acid (EPA, 20: 5n-3) from wet *Phaeodactylum tricornutum* UTEX 640 biomass. *J. Appl. Phycol.* 8, 359-367.
- Moreno, V.J., De Moreno, J.E.A., Brenner, R.R., 1979. Biosynthesis of unsaturated fatty acids in the diatom *Phaeodactylum tricornutum*. *Lipids* 14, 15-19.
- Navarro, E., Esteve, M., Olive, A., 2000. Abnormal fatty acid pattern in rheumatoid
- 25 arthritis. A rationale for treatment with marine and botanical lipids. *J Rheumatol.* 27, 298-303.
- Nugent, A.P., 2004. The metabolic syndrome, *Nutr Bull*, 29, 36-43.
- Onta S., Chang, T., Aozasa, O., Ikegami, N., Miyata, H., 1993. Alterations in fatty acid composition of marine red alga *Porphyridium purpureum* by environmental factors. *Bot.*
- 30 *Mar*, 36, 103-107.
- Qiang, H., Zhengyu, H., Cohen, Z., Richmond, A., 1997. Enhancement of eicosapentaenoic acid (EPA) and Klinolenic acid (GLA) production by manipulating algal density of outdoor cultures of *Monodus subterraneus* (Eustigmatophyta) and *Spirulina platensis* (Cyanobacteria). *Eur. J. Phycol* 32, 81-86.

- Qiu, X., Hong, H., MacKenzie, S.L., 2001. Identification of a  $\Delta^4$  fatty acid desaturase from *Thraustochytrium* sp. involved in the biosynthesis of docosahexanoic acid by heterologous expression in *Saccharomyces cerevisiae* and *Brassica juncea*J. Biol.Chem 276, 31561-6.
- 5 Radakovits, R., Eduafo, P., Posewitz M., 2011. Genetic engineering of fatty acid chain length in *Phaeodactylum tricornutum*. Metab.Eng 13, 89-95.
- Renaud, S.M., Parry, D.L., Thinh, L.V., 1994. Microalgae for use in tropical aquaculture: I. Gross chemical and fatty acid composition of twelve species of microalgae from the North Territory, Australia. J. Appl. Phycol 6, 337-345.
- 10 Renaud, S.M., Thinh, L.V., Parry, D.L., 1999. The gross chemical composition and fatty acid composition of 18 species of tropical Australian microalgae for possible use in mariculture. Aquaculture 170, 147-159.
- Roessler, P.G., 1990. Environmental control of glycerolipid metabolism in microalgae: commercial implications and future research directions. J. Phycol 26, 393-399.
- 15 Ruiz-Lopez N., Haslam R.P., Venegas-Caleron M., Li T., Bauer J., Napier J.A., 2012. Enhancing the accumulation of omega-3 long chain polyunsaturated fatty acids in transgenic *Arabidopsis thaliana* via iterative metabolic engineering and genetic crossing. Transgenic Res 18.
- Sayanova, O., Smith, M.A., Lapinskas, P., Stobart, A.K., Dobson, G., Christie, W.W., Shewry, P.R., Napier, J.A., 1997. Expression of a borage desaturase cDNA containing an N-terminal cytochrome  $b_5$  domain results in the accumulation of high levels of  $\Delta^6$ -desaturated fatty acids in transgenic tobacco. Proc. Natl. Acad. Sci. USA 94, 4211-6.
- 20 Sayanova, O., Beaudoin, F., Michaelson, L., Shewry, P., Napier, J.A., 2003. Identification of *Primula* fatty acid  $\Delta^6$ -desaturases with  $n-3$  substrate preferences. FEBS Lett 542, 100-104.
- 25 Sayanova O., Ruiz-Lopez N., Haslam R.P., Napier J.A., 2012. The role of  $\Delta^6$ -desaturase acyl-carrier specificity in the efficient synthesis of long-chain polyunsaturated fatty acids in transgenic plants. Plant Biotechnology Journal 10, 195-206.
- 30 Seto, A., Wang, H.L., Hesseitine C.W., 1984. Culture conditions affect eicosapentaenoic acid content of *Chlorella minutissima*. J. Am. Oil Chem. Soc 61, 892-894.
- Siaut, M., Heijde, M., Mangogna, M., Montsant, A., Coesel, S., Allen, A., Manfredonia, A., Falciatore, A., Bowler, C., 2007. Molecular toolbox for studying diatombiology in *Phaeodactylum tricornutum*. Gene 406, 23-35.
- 35

- Springer, M., Franke, H., Putz, O., 1994. Increase of the content of polyunsaturated fatty acids in *Porphyridium cruentum* by low-temperature stress and acetate supply. J. Plant Physiology 143, 534-537.
- Stinson, E.E., Kwoczak, R., Kurantz, M., 1991. Effect of culture conditions on production of eicosapentaenoic acid by *Pythium irregular.* Ind. Microbiol 8, 171-178.
- 5 Sukenik A., 1991, Ecophysiological considerations in the optimization of eicosapentaenoic acid production by *Nannochloropsis* sp. (Eustigmatophyceae) Bioresour. Technol 35, 263-269.
- Tan, C.K., Johns, M.R., 1996. Screening of diatoms for heterotrophic eicosapentaenoic acid production. J. Appl. Phycol 8, 59-64.
- 10 Tonon T., Harvey D., Tony R. Larson T.R, Graham I.A., 2002. Long chain polyunsaturated fatty acid production and partitioning to triacylglycerols in four microalgae. Phytochemistry 61, 15-24.
- Venegas-Caleron M., Sayanova O., Napier J.A., 2010. An alternative to fish oils: metabolic engineering of oil-seed crops to produce omega-3 long chain polyunsaturated fatty acids. Prog Lipid Res 49, 108-119.
- 15 Voigt, R.G., Jensen, C.L, Fraley, J.K., Rozelle, J.C., Brown, F.R., Heird, W.C., 2000. Relationship between omega-3 long-chain polyunsaturated fatty acid status during early infancy and neurodevelopmental status at 1 year of age. J Hum Nutr Diet 15, 111-120.
- 20 Wagner, M., Hoppe, K., Czabany, T., Heilmann, M., Daum, G., Feussner, I., Fulda, M., 2010. Identification and characterization of an acyl-CoA:diacylglycerol acyltransferase 2 (DGAT2) gene from the microalga *O. tauri*. Plant Physiology and Biochemistry 48, 407-416.
- 25 Welti, R., Li, W., Li, M., Sang, Y., Biesiada, H, Zhou, H.E., Rajashekar, C.B., Williams, T.D., Wang, X., 2002. Profiling membrane lipids in plant stress responses. Role of phospholipase D alpha in freezing-induced lipid changes in Arabidopsis. J Biol Chem. 30, 277, 35, 31994.
- Wen, Z.Y., Chen, F., 2001. Optimization of nitrogen sources for heterotrophic production of eicosapentaenoic acid by the diatom *Nitzschia laevis*. Enzyme Microb. Technol 29, 341-347.
- 30 Xiao, S., Gao, W., Chen, Q.F., Chan, S.W., Zheng, S.X., Ma, J., Wang, M., Welti, R., Chye, M.L., 2010. Overexpression of Arabidopsis acyl-CoA binding protein ACBP3 promotes starvation-induced and age-dependent leaf senescence. Plant Cell 22, 5, 1463-82.
- 35

Yongmanitchai, W., Ward, O.P., 1989. Omega-3 fatty acids: alternative sources of production. *Process Biochem* 24, 117-125.

Yongmanitchai, W., Ward, O., 1991. Growth and omega-3 fatty acid production by the *Phaeodactylum tricornutum* under different culture conditions. *Applied and Environmental Microbiology* 419-425.

Yongmanitchai, W., Ward, O.P., 1993. Positional distribution of fatty acids, and molecular species of polar lipids, in the diatom *Phaeodactylum tricornutum* *J Gen Microbiol* 139,465-472.

Yu, E.T., Zendejas, F.J., Lane P.D., Gaucher, S., Simmons B.A., Lane, T.W., 2009. Triacylglycerol accumulation and profiling in the model diatoms *Thalassiosira pseudonana* and *Phaeodactulum tricornutum* (Baccilariophyceae) during starvation. *J Appl Phycol* 21, 669-681 .

Zaslavskaja, LA., Lippmeier, J.C., Kroth, P.G., Grossman, A.R., Apt, K.E., 2000. Transformation of the diatom *Phaeodactylum tricornutum* (Bacillariophyceae) with a variety of selectable marker and reporter genes. *J. Phycol* 36, 379-986.

### Sequence listing

Nucleic acids analogous to cDNA are shown.

20 SEQ ID No 1 Nucleic acid sequence OtElo5

```

atgagcgcctccgggtgcgctgctgcccgcgatcgcgctccgccggtacgcgtacgcgacg
tacgcctacgcctttgagtggtcgcacgcgaatggcatcgacaacgtcgacgcgcgcgag
tggatcgggtgcgctgtcgttgaggctcccggcgatcgcgacgcgacgatgtacctgttctc
tgccctggtcggaccgaggttgatggcgaagcgcgagggcgttcgacccgaaggggttcag
25 ctggcgtacaatgcgtatcagacggcgttcaacgtcgtcgtgctcgggatgttcgcgcga
gagatctcggggctggggcagcccgtgtgggggtcaaccatgccgtggagcgatagaaaa
tcgtttaagatcctcctcgggggtgtggttgactacaacaacaatatttgagctattg
gacactgtgttcattggttgccgcgcaagaagacgaagcagttgagcttcttgacgcttat
catcacgccctggtgatctgggcgtggtggttggtgtgtcacttgatggccacgaacgat
30 tgtatcgatgcctacttcggcgcggcgtgcaactcgttcattcacatcgtgatgtactcg
tattatctcatgtcggcgcctcggcattcgatgcccggtggaagcgatacatcaccaggct
caaatgctccaattcgtcattgtcttcgcgcacgcctgttcgtgctgcgtcagaagcac
tgcccggtcacccttccttgggcgcaaattggtcgtcatgacgaacatgctcgtgctcttc
gggaacttctacctcaaggcgtactcgaacaagtcgcgcggcgacggcgcgagttccgtg
35 aaaccagccgagaccacgcgcgcgccagcgtgacgcacgcgcgatctcgaaaaattgac
taa

```

SEQ ID No 2 Amino acid sequence OtElo5

MSASGALLPAIASAAYAYATYAYAFEWSHANGIDNVDAREWIGALSLRLPAIATT  
 MYLLFCLVGPRLMAKREAFDPKGFMLAYNAYQTAFNVVVLGMFAREISGLGQPVW  
 GSTMPWSDRKSFKILLGVWLHYNNKYLELLD TVFMVARKKTKQLSFLHVYHHALL  
 IWAWWL VCHLMATNDCIDAYFGAACNSFIHIVMYSYLLMSALGIRCPWKRYITQA  
 5 QMLQFVIVFAHAVFVLRQKHCPVTL PWAQMFVMTNMLVLFGNFYLKAYSNKSREGD  
 GAS SVKPAETT RAPS VRRTRSRKI D\*

SEQ ID No 3 OtD6 nucleic acid sequence

10 atgtgCGTGGAGACGGAAAATAACGATGGGATCCCCACGGTGGAGATCGCGTTCGACGGT  
 GAGCGGAGCGGGCGGAGGCAAACGTGAAGCTGTCCGCGGAGAAGATGGAGCCGGCGGC  
 CTGGCGAAGACGTTCCGCGAGGCGGTACGTCGTGATCGAGGGGGTGGAGTACGATGTGACG  
 GATTTTAAGCACCCGGGAGGAACGGTTATTTTCTATGCGTTGTCAAACACCGGGGCGGAC  
 GCGACGGAAGCGTTC AAGGAGTT TEAT CAT CGGTCGAGAAAGGCGAGGAAAGCCTTGGCG  
 15 GCGCTCCCGTCTCGACCGGCCAAGACGGCCAAGGTGGACGACGCGGAGATGCTCCAAGAT  
 TTCGCCAAGTGGCGGAAAGAATTGGAGAGAGATGGATTCTTCAAGCCCTCTCCGGCGCAC  
 GTGGCGTATCGCTTCGCCGAGCTCGCGGCGATGTACGCTCTCGGGACGTACCTGATGTAC  
 GCTCGATACGTGCTCTCCTCGGTGCTCGTGTACGCTTGCTTTTTCGGCGCCCGATGCGGT  
 TGGGTGCAGCACGAGGGCGGACACAGCTCGCTGACGGGCAACATTTGGTGGGACAAGCGC  
 20 ATCCAGGCCTTACAGCCGGTTCGGTCTCGCCGGTAGCGGCGACATGTGGAACCTCGATG  
 CACAACAAGCATCACGCGACGCCTCAAAAGGTTCTGTCACGACATGGATCTGGACACCACC  
 CCCGCGGTGGCGTCTCTCAACACCGCGGTGGAAGACAATCGTCCCCTGGCTTTAGCAAG  
 TACTGGTTGCGCCTTCAGGCGTGGACCTTCATCCCCGTGACGTCGGGCTTGGTGTCTCTT  
 TCTGGATGTTTTCTCCACCCTCCAAGGCTTTGAAGGGTGGCAAGTACGAAGAGTTG  
 25 GTGTGGATGCTCGCCGCGCACGTCATCCGACGTTGGACGATCAAGGCGGTGACCGGATTC  
 ACCGCGATGCAGTCTACGGCTTATTTTGGCGACGAGCTGGGTGAGCGGCTGCTATCTG  
 TTTGCACACTTCTCCACGTCGCACACGCACCTGGATGTGGTGCCCGCGGACGAGCATCTC  
 TCCTGGGTTCGATACGCCGTGATCACACGATCGACATCGATCCGAGTCAAGGTTGGGTG  
 AACTGGTTGATGGGCTACCTCAACTGCCAAGTCATCCACCACCTCTTCCGAGCATGCCG  
 30 CAGTTCGCCAGCCCGAGGTATCTCGCCGCTTCGTGCTCTTTCGAAAAAGTGGAACTC  
 AACTACAAGGTCATGACCTACGCCGGTGCCTGGAAGGCAACGCTCGGAAACCTCGACAAC  
 GTGGGTAAGCACTACTACGTGCACGGCCAACACTCCGGAAAGACGGCGTAA

SEQ ID No 4 OtD6 amino acid sequence

35 MCVET ENNDGI PTVE IAFDGE RERAE ANVKLSA EKME PAALAKT FARRYVVI EGVE YDVT  
 DFKHPGGTVI FYALSNTGADATEAFKEFHRSRKARKALAALPSRPAKTAKVDDAEMLQD  
 FAKWRKELERDGF FKPSPAHVAYRFAELAAMYALGTYL MYARYVVS SVLVYACFFGARCG  
 WVQHEGGHSSLTGNIIWWDKRIQAF TAGFGLAGSGDMWNSMHNKHHATPQKVRHMDLDTT  
 40 PAVAFFNTAVEDNRPRGFSKYWLRLQAWTFIPVTSGLVLLFWMFFLHPSKALKGGKYEEL  
 VWMLAAHVIRTWTIKAVTGFTAMQSYGLFLATSWVSGCYLFAHFSTSHTHLDVVPADDEHL  
 SWVRYAVDHTIDIDPSQGWVNWLMGYLNCQVIHHLFP SMPQFRQPEVSRRFVAF AKKWNL  
 NYKVM T YAGAWKAT LGNL DNVGKHYYVHGQHS GKTA\*

SEQ ID No 5 OtD6Pt nucleic acid sequence optimised codon

5 ggtaccaagcttgatatacaccaaaatggtggtcgaaacggaaaacaacgatggaatccccacgg  
 tcgaaattgcctttgatggagaacgcgaacgcgcgccaagccaacgtcaagctctccgcccga  
 gatggaaccgcccgccttgccaagacctcgcccgtcgctacgtcgtcattgaagggtcgaa  
 tacgatgtcaccgacttcaagcaccgggaggtacgggtcatcttttacgccctctccaacaccg  
 gagccgacgccacggaagccttcaaggaattcaccaccgttcccgcaaggcccgtgaaggcct  
 cgccgccttgccctcgcgcccggccaagaccgccaagggtcgacgatgccgaaatgcttcaggat  
 10 ttcgccaagtggcgtaaggaactcgaacgcgacggcttctttaagccctccccggcccacgtcg  
 cctaccgttttgccgaactcgccgccatgtacgcccttggaaacctacctcatgtacgcccgtta  
 cgtcgtctcctcgggtcttggctacgcctgcttctttgggtgcccgctgtggatgggtccagcac  
 gaaggcggacactcctcgctcaccggaacatattgggtgggataagcgtatccaagccttcacgg  
 ccggatttggtttggccggctccggagacatgtggaactcgatgcacaacaagcaccacgccac  
 cccccagaaggtccgctcacgacatggatctcgacaccacgccggccgtcgcccttctttaacacc  
 15 gccgtcgaagataaccgtccccgcggattctccaagtaactggcttcgtctccaagcctggacct  
 tcatccccgtcacgtccggtttggctcctcttgttttggatgttctttcttcaccgcgtgaaggc  
 cctcaaggggtggcaagtaacgaagaattgggtctggatgcttggccgccacgtcattcgtacctgg  
 acgatcaaggccgctcaccggtttcacggccatgcagtcctacggcttggtttcttgccacctcct  
 20 ggggtctcgggttgcctacctcttcgccacttttccacctcgcacacgcacttggatgtcgtccc  
 cgccgacgaacacctttcctgggtccgctacgcgctcgaccacaccattgacattgaccgcgtcg  
 cagggatgggtcaactggctcatgggttacttgaactgtcaagtcatccaccacctcttcccct  
 ccatgccgcagtttcgtcaaccgaagtctcgcgctcgcttctcgtcgcctttgccagaagtggaa  
 cttgaactacaaggtcatgacctacgccggagcctggaaggccacgcttggaaaccttgataac  
 25 gtcggaaagcactactacgtccacggccagcactcgggaaagaccgcctaagagctcggtagcc  
 tcgag

SEQ ID No 6 OtD6 amino acid sequence optimised codon

30 MCVET ENNDGI PTVE IAFDGE RERAE ANVKLSAEKME PAALAKT FARRYVVI EGVE YDVT  
 DFKHPGGTVIFYALSNTGADATEAFKEFHHRSRKARKALAALPSRPAKTAKVDDAEMLQD  
 FAKWRKELERDGF FKPSPAHVAYRFAELAAMYALGTYLMYARYVVSSVLVYACFFGARGC  
 WVQHEGGHSSLTGNIWWDKRIQAF TAGFGLAGSGDMWNSMHNKHHATPQKVRHMDLDTT  
 PAVAFFNTAVEDNRPRGFSKYWLRLQAWTFIPVTSGLVLLFWMFFLHPSKALKGGKYEEL  
 VWMLAAHVIRTWTIKAVTGFTAMQSYGLFLATSWVSGCYLFAHFSTSHTHLDVVPAD EHL  
 35 SWVRYAVDHTIDIDPSQGWVNWLMGYLNCQVIHHLFSPMPQFRQPEVSRRFVAFKKNL  
 NYKVMTYAGAWKATLGNLDNVGHYVHGQHSKTA

SEQ ID No 7 A6-desaturase nucleic acid from *Ostreococcus RCC809*

40 atgctcgcgtcgaaacggaggacgacaacgttccgacgggtcaccgctcggactgtcggaggag  
 agcgacgggatgaagggggcgagaaacccccggggcgcgggcggtggaaatcgacgctcgag  
 ccgcacgcgggtggccaagtcttcgatcgacgggtgggtcaaggttgacggcgtcgagtac  
 gacgtcacggat ttttaagcatccgggtggatctgtgattttattacatgctgtcgaacacc

ggagcggacgcgacggaggcggttcaaagagtt teat tat cggtcgaaaaaggcgagaaag  
 gcgttggcggcggttgccgcagcgcgagccggaggacgcgtcgccagtggaagacgcgaat  
 atgttgaaaggatttcgcgaaatggcgcaaagatttgagcgcgaggggtttctttaaacgcg  
 tcgccggcgcacgtggcggtacagattcgcggaactcgcggccatggttcgcgctcgggacg  
 5 gcgttgatgtacgctcgatggcacgccacctcagtcttcgtcaccgcgctgcttttcggc  
 gcgcgggtgcggttgggtgcaacacgaggggtggtcacagctcgctgacggggagcatttgg  
 tgggacaagcgaatccaagcgttcaccgcgggttcggattagcatcgagcggcgacatg  
 tggaacctcatgcacaacaagcaccacgccactccgcaaaagggtgcgacacgacatggac  
 ctcgacaccacgcggcggtggccttcttcaacactgcggtcgaggaaaaccgtccgcgc  
 10 aagttcagtaagttatggttgccgctgcagggcgtggacggttcgtcccggtcacctctggt  
 ttggtggtgctcgctggatgtacctcttgcatccgagacacattgctcgccgtaaaaac  
 tacgaagaggctgcgtggatcgctcgccgcgcacgtcatccgcacgtcggtcatcaaagcc  
 gtgaccggttactcctggatcacgtgctacggtttgttcttgtccaccatgtgggtgagc  
 ggctgctacctctttgcgacttctccacgtctcacacgcacctcgacgtcggtccgagc  
 15 gataagcatctctcttgggtgcgatacgccgctcgaccacaccatcgacatcgaccgagc  
 aagagcgtcgtcaactgggtgatgggttacctgaactgccaggtcatccatcacttgttt  
 ccggacatgcctcagttccgctcagcccgaagtctctcgcgcgcttcgctctcttttgcgaaa  
 aagtggaacctcaattacaaggtcatgagctactacggcgcggtggaaggccaccttcggt  
 aacttgaacgaggtcggcaagcactattacatccaaggttctcaaatcacgaagaagacg  
 20 gtgtaa

SEQ ID No 8 A6-desaturase amino acid from *Ostreococcus RCC809*

MRVETEDDNVPTVTVGLSEESDGMKGARNPGARAWKSTLEPHAVAKSFDRRWVKVDGVEYDVTD  
 25 FKHPGGSVIYYMLSNTGADATEAFKEFHYSKARKALALPQREPEDASPVEDANMLKDFAKW  
 RKDLEREGFFKPSPAHVAYRFAELAAMFALGTALMYARWHATSVFVTACFFGARCGWVQHEGGH  
 SSSLTGSIWWDKRIQAFTAGFGLASSGDMWNLMHKHHATPQKVRHMDLDLTPAVAFFNTAVEE  
 NRPRKFSKLWLRVQAWTFVPVTSGLVLLAWMYLLHPRHIARRKNYEEAAWIVAHHVIRTSVIKA  
 VTGYSWITCYGLFLSTMWVSGCYLFAHFSTSHTHLDVVP SDKHLSWVRYAVDHTIDIDPSKSVV  
 30 NWLMGYLNCQVIHHLFPDMPQFRQPEVSRRFVSFAKKWNLNYKVMSYYGAWKATFGNLNEVGKH  
 YIIQGSQITKKTV

SEQ ID No 9 A6-desaturase (Ost809A6) nucleic acid from *Ostreococcus RCC809*  
 codon optimised for expression in *T.pseudonana*

atgcggtggaaccgaagacgataatgtgccaaactgttactgtgggattgtcagaggagtcg  
 atggaatgaaggagcaaggaaccccgagcacgtgcttggaaagtcgacgttgagccgcacgc  
 cgtggcaagtcattcgatcgtaggtgggttaaggttgacggagtcgaatacgacgtaactgat  
 ttcaagcatcccggaggatcagttatctactatgtcttctaacaccggagctgatgccactg  
 40 aggctttcaaggaatctcactatcgtagtaagaaggccaggaaggcacttgctgcctcccaca

acgtgagcctgaagacgcttcgccagtcgaggatgccaatatgctcaaggacttcgcaaagtgg  
 cgtaaggatttggagaggggaaggattctttaagccaagtcctgctcacgtggcctaccgtttcg  
 ccgaactcgcagctatgtttgctttgggaactgcccttatgtatgcacgttggcatgctacgtc  
 5 tgtcttcgtaacagcctgtttctttggagcaagggtgtggatgggtgcaacacgagggaggacat  
 tcttccttgaccggatccatctgggtgggataagcgtattcaggcattcactgctggatttggac  
 ttgccagttcgggagacatgtggaacctcatgcacaataagcaccatgcaacgccacaaaaagt  
 taggcatgatatggacctcgataccactcctgcagtggtttctttaacacagctgttgaggaa  
 aatcgtcctaggaagtctctaagttgtggcttcgtgtccaggcctggacctttgtgcccgta  
 10 ctcccgattggtactcttggcatggatgtaccttctccaccgcgcatatcgctcgtaggaa  
 gaactatgaggaagccgcatggattgtggctgcccatgttatcaggacctccgctcattaaggct  
 gtaacgggatacagttggatcacatgttatggactcttcttgcgactatgtgggtctcaggat  
 gctacctcttcgctcacttttcaacgtctcacacacatttggacgtggttccatctgataagca  
 cctttcctgggtgcggttacgccgttgatcataccatcgacattgatccttccaagagtgtcgta  
 15 aactggctcatgggatatttgaactgtcaggttatccaccatttgttccccgacatgccgcaat  
 ttcgtcagcccgaagtcagtcgtaggttcgtatcgtttgccaagaagtggaaccttaattaaa  
 ggtcatgtcttactatggagcctggaaggcaaccttcggaaatctcaacgaagtcggaagcac  
 tactacatccaaggaagtcaaatcacaagaagacggtttag

20 SEQ ID No 10 A6-desaturase amino acid from *Ostreococcus RCC809* codon optimised

MRVETEDDNVPTVTVGLSEESDGMKGARNPGARAWKSTLEPHAVAKSFDRRWVKVDGVEY  
 DVTDFKHPGGSVIYYMLSNTGADATEAFKEFHYSKKARKALALPQREPEDASPVEDAN  
 MLKDFAKWRKDLEREGFFKPSPAHVAYRFAELAAMFALGTALMYARWHATSVFVTACFFG  
 ARCGWVQHEGGHSSLTGSIWWDKRIQAFTAGFGLASSGDMWNLMHNKHHATPQKVRHDMD  
 25 LDTPAVAAFFNTAVEENRPRKFSKLWLRVQAWTFVPVTSGLVLLAWMYLLHPRHIARRKN  
 YEEAAWIVA AHVIRTSVIKAVTGYSWITCYGLFLSTMWVSGCYLFAHFSTSHTHLDVVP  
 DKHLSWVRYAVDHTIDIDPSKSVNWLGMYLNCQVIHHLFPDMPQFRQPEVSRRFVSFAK  
 KWNLNYKVMSY YGAWKAT FGNLNE VGKHYY IQGSQ ITKKTV

30 SEQ No. 11 A4-desaturase from *E. huxleyi* (EhD4) codon-optimized for expression in Arabidopsis

atgggagggcgccggcgcgagcgaggctgaacggcccaagtggaccacgatccacgggcggcacg  
 tcgatgtgtcaaagttccgccaccgggtgggaacatcatcgagctcttctatggcatggactc  
 35 gacgagcgcgttcgagcagttccacggccaccacaagggcgctggaagatgctcaaggcgtg  
 ccgaccaaggaggtcgaccccgccgacgtgccgcagcagccgcaggagcacgttgccgagatga  
 cgcggtgatgacgtcgtggcgcgagcgcggtctcttaagccgccccgctgcctcgggcat  
 ctacggctcgcgctcgtcgtgccatcgtcgcgtgcacgcctgcgcgcgcacgcgcgggtg  
 ctgagcgggatcgggct egg cage tgctgggcgagtcgggttccctgcagcacatgggccccg  
 40 accgcgagtggggggtgcggtactccttcctcctgcagcacttcttcgagggcctcctcaaggg  
 cgggtccgctcgtggtggcgcaaccgccacaacaagcatcacgcaagactaacgtgctcggc  
 gaggacggcgacctgcgacgactcccttcttcgctgggacccgacgctcgccaagaaggttc  
 cagactggtcgtcaagacgcaggccttcacctcctccccgcctcggagcgtacgtctttgt  
 ctttgcttcacgatccgcaagtatgccgtcgtcaagaagctctggcacgagctcgactcatg  
 45 atcgcgcactacgcgatgttctactacgcgctgcagctcgccggtgcgtcgtcggcagcggcc



tcgccttttactgcaccggctacgcctggcaaggcatctacctcggcttcttcttcggcctgtc  
 ccacttcgcggtcgagcgagtcccctccaccgccacctggctcgagtcgtccatgatcggcacc  
 gtgcactggggaggctcctccgccttttgcggctacgtctccggcttccctcaacatccagatcg  
 agcaccacatggcgccgcagatgccgatggagaacctgcgccagatccgcgcccactgcaaggc  
 5 gagcgcgagaagctcgggcttccctatcgcgagctctccttcgccggcgcggtcaagctgatg  
 atggctcggcctctggcgcacggggaggagcagctgcagctgcgctccgacaggcgcaagtact  
 cgcgcaccaggcctacatggcgccgcctcggcggtggtggagaacctcaaggcggactag

10 SEQ No. 12 A4-desaturases from *E. huxleyi* codon-optimized for expression in  
 Arabidopsis

MNGNLPASTAQLKSTSKPQQQHEHRTISKSELAQHNTPKSAWCAVHSTPATDPSHSNNKQHAH  
 LVLDITDFASRHPGGDLILLASGKDASVLFETYHPRGVPTSLIQKLQIGVMEEEAFRDSFYSWT  
 DSDFYTVLKRVRVERLEERGLDRRGSKEIWIKALFLLVGFWYCLYKMYTTSIDIDQYGIAYSI  
 15 GMGTFAAFIGTICIQHDGNHGAFQAQNKLLNKLAWTLDMIGASAFWELQHMLGHPYTNVLDGV  
 EEERKERGEDVALEEKDQESDPDVFSSPFLMRMHPHHTTSWYHKYQHLYAPPLFALMTLAKVFQ  
 QDFEVATSGRLYHIDANVRYGSVWNVMRFWAMKVI TMGYMMGLPIYFHGVLRGVGLFVIGHLAC  
 GELLATMFIVNHVIEGVSYGTKDLVGGASHGDEKKIVKPTTVLGDTPMEKTRREEALKSNSNNNK  
 20 KKGEKNSVPSVPFNDWAAVQCQTSVNWSPGSWFNHFSGGLSHQIEHHLFPSICHTNYCHIQDV  
 VESTCAEYGVYPYQSESNLFFVAYGKMISHLKFGLKAKCE\*

SEQ ID No. 13 D4-desaturase from *Thalassiosira pseudonana* nucleic acid

atgggcaacggcaacctcccagcatccaccgcacagctcaagtccacctcgaagccccagcagc  
 25 aacatgagcatcgcaccatctccaagtcgagctcgcccaacacaacacgcccacatcagcatg  
 gtgtgccgtccactccactcccgccaccgacccatccactccaacaacaacaacacgcacac  
 ctagtctcgacattaccgactttgctcccggccatccagggggagacctcatcctcctcgctt  
 ccggcaaagacgcctcgggtgctgtttgaaacataccatccacgtggagttccgacgtctctcat  
 tcaaaagctgcagattggagtgatggaggaggagcgtttcgggattcgttttacagttggact  
 30 gattctgactttttatactgtgttgaaagaggaggttgtggagcgggtggaggagagggggttg  
 acaggaggggatcgaaagagatttgatcaaggctttgttcttgttggttgattttggtactg  
 tttgtacaagatgtatactacgtcgatattgatcagtagcgtattgccattgcctattctatt  
 ggaatgggaacctttgcgcatcctcggcacgtgtattcaacacgatggaaatcacggtgcat  
 tcgctcagaacaagtactcaacaagttggctgggtggacggttgatgatgggtgcgagtg  
 35 gtttacgtgggagcttcagcacatgctggggcatcatccatatacgaatgtgttgatgggggtg  
 gaggaggagaggaaggagaggggggaggatgttgctttggaagaaaaggatcaggaatcagatc  
 cagacgtattctcctccttccctctcatgagaatgcatccccaccatacaacctcatggtatca  
 taaataccaacacctctacgctccaccctctttgcattgatgacacttgccaaagtattccaa  
 caggattttgaagttgccacatccggacgattatatcatattgatgccaatgtacggttatgggt  
 40 cggatggaatgtcatgaggttttgggctatgaaggtcattacgatgggatatatgatgggatt  
 accaatctactttcatggagtactgaggggagttggattgtttgttattgggcatttggcggtg  
 ggagagttggtggcgacgatgtttattgtgaatcacgtcattgaggggtgtgagttatggaacga

aggatttggttggtggtgctgagtcacggagatgagaagaagattgtcaagccaacgactgtatt  
 gggagatacaccaatggaaaagactcgcgaggaggcattgaaaagcaacagcaataacaacaag  
 aagaagggagagaagaactcgggtaccatccggttccattcaacgactgggcagcagtcctaatgcc  
 agacctccgtgaattggtctccaggctcatggttctggaatcacttttctgggggactctctca  
 5 ttagattgagcatcacttggtccccagcatttgtcatacaaaactactgtcatatccaggatggt  
 gtggagagtagctgtgctgagtagcggagttccgtagcagagtgagagtaatttggttggtgctt  
 atggaaagatgattagtcatttgaagttttgggtaagccaagtgtgagtag

10 SEQ ID No. 14 D4-desaturase from *Thalassiosira pseudonana* amino acid acid

MGGAGASEAERPkwTtIHGRHVDVSKFRHPGGNI IELFYGMDSTSAFEQFHGHHKGAWKM  
 LKALPTKEVDPADVPQQPQEHVAEMTRLMTSWRERGLFKPRPVASGIYGLAVVAIVACI  
 ACAPHAPVLSGIGLGSWAQCGLQHMGHREWGVRYSFLLQHFFEGLLKGSASWWRNR  
 HNKHHAKTNVLGEDGLRTPFFAWDPTLAKKVPDWSLKTQAFT FLPALGAYVFVFAFTI  
 15 RKYAVVKKLWHELALMIAHYAMFYALQLAGASLGSGLAFYCTGYAWQGIYLGFFFGLSH  
 FAVERVPSTATWLESSMIGTVDWGGSSAFCGYVSGFLNIQIEHHMAPQMPMENLRQIRAD  
 CKASAEKLGLPYRELSFAGAVKLMVGLWRTGRDELQLRSDRRKYSRTQAYMAAASAVVE  
 NLKAD\*

20 SEQ ID No. 15 A4-desaturase *Ostreococcus* RCC809 nucleic acid

atgccgacgactcgatcgcgcgcgcgctgacgacgccccctcgcgagacgccgacgagagcga  
 acaccgtcgccgcgctcgatcccgagcgcaagtacacgcgcatcgcggcgtcgtgtacgacgt  
 cacggatttcgccagccgctcatccgggtggcgcgcaattgttatcgctgtgctggtgggagagac  
 25 gccaccatcctggtggagagtcacaccttcgtccggaggtggtgcaaaagtacctgaagacgc  
 ttcctcggtggtagggcgcgccggggcggttcgggcccaggagacggttcggaaaccgctcga  
 ctcgatttgtaccgaaagattcagggcgcgcttcgtaaagagatcgtcgaaccggtgaagatg  
 acgcgcgacgcgagccgcacgggagggctggtgctgttggaacgccccgggtggtgtggctt  
 tcttcgcttcgcttgggagtcatttgaagacgcccagcgggtggcgacgggggtgcctgtggg  
 30 gctcgccgggtactggagcggcaccggattgcaacacacggcgaaccacgggtggattggcgaag  
 agtgggttttggaaatcagttttggggatggctcgggaacgacgctcgccatcgggaagagctcgg  
 tggagtgagatatcatcacatgggtgagccaccactcgtattgcaacgacgcggacctcgatca  
 agacgtgtacaccgcgctgccgcttcttcggttggaccgctcccaggagttgaagtgggtccac  
 cgctaccaagcgttctacgccccgctgatgtggccgatggtgtggctcgccgcgacggttggcg  
 35 acgcgcaaaatatatttagtgataaggcgtctccggcgctcgagtacaagggcctcatgaagct  
 cgaagtcgctgtacgcttctcggaaagttttgcatttttagcttgttgctcggcgctaccggcc  
 tacttgcacgggtttgcaacgccatcgtgccgttcatcgcgtacggtgcttccggttcggttcg

tccctgtgctgggtttttcatcgtcagtcacaacttggaggcgttgacccaatcaatctgagcaa  
 atccacgaagaatgactggggcgcggtggcaaatcgaaacttccgcgtcctggggcaacggcttc  
 tggagctttttctccggcggttgaatttgcaaatcgagcaccacttgttcccggttgcgcgc  
 acaacttgtacccgaagatggttcccatcatcaaggaagagtgcgaaaaggctggcgtcacgta  
 5 caccggttacggtgggtactttgggtctccttcccatcactcgggacatgttcgcgtacttgta  
 aaaatgggcccgaagcaaaaagtcggcgtaa

SEQ ID No. 16 A4-desaturase *Ostreococcus* RCC809 amino acid

10 MPTTRSRRARVTPPRETPTRANTVAALDPERKYTRIRGVVYDVTFASRHPGGAQLLSLCVGRD  
 ATILVESHHLRPEVVQKYLKTLPVVEGAAGAFGPEETFPKPLDSDLYRKIQGRVRKEIVEPLKM  
 TRGREPHGRGWCVLDAGVVLAFFAFALGVYWKTPTVATGCLLGLAGYWSGTGLQHTANHGGLAK  
 SGFWNQFWGWLGNDAIGKSSVEWRYHHMVSHHSYCNDADLDQDVYTALPLLRLDPSQELKWFH  
 RYQAFYAPLMWPMLWLAQFGDAQNILVDKASPGVEYKGLMKLEVALYVLGKFLHFSLLLGVPA  
 15 YLHGFANAIVPFIAYGAFGSFVLCWFFIVSHNLEALTPINLSKSTKNDWGAWQIETSASWNGNF  
 WSFFSGGLNLQIEHHLFPGCAHNLYPKMVPIIKEECEKAGVTYTYGGYFGLLPITRDMFAYLY  
 KMGRQSKKSA\*

SEQ ID No. 17 A4-desaturase *Ostreococcus* RCC809 nucleic acid codon optimised  
 acid for expression in Pt

20 ggatccggtaccaagcttgatatacctcaaaatgccaaactactcgttctcgtgctcgtgttacta  
 ctccacctcgtgaaactcctactcgtgctaatactggtgctgcttagatccagaacgtaaata  
 tacacgtattcgaggtggtgtatgatggtactgattttgctagtcgacatccaggtggtgca  
 25 caattattatctttatgtggtggtcgtgatgctacaattttagtagaatcacatcatttacgac  
 cagaagttgtacaaaaatatttaaaaacattacctggtgtagaaggtgctgctgggtgcatttg  
 tccagaagaaacttttccaaaacttttagatagtgatttatatcgtaaaattcaaggtcgtggt  
 cgaaaagaaattgtagaaccattaaaaatgacacgtggctcgagaacctcatggctcgtggtggt  
 gtgtttagatgctggtggtgatttagctttctttgcttttgcattaggtggttattggaaaac  
 30 accaactgtagctactggtggttatttaggttttagcagggttattggctcgttacagggtttacaa  
 catactgctaatacatgggtggttttagcaaaatcagggtttggaatcaattttgggggtggttagg  
 aatgatggtgctattggtaaatcaagtgtagaatggcgttatcatcatatgggtttcacatcat  
 agttattgtaatgatgctgatttagatcaagatggttatacagcattaccattattacgttttag  
 atccttcacaagaattaaaatgggtttcatcgttatcaagcattttatgcacctttaatgtggcc  
 35 tatggtatgggttagctgcacaatttggtgatgctcaaaatatttttagttgataaagcaagtc  
 ggtgtagaatataaagggtttaatgaaattagaagttgctttatatgtattaggaaaattttta

catttttctttattatttaggtggttcctgcatatttacatgggttttgctaatgcaattgtacat  
 ttattgcttatggtgcatattggttcatttggtttatggtgggtttttcattgtaagtcataattt  
 agaagcattaacaccaattaatctctaaatcaactaaaaatgattgggggtgcttggcaaatt  
 gaaactagtgcattcttggggtaatgggttttggtcatttttctcaggtgggtttaaatttcaaaa  
 5 ttgaacatcatttatttctggttggtgctcataatctatccaaaaatggttcctattattaa  
 agaagaatgtgaaaaagcaggtggttacatatactgggttatgggtggttattttgggtttattacca  
 attactcgtgatatggttgcttattatataaaaatgggtcgtcaatctaaaaaatctgcttaag  
 agctcggtagcctcaggtctaga

10 SEQ ID No. 18 A4-desaturase *Ostreococcus* RCC809 amino acid codon optimised  
 acid for expression in Pt

MPTRSRARVTTPPRETPTRANTVAALDPERKYTRIRGVVYDVTDFASRHPGGAQLLSLCVGRD  
 ATILVESHHLRPEVVQKYLKTLPVVEGAAGAFGPEETFPKPLDSDLYRKIQGRVRKEIVEPLKM  
 15 TRGREPHGRGWCVLDAVVLAFFAFALGVYKTPTVATGCLLGLAGYWSGTGLQHTANHGGLAK  
 SGFWNQFWGLGNDVAIGKSSVEWRYHHMVSHHSYCNDADLDQDVYTALPLLRLDPSQELKWFH  
 RYQAFYAPLMWMLWLAAQFGDAQNILVDKASPGVEYKGLMKLEVALYVLGKFLHFSLLLGVPA  
 YLHGFANAIVPFIAYGAFGSFVLCWFFIVSHNLEALTPINLSKSTKNDWGAWQIETSASWGNF  
 WSFSGGLNLQIEHHLFPGCAHNLYPKMVPIIKEECEKAGVTYTYGGY FGLLPITRDMFAYLY  
 20 KMGRQSKKSA\*

SEQ ID No. 19 Δ6-elongase from *Fragilariopsis cylindrus* nucleic acid

ccatggggtagcagatcaccaaaatggacgagtacaaagcaactcctgaatctgt  
 25 tggggatgctat cat ccaatgggcagatcctgaaagtcagttcaccgggtt caeca  
 agggatggttcttgacagatctcacatctgcgttttagtattgcacttgatatacgtc  
 ttatttgatcattggttctcaagtgatgaaagtcctacctgctattgatccgta  
 cccaatcaagttttttacaatgtatcacaattatgctgtgtgcttacatgacga  
 ttgaagcatgtctgtagcgtaccgtaacggatacactatcatgcatgtgtcgga  
 30 tacaatagagatgatccagcaattggaaatcttttatgggtattttatggtttcaa  
 agtttgggatttttgggataccatctttatcgttttggggaagaagtgagacaac  
 tttctttccttcacgtttaccatcataccaccatctttttggttctactggcttaac  
 gcgaatgtcttttatgatgggtgatatttatcttaccattgctctgaatgggtttcat  
 ccatactggtatgtacacatactactttatctgtatgcatactaaagacaagaaaa  
 35 ctggaaaatcgcttctatctggtggaaatcatctttgactttggtgcaattggtt  
 cagttcattaccatgatgtcacagggcttataccttatcatttttgggtgtgaatc

actttctatccgagtcactgcgacatacgttgtttacatattgtcactttttctttt  
 tgtttgcgcaattcttcggtgcatcttacatgcaacctaagaaatcgaagactgcc  
 taagagctcggtagcttaattaa

5 SEQ ID No. 20 A6-elongase from *Fragilariopsis cylindrus* amino acid

MDEYKATLESVGDAL IQWADPESQFTGFTKGWFLTDFTSAFSIALVYVLFVI IGSQVMKVL PAI  
 DPYPIKFFYNVSQIMLCAYMTIEACLLAYRNGYTIMPCVGYNRDDPAIGNLLWLFYVSKVWDFW  
 DTIFIVLGKKWRQLSFLHVYHHTTI FLFYWLNAVFDGDIYLTIALNGFIHTVMYTYFYFICMH  
 10 TKDKKTGKSLPIWKKSSLTLLQLFQFITMMSQGLYLII FGCELSLIRVTATYVVYILSLFFLFA  
 QFFVASYMQPKKSKTA

SEQ ID No. 21  $\Delta\delta$ -desurase from *Fragilariopsis cylindrus* nucleic acid

1 ATGGCACCCGACGCCGATCACAAGCTGAGACAGCGCCGTCTAAAAGGCGACGAAGTTTGT  
 15 61 ATCGATGGAATTATCTATGATATATCAT CCTTCGAGCAT CCGGGTGGT GACTACTCAAC  
 121 GTATTTGGTGGAAAC GATGCAACAATTCAGTACAAAAT GATTCACCCGTACCATACCACG  
 181 AAGCATTTAGAAAAAAT GAAGGTAGT TGGTAAAGTTCCAGAC TACTACTCAGAATACAAA  
 241 TGGGATACACCCTTCGAACGTGAAATGAAACGTGAGGTATTTAAATTGTACGACGTGGA  
 301 CAAGAATTTGGTACAAATGGATATTTTTCCGTGCCATTTCTGTATATTGCTATGTTTTTT  
 20 361 TATCTGCAATATTTATGGATGCAAGAATCTTCCTACACGTTAGCCATCGTATACGGGAT T  
 421 AGTATGGGATTGATTGGACTGAATGTCCAGCATGATGCGAACCACGGAGCTGCATCGAAA  
 481 AAAGTGTGGGTGAATGACCTCCTAGGATTGGGAGCAGACTTTATCGGAGGATCGAAATGG  
 541 TTGTGGATGGAAAAACATTGGACGCATCATGCTTTTACAAA C CATCGAGAAAAGGATCCA  
 601 GATGGGTTAGCAGCGGAACCTTTCCTATTGTTCAACGACTACGACTTGTTCGAGTTCCAAA  
 25 661 CGTGCTGGATATCATGCATACCAAGGAATTTATTTAGTCCTATTATTGTGTGGGTATTGG  
 721 CTTTCGGCAATTATTGATATACCTGTAATTTGGAATCTACAAGATCGTGGTGCCTTACG  
 781 GTAGGAATCCAGCTGGATAACGATTGGATTGCTAGTCGAAGAAAGTACGCGGTAGTCTT  
 841 CGAATCTTATACCTCTTTTGTAAACATCGTCGTTCTCTCTATAACAATTTCTCTGGACA  
 901 ACCGTGAGTCATATCAATGTAATGGGAATTTGTGGTAGCCTTACATTAGGACTACTTTTT  
 30 961 ACCTTGTCG CACAATTTT GAGAATGTAGATCGAGATCCTACCAATCTGAACTTAAATGAA  
 1021 ACAGAAGAACCCTGTTTGCTGGTTCAAATCTCAAGTAGAAACTTCTTCAACATACGGGGC  
 1081 ATGATATCCGGATGGTTAACCGGCGGATTAAACTTTTCAGGTTGAGCACCATTTATCCCG  
 1141 AGAATGTCTAGTGCCTGGTATCCATTTATTGCACCAAAAAGTTCGTGAAATTTGCAAAAAG  
 1201 CACGGAGTTCGTTACGTATACTATCCATGGTGTGCAAAAATATGTATTCGACGTTGAAG  
 35 1261 TACACCACGAGGTTGGTGTGCGGCTCACATTTGGAAGGATAATCCTTTTAAGGGTGAATG  
 1321 TAG

SEQ ID No. 22  $\Delta\delta$ -desurase from *Fragilariopsis cylindrus* amino acid

1 MAPDADHKLQRRLKGDEVCIDGIIYDISSFEHPGGDTINVFGGNDATIQYKMIHPYHTT  
 61 KHLEKMKWKGKVPDYSEYKWDTPFEREMKREVKIVRRGQEFGTNGYFFRAISYIAMFF  
 121 YLQYLWMQESSYTLAIVYGISMGLIQLNVQHDANHGAASKKVVWDLGLGADFIGGSKW  
 5 181 LWMEKHWTHHAFTHNREKDPDGLAAEPFLLFNDYDLSSSKRAGYHAYQGIYLVLLCCGYW  
 241 LSAIIDIPVIWNLQDRGALTVGIQLDNDWIASRRKYAVSLRILYLCNIWPLYNFSWT  
 301 TVSHINVMGICGSLTLGLLFTLSHNFENVDRDPTNLNLNETEEPVCWFKSQVETSSTYGG  
 361 MISGWL TGGLNFQVEHHLFPRMSSAWYFPIAPKVREICKKHGVRVYVYPWLLQNMYSTLK  
 421 YTHEVGVGSHWKDNPFGEM-

10

SEQ ID No. 23 *P. patens* PpHUPI L codon-optimised for expression in *Phaeodactylum tricornutum*

15 1  
 ATGGCAGGGGGGGTGTCTGTTACGGCGGGGAGATCAAGCACTACCCCGGCCGAACAACC  
 61  
 TTCTTTGTGATTATGGTCTGTATAGTGGCGGCATCCGGAGGTCTCATGTTCCGATACGAT  
 121  
 20 GTCGGAATTCAGGGGGTGTACGTCTATGGACGAATTTTTGGCGAAATTTTTCTCTGCG  
 181  
 GTGTTGGCGAAGAAGCGAGCAGAGGCAGCTTCGGAGAGCGCCTACTGCAAGTATGATGAC  
 241  
 CAGAAGCTGCAAGCCTTCACATCGTCGCTGTACATTTCCGCACTCGTGTGACATTCTTC  
 25 301  
 TCGTCGTACACCACCAGGCACTACGGCCGTAAATTTACCATGCTCATAGCTGGTTTCGCC  
 361  
 TTCTGCTTCGGCGTCATCTTCACCGCCGCTGCGCAAGAAATCATCATGCTAATCATAGGG  
 421  
 30 CGCGTCTCTGTTGGGGTGTGCGATTGCTAACCAGGCTGTTCCGTTGTACCTCTCC  
 481  
 GAAATGGCACCCCTCCAAGTGGCGAGGTGCGCTCAACATCCTCTTCCAATTGGCGGTGACC  
 541  
 ATTGGCATCCTGTTCCGAGTCTCGTGAACACTACGGCACAGAGAAGATGGCTCGCAACGGG  
 35 601  
 TGGCGTGTTCCTCGCCATCGCCGGCCTGCCTGCGATCTTCATCACCCCTCGGAGGATTA  
 661  
 CTCCTGCCAGACACACCGAATTCCTCGTGAACGCGGCAAGCACGAGAGCGCCCGCCAG  
 721  
 40 GTCCTACGCAGGATTGTTGGCGTCGACAACATTGAGGAAGAGTTCGACGACATCCTCAT  
 781  
 GCCAGTAACGAAGCCGCCTCCGTGAAGCACCCCTTCGCAATATCTTGAAACGCCGCAAC  
 841  
 CGCCCTCAGCTGGTCATCTCCATGGCTCTTCAGTTTTCCAGCAATTCCTGGAATTAAT  
 45 901  
 GCTATTATGTTTTACGCGCCTGTCTTGTTCAGACGCTGGGATTCCGGAGTTCGCTTCA  
 961  
 CTTTACTCTGCTGTCATCGTTGGAGCCGTGAATGTGCTGGCCACTTGCGTGCCTATCGCT  
 1021  
 50 GTTGTGGATCGATTCCGGTTCGACGATGGTTGCTCTTGAAGCTTGCATCCAAATGTTCTTA

1081  
 GCACAGACGGCGATTGCAATTATCCTGGCGGCGGGATTGAAGGGGACCGAGATGCCGGAG  
 1141  
 TATCTGGGATGGATCGCGGTGGTATTGATTTGCGTGTACGTGTCTTCTTTTCGCGTGGTCT  
 5 1201  
 TGGGGTCCACTTGGATGGTTGATTCCAAGTGAGATTTTCCCCTTGGAGACGCGTTCAGCA  
 1261  
 GGGCAAGCCATCACGGTGTGACCAACATGGTCTTCACCTTCCTCATCGCGCAAGTGTTTC  
 1321  
 10 CTGTCAATGTTGTGCGCGTTCAAGTGGGGCATCTTCTCTTCTTCGCCGCGTGGGTGGTG  
 1381  
 GTGATGTTCTTTTACGTACTTTTTAATCCCCGAGACGAAGGGCATCCCCATCGAGGAG  
 1441  
 ATGGATCTCGTGTGGACCAAGCACTGGTCTGGAAGCGCTACGTCCCCTACCCTGAGACT  
 15 1501  
 CTCGCTCACACCAGCGGCATCCCCATGGGAGATATGAAGGTCAGCAAGCTGGAGAATGGC  
 1561 TCCGCAAATGGCCACAACTGTAA

20 SEQ ID No. 24 Deduced polypeptide sequence of **PpHUP1L**

1 MAGGGVVTAGEIKHYPRRTTFFVIMVCIVAASGGLMFGYDVGISGGVTSMDDEF LAKFFPA  
 61  
 VLAKKRAEAAASESAYCKYDDQKLQAF TSSLYISALVSTFFSSYTTRHYGRKFTMLIAGFA  
 25 121  
 FCFGVIFTAAAQEIIMLIIGRVLLGWGVGFANQAVPLYLSEMAPSKWRGALNILFQLAVT  
 181  
 IGILFASLVNYGTEKMARNGWVSLAIAGLPAIFITLGGLLLPDTPNLSLVQRGKHESARQ  
 241  
 30 VLRRIRGVDNIEEFDDILIASNEAASVKHPFRNILKRRNRPQLVISMALQFFQOFTGIN  
 301  
 AIMFYAPVLFQTLGFGSSASLYSAVIVGAVNVLATCVAIAVVDRFGRRWLLEACIQMFL  
 361  
 AQTALAI ILAAGLKGTEMPEYLGWIAVVLICVYVSSFAWSWGPLGWLIPSEI FPLETRSA  
 35 421  
 GQAITVSTNMVFT FLIAQVFLSMLCAFKWGIFLFFAAWVVVMFLFTYFLIPETKGIPIEE  
 481 MDLVWTKHFWKRYVPYPETLAHTSGIPMGDMKVSKLENGSANGHKL-

40 SEQ ID No. 25 Homo sapiens HsGLUT1 **codon-optimised for expression in**  
*Phaeodactylum tricornutum*

1  
 ATGGAGCCCAGCAGCAAGAAGCTGACGGGTCGCCTCATGCTGGCTGTGGGAGGAGCAGTG  
 45 61  
 CTTGGCTCCCTGCAGTTTGGCTACAACACTGGAGTCATCAATGCCCCCAGAAGGTGATC  
 121  
 GAGGAGTTCTACAACCAGACATGGGTCCACCGCTATGGGGAGAGCATCCTGCCACCACG  
 181  
 50 CTCACCACGCTCTGGTCCCTCTCAGTGGCCATCTTTTCTGTTGGGGGCATGATTGGCTCC

241  
 TTCTCTGTGGGCCTTTTTCGTTAACCGCTTTGGCCGGCGGAATTCAATGCTGATGATGAAC  
 301  
 CTGCTGGCCTTCGTGTCCGCCGTGCTCATGGGCTTCTCGAAACTGGGCAAGTCCTTTGAG  
 5 361  
 ATGCTGATCCTGGGCCGCTTCATCATCGGTGTGTACTGCGGCCTGACCACAGGCTTCGTG  
 421  
 CCCATGTATGTGGGTGAAGTGTACCCACAGCCTTTTCGTGGGGCCCTGGGCACCCTGCAC  
 481  
 10 CAGCTGGGCATCGTCGTCCGCATCCTCATCGCCAGGTGTTTCGGCCTGGACTCCATCATG  
 541  
 GGCAACAAGGACCTGTGGCCCCTGCTGCTGAGCATCATCTTCATCCCGGCCCTGCTGCAG  
 601  
 TGCATCGTGCTGCCCTTCTGCCCCGAGAGTCCCCGCTTCTGCTCATCAACCGCAACGAG  
 15 661  
 GAGAACC GGCCAAGAGTGTGCTAAAGAAGCTGCGCGGGACAGCTGACGTGACCCATGAC  
 721  
 CTGCAGGAGATGAAGGAAGAGAGTCGGCAGATGATGCGGGAGAAGAAGGTCACCATCCTG  
 781  
 20 GAGCTGTTCCGCTCCCCCGCCTACCGCCAGCCATCCTCATCGCTGTGGTGCTGCAGCTG  
 841  
 TCCCAGCAGCTGTCTGGCATCAACGCTGTCTTCTATTACTCCACGAGCATCTTCGAGAAG  
 901  
 GCGGGGGTGCAGCAGCCTGTGTATGCCACCATTGGCTCCGGTATCGTCAACACGGCCTTC  
 25 961  
 ACTGTCGTGTCGCTGTTTGTGGTGGAGCGAGCAGGCCGGCGGACCCTGCACCTCATAGGC  
 1021  
 CTCGCTGGCATGGCGGGTTGTGCCATACTCATGACCATCGCGCTAGCACTGCTGGAGCAG  
 1081  
 30 CTACCCTGGATGTCCTATCTGAGCATCGTGGCCATCTTTGGCTTTGTGGCCTTCTTTGAA  
 1141  
 GTGGGTCTGGCCCCATCCCATGGTTCATCGTGGCTGAACTCTTCAGCCAGGGTCCACGT  
 1201  
 CCAGCTGCCATTGCCGTTGCAGGCTTCTCCAAGTTCCTGAGACTAAAGGCCGGACCTTC  
 35 1261  
 TGCTTCCAGTATGTGGAGCAACTGTGTGGTCCCTACGTCTTCATCATCTTCACTGTGCTC  
 1321  
 CTGTTCTGTTCTTCATCTTCACCTACTTCAAAGTTCCTGAGACTAAAGGCCGGACCTTC  
 1381  
 40 GATGAGATCGCTTCCGGCTTCCGGCAGGGGGAGCCAGCCAAAGTGATAAGACACCCGAG  
 1441 GAGCTGTTCCATCCCCCTGGGGGCTGATCCCAAGTGTGA

**SEQ ID No. 26 Deduced polypeptide sequence of HsGLUT1**

45 1  
 MEPSSKLLTGRMLAVGGAVLGS LQFGYNTGVINAPQKVIEEFYNQTWVHRYGESILPTT  
 61  
 LTTLWLSVAI FSVGGMIGSFVGLFVNRFGRNSMLMMNLLAFVSAVLMGF SKLGKSFE  
 121  
 50 MLILGRFIIIGVYCGLTTGFVPMYVGEVSPTAFRGALGTLHQLGIVVGILIAQVFG LDSIM  
 181  
 GNKDLWPLLLSII FIPALLQCIVLPFCPE SPRFLLINRNEENRAKSVLKKLRGTADVTHD  
 241  
 LQEMKEESRQMMREKKVTILELFRSPAYRQPILIAVVLQLSQQLSGINAVFYYSTSI FEK



301  
AGVQQPVYATIGSGIVNTAFTVVSLFVVERAGRRTLHLIGLAGMAGCAILMTIALALLEQ  
361  
LPWMSYLSIVAIFGFVAFFEVGPGPIPWFIVAELFSQGPRPAAIAVAGFSNWTSNFIVGM  
5 421  
CFQYVEQLCGPYVFI IFTVLLVLFFI FTYFKVPETKGRTFDEIASGFRQGGASQSDKTPE  
481 ELFHPLGADSQV-

10

**CLAIMS:**

1. A transgenic microalgae with increased production of omega-3 LC-PUFAs.
2. A transgenic microalgae according to claim 1 wherein the microalgae is a diatom.
3. A transgenic microalgae according to claim 2 wherein the diatom is *P. tricornutum*.
4. A transgenic microalgae according to a preceding claim wherein the omega-3 LC-PUFA is selected from EPA or DHA.
5. A transgenic microalgae according to claim 4 wherein the omega-3 LC-PUFA is DHA.
6. A transgenic microalgae according to claim 5 wherein the increased DHA content as a percentage of total fatty acids is increased by at least 1%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15% or more compared to a control microalgae.
7. A transgenic microalgae according to a preceding claim expressing a nucleic acid encoding a  $\Delta\delta$ -elongase.
8. A transgenic microalgae according to claim 7 wherein said nucleic acid comprises SEQ ID No. 1 or a sequence that encodes for a  $\Delta\delta$ -elongase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 2.
9. A transgenic microalgae according to any of claims 7 to 9 wherein the transgenic microalgae further comprises one or more nucleic acids encoding for a polypeptide involved in regulation of the LC-PUFA pathway.
10. A transgenic microalgae according to claim 9 wherein said nucleic acid encodes a  $\Delta 6$ -desaturase comprising SEQ ID No. 4 or 6 or a sequence that encodes for a  $\Delta\delta$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 4 or 6.
11. A transgenic microalgae according to claim 9 wherein said nucleic acid encodes a  $6\Delta$ -desaturase comprising SEQ ID No. 8 or 10 or a sequence that encodes for a  $6A$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 8 or 10.

12. A transgenic microalgae according to any of claims 1 to 4 wherein the omega 3-fatty acid is EPA.
13. A transgenic microalgae according to claim 12 wherein said microalgae expresses a nucleic acid that encodes a  $\Delta 6$ -desaturase.
- 5 14. A transgenic microalgae according to claim 12 wherein said nucleic acid encodes a  $\Delta 6$ -desaturase comprising SEQ ID No. 4 or 6 or a sequence that encodes for a  $\Delta 6$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 4 or 6.
- 10 15. A transgenic microalgae according to claim 12 wherein said nucleic acid encodes a  $\Delta 6$ -desaturase comprising SEQ ID No. 8 or 10 or a sequence that encodes for a  $\Delta 6$ -desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% homology to SEQ ID No. 8 or 10.
- 15 16. A transgenic microalgae according to any of claims 7 to 15 wherein said nucleic acid further comprises a regulatory sequence.
17. The use of a transgenic microalgae according to a preceding claim in producing omega-3 LC-PUFAs or increasing one or more omega-3 LC-PUFAs.
18. The use according to claim 17 wherein the omega-3 LC-PUFA is EPA or DHA.
- 20 19. The use according to claim 17 wherein the omega-3 LC-PUFAs is DHA.
20. A method for producing transgenic microalgae with increased omega-3 LC-PUFAs content.
21. A method according to claim 20 wherein the omega-3 LC-PUFA is DHA and the method comprises transforming a microalgae with a nucleic acid encoding a  $\Delta 5$ -elongase.
- 25 22. A method according to claim 20 wherein the omega-3 LC-PUFA is EPA and the method comprises transforming a microalgae with a nucleic acid encoding a  $\Delta 6$ -desaturase.
23. A method for increasing production of one of more omega-3 LC-PUFA in microalgae comprising
- 30 a) cultivating a transgenic microalgae according to any of claims 1 to 16 under conditions which allow for the production of one of more omega-3 LC-PUFAs and
- b) obtaining said one of more omega-3 LC-PUFA from the transgenic
- 35 microalgae.

24. A method according to claim 23 wherein said omega-3 LC-PUFA is DHA and the method comprises
- a) cultivating a transgenic microalgae according to any of claims 5 to 11 or 16 under conditions which allow for the production of DHA and
  - 5 b) obtaining said DHA from the transgenic microalgae.
25. A method according to claim 20 wherein said omega- 3 LC-PUFA is EPA and the method comprises
- a) cultivating a transgenic microalgae according to any of claims 12 to 16 under conditions which allow for the production of EPA and
  - 10 b) obtaining said EPA from the transgenic microalgae.
26. An oil isolated from a microalgae according to any of claims 1 to 16 or a foodstuff, feedstuff, nutraceutical or cosmetic obtained from a microalgae according to any of claims 1 to 16.
27. A composition comprising a transgenic microalgae according to any of claims 1
- 15 to 16 or an oil according to claim 26.
28. A composition comprising a transgenic microalgae according to any of claims 1 to 16 for use as a medicament.
29. A composition comprising a transgenic microalgae according to any of claims 1 to 16 for use in the treatment or prevention of cardiovascular conditions,
- 20 including atherosclerosis, thrombosis, high blood pressure, myocardial infarction and atherosclerosis, inflammatory conditions, depression, cognitive decline, arthritis, eczema, metabolic syndrome and type II diabetes.
30. A transgenic microalgae according to any of claims 1 to 16 or a composition comprising a transgenic microalgae according to any of claims 1 to 16 for use
- 25 as a foodstuff, feedstuff, nutraceutical or cosmetic.
31. A method for making a feedstuff comprising
- a) cultivating a transgenic microalgae comprising a heterologous transgene as defined in any of claims 1-16 under conditions which allow for the production of one or more omega-3 LC-PUFAs and
  - 30 b) obtaining said one or more omega-3 LC-PUFA from the transgenic microalgae.
32. An isolated nucleic acid comprising SEQ ID No. 7 or 9 encoding A6-desaturase (Ost809A6) comprising SEQ ID No. 8 or 10, a functional variant thereof or a  $\Delta$ 6-desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at

least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 8 or 11.

- 5 33. An isolated nucleic acid comprising SEQ ID No. 15 or 17 encoding a  $\Delta 4$ -desaturase (Ost809A4) comprising SEQ ID No. 16 or 18, a functional variant thereof or a A4-desaturase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 16 or 18.
- 10 34. An isolated nucleic acid comprising SEQ ID No. 19 encoding A6-elongase (FcEL06) comprising ID No. 20 a functional variant thereof or a A6-elongase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 20.
- 15 35. An isolated nucleic acid comprising SEQ ID No. 21 encoding  $\Delta \delta$ -desaturase comprising ID No. 22 a functional variant thereof or a A6-elongase that has at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% homology to SEQ ID No. 22.
- 20 36. A vector comprising an isolated nucleic according to claim 32, 33, 34 and/or 35.
37. A host cell comprising a vector according to claim 36.
38. A host cell according to claim 37 wherein the host cell is an algae or higher plant cell.
- 20 39. The use of an isolated nucleic according to claim 32, 33, 34 and/or 34 in increasing the production of omega-3 LC-PUFAs in microalgae, the preparation of a foodstuff, feedstuff, nutraceutical, cosmetic or medicament.
40. The use according to claim 39 wherein the omega-3 LC-PUFAs is EPA or DHA.
- 25 41. A method for increasing production of one of more omega-3 LC-PUFA in microalgae comprising
- 30 a) cultivating a transgenic microalgae comprising a heterologous transgene comprising one or more of the nucleic acids defined in claims 30, 31 or 32 under conditions which allow for the production of one of more omega-3 LC-PUFAs and
- b) obtaining said one of more omega-3 LC-PUFA from the transgenic microalgae.
42. A transgenic organism with increased DHA levels expressing a heterologous  $\Delta \delta$ -elongase.
- 35 43. A transgenic organism according to claim 42 wherein the  $\Delta \delta$ -elongase is a  $\Delta 5$ -elongase from *Ostreococcus tauri*.

44. A transgenic organism according to claim 42 and 43 wherein no other heterologous transgenes involved in the regulation of the LC-PUFAs biosynthetic pathway are expressed in said organism.

FIGURE 1




FIGURE 2a




FIGURE 2b






FIGURE 3a




FIGURE 3b




FIGURE 4a




FIGURE 4b




FIGURE 5a




FIGURE 5b




FIGURE 6a




FIGURE 6b




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10






FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17





FIGURE 18



Dark grown plates +/- glucose 10 days after single colonies were streaked on to plates  
 WT cells cannot grow in the dark (top of plates)

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/GB2013/052553

A. CLASSIFICATION OF SUBJECT MATTER  
INV. C12N1/Q0  
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal , WPI Data, BIOSIS, MEDLINE, EMBASE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2010/057246 AI (COMMW SCIENT IND RES ORGANISAT [AU] ; PETRIE JAMES ROBERTSON [AU] ; MACK) 27 May 2010 (2010-05-27) page 1 - page 170; claims 1-175 -----	1-44
X	wo 2011/161678 A2 (UNIV BEN GURION [IL] ; HACHEN ZVI [IL] ; KHOZIN GOLDBERG INNA [IL] ; UMI) 29 December 2011 (2011-12-29) page 1 - page 53; claims 1-21 ----- - / - -	1-44

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 November 2013

Date of mailing of the international search report

28/11/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040,  
Fax: (+31-70) 340-3016

Authorized officer

Brochado Garganta, M

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/GB2013/052553

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>HUANG Y-S ET AL: "Enzymes for transgenic biosynthesis of long-chain polyunsaturated fatty acids", BIOCHIMIE, MASSON, PARIS, FR, vol. 86, no. 11, 1 November 2004 (2004-11-01), pages 793-798, XPQ04689088, ISSN: 0300-9084, DOI: 10.1016/J.BIOCHI.2004.09.019 page 793 - page 796</p> <p style="text-align: center;">-----</p>	1-44
X	<p>MEYER A ET AL: "NOVEL FATTY ACID ELONGASES AND THEIR USE FOR THE RECONSTITUTION OF DOCOSAHEXAENOIC ACID BIOSYNTHESIS", JOURNAL OF LIPID RESEARCH, AMERICAN SOCIETY FOR BIOCHEMISTRY AND MOLECULAR BIOLOGY, INC, US, vol. 45, no. 10, 1 October 2004 (2004-10-01), pages 1899-1909, XP009046591, ISSN: 0022-2275, DOI: 10.1194/JLR.M400181-JLR200 page 1899 - page 1908</p> <p style="text-align: center;">-----</p>	1-44
X	<p>ZHOU ET AL: "Isolation and characterization of genes from the marine microalga Pavlova salina encoding three front-end desaturases involved in docosahexaenoic acid biosynthesis", PHYTOCHEMISTRY, PERGAMON PRESS, GB, vol. 68, no. 6, 3 March 2007 (2007-03-03), pages 785-796, XP005912642, ISSN: 0031-9422, DOI: 10.1016/J.PHYTOCHEM.2006.12.016 page 785 - page 795</p> <p style="text-align: center;">-----</p>	1-44
A	<p>WARD OP ET AL: "Omega-3/6 fatty acids: Alternative sources of production", PROCESS BIOCHEMISTRY, ELSEVIER, NL, vol. 40, no. 12, 1 December 2005 (2005-12-01), pages 3627-3652, XP027794053, ISSN: 1359-5113 [retrieved on 2005-12-01] the whole document</p> <p style="text-align: center;">-----</p> <p style="text-align: center;">-/--</p>	1-44

## INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2013/052553

(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>HARWOOD J L ET AL: "The versatility of algae and their lipid metabolism", BIOCHIMIE, MASSON, PARIS, FR, vol . 91, no. 6, 1 June 2009 (2009-06-01) , pages 679-684, XP026119774, ISSN : 0300-9084, DOI : 10.1016/J.BIOCHI.2008.11.004 [retrieved on 2008-11-27] the whole document</p> <p style="text-align: center;">-----</p>	1-44
A	<p>RUBIO-RODRIGUEZ N ET AL: "Production of omega-3 polyunsaturated fatty acid concentrates : A review" ; INNOVATIVE FOOD SCIENCE AND EMERGING TECHNOLOGIES, ELSEVIER, AMSTERDAM, NL, vol . 11, no. 1, 1 January 2010 (2010-01-01) , pages 1-12, XP026825026, ISSN : 1466-8564 [retrieved on 2010-01-05] the whole document</p> <p style="text-align: center;">-----</p>	1-44
A	<p>wo 2011/054800 AI (DSM IP ASSETS BV [NL] ; VERKOEIJEN DANIEL [US] ; BIJL HENDRIK LOUIS [NL] ) 12 May 2011 (2011-05-12) the whole document</p> <p style="text-align: center;">-----</p>	1-44



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/GB2013/052553
---

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2010057246	AI	27-05-2010	AR 074364 AI 12-01 -2011
			AU 2009317860 AI 27-05 -2010
			CA 2743880 AI 27-05 -2010
			CN 102317459 A 11-01 -2012
			EP 2358882 AI 24-08 -2011
			JP 2012509059 A 19-04 -2012
			NZ 593097 A 22-02 -2013
			US 2012016144 AI 19-01 -2012
			WO 2010057246 AI 27-05 -2010
WO 2011161678	A2	29-12-2011	NONE
WO 2011054800	AI	12-05-2011	AU 2010317139 AI 24-05-2012
			CA 2779551 AI 12-05-2011
			CN 102665431 A 12-09-2012
			JP 2013509860 A 21-03-2013
			KR 20120092145 A 20-08-2012
			US 2013129902 AI 23-05-2013
			WO 2011054800 AI 12-05-2011