

Rothamsted Repository Download

A - Papers appearing in refereed journals

Greenop, A., Cecelja, A., Woodcock, B. A., Wilby, A., Cook, S. M. and Pywell, R. F. 2019. Two common invertebrate predators show varying responses to different types of sentinel prey in a mesocosm study.
Journal of Applied Entomology.

The publisher's version can be accessed at:

- <https://dx.doi.org/10.1111/jen.12612>

The output can be accessed at: <https://repository.rothamsted.ac.uk/item/8w8v6>.

© Rothamsted Research. Licensed under the Creative Commons CC BY.

Title: Two common invertebrate predators show varying responses to different types of sentinel prey in a mesocosm study

Authors

Arran Greenop.^{1,2}, Andreas Cecelja., Ben A Woodcock¹, Andrew Wilby², Samantha M Cook³ & Richard F Pywell¹.

¹. NERC Centre for Ecology & Hydrology, Maclean Building, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB, UK. Tel. +44(0)1491692415

². Lancaster Environment Centre, Library Avenue, Lancaster University, Lancaster LA1 4YQ

³. Biointeractions and Crop Protection Department, Rothamsted Research, Harpenden, Herts, AL5 2JQ, UK.

Contact Author: Arran Greenop, e-mail:arrgre@ceh.ac.uk, 07538808379

Acknowledgements

This study was supported by the Centre for Ecology & Hydrology, Rothamsted Research and Lancaster University. The research was also supported by the Natural Environment Research Council (NERC) and the Biotechnology and Biological Sciences Research Council (BBSRC) under research programme NE/N018125/1 LTS-M ASSIST – Achieving Sustainable Agricultural Systems www.assist.ceh.ac.uk.

Abstract

Sentinel prey (an artificially manipulated patch of prey) are widely used to assess the level of predation provided by natural enemies in agricultural systems. While a number of different methodologies are currently in use, little is known about how arthropod predators respond to artificially-manipulated sentinel prey in comparison to predation on free-living prey populations. We assessed how attack rates on immobilised (aphids stuck to cards) and artificial (plasticine lepidopteran larvae mimics) sentinel prey differed to predation on free moving live prey (aphids). Predation was assessed in response to density of the common invertebrate predators, a foliar active ladybird *Harmonia axyridis* (Coleoptera: Coccinellidae), and a ground active beetle *Pterostichus madidus* (Coleoptera: Carabidae). Significant increases in attack rates were found for the immobilised and artificial prey between the low and high predator density treatments. However, an increased predator density did not significantly reduce numbers of free living live aphids included in the mesocosms in addition to the alternate prey. We also found no signs of predation on the artificial prey by the predator *H. axyridis*. These findings suggest that if our assessment of predation had been based solely on the foliar artificial prey then no increase in predation would have been found in response to increased predator density. Our results demonstrate that predators differentially respond to sentinel prey items which could affect the level of predation recorded where target pest species are not being used.

Key words: ecosystem services, biological control, artificial caterpillars, aphids, ladybirds, ground beetle

52 Introduction

53 Biodiversity and ecosystem functioning studies are central to understanding how humans
54 can manage the natural environment to maximise ecosystem services including pollination
55 and pest control (Bianchi et al., 2006; Cardinale et al., 2006; Foster et al., 2011). Of these
56 ecosystem services, pest regulation has received considerable attention, much of it relating
57 to the potential of natural enemies to reduce crop pest populations (Snyder et al., 2008;
58 Gardiner et al., 2009; Rusch et al., 2016; Begg et al., 2017; Greenop et al., 2018).

59

60 Fundamental to understanding the value of natural pest control in agricultural ecosystems is
61 an accurate measure of service delivery (Macfadyen et al., 2015). Several methods exist to
62 assess the suitability and function of pest control provided by natural enemies, ranging from
63 carefully-selected species assemblages in mesocosm studies conducted under laboratory
64 conditions (Straub & Snyder, 2006; Northfield et al., 2010) to the exclusion of entire
65 functional groups under real-world agricultural conditions (Gardiner et al., 2009; Holland et
66 al., 2012; Woodcock et al., 2016; Mansion-Vaquié et al 2017). The current body of literature
67 is dominated by studies that either use natural enemy abundances as a proxy for pest
68 control (Elliott et al. 1999; Schmidt et al. 2005; Bianchi, Booij & Tscharnke 2006) or infer
69 predation rates based on pest abundances (Chaplin-Kramer et al. 2011). However, both
70 approaches have associated problems that may result in the misrepresentation of the true
71 levels of pest control. For example, inferring predation based on natural enemy abundances
72 provides no direct measure of prey suppression (Macfadyen et al., 2015). Additionally pest
73 abundances are often patchy in distribution (Ferguson & Stiling 1996; Winder, Perry &
74 Holland 1999; Wan *et al.* 2018) and are influenced by bottom up as well as top-down factors
75 (Chaplin-Kramer et al., 2011). Ultimately, over- or under-estimating the efficacy of natural
76 pest control limits our capacity to manage and enhance this service to support sustainable
77 intensification of agricultural systems (Macfadyen et al., 2015; Zalucki et al., 2015).

78

79 To address this issue the use of sentinel prey has become a widely used methodology to
80 infer rates of natural pest control, particularly for agro-ecosystems (Lövei & Ferrante 2017).
81 These approaches use an artificially manipulated patch of prey that can be directly
82 monitored to assess rates of predation under field conditions (Howe et al., 2009; Winqvist et
83 al., 2011; Roslin et al., 2017). As such they provide a quantitative measure of the number of
84 prey consumed or parasitized, which is comparable between experimental treatments
85 (Birkhofer et al., 2017; Lövei & Ferrante, 2017). Two of the most common types of sentinel
86 prey currently used are: 1) live prey that have been immobilised, either by attaching them to
87 sticky labels (Winqvist et al., 2011), cards (Bianchi et al., 2005) or tethering the prey item
88 (Mathews et al., 2004); and 2) artificial prey items that act as lures and elicit a bite response
89 by predators that can then be observed as marks on the lure surfaces. These are
90 constructed out of materials such as modelling clay (Howe et al., 2009; Roslin et al., 2017;
91 Mansion-Vaquié et al., 2017). Both approaches have the practical advantage of allowing
92 studies to control prey densities and as such produce standardised assessments of
93 predation that can be replicated a large number of times at a relatively low cost.

94

95 Several concerns have been raised about the different sentinel prey approaches. One of the
96 most important is that immobilised or artificial prey no longer exhibit ecological mechanisms
97 that play important roles in predation rates. For example, certain aphids show a dropping
98 escape response to foliar-active natural enemies that can reduce predation rates (Losey &
99 Denno, 1998a; Dixon, 1958). Additionally, the state (live, wounded, artificial or dead) of prey
100 items has also been found to influence their attractiveness to predators (Zou et al., 2017;
101 Ferrante et al., 2017). Such ecological mechanisms therefore have potential to impact on the
102 level of predation recorded and consequently our capacity to infer pest control ecosystem
103 services.

104

105 While different sentinel prey methods are currently widely used to infer predation rates, little
106 is known about how many common predators respond to sentinel prey items and the manner
107 in which they are presented. In this study we aim to address this issue by comparing the
108 attack rates by two common predators: the Harlequin ladybird *Harmonia axyridis*
109 (Coleoptera: Coccinellidae), and a carabid beetle *Pterostichus madidus* (Coleoptera:
110 Carabidae) on immobilised prey aphids (*Sitobion avenae* (Hemiptera: Aphididae)) glued to
111 card, and artificial prey (plasticine caterpillars). We compare attack rates on the sentinel
112 methods and also assess how this differs to predation on live pest populations (free moving
113 *S. avenae*) at two different predator densities in a mesocosm study system. We predicted:
114 1) that an increase in attack rates on both sentinel prey (artificial caterpillars and immobilised
115 aphids stuck to cards) and a reduction in live free moving aphid numbers (live pest
116 population) in response to increasing predator density, under assumption that predator
117 attack rates are a linear function of predator density (Abrams & Ginzburg 2000); 2) attack
118 rates on immobilised aphids will be higher than on artificial prey, as the artificial prey do not
119 possess any chemical cues used by both predator species to locate prey and do not
120 represent a valid food item (Lövei & Sunderland, 1996; Kiehl et al., 1996; Abassi et al.,
121 2000); and, 3) the ground beetle will be more likely than the ladybird to attack artificial prey
122 as they have been shown to be highly opportunistic and generalist visual hunters (Lang &
123 Gsödl, 2008; Ferrante et al., 2017). In contrast *H. axyridis* is highly dependent on olfactory
124 as well as visual cues to locate prey (Koch, 2003).

125

126 Methods

127 Experimental system

128 We used an experimental mesocosm design to control predator density and composition
129 between treatments. Each mesocosm comprised a 10L plant pot (28.5cm diameter / 22.5cm

deep), filled with peat-free compost and three wheat plants (*Triticum aestivum* L. Em. Fiori & Paol. Variety: KWS Dacanto), enclosed within a porous plastic mesh (height 36.5cm / diameter 28.5cm, pore size 0.05mm, held under standardised environmental conditions of $19.5 \pm 1^\circ\text{C}$ and LD 16:8 h). The combination of a ground-foraging predator, *P. madidus* and a foliar-foraging predator, *H. axyridis* was used as model predator community. Both species are predators of aphids, though have spatially segregated hunting niches (ground vs. canopy, respectively) (Schmitz, 2007; Woodcock & Heard, 2011). Adult *P. madidus* were collected through dry pitfall trapping and then stored in a controlled temperature facility ($19.5 \pm 1^\circ\text{C}$ and LD 16:8 h) in plastic cups containing moist soil, and were fed with dog food *ad libitum*. Within the same environment, adult *H. axyridis* were collected by hand from the field and stored in plastic 10L pots (28.5cm diameter / 22.5cm deep) covered with a porous plastic mesh (pore size 0.05mm) and were fed *ad libitum* with live aphids. Predators were kept for a maximum of four weeks in the laboratory. The pest species on which predation was assessed was *S. avenae*, an important aphid pest of wheat frequently used as a model prey item for measuring pest control (Mansion-Vaquié et al. 2017; Bøsem Baillod et al. 2017). This aphid species shows a dropping behaviour in response to predator attacks (Winder 1990).

We tested two forms of sentinel prey commonly used to assess the delivery of natural pest control ecosystem services under field conditions. Immobilised prey represented by 10 aphids glued using superglue (Loctite Super Glue, Henkel, Düsseldorf, Germany) to 4 x 6cm pieces of green card; aphids were placed approx., 0.5 cm apart. This reflects methodologies established by Winqvist et al. (2011). Within each mesocosm we suspended one card in the canopy of the wheat using a pin, and placed another on the soil surface of the plant pot (adapted from Winqvist et al. (2011)). We also used artificial prey designed to mimic lepidopteran caterpillars. Whilst the focus of our experiment was aphid prey, the use of artificial caterpillars has been widely used to infer predation rates in agricultural settings where the target pest species is not necessarily lepidopteran (Mansion-Vaquié et al. 2017).

Following approaches described in Howe, Lövei & Nachman (2009), caterpillars were made of non-toxic green plasticine (Newplast, Newclay, Devon, UK) and were 2cm × 0.5cm in diameter (Supplementary material: Appendix S1; Figure S1 & S2). Caterpillars were glued using superglue (Loctite) in pairs to 3x3cm pieces of green card. This ensured once constructed, no further handling of individual caterpillars occurred, avoiding the risk of accidental marks (important as marks were used as a measure of predation). A total of 10 artificial prey items were suspended in the canopy by pinning the card with the caterpillars attached to the wheat foliage and 10 caterpillars placed on the soil surface, so the method could be quantitatively compared to the immobilised prey. In each mesocosm we also included live prey so that attack rates on the sentinel prey could be compared to live prey populations. Live prey populations were established as 20 free-moving adult *S. avenae* aphids evenly distributed on the leaves of each wheat plant. Aphids were allowed to settle for 4 hours, after which the two predator species were introduced. In addition to the two sentinel prey treatments, we also included a control treatment for each sentinel prey type that contained no predators. The control treatments were established following the same experimental protocol as above.

Using this model system, we assessed whether an increased density of predators resulted in higher attack rates on the sentinel prey and lower numbers of live aphids. We prepared a low-density treatment comprising two *H. axyrdis* and two *P. madidus*, and a high-predator density treatment with four individuals each of *H. axyrdis* and *P. madidus*. Each treatment was replicated seven times. All treatments were run at the same time with predators that were starved 24h prior to the experiment (predators were used only once i.e. a total of 84 individuals of each species were used over the whole experiment). The proportion of immobilised aphids and the proportion of plasticine caterpillars showing evidence of attack were recorded out of 20 and the number of live aphids were counted after 24h from the point where predators were added.

Statistical analysis

We wanted to determine whether prey location (ground vs. foliar) affected predation rates at the two predator densities (low vs. high) and whether these attack rates differed between the sentinel prey methods (immobilised vs. artificial). However, we found no signs of predation by the ladybird on the artificial prey. This resulted in zero variation for this parameter which can lead to unreliable results in generalized linear models (Kuhn & Johnson 2013). Therefore, we first analysed the immobilised prey separately to determine whether attack rates differed between the ground and foliar predators at the different predator densities. Prey items were not analysed individually as statistically independent units, but rather a proportional attack rate across all 10 prey items at either the ground or foliage. We used a binomial distribution reflecting the bounding (0-1) of data. The response variable was attack rate (proportion of prey attacked out of 10) and the explanatory variables were predator density (low and high) and predator feeding location (ground or foliar) and the interaction between these two factors. As there were no predators in the controls for the artificial and immobilised prey treatments we found no signs of attack on the plasticine caterpillars or the aphids glued to cards (except one missing aphid from a card). This again meant that there was near zero variation for the controls and they were excluded from analysis. We then analysed the ground sentinel prey separately to determine whether *P. madidus* had higher attack rates on the plasticine caterpillars in comparison to the immobilised live prey, as it actively attacked both prey types. We used a binomial GLM with attack rate as the response variable and the explanatory variables predator density and prey type and the interaction between these two factors. Significance was assessed against a chi distribution.

To determine how predator density affected predation on live aphids we used a negative binomial GLM implemented in the MASS package (Venables & Ripley, 2002). A negative binomial GLM was used to account for overdispersion in the count data and for the fact that pest populations have the capacity to reproduce, even over a 24h time frame. The response

variable was the number of aphids counted in the mesocosm at the end of the experiment and the explanatory variables were predator density (control (no predators), low and high) and alternative prey type (artificial and immobilised) and the interaction between these two factors. Significance was assessed against a chi distribution. Where the interaction was significant, orthogonal post-hoc contrasts were carried out. All analyses were carried out in R (R Core Team, 2017).

Results

For both sentinel prey methods, evidence of attack was recorded after the 24h foraging period, suggesting that immobilised prey stuck to cards and artificial caterpillars elicited a predation response in the predators. However, we found no signs of predation on the artificial caterpillars in the canopy and found no jaw marks from the predator *H. axyridis* on any of the artificial caterpillars placed on the soil surface; these showed predation only by *P. madidus*. Of the immobilised aphid prey, we found no significant interaction between predator feeding location and predator density on predator attack rates ($\chi^2 = 0.210$, $df = 1$, $p = 0.647$). Predator feeding location was also not found to have a significant effect on attack rates on the immobilised prey ($\chi^2 = 1.981$, $df = 1$, $p = 0.159$), however there was a significant effect of predator density ($\chi^2 = 10.407$, $df = 1$, $p = 0.002$). Attack rates were significantly higher at the high predator density compared to the low predator density (proportion of prey attacked out of 10 on immobilised prey: low predator density = $0.207 \pm 1SE\ 0.046$; high predator density = $0.779 \pm 1SE\ 0.094$). Where predation was compared between sentinel prey types for *P. madidus* we found there was no significant interaction between prey type and predator density ($\chi^2 = 0.269$, $df = 1$, $p = 0.604$). Prey type was also not significant ($\chi^2 = 0.020$, $df = 1$, $p = 0.887$), however there was a significant increase in attack rates by *P. madidus* between predator densities ($\chi^2 = 10.080$, $df = 1$, $p = 0.001$) (low predator density mean = $0.114 \pm 1SE\ 0.038$; high predator density mean = $0.679 \pm 1SE\ 0.070$).

There was no significant interaction between predator density and alternative prey type on the number of live prey in each treatment ($\chi^2 = 1.110$, $df = 1$, $p = 0.574$), however both main effects predator density and alternative prey type were significant (Alternative prey type $\chi^2 = 6.066$, $df = 1$, $p = 0.014$; Predator density $\chi^2 = 21.813$, $df = 2$, $p = <0.001$). Post hoc comparisons showed that there was a significant difference between the number of live aphids in the control treatments and the predation treatments ($z = -4.521$, $p = <0.001$) (Table 1). However, there was no significant difference between the low predator density treatment and high predator density treatment ($z = 1.100$, $p = 0.271$). The number of live aphids in the artificial prey treatment was significantly lower than the immobilised prey treatment (Table 1).

Discussion

Effect of predator density on attack rates

In accordance with our first prediction, both the immobilised and artificial prey detected increased attack rates in response to a higher predator density. However, in the case of the live aphids there was no evidence of increased consumption at the higher predator densities. This contrasts with the higher attack rates seen for the sentinel prey under the same conditions. The sentinel prey represented both aggregated and highly conspicuous prey that, in contrast to the live aphids, were unable to escape from predators. In this situation, once the predators located the prey the two predominant limiting effects on attack rates would be predator satiation or negative intraguild interactions (Gagnon, Heimpel & Brodeur 2011). Immobilised prey were viable food items, so would contribute to predator satiation, which could have reduced predation on the free moving prey (the number of free moving aphids was still significantly lower in the predation treatments compared to the control, indicating that predation did occur on the live pest populations). In contrast, the artificial prey is unlikely to contribute to predator satiation as it offers no nutrition, which could lead to an inflation of attack rates on artificial prey (where predators continually attack the prey due to a lack of satiation) or cause them to search for alternative prey (Lövei & Ferrante 2017). We found that significantly more free-moving aphids were consumed in the artificial prey treatment

compared to the immobilised prey treatment, suggesting the predators were attacking the live prey to gain food (although the number of aphids consumed did not change as a function of predator density). However, there was strong evidence that at the higher predator densities artificial caterpillars were often attacked multiple times (i.e. >40% of caterpillars were attacked). This relatively high attack rate for the artificial caterpillars on the ground may have reduced predation by the ground foraging beetles on the live aphids. A final point is that in comparison to the sentinel prey, the free-moving aphids would be able to avoid predators through either escape responses such as dropping from the plant when attacked, or persisting in refuges where they are less vulnerable to predation; both mechanisms have been found to reduce predation rates (Losey & Denno, 1998a; b; Berryman et al., 2006; Bommarco et al., 2007). This could also explain the lower levels of predation on the free-moving aphid populations also included in the mesocosm.

Predation responses to the different sentinel prey methods

We found equivocal evidence in support of our second prediction that attack rates were lower on the artificial prey compared to the immobilised prey, with no significant difference in attack rates by *P. madidus* being identified between the sentinel prey. However, if predation assessments were based only on the artificial caterpillars located in the plant canopy, then no difference in predation would have been detected as *H. axyridis* was not seen to bite these artificial prey items. This agrees with the findings of Lövei and Ferrante, (2017) who demonstrated lower predation on artificial sentinel prey compared to real sentinel prey. Our results suggest this is due to individual predator feeding preferences. The lack of predation by *H. axyridis* supported our third prediction that ground beetles would be more likely to attack artificial prey. Both ground beetles and ladybirds have been found to use visual cues when selecting feeding patches (e.g. attracted to high prey densities) (Lövei & Sunderland, 1996; Osawa, 2000; Lee & Kang, 2004) and both groups have also been found to respond to and locate prey based on aphid volatiles (Lövei & Sunderland, 1996; Kielty et al., 1996;

Koch, 2003). However, our results suggest that either *H. axyridis* does not view plasticine caterpillars as a prey item, or demonstrates preferences for live aphids over lepidopteran prey. It is worth noting here that *H. axyridis* is polyphagous and will feed on juvenile stages of Lepidoptera (Koch et al., 2003). For this reason other factors may also contribute to the effective avoidance of the artificial caterpillars by *H. axyridis*. For example, *H. axyridis* relies more on olfactory cues and has been shown to be highly attracted to the chemical (*E*)- β -farnesene a key component of the alarm pheromone for most aphid species including *S. avenae* (Verheggen et al., 2007). In contrast, ground beetles are more opportunistic predators and may base feeding choices on prey vulnerability (Lang & Gsödl, 2008), which could increase the likelihood of ground beetles attacking artificial prey items. The use of plasticine caterpillars may therefore be a poor measure of predation where the dominant predators in the ecosystem are Coccinellidae or other taxa that show similar hunting strategies.

Sentinel methods are rarely used to calculate absolute predation and are more frequently used to compare the relative amount of predation between experimental units (Lövei & Ferrante, 2017). When combined with information on crop yield, direct measures of pest damage and conventional quantification of both pest and predator densities, sentinel prey approaches have the potential to provide valuable insights into pest control dynamics in agro-ecosystems. Whilst understanding relative changes in predation between experimental units is useful in elucidating ecosystems dynamics, being able to use sentinel prey items to provide a surrogate measure of pest control for target pest species could be developed into a standardised measure of pest control that can be applied to a range of farming systems. Our study provides a basic demonstration that live and sentinel prey items detect varying levels of predation in response to different predator species and predator densities, which highlights potential limitations of using sentinel prey as proxies for pest suppression. However, as live prey populations are able to reproduce and move, dynamics which cannot

be replicated in sentinel prey, the measure of success for real prey is often based on pest threshold densities. As such it is very difficult to draw parallels between predation on live and sentinel prey items.

There are limitations in this study to the sentinel prey approaches used to evaluate natural pest control that merit consideration if the findings of this research are to inform future work. Firstly, the number of artificial caterpillars may have been unrealistically high as this prey item was included in mesocosms at the same density as the immobilised aphids. This was done in part for practical reasons; if the number of prey were too small then detectable differences between experimental units would be hard to observe, particular where all the prey were consistently attacked or consumed (a problem akin to the “ceiling effect” in statistics) (Austin & Brunner 2003). However, comparable densities to the immobilised prey used in this study are not uncommon for pest populations under field conditions. For example, caterpillars such as *Artogeia rapae* (small cabbage white) can reach similar densities (Hooks, Pandey & Johnson 2003), while aggregations of aphids will normally exceed those used in this study (Sunderland & Vickerman 1980; Sopp, Sunderland & Coombes 1987). Secondly, the sentinel prey types in our study could have been assessed in isolation without alternative real prey. This would have enabled the relative differences in predation between methods to be directly compared more easily. However, to understand how these methods perform in the real-world, where predators are exposed to both naturally occurring free-moving and experimental sentinel prey, using more than one prey species provides a more realistic comparison. In mesocosms a predator may attack the sentinel prey (where it is the only prey) out of necessity (starvation), which directly contrasts to an agroecosystem where alternative prey are likely to be available. Accordingly, this could falsely represent predation by that species on sentinel prey. A similar criticism could also be made where studies use a single real prey species to assess natural pest control. However, typically these studies focus on a model prey species deliberately selected as it represents a

pest of economic importance to that crop. In this situation avoidance of that key pest species in preference to alternative prey still provides key biologically relevant information in terms of assessing pest control, at least for that key pest. Finally, it is possible that due to the close spacing of the prey, that the free-moving aphids could walk on the caterpillars and potentially leave prey-related chemical cues on them increasing their level of attraction to the predators. However, we found that more aphids were consumed where the alternate prey were the plasticine caterpillars as opposed to the immobilised prey. This would suggest that the predators were distinguishing between the artificial prey and real prey in the mesocosm without being affected by such chemical cues.

Conclusion

Sentinel prey methods offer a simple way to measure predation that have significant advances over surrogate measures that rely on variation in prey or predator abundances (Lövei & Sunderland, 1996; Chaplin-Kramer et al., 2011; Lövei & Ferrante, 2017). However, when using sentinel prey our results highlight the importance of considering the effects of predator and pest species life-history traits and the influence these have on observed predation. A sensible approach would be to consider more than one measure of prey suppression and tailor it to the desired measure of the study (Macfadyen et al., 2015). For example, using plasticine caterpillars in conjunction with live, free moving prey (of a known focal pest species) would allow a practitioner to record potential key predators within an ecosystem based on the detection of bite marks in the plasticine, whilst also giving an indication of actual pest suppression on the live prey. Correlation in predation rates between the two methods could be used to determine whether the predators revealed by the artificial prey method are the dominant predators responsible for pest control within that particular agroecosystem. Accounting for variation in the attractiveness of different prey items to predators, the effects of inhibiting important ecological escape mechanisms and the effects

of different sentinel prey densities will improve estimates of prey suppression. Ultimately, this will improve our understanding of how natural pest control is delivered under field conditions.

Author contribution

- All authors conceived the research
- Author 1 and author 2 designed and conducted experiments
- Author 1 conducted statistical analyses
- Author 1 wrote the draft manuscript
- All authors contributed to writing the final and revised manuscripts
- All authors approved the final version of the manuscript

References

- Abassi SA, Birkett MA, Pettersson J, Pickett JA, Wadhams LJ & Woodcock CM. (2000). Response of the Seven-spot Ladybird to an Aphid Alarm Pheromone and an Alarm Pheromone Inhibitor is Mediated by Paired Olfactory Cells. *Journal of Chemical Ecology*, 26:1765-1771
- Abrams, P.A. & Ginzburg, L.R. (2000) The nature of predation: prey dependent, ratio dependent or neither? *Trends in Ecology & Evolution*, 15, 337–341.
- Austin, P.C. & Brunner, L.J. (2003) Type I error inflation in the presence of a ceiling effect. *The American Statistician*, 57, 97–104.
- Begg, G.S., Cook, S.M., Dye, R., Ferrante, M., Franck, P., Lavigne, C., Lövei, G. Mansion-Vaquie, A., Pell, J., K., Petit, S., Quesada, N., Ricci, B. & Birch, A.N.E (2017) A functional overview of conservation biological control. *Crop Protection* 97: 145-158
- Berryman AA, Hawkins BA & Hawkins BA. (2006). The refuge as an integrating concept in ecology and evolution. *Oikos*, 115:192–196.
- Bianchi FJJA, Booij CJH & Tscharntke T. (2006). Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings. Biological Sciences / the Royal Society*, 273:1715–1727.
- Bianchi FJJA, van Wingerden WKRE, Griffioen AJ, van der Veen M, van der Straten MJJ, Wegman RMA & Meeuwsen HAM. (2005). Landscape factors affecting the control of

406 *Mamestra brassicae* by natural enemies in Brussels sprout. *Agriculture, Ecosystems &*
407 *Environment*, 107:145–150.

408 Birkhofer K, Bylund H, Dalin P, Ferlian O, Gagic V, Hambäck PA, Klapwijk M, Mestre L,
409 Roubinet E, Schroeder M, Stenberg JA, Porcel M, Björkman C & Jonsson M. (2017).
410 Methods to identify the prey of invertebrate predators in terrestrial field studies. *Ecology and*
411 *Evolution*, 7:1942–1953.

412 Bommarco R, Firling SO & Ekblom B. (2007). Outbreak suppression by predators depends on
413 spatial distribution of prey. *Ecological Modelling*, 201:163–170.

414 Bosem Baillod, A., Tscharnke, T., Clough, Y. and Batáry, P. (2017). Landscape-scale
415 interactions of spatial and temporal cropland heterogeneity drive biological control of cereal
416 aphids. *Journal of Applied Ecology*, 54:1804-1813.

417 Cardinale BJ, Srivastava DS, Duffy JE, Wright JP, Downing AL, Sankaran M & Jouseau C.
418 (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*,
419 443:989–992.

420 Chaplin-Kramer R, O'Rourke ME, Blitzer EJ & Kremen C. (2011). A meta-analysis of crop
421 pest and natural enemy response to landscape complexity. *Ecology Letters*, 14:922–932.

422 Dixon AFG. (1958). The escape responses shown by certain aphids to the presence of the
423 coccinellid *Adalia decempunctata* (L.). *Transactions of the Royal Entomological Society of*
424 *London*, 110:319–334.

425 Elliott NC, Kieckhefer RW, Lee J-H & French BW. (1999). Influence of within-field and
426 landscape factors on aphid predator populations in wheat. *Landscape Ecology* 14:239-252.

427 Ferguson KI & Stiling P. (1996). Non-additive effects of multiple natural enemies on aphid
428 populations. *Oecologia*, 108:375–379.

429 Ferrante M, Barone G & Lövei GL. (2017). The carabid *Pterostichus melanarius* uses
430 chemical cues for opportunistic predation and saprophagy but not for finding healthy prey.
431 *Biocontrol*, 62:741–747.

432 Foster WA, Snaddon JL, Turner EC, Fayle TM, Cockerill TD, Ellwood MDF, Broad GR,
433 Chung AYC, Eggleton P, Khen CV & Yusah KM. (2011). Establishing the evidence base for
434 maintaining biodiversity and ecosystem function in the oil palm landscapes of South East
435 Asia. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
436 *Sciences*, 366:3277–3291.

437 Gagnon, A.-È., Heimpel, G.E. & Brodeur, J. (2011) The ubiquity of intraguild predation
438 among predatory arthropods. *Plos One*, 6, e28061.

439 Gardiner MM, Landis DA, Gratton C, DiFonzo CD, O'Neal M, Chacon JM, Wayo MT,
440 Schmidt NP, Mueller EE & Heimpel GE. (2009). Landscape diversity enhances biological
441 control of an introduced crop pest in the north-central USA. *Ecological Applications*, 19:143–
442 154.

443 Greenop A, Woodcock BA, Wilby A, Cook SM & Pywell RF. (2018). Functional diversity
444 positively affects prey suppression by invertebrate predators: a meta-analysis. *Ecology*,
445 doi.org/10.1002/ecy.2378

446 Holland JM, Oaten H, Moreby S, Birkett T, Simper J, Southway S & Smith BM. (2012). Agri-
 447 environment scheme enhancing ecosystem services: A demonstration of improved biological
 448 control in cereal crops. *Agriculture, Ecosystems & Environment*, 155:147–152.

449 Hooks, C.R.R., Pandey, R.R. & Johnson, M.W. (2003) Impact of avian and arthropod
 450 predation on lepidopteran caterpillar densities and plant productivity in an ephemeral
 451 agroecosystem. *Ecological Entomology*, 28, 522–532.

452 Howe A, Lövei GL & Nachman G. (2009). Dummy caterpillars as a simple method to assess
 453 predation rates on invertebrates in a tropical agroecosystem. *Entomologia experimentalis et*
 454 *applicata*, 131:325–329.

455 Kielly JP, Allen-Williams LJ, Underwood N & Eastwood EA. (1996). Behavioral responses of
 456 three species of ground beetle (Coleoptera: Carabidae) to olfactory cues associated with
 457 prey and habitat. *Journal of Insect Behavior*, 9:237–250.

458 Koch RL, Hutchison WD, Venette RC & Heimpel GE. (2003). Susceptibility of immature
 459 monarch butterfly, *Danaus plexippus* (Lepidoptera: Nymphalidae: Danainae), to predation by
 460 *Harmonia axyridis* (Coleoptera: Coccinellidae). *Biological Control*, 28:265–270.

461 Koch RL. (2003). The multicolored Asian lady beetle, *Harmonia axyridis*: A review of its
 462 biology, uses in biological control, and non-target impacts. *Journal of Insect Science*, 3:1–16.

463 Kuhn, M. & Johnson, K. (2013) *Applied Predictive Modeling*. Springer New York, New York,
 464 NY.

465 Lang A & Gsödl S. (2008). Prey vulnerability and active predator choice as determinants of
 466 prey selection: a carabid beetle and its aphid prey. *Journal of Applied Entomology*, 125:53–
 467 61.

468 Lee J-H & Kang T-J. (2004). Functional response of *Harmonia axyridis* (Pallas) (Coleoptera:
 469 Coccinellidae) to *Aphis gossypii* Glover (Homoptera: Aphididae) in the Laboratory. *Biological*
 470 *Control*, 31:306–310.

471 Losey JE & Denno RF. (1998a). The escape response of pea aphids to foliar-foraging
 472 predators: factors affecting dropping behaviour. *Ecological Entomology* 23:53–61.

473 Losey JE & Denno RF. (1998b). Interspecific variation in the escape responses of aphids:
 474 effect on risk of predation from foliar-foraging and ground-foraging predators. *Oecologia*,
 475 115:245–252.

476 Lövei GL & Ferrante M. (2017). A review of the sentinel prey method as a way of quantifying
 477 invertebrate predation under field conditions. *Insect Science*, 24:528–542.

478 Lövei GL & Sunderland KD. (1996). Ecology and behavior of ground beetles (Coleoptera:
 479 Carabidae). *Annual Review of Entomology*, 41:231–256.

480 Macfadyen S, Davies AP & Zalucki MP. (2015). Assessing the impact of arthropod natural
 481 enemies on crop pests at the field scale. *Insect Science*, 22:20–34.

482 Mansion-Vaquié A, Ferrante M, Cook SM, Pell JK & Lövei GL. (2017). Manipulating field
 483 margins to increase predation intensity in fields of winter wheat (*Triticum aestivum*). *Journal*
 484 *of Applied Entomology*, 141:600–611.

485 Mathews CR, Bottrell DG & Brown MW. (2004). Habitat manipulation of the apple orchard
 486 floor to increase ground-dwelling predators and predation of *Cydia pomonella* (L.)
 487 (Lepidoptera: Tortricidae). *Biological Control*, 30:265–273.

488 Northfield TD, Snyder GB, Ives AR & Snyder WE. (2010). Niche saturation reveals resource
489 partitioning among consumers. *Ecology Letters*, 13:338–348.

490 Osawa N. (2000). Population field studies on the aphidophagous ladybird beetle *Harmonia*
491 *axyridis* (Coleoptera: Coccinellidae): resource tracking and population characteristics.
492 *Population Ecology*, 42:115–127.

493 R Core Team (2017). R: A language and environment for statistical computing. R
494 Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

495 Roslin T, Hardwick B, Novotny V, Petry WK, Andrew NR, Asmus A, Barrio IC, Basset Y,
496 Boesing AL, Bonebrake TC, Cameron EK, Dáttilo W, Donoso DA, Drozd P, Gray CL, Hik
497 DS, Hill SJ, Hopkins T, Huang S, Koane B, Laird-Hopkins B, Laukkanen L, Lewis OT, Milne
498 S, Mwesige I, Nakamura A, Nell CS, Nichols E, Prokurat A, Sam K, Schmidt NM, Slade A,
499 Slade V, Suchanková A, Teder T, van Nouhuys S, Vandvik V, Weissflog A, Zhukovich V &
500 Slade EM. (2017). Higher predation risk for insect prey at low latitudes and elevations.
501 *Science*. 356:742–744.

502 Rusch A, Chaplin-Kramer R, Gardiner MM, Hawro V, Holland J, Landis D, Thies C,
503 Tscharnkte T, Weisser WW, Winqvist C, Woltz M & Bommarco R. (2016). Agricultural
504 landscape simplification reduces natural pest control: A quantitative synthesis. *Agriculture,*
505 *Ecosystems & Environment*, 221:198–204.

506 Schmidt MH, Roschewitz I, Thies C & Tscharnkte T. (2005). Differential effects of landscape
507 and management on diversity and density of ground-dwelling farmland spiders. *Journal of*
508 *Applied Ecology*, 42:281–287.

509 Schmitz OJ. (2007). Predator diversity and trophic interactions. *Ecology*, 88:2415–2426.

510 Snyder GB, Finke DL & Snyder WE. (2008). Predator biodiversity strengthens aphid
511 suppression across single- and multiple-species prey communities. *Biological Control*,
512 44:52–60.

513 Sopp, P.I., Sunderland, K.D. & Coombes, D.S. (1987) Observations on the number of cereal
514 aphids on the soil in relation to aphid density in winter wheat. *Annals of Applied Biology*, 111,
515 53–57.

516 Straub CS & Snyder WE. (2006). Species identity dominates the relationship between
517 predator biodiversity and herbivore suppression. *Ecology*, 87:277–282.

518 Sunderland, K.D. & Vickerman, G.P. (1980) Aphid feeding by some polyphagous predators
519 in relation to aphid density in cereal fields. *The Journal of Applied Ecology*, 17, 389.

520 Venables WN & Ripley BD. (2002). Modern Applied Statistics with S. Statistics and
521 Computing. (ed by J Chambers, W Eddy, W Härdle, S Sheather & L Tierney) 4th edn.
522 Springer New York, New York, NY,.

523 Verheggen FJ, Fagel Q, Heuskin S, Lognay G, Francis F & Haubruge E. (2007).
524 Electrophysiological and behavioral responses of the multicolored Asian lady beetle,
525 *Harmonia axyridis* pallas, to sesquiterpene semiochemicals. *Journal of Chemical Ecology*,
526 33:2148–2155.

527 Wan, N.-F., Ji, X.-Y., Kiær, L.P., Liu, S.-S., Deng, J.-Y., Jiang, J.-X. & Li, B. (2018) Ground
528 cover increases spatial aggregation and association of insect herbivores and their predators
529 in an agricultural landscape. *Landscape Ecology*, 33, 799–809.

- Winder, L. (1990). Predation of the cereal aphid *Sitobion avenae* by polyphagous predators on the ground. *Ecological Entomology*, 15:105-110.
- Winder, L., Perry, J.N. & Holland, J.M. (1999) The spatial and temporal distribution of the grain aphid *Sitobion avenae* in winter wheat. *Entomologia experimentalis et applicata*, 93, 275–288.
- Winqvist C, Bengtsson J, Aavik T, Berendse F, Clement LW, Eggers S, Fischer C, Flohre A, Geiger F, Liira J, Pärt T, Thies C, Tschardt T, Weisser WW & Bommarco R. (2011). Mixed effects of organic farming and landscape complexity on farmland biodiversity and biological control potential across Europe. *Journal of Applied Ecology*, 48:570–579.
- Woodcock B., Bullock J., McCracken M, Chapman R., Ball S., Edwards M., Nowakowski M & Pywell R. (2016). Spill-over of pest control and pollination services into arable crops. *Agriculture, Ecosystems & Environment*, 231:15–23.
- Woodcock BA & Heard MS. (2011). Disentangling the effects of predator hunting mode and habitat domain on the top-down control of insect herbivores. *The Journal of Animal Ecology*, 80:495–503.
- Zalucki MP, Furlong MJ, Schellhorn NA, Macfadyen S & Davies AP. (2015). Assessing the impact of natural enemies in agroecosystems: toward “real” IPM or in quest of the Holy Grail? *Insect Science*, 22:1–5.
- Zou Y, de Kraker J, Bianchi FJJA, van Telgen MD, Xiao H & van der Werf W. (2017). Video monitoring of brown planthopper predation in rice shows flaws of sentinel methods. *Scientific Reports*, 7:42210.

Table 1. Number of live aphids (*Sitobion avenae*) recorded after 24h exposure to the predators *Pterostichus madidus* and *Harmonia axyridis* in a mesocosm where either immobilised prey (20 *S. avenae* aphids glued to card) or artificial prey (20 plasticine caterpillars) were also available. Predator densities control (no predators), low (two *H. axyridis* and two *P. madidus*) and high (four *H. axyridis* and four *P. madidus*) are the mean across both alternate prey types. Artificial prey treatment and immobilised prey treatment are the mean across all predator densities.

Treatment	Number of live aphids (mean \pm 1SE)
Control	28.786 \pm 2.823
Low predatory density	18.429 \pm 2.336
High predator density	15.643 \pm 1.561
Artificial prey treatment	18.190 \pm 2.102
Immobilised prey treatment	23.714 \pm 2.212