



US 20170067070A1

(19) **United States**(12) **Patent Application Publication**  
**Bauer et al.**(10) **Pub. No.: US 2017/0067070 A1**(43) **Pub. Date: Mar. 9, 2017**(54) **NOVEL FATTY ACID DESATURASES AND  
ELONGASES AND USES THEREOF**(71) Applicant: **BASF Plant Science Company GmbH,**  
Ludwigshafen (DE)(72) Inventors: **Jörg Bauer,** Teltow (DE); **Johnathan**  
**A. Napier,** Preston (GB); **Olga**  
**Sayanova,** Redbourn (GB)(21) Appl. No.: **15/351,962**(22) Filed: **Nov. 15, 2016****Related U.S. Application Data**(63) Continuation of application No. 13/384,277, filed on  
Jan. 16, 2012, now Pat. No. 9,493,520, filed as  
application No. PCT/EP10/60178 on Jul. 15, 2010.(60) Provisional application No. 61/226,301, filed on Jul.  
17, 2009.(30) **Foreign Application Priority Data**

Jul. 17, 2009 (EP) ..... 09165752.8

**Publication Classification**(51) **Int. Cl.****C12N 15/82** (2006.01)**C12N 9/00** (2006.01)**C12N 9/02** (2006.01)(52) **U.S. Cl.**CPC ..... **C12N 15/8247** (2013.01); **C12N 9/0071**  
(2013.01); **C12N 9/93** (2013.01); **C12Y**  
**114/19003** (2013.01); **C12Y 114/19** (2013.01);  
**C12Y 114/19006** (2013.01); **C12Y 114/19004**  
(2013.01); **C12Y 602/01003** (2013.01)

## (57)

**ABSTRACT**

The invention provides isolated nucleic acid molecules which encode novel fatty acid desaturases and elongases from the organism *Emiliana huxleyi*. The invention also provides recombinant expression vectors containing desaturase or elongase nucleic acid molecules, host cells into which the expression vectors have been introduced, and methods for large-scale production of long chain polyunsaturated fatty acids (LCPUFAs), e.g. arachidonic acid (ARA), eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA).

**ω3- pathway**

**aerobic**

**ω6- pathway**

**PKS (anaerobic)  
e.g. bacteria**

**Sprecher pathway  
e.g. mammals**

**β-oxidation**

18:2 Δ 9,12 → Δ 9-elongase → 20:2 Δ 11,14 → Δ 8-desaturase → 20:3 Δ 8,11,14 → Δ 5-desaturase → 20:4 Δ 6,8,11,14 → Δ 5-elongase → 22:4 Δ 7,10,13,16 → Δ 4-desaturase → 22:5 Δ 4,7,10,13,16

18:3 Δ 9,12,15 → Δ 6-desaturase → 18:4 Δ 6,9,12,15 → Δ 6-elongase → 20:4 Δ 8,11,14,17 → Δ 5-desaturase → 20:5 Δ 5,8,11,14,17 → Δ 5-elongase → 22:5 Δ 7,10,13,16,19 → Δ 4-desaturase → 22:6 Δ 4,7,10,13,16,19

20:3 Δ 11,14,17 → Δ 9-elongase → 20:3 Δ 11,14,17 → Δ 8-desaturase → 20:4 Δ 8,11,14,17 → Δ 5-desaturase → 20:5 Δ 5,8,11,14,17 → Δ 5-elongase → 22:5 Δ 7,10,13,16,19 → Δ 4-desaturase → 22:6 Δ 4,7,10,13,16,19

24:5 Δ 9,12,15,18,21 → Δ 6-desaturase → 24:6 Δ 6,9,12,15,18,21

Acetyl-CoA  
Malonyl-CoA → 20:5 Δ 5,8,11,14,17 → 22:6 Δ 4,7,10,13,16,19

**ARA** 20:4 Δ 5,8,11,14

**EPA** 20:5 Δ 5,8,11,14,17

**DHA** 22:6 Δ 4,7,10,13,16,19




Figure 2 of 13: Gas chromatograph of yeast expression experiment with feeding of 22:5n-3 in the presence (A) and absence (B) of d4Des(Eh). The arrow indicates the formation of 22:6, the product of d4Des(Eh) activity.

### Expression in yeast




Figure 3 of 13: Gas chromatograph of yeast expression experiment with feeding of 20:3n-3 in the presence (A) and absence (B) of d8Des(Eh). The arrow indicates the formation of 20:4n-3, the product of d8Des(Eh) activity.




Figure 4 of 13: Gas chromatograph of yeast expression experiment with feeding of 18:3n-3 (ALA) in the presence (A) and absence (B) of d9Elo(Eh). The arrow indicates the formation of 20:3n-3, the product of d9Elo(Eh) activity.




Figure 5 of 13: Gas chromatograph of yeast expression experiment with feeding of 18:3n-6 (GLA) and 18:4n-3 (SDA) in the presence (A) and absence (B) of d5Elo(Eh). The arrow indicates the formation of 20:3n-6 or 20:4n-3, respectively, the products of d5Elo(Eh) activity




Figure 6 of 13: Gas chromatograph of yeast expression experiment with feeding of 20:4n-6 (ARA) and 20:5n-3 (EPA) in the presence (A) and absence (B) of d5Elo(Eh). The arrow indicates the formation of 22:4n-6 or 22:5n-3, respectively, the products of d5Elo(Eh) activity.




Figure 7 of 13: Gas chromatograph analysis of mature Arabidopsis seeds. Peaks were quantified and listed in the table below. EmiElo91 and EmiElo92 are two selected events transformed with d9Elo(Eh). WT is a non-transformed control. The products of d9Elo(Eh) activity are 20:2 and 20:3n3, which are 10fold increased compared to the levels of the non-transformed control.

Fatty acids	16:0	18:0	18:1	18:2	ALA	18:4	20:0	20:1	20:2	20:3n3	22:1
EmiElo91	4.64	3.43	13.34	23.88	12.64	0.18	2.55	22	11.26	3.73	1.69
EmiElo92	4.21	3.04	12.63	24.63	14.57	0.34	2.28	21.56	10.94	4.02	1.8
WT	6.22	3.21	16.17	27.75	16.63	0.16	2.4	22.2	1.95	0.46	2.15



Figure 8 of 13: Acyl-CoA analysis of mature Arabidopsis seeds of event EmiElo91 transformed with d9Elo(Eh). Col0 is seed material from a non-transformed plant.  $\Delta 9\text{elo}$  is seed material from EmiElo91. The product of d9Elo(Eh) activity is marked with a star (20:2), which is massively increased compared to the control, indicating the functional expression of d9Elo(Eh).




Figure 9 of 13: Gas chromatographic analysis of mature *Arabidopsis* seeds transformed with the construct AP2 (gene combinations d9Elo(Eh)\_d8Des(Eh)\_d5Des(Eh)\_d12Des(Ps)\_o3Des(Pi). Peaks were quantified and listed in the table below. The products of d9Elo(Eh) activity are 20:2 and 20:3n3.

FA	16:0	18:0	18:1c9	18:1c11	18:2	18:3n3	20:0	20:1	20:2	20:3n3
AP2 1	6.5	3.4	12.9	1.8	27.4	13.2	2.0	15.9	7.5	4.5
AP2 5	5.7	3.2	14.9	1.5	26.4	13.1	2.0	17.9	6.8	3.2
AP2 11	5.9	3.2	13.5	1.6	26.6	13.3	2.0	17.1	7.7	4.0
AP2 16	6.5	3.1	11.9	1.7	26.9	13.5	2.0	16.2	7.8	5.0
AP2 17	7.6	3.4	12.2	3.1	30.6	13.8	2.0	14.9	3.8	2.2
AP2 18	6.5	3.3	11.4	1.9	27.2	14.2	2.3	16.7	6.7	3.9
AP2 21	6.6	3.1	10.9	1.8	24.9	15.5	2.2	15.9	7.4	5.7
AP2 22	6.2	3.2	9.7	1.7	25.9	14.4	2.1	15.9	8.7	6.4
AP2 23	6.3	3.1	10.8	1.9	27.1	14.6	2.1	16.0	7.5	5.2
AP2 24	5.9	3.1	10.3	1.6	25.4	14.2	2.1	16.1	9.4	5.9
AP2 25	6.2	3.0	12.0	1.6	26.0	14.8	2.0	17.1	7.7	3.9
AP2 26	6.2	3.3	11.1	1.8	26.6	13.1	2.1	16.0	8.6	5.7
AP2 27	5.8	3.1	9.8	2.0	24.1	12.2	2.2	15.8	9.5	8.6
AP2 28	6.0	3.2	9.8	1.8	25.0	13.0	2.2	15.7	8.9	7.6
AP2 29	6.6	3.3	10.1	2.0	26.5	14.1	2.2	15.8	7.9	5.8
AP2 30	5.6	3.0	10.7	1.5	26.7	13.2	2.0	16.6	10.5	4.7
AP2 31	5.5	3.2	11.2	1.8	24.5	12.6	2.1	16.5	10.6	6.0
AP2 32	5.7	3.2	11.1	1.6	25.5	13.5	2.2	17.4	8.8	4.9
AP2 33	6.5	3.2	9.2	1.9	26.7	15.4	2.3	15.8	7.6	5.5
AP2 34	6.4	3.3	10.1	2.0	26.1	13.7	2.3	16.7	8.0	5.7
AP2 42	5.7	3.4	13.1	1.8	25.6	12.8	2.3	18.0	7.6	4.2
AP2 46	6.3	3.3	10.3	2.4	27.2	12.8	2.4	17.0	6.7	5.0
AP2 47	5.7	3.5	13.3	2.0	26.7	12.4	2.3	17.5	7.3	3.8
AP2 48	5.6	3.5	12.0	2.0	25.5	12.2	2.3	16.6	8.9	5.7
AP2 49	6.0	2.9	12.9	1.6	26.7	14.2	2.2	18.4	6.5	3.2
AP2 50	5.8	3.0	10.8	2.5	26.6	13.9	2.2	17.1	6.7	5.1
average	6.1	3.2	11.4	1.9	26.3	13.6	2.2	16.6	7.9	5.1
std	0.5	0.2	1.4	0.3	1.2	0.9	0.1	0.8	1.4	1.4

Figure 10 of 13:

FA	16:0	18:0	18:1	18:1 Δ9	18:1 Δ11	18:2 Δ9,12	GLA 18:3n6	18:4	20:0	20:1 Δ11	20:2 Δ11,14	DHGLA 20:3n6	ARA 20:4n3	20:3n3	ETA 20:4n3	EPA 20:5n3	22:0	22:1	DPA	DHA 22:5	22:6
OstELO5EmD4_1	6	3	10	1.9	2.5	0.7	0.3	2	17.9	2.0	0	0	1	0	1	2	0	1	1	2.3	
OstELO5EmD4_2	7	3	13	2.4	27.4	2.7	1.5	2	16.0	1.1	0	0	0	0	0	1	0	1	0	1.2	
OstELO5EmD4_3	6	3	10	2.3	26.2	1.9	1.0	2	15.5	1.4	0	0	1	0	0	3	0	1	0	2.7	
OstELO5EmD4_4	7	3	8.7	2.8	23.7	5.7	3.2	2	15.2	1.1	0	0	0	1	0	2	0	1	0	1.6	
OstELO5EmD4_5	7	3	10	2.4	24.6	3.2	1.8	2	16.3	1.1	0	0	0	0	1	2	0	1	0	2.0	
OstELO5EmD4_6	8	3	9.0	2.2	23.2	6.1	4.0	2	15.7	1.3	0	0	0	0	0	1	0	1	0	1.4	
OstELO5EmD4_7	7	3	7.9	2.2	24.0	1.0	0.7	2	14.6	2.1	0	0	1	1	1	3	0	1	1	4.7	
OstELO5EmD4_8	9	3	10	1.7	26.8	1.1	0.5	1	13.7	1.6	0.7	0.7	1	0	1	2	0	1	0	2	
OstELO5EmD4_14	8	3	12	1.6	25.4	4.8	2	2	15.7	1.2	0.1	0	0	0	0	0	0	1	0	0	
OstELO5EmD4_15	9	3	11	1.7	25.8	3.9	2	1	13.6	1.2	0.1	0	0	0	0	2	0	0	0	1	
OstELO5EmD4_16	7	3	9.7	2.2	24.3	6.7	3	2	14.2	0.9	0.1	0	0	1	0	1	0	1	0	1	
OstELO5EmD4_18	7	3	10	2.1	24.4	5.1	3	2	15.7	1.3	0.5	0	0	0	0	2	0	1	0	1	
OstELO5EmD4_20	6	3	8.8	2.0	24.4	1.0	0	2	16.5	1.8	0.7	1	1	1	1	3	0	1	0	3	
OstELO5EmD4_21	5	3	11	2.0	26.4	0.6	0	2	18.3	1.9	0.7	1	0	1	1	2	0	1	0	1	
OstELO5EmD4_25	6	3	10	1.9	24.9	2.1	1	2	17.4	1.9	0.5	0	0	0	0	1	0	1	0	1	
OstELO5EmD4_26	6	3	11	2.1	25.5	5.0	3	1	15.6	1.3	0.2	0	0	1	0	1	0	1	0	1	
OstELO5EmD4_27	6	3	12	1.9	24.2	4.2	2	2	16.4	1.3	0.4	0	0	0	0	2	0	1	0	1	
OstELO5EmD4_28	8	3	8.2	2.4	23.5	3.7	2	2	14.2	1.1	0.8	1	1	1	1	4	0	1	1	2	
OstELO5EmD4_101	6	3	8.2	2.1	23.1	1.4	0	3	17.4	1.7	0.6	1	1	1	1	3	0	1	1	3	
OstELO5EmD4_103	6	3	10	2.2	26.7	1.1	0	2	17.1	2.0	0.2	0	0	1	0	1	0	1	0	1	
OstELO5EmD4_104	6	3	9.2	2.4	25.3	1.2	0	2	15.9	1.6	0.7	1	1	1	1	4	0	1	1	3	
OstELO5EmD4_105	6	3	9.3	2.0	24.9	3.5	1	2	15.1	1.4	0.7	2	0	0	0	3	0	1	1	2	
OstELO5EmD4_106	7	3	7.2	2.3	19.5	3.1	2	2	13.8	1.3	0.4	1	1	1	1	7	0	1	1	4	
OstELO5EmD4_109	7	3	7.5	2.4	21.7	1.8	1	2	14.7	1.7	0.3	1	1	1	0	5	0	1	2	4	
OstELO5EmD4_111	7	3	10	2.2	25.6	2.5	1	2	14.5	1.3	0.3	0	0	0	0	3	0	1	1	2	
OstELO5EmD4_112	7	3	6.3	3.5	20.1	2.1	1	2	13.3	1.7	1.0	0	0	1	2	6	0	1	1	3	
OstELO5EmD4_115	8	3	7.8	2.7	24.1	2.6	1	2	13.8	1.5	0.3	1	1	1	0	4	0	1	2	3	
OstELO5EmD4_116	7	3	10	2.5	24.8	5.7	3	2	13.7	0	0.2	0	0	1	0	2	0	1	0	1	
OstELO5EmD4_117	6	3	9	2	26	1	0	2	15.2	1	0.9	1	1	1.3	1	3	0	1	1	3	
OstELO5EmD4_118	6	3	10.4	2	23	3	2	2	15.3	1	1.2	0	0	1.1	2	3	0	1	1	1	
OstELO5EmD4_119	7	3	9	2	25	4	2	2	14.2	1	0.2	0	0	1.0	0	2	0	0	0	2	
OstELO5EmD4_120	6	3	10.0	2	27	1	0	2	15.1	1	0.8	1	1	0.9	1	2	0	1	1	2	
average	7	3	9	2.2	24	3.0	1	2	15.4	1	0	0	0	1.0	0	3	0.4	1.3	1	2	
std	0	0	1	0.4	1	1.8	1	0	1	0	0	0	0	0.2	0	1	0.1	0.2	0	1	

Figure 11 of 13: Gas chromatographic analysis of mature *Arabidopsis* seeds transformed with different constructs. Values are generated from the best individual *Arabidopsis* line. Following constructs are compared: BBC, OstELO5TcD4 and OstELO5EmD4. The construct with d4Des(Eh) delivered highest levels of DHA and highest ratio of DHA:DPA (2,9).

Line	16:	18:	18:	18: n-6	GL	18: n-3	SD	20: n-9	A	EP	DP	DH
BBC	3.9	4.1	9.9	21.0	1.0	10.5	0.6	22.	1.6	10.	-	-
OstELO TcD4	6.2	3.8	6.2	21.8	2.0	11.6	1.1	14.	1.5	5.4	4.4	3.8
OstELO EmD4	7.1	3.3	7.9	24.0	1.0	13.6	0.7	14.	1.2	3.8	1.6	4.7
BBC-OTelo5-EhD4-2	6.9	3.2	7.1	21.5	1.2	11.8	0.6	13.	1.3	4.6	8.5	0.2

Figure 12 of 13: Gas chromatographic analysis of mature *Arabidopsis* seeds transformed with the construct EmELO5Tcd4. The production of 22:5 and 22:6 demonstrate the activity of d5Elo(Eh) in seeds.

	16:0	18:0	18:1	18:1	18:2	GLA	ALA	18:4	20:0	20:1	20:1	20:2	DHGLA	ARA	20:3n3	20:4n3	20:5n3	EPA	22:1	DPA	DHA
FA			$\Delta 9$	$\Delta 11$	$\Delta 9,12$	18:3n6	18:3n3			$\Delta 11$	$\Delta 13$	$\Delta 11,14$	20:3n6	20:4n3	20:3n3	20:4n3	20:5n3				
EmELO5Tcd4_1	6.5	3.4	10.5	2.5	24.2	3.3	12.7	2.0	2.3	17.5	3.3	1.4	0.2	0.5	1.0	0.5	4.1	1.7	0.2	0.2	
EmELO5Tcd4_2	6.4	3.6	10.0	1.9	25.2	3.1	13.1	1.8	2.3	16.1	2.4	1.4	0.2	1.0	1.2	0.5	5.6	1.3	0.2	0.2	
EmELO5Tcd4_3	6.1	3.5	14.3	1.8	27.0	0.0	16.8	0.0	2.5	18.4	2.3	3.6	0.0	0.0	1.6	0.0	0.0	1.6	0.0	0.0	
EmELO5Tcd4_4	7.0	3.6	10.1	2.4	26.2	3.4	11.9	2.1	2.3	15.9	2.9	1.1	0.3	0.6	1.2	0.5	4.3	1.2	0.3	0.4	
EmELO5Tcd4_5	6.3	3.5	8.3	2.1	23.1	1.5	12.0	0.8	2.2	15.6	2.7	1.4	1.1	1.3	1.4	2.5	8.7	1.2	0.2	0.3	
EmELO5Tcd4_6	6.1	3.6	10.6	2.1	24.9	1.6	13.3	0.8	2.3	16.6	2.5	1.8	0.9	0.7	1.3	1.7	4.8	1.3	0.2	0.2	
EmELO5Tcd4_7	5.5	3.4	10.3	1.8	25.7	0.5	12.6	0.2	2.5	18.2	2.3	2.0	0.9	1.4	1.1	1.4	6.0	1.5	0.2	0.3	
average	6.3	3.5	10.6	2.1	25.2	1.9	13.2	1.1	2.4	16.9	2.6	1.8	0.5	0.8	1.2	1.0	4.8	1.4	0.2	0.2	
std	0.5	0.1	1.8	0.3	1.3	1.4	1.7	0.9	0.1	1.1	0.4	0.9	0.4	0.5	0.2	0.9	2.6	0.2	0.1	0.1	

Figure 13 of 13: Gas chromatographic analysis of mature *Arabidopsis* seeds transformed with four different constructs (A). The production of ARA and EPA demonstrates the activity of d5Des(Eh) in seeds (B). It could be shown that d5Des(Eh) has a preference for n-6 fatty acids (ARA).

A)

Constructs	
Borage Δ6=	Glycinin promoter + Borage Δ6
Borage Δ6: PpELO6=	Glycinin promoter + Borage Δ6 Glycinin promoter + PpELO6
Ostr Δ6=	Glycinin promoter + Ostr Δ6
Ostr Δ6: PpELO6=	Glycinin promoter + Ostr Δ6 Glycinin promoter + PpELO6

B)

Construct	16:0	16:1	16:2	GLA	18:3	SDA	20:1	20:2	DHGLA	ARA	20:3+3	20:4+3	EPA	
Bd6+elo6	6.2	4.1	17.3	24.8	4.4	9.5	1.8	15.4	4.3	2.6	-	1.3	0.9	-
Od6+elo6	6.6	4.3	12.9	26.3	0.9	9.5	0.4	11.4	1.7	9.1	-	0.5	4.2	-
B6/elo6/Ed5	10.8	4.3	9.3	26.0	4.4	10.7	2.0	10.6	5.0	2.0	1.4	1.7	1.0	0.5
O6/elo6/Ed5	10.3	3.9	9.5	23.9	1.0	10.5	0.0	11.5	2.3	5.2	4.7	0.5	3.9	1.4

## NOVEL FATTY ACID DESATURASES AND ELONGASES AND USES THEREOF

### RELATED APPLICATIONS

**[0001]** This application is a continuation of U.S. application Ser. No. 13/384,277 filed Jan. 16, 2012, which is a national stage application (under 35 U.S.C. §371) of PCT/EP2010/060178, filed Jul. 15, 2010, which claims benefit of U.S. Provisional Application No. 61/226,301, filed Jul. 17, 2009, and European Application No. 09165752.8, filed Jul. 17, 2009. The entire contents of each of these applications are hereby incorporated by reference herein in their entirety.

### SUBMISSION OF SEQUENCE LISTING

**[0002]** The Sequence Listing associated with this application is filed in electronic format via EFS-Web and hereby incorporated by reference into the specification in its entirety. The name of the text file containing the Sequence Listing is Sequence\_Listing\_074021\_0167\_01. The size of the text file is 58 KB, and the text file was created on Nov. 15, 2016.

**[0003]** The invention in principle pertains to the field of recombinant manufacture of fatty acids. It provides nucleic acid molecules which encode novel fatty acid desaturases and elongases. The invention also provides recombinant expression vectors containing desaturase and elongase nucleic acid molecules, host cells into which the expression vectors have been introduced, and methods for large-scale production of long chain polyunsaturated fatty acids (LCPUFAs), e.g. arachidonic acid (ARA), eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA).

**[0004]** Fatty acids are carboxylic acids with long-chain hydrocarbon side groups that play a fundamental role in many biological processes. Fatty acids are rarely found free in nature but, rather, occur in esterified form as the major component of lipids. As such, lipids/fatty acids are sources of energy (e.g.,  $\beta$ -oxidation). In addition, lipids/fatty acids are an integral part of cell membranes and, therefore, are indispensable for processing biological or biochemical information.

**[0005]** Fatty acids can be divided into two groups: saturated fatty acids formed of single carbon bonds and the unsaturated fatty acids which contain one or more carbon double bonds in *cis*-configuration. Unsaturated fatty acids are produced by terminal desaturases that belong to the class of nonheme-iron enzymes. Each of these enzymes are part of an electron-transport system that contains two other proteins, namely cytochrome  $b_5$  and NADH-cytochrome  $b_5$  reductase. Specifically, such enzymes catalyze the formation of double bonds between the carbon atoms of a fatty acid molecule, for example, by catalyzing the oxygen-dependent dehydrogenation of fatty acids (Sperling et al., 2003). Human and other mammals have a limited spectrum of desaturases that are required for the formation of particular double bonds in unsaturated fatty acids and thus, have a limited capacity for synthesizing essential fatty acids, e.g., long chain polyunsaturated fatty acids (LCPUFAs). Thus, humans have to take up some fatty acids through their diet. Such essential fatty acids include, for example, linoleic acid (C18:2), linolenic acid (C18:3). In contrast, insects, microorganisms and plants are able to synthesize a much larger

variety of unsaturated fatty acids and their derivatives. Indeed, the biosynthesis of fatty acids is a major activity of plants and microorganisms.

**[0006]** Long chain polyunsaturated fatty acids (LCPUFAs) such as docosahexaenoic acid (DHA, 22:6(4,7,10,13,16,19)) are essential components of cell membranes of various tissues and organelles in mammals (nerve, retina, brain and immune cells). For example, over 30% of fatty acids in brain phospholipid are 22:6 (n-3) and 20:4 (n-6) (Crawford, M. A., et al, (1997) *Am. J. Clin. Nutr.* 66:1032S-1041S). In retina, DHA accounts for more than 60% of the total fatty acids in the rod outer segment, the photosensitive part of the photoreceptor cell (Giusto, N. M., et al. (2000) *Prog. Lipid Res.* 39:315-391). Clinical studies have shown that DHA is essential for the growth and development of the brain in infants, and for maintenance of normal brain function in adults (Martinetz, M. (1992) *J. Pediatr.* 120:S129-S138). DHA also has significant effects on photoreceptor function involved in the signal transduction process, rhodopsin activation, and rod and cone development (Giusto, N. M., et al. (2000) *Prog. Lipid Res.* 39:315-391). In addition, some positive effects of DHA were also found on diseases such as hypertension, arthritis, atherosclerosis, depression, thrombosis and cancers (Horrocks, L. A. and Yeo, Y. K. (1999) *Pharmacol. Res.* 40:211-215). Therefore, appropriate dietary supply of the fatty acid is important for human health. Because such fatty acids cannot be efficiently synthesized by infants, young children and senior citizens, it is particularly important for these individuals to adequately intake these fatty acids from the diet (Spector, A. A. (1999) *Lipids* 34:S1-S3).

**[0007]** Currently the major sources of DHA are oils from fish and algae. Fish oil is a major and traditional source for this fatty acid, however, it is usually oxidized by the time it is sold. In addition, the supply of fish oil is highly variable, particularly in view of the shrinking fish populations. Moreover, the algal source of oil is expensive due to low yield and the high costs of extraction.

**[0008]** EPA and ARA are both 45 essential fatty acids. They form a unique class of food and feed constituents for humans and animals. EPA belongs to the n-3 series with five double bonds in the acyl chain. EPA is found in marine food and is abundant in oily fish from North Atlantic. ARA belongs to the n-6 series with four double bonds. The lack of a double bond in the  $\omega$ -3 position confers on ARA different properties than those found in EPA. The eicosanoids produced from AA have strong inflammatory and platelet aggregating properties, whereas those derived from EPA have anti-inflammatory and anti-platelet aggregating properties. ARA can be obtained from some foods such as meat, fish and eggs, but the concentration is low.

**[0009]** Gamma-linolenic acid (GLA) is another essential fatty acid found in mammals. GLA is the metabolic intermediate for very long chain n-6 fatty acids and for various active molecules. In mammals, formation of long chain polyunsaturated fatty acids is rate-limited by  $\Delta$ 6 desaturation. Many physiological and pathological conditions such as aging, stress, diabetes, eczema, and some infections have been shown to depress the  $\Delta$ 6 desaturation step. In addition, GLA is readily catabolized from the oxidation and rapid cell division associated with certain disorders, e.g., cancer or inflammation. Therefore, dietary supplementation with GLA can reduce the risks of these disorders. Clinical studies have shown that dietary supplementation with GLA is effective in

treating some pathological conditions such as atopic eczema, premenstrual syndrome, diabetes, hypercholesterolemia, and inflammatory and cardiovascular disorders.

**[0010]** A large number of beneficial health effects have been shown for DHA or mixtures of EPA/DHA. DHA is a n-3 very long chain fatty acid with six double bonds.

**[0011]** Although biotechnology offers an attractive route for the production of specialty fatty acids, current techniques fail to provide an efficient means for the large scale production of unsaturated fatty acids. Accordingly, there exists a need for an improved and efficient method of producing unsaturated fatty acids, such as DHA, EPA and ARA.

**[0012]** Thus, the present invention relates to a polynucleotide comprising a nucleic acid sequence elected from the group consisting of:

**[0013]** a) a nucleic acid sequence having a nucleotide sequence as shown in SEQ ID NOs: 1, 3, 5, 7 or 9;

**[0014]** b) a nucleic acid sequence encoding a polypeptide having an amino acid sequence as shown in SEQ ID NOs: 2, 4, 6, 8 or 10;

**[0015]** c) a nucleic acid sequence being at least 70% identical to the nucleic acid sequence of a) or b), wherein said nucleic acid sequence encodes a polypeptide having desaturase or elongase activity;

**[0016]** d) a nucleic acid sequence encoding a polypeptide having desaturase or elongase activity and having an amino acid sequence which is at least 82% identical to the amino acid sequence of any one of a) to c); and

**[0017]** e) a nucleic acid sequence which is capable of hybridizing under stringent conditions to any one of a) to d), wherein said nucleic acid sequence encodes a polypeptide having desaturase or elongase activity.

**[0018]** The term "polynucleotide" as used in accordance with the present invention relates to a polynucleotide comprising a nucleic acid sequence which encodes a polypeptide having desaturase or elongase activity. Preferably, the polypeptide encoded by the polynucleotide of the present invention having desaturase or elongase activity upon expression in a plant shall be capable of increasing the amount of PUFA and, in particular, LCPUFA in, e.g., seed oils or the entire plant or parts thereof. Such an increase is, preferably, statistically significant when compared to a LCPUFA producing transgenic control plant which expresses the present state of the art set of desaturases and elongases required for LCPUFA synthesis but does not express the polynucleotide of the present invention. Whether an increase is significant can be determined by statistical tests well known in the art including, e.g., Student's t-test. More preferably, the increase is an increase of the amount of triglycerides containing LCPUFA of at least 5%, at least 10%, at least 15%, at least 20% or at least 30% compared to the said control. Preferably, the LCPUFA referred to before is a polyunsaturated fatty acid having a C-20 or C-22 fatty acid body, more preferably, ARA, EPA or DHA. Suitable assays for measuring the activities mentioned before are described in the accompanying Examples.

**[0019]** The term "desaturase" or "elongase" as used herein refers to the activity of a desaturase, introducing a double bond into the carbon chain of a fatty acid, preferably into fatty acids with 18, 20 or 22 carbon molecules, or an elongase, introducing two carbon molecules into the carbon chain of a fatty acid, preferably into fatty acids with 18, 20 or 22 carbon molecules

**[0020]** More preferably, polynucleotides having a nucleic acid sequence as shown in SEQ ID NOs: 1, 3, 5, 7 or 9 encoding polypeptides having amino acid sequences as shown in SEQ ID NOs: 2, 4, 6, 8 or 10 or variants thereof, preferably, exhibit desaturase or elongase activity.

**[0021]** Polynucleotides encoding a polypeptide having desaturase or elongase activity as specified above has been obtained in accordance with the present invention, preferably, from *Emiliana huxleyi*. However, orthologs, paralogs or other homologs may be identified from other species. Preferably, they are obtained from plants such as algae, for example *Isochrysis*, *Mantoniella*, *Ostreococcus* or *Cryptocodinium*, algae/diatoms such as *Phaeodactylum*, *Thalassiosira* or *Thraustochytrium*, mosses such as *Physcomitrella* or *Ceratodon*, or higher plants such as the Primulaceae such as *Aleuritia*, *Calendula stellata*, *Osteospermum spinescens* or *Osteospermum hyoseroides*, microorganisms such as fungi, such as *Aspergillus*, *Phytophthora*, *Entomophthora*, *Mucor* or *Mortierella*, bacteria such as *Shewanella*, yeasts or animals. Preferred animals are nematodes such as *Caenorhabditis*, insects or vertebrates. Among the vertebrates, the nucleic acid molecules may, preferably, be derived from *Euteleostomi*, *Actinopterygii*; *Neopterygii*; *Teleostei*; *Euteleostei*, *Protacanthopterygii*, *Salmoniformes*; *Salmonidae* or *Oncorhynchus*, more preferably, from the order of the Salmoniformes, most preferably, the family of the Salmonidae, such as the genus *Salmo*, for example from the genera and species *Oncorhynchus mykiss*, *Trutta trutta* or *Salmo trutta fario*. Moreover, the nucleic acid molecules may be obtained from the diatoms such as the genera *Thalassiosira* or *Phaeodactylum*.

**[0022]** Thus, the term "polynucleotide" as used in accordance with the present invention further encompasses variants of the aforementioned specific polynucleotides representing orthologs, paralogs or other homologs of the polynucleotide of the present invention. Moreover, variants of the polynucleotide of the present invention also include artificially generated muteins. Said muteins include, e.g., enzymes which are generated by mutagenesis techniques and which exhibit improved or altered substrate specificity, or codon optimized polynucleotides. The polynucleotide variants, preferably, comprise a nucleic acid sequence characterized in that the sequence can be derived from the aforementioned specific nucleic acid sequences shown in any one of SEQ ID NOs: 1, 3, 5, 7 or 9 or by a polynucleotide encoding a polypeptide having an amino acid sequence as shown in any one of SEQ ID NOs: 2, 4, 6, 8 or 10 by at least one nucleotide substitution, addition and/or deletion, whereby the variant nucleic acid sequence shall still encode a polypeptide having a desaturase or elongase activity as specified above. Variants also encompass polynucleotides comprising a nucleic acid sequence which is capable of hybridizing to the aforementioned specific nucleic acid sequences, preferably, under stringent hybridization conditions. These stringent conditions are known to the skilled worker and can be found in Current Protocols in Molecular Biology, John Wiley & Sons, N. Y. (1989), 6.3.1-6.3.6. A preferred example for stringent hybridization conditions are hybridization conditions in 6× sodium chloride/sodium citrate (=SSC) at approximately 45° C., followed by one or more wash steps in 0.2×SSC, 0.1% SDS at 50 to 65° C. The skilled worker knows that these hybridization conditions differ depending on the type of nucleic acid and, for example when organic solvents are present, with regard to the tem-



perature and concentration of the buffer. For example, under "standard hybridization conditions" the temperature differs depending on the type of nucleic acid between 42° C. and 58° C. in aqueous buffer with a concentration of 0.1 to 5×SSC (pH 7.2). If organic solvent is present in the above-mentioned buffer, for example 50% formamide, the temperature under standard conditions is approximately 42° C. The hybridization conditions for DNA:DNA hybrids are, preferably, 0.1×SSC and 20° C. to 45° C., preferably between 30° C. and 45° C. The hybridization conditions for DNA:RNA hybrids are, preferably, 0.1×SSC and 30° C. to 55° C., preferably between 45° C. and 55° C. The above-mentioned hybridization temperatures are determined for example for a nucleic acid with approximately 100 bp (=base pairs) in length and a G+C content of 50% in the absence of formamide. The skilled worker knows how to determine the hybridization conditions required by referring to textbooks such as the textbook mentioned above, or the following textbooks: Sambrook et al., "Molecular Cloning", Cold Spring Harbor Laboratory, 1989; Hames and Higgins (Ed.) 1985, "Nucleic Acids Hybridization: A Practical Approach", IRL Press at Oxford University Press, Oxford; Brown (Ed.) 1991, "Essential Molecular Biology: A Practical Approach", IRL Press at Oxford University Press, Oxford. Alternatively, polynucleotide variants are obtainable by PCR-based techniques such as mixed oligonucleotide primer-based amplification of DNA, i.e. using degenerated primers against conserved domains of the polypeptides of the present invention. Conserved domains of the polypeptide of the present invention may be identified by a sequence comparison of the nucleic acid sequences of the polynucleotides or the amino acid sequences of the polypeptides of the present invention. Oligonucleotides suitable as PCR primers as well as suitable PCR conditions are described in the accompanying Examples. As a template, DNA or cDNA from bacteria, fungi, plants or animals may be used. Further, variants include polynucleotides comprising nucleic acid sequences which are at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the nucleic acid sequences shown in any one of SEQ ID NOs: 1, 3, 5, 7 or 9, preferably, encoding polypeptides retaining a desaturase or elongase activity as specified above. Moreover, also encompassed are polynucleotides which comprise nucleic acid sequences encoding a polypeptide having an amino acid sequences which are at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the amino acid sequences shown in any one of SEQ ID NOs: 2, 4, 6, 8 or 10, wherein the polypeptide, preferably, retains desaturase or elongase activity as specified above. The percent identity values are, preferably, calculated over the entire amino acid or nucleic acid sequence region. A series of programs based on a variety of algorithms is available to the skilled worker for comparing different sequences. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch algorithm (Needleman 1970, J. Mol. Biol. (48):444-453) which has been incorporated into the needle program in the EMBOSS software package (*EMBOSS: The European Molecular Biology Open Software Suite*, Rice, P., Longden, I., and Bleasby, A, Trends in Genetics 16(6), 276-277, 2000), using either a BLOSUM

45 or PAM250 scoring matrix for distantly related proteins, or either a BLOSUM 62 or PAM160 scoring matrix for closer related proteins, and a gap opening penalty of 16, 14, 12, 10, 8, 6, or 4 and a gap extension penalty of 0.5, 1, 2, 3, 4, 5, or 6. Guides for local installation of the EMBOSS package as well as links to WEB-Services can be found at [emboss.sourceforge.net](http://emboss.sourceforge.net). A preferred, non-limiting example of parameters to be used for aligning two amino acid sequences using the needle program are the default parameters, including the EBLOSUM62 scoring matrix, a gap opening penalty of 10 and a gap extension penalty of 0.5. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the needle program in the EMBOSS software package (*EMBOSS: The European Molecular Biology Open Software Suite*, Rice, P., Longden, I., and Bleasby, A, Trends in Genetics 16(6), 276-277, 2000), using the EDNAFULL scoring matrix and a gap opening penalty of 16, 14, 12, 10, 8, 6, or 4 and a gap extension penalty of 0.5, 1, 2, 3, 4, 5, or 6. A preferred, non-limiting example of parameters to be used in conjunction for aligning two amino acid sequences using the needle program are the default parameters, including the EDNAFULL scoring matrix, a gap opening penalty of 10 and a gap extension penalty of 0.5. The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the BLAST series of programs (version 2.2) of Altschul et al. (Altschul 1990, J. Mol. Biol. 215:403-10). BLAST using acyltransferase nucleic acid sequences of the invention as query sequence can be performed with the BLASTn, BLASTx or the tBLASTx program using default parameters to obtain either nucleotide sequences (BLASTn, tBLASTx) or amino acid sequences (BLASTx) homologous to acyltransferase sequences of the invention. BLAST using acyltransferase protein sequences of the invention as query sequence can be performed with the BLASTp or the tBLASTn program using default parameters to obtain either amino acid sequences (BLASTp) or nucleic acid sequences (tBLASTn) homologous to acyltransferase sequences of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST using default parameters can be utilized as described in Altschul et al. (Altschul 1997, Nucleic Acids Res. 25(17):3389-3402).

TABLE 1

Relation of sequence types of query and hit sequences for various BLAST programs				
Input query sequence	Converted Query	Algorithm	Converted Hit	Actual Database
DNA		BLASTn		DNA
PRT		BLASTp		PRT
DNA	PRT	BLASTx		PRT
PRT		tBLASTn	PRT	DNA
DNA	PRT	tBLASTx	PRT	DNA

**[0023]** A polynucleotide comprising a fragment of any of the aforementioned nucleic acid sequences is also encompassed as a polynucleotide of the present invention. The fragment shall encode a polypeptide which still has desaturase and elongase activity as specified above. Accordingly, the polypeptide may comprise or consist of the domains of

the polypeptide of the present invention conferring the said biological activity. A fragment as meant herein, preferably, comprises at least 50, at least 100, at least 250 or at least 500 consecutive nucleotides of any one of the aforementioned nucleic acid sequences or encodes an amino acid sequence comprising at least 20, at least 30, at least 50, at least 80, at least 100 or at least 150 consecutive amino acids of any one of the aforementioned amino acid sequences.

**[0024]** The variant polynucleotides or fragments referred to above, preferably, encode polypeptides retaining desaturase or elongase activity to a significant extent, preferably, at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90% of the desaturase and elongase activity exhibited by any of the polypeptide shown in any one of SEQ ID NOs: 2, 4, 6, 8 or 10. The activity may be tested as described in the accompanying Examples.

**[0025]** The polynucleotides of the present invention either essentially consist of the aforementioned nucleic acid sequences or comprise the aforementioned nucleic acid sequences. Thus, they may contain further nucleic acid sequences as well. Preferably, the polynucleotide of the present invention may comprise in addition to an open reading frame further untranslated sequence at the 3' and at the 5' terminus of the coding gene region: at least 500, preferably 200, more preferably 100 nucleotides of the sequence upstream of the 5' terminus of the coding region and at least 100, preferably 50, more preferably 20 nucleotides of the sequence downstream of the 3' terminus of the coding gene region. Furthermore, the polynucleotides of the present invention may encode fusion proteins wherein one partner of the fusion protein is a polypeptide being encoded by a nucleic acid sequence recited above. Such fusion proteins may comprise as additional part other enzymes of the fatty acid or PUFA biosynthesis pathways, polypeptides for monitoring expression (e.g., green, yellow, blue or red fluorescent proteins, alkaline phosphatase and the like) or so called "tags" which may serve as a detectable marker or as an auxiliary measure for purification purposes. Tags for the different purposes are well known in the art and comprise FLAG-tags, 6-histidine-tags, MYC-tags and the like.

**[0026]** The polynucleotide of the present invention shall be provided, preferably, either as an isolated polynucleotide (i.e. purified or at least isolated from its natural context such as its natural gene locus) or in genetically modified or exogenously (i.e. artificially) manipulated form. An isolated polynucleotide can, for example, comprise less than approximately 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in the genomic DNA of the cell from which the nucleic acid is derived. The polynucleotide, preferably, is provided in the form of double or single stranded molecule. It will be understood that the present invention by referring to any of the aforementioned polynucleotides of the invention also refers to complementary or reverse complementary strands of the specific sequences or variants thereof referred to before. The polynucleotide encompasses DNA, including cDNA and genomic DNA, or RNA polynucleotides.

**[0027]** However, the present invention also pertains to polynucleotide variants which are derived from the polynucleotides of the present invention and are capable of interfering with the transcription or translation of the polynucleotides of the present invention. Such variant poly-

nucleotides include anti-sense nucleic acids, ribozymes, siRNA molecules, morpholino nucleic acids (phosphorodiamidate morpholino oligos), triple-helix forming oligonucleotides, inhibitory oligonucleotides, or micro RNA molecules all of which shall specifically recognize the polynucleotide of the invention due to the presence of complementary or substantially complementary sequences. These techniques are well known to the skilled artisan. Suitable variant polynucleotides of the aforementioned kind can be readily designed based on the structure of the polynucleotides of this invention.

**[0028]** Moreover, comprised are also chemically modified polynucleotides including naturally occurring modified polynucleotides such as glycosylated or methylated polynucleotides or artificial modified ones such as biotinylated polynucleotides.

**[0029]** In the studies underlying the present invention, advantageously, polynucleotides where identified encoding desaturase and elongases from *Emiliana huxleyi*. In particular, the *Emiliana huxleyi* desaturases  $\Delta 4\text{Des(Eh)}$ ,  $\Delta 8\text{Des(Eh)}$  and  $\Delta 5\text{Des(Eh)}$  and elongases  $\Delta 9\text{Elo(Eh)}$  and  $\Delta 5\text{Elo(Eh)}$  have been identified. Each of the desaturases are capable of introducing a double bond into fatty acids. For example, the expression of the  $\Delta 8\text{Des(Eh)}$  leads to introduction of a double bond at position eight into C20:2n-6 fatty acid. The polynucleotides of the present invention are particularly suitable for the recombinant manufacture of LCPUFAs and, in particular, ARA, EPA and/or DHA.

**[0030]** In a preferred embodiment of the polynucleotide of the present invention, said polynucleotide further comprises an expression control sequence operatively linked to the said nucleic acid sequence.

**[0031]** The term "expression control sequence" as used herein refers to a nucleic acid sequence which is capable of governing, i.e. initiating and controlling, transcription of a nucleic acid sequence of interest, in the present case the nucleic sequences recited above. Such a sequence usually comprises or consists of a promoter or a combination of a promoter and enhancer sequences. Expression of a polynucleotide comprises transcription of the nucleic acid molecule, preferably, into a translatable mRNA. Additional regulatory elements may include transcriptional as well as translational enhancers. The following promoters and expression control sequences may be, preferably, used in an expression vector according to the present invention. The cos, tac, trp, tet, trp-tet, lpp, lac, lpp-lac, lacIq, T7, T5, T3, gal, trc, ara, SP6,  $\lambda$ -PR or  $\lambda$ -PL promoters are, preferably, used in Gram-negative bacteria. For Gram-positive bacteria, promoters amy and SPO2 may be used. From yeast or fungal promoters ADC1, AOX1r, GAL1, MF $\alpha$ , AC, P-60, CYC1, GAPDH, TEF, rp28, ADH are, preferably, used. For animal cell or organism expression, the promoters CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer are preferably used. From plants the promoters CaMV/35S (Franck 1980, Cell 21: 285-294), PRP1 (Ward 1993, Plant. Mol. Biol. 22), SSU, OCS, lib4, usp, STLS1, B33, nos or the ubiquitin or phaseolin promoter. Also preferred in this context are inducible promoters, such as the promoters described in EP 0 388 186 A1 (i.e. a benzylsulfonamide-inducible promoter), Gatz 1992, Plant J. 2:397-404 (i.e. a tetracyclin-inducible promoter), EP 0 335 528 A1 (i.e. a abscisic-acid-inducible promoter) or WO 93/21334 (i.e. a ethanol- or cyclohexenol-inducible promoter). Further suitable plant promoters are the promoter of

cytosolic FBPase or the ST-LSI promoter from potato (Stockhaus 1989, EMBO J. 8, 2445), the phosphoribosyl-pyrophosphate amidotransferase promoter from *Glycine max* (Genbank accession No. U87999) or the node-specific promoter described in EP 0 249 676 A1. Particularly preferred are promoters which enable the expression in tissues which are involved in the biosynthesis of fatty acids. Also particularly preferred are seed-specific promoters such as the USP promoter in accordance with the practice, but also other promoters such as the LeB4, DC3, phaseolin or napin promoters. Further especially preferred promoters are seed-specific promoters which can be used for monocotyledonous or dicotyledonous plants and which are described in U.S. Pat. No. 5,608,152 (napin promoter from oilseed rape), WO 98/45461 (oleosin promoter from *Arabidopsis*, U.S. Pat. No. 5,504,200 (phaseolin promoter from *Phaseolus vulgaris*), WO 91/13980 (Bce4 promoter from *Brassica*), by Baeumlein et al., Plant J., 2, 2, 1992:233-239 (LeB4 promoter from a legume), these promoters being suitable for dicots. The following promoters are suitable for monocots: lpt-2 or lpt-1 promoter from barley (WO 95/15389 and WO 95/23230), hordein promoter from barley and other promoters which are suitable and which are described in WO 99/16890. In principle, it is possible to use all natural promoters together with their regulatory sequences, such as those mentioned above, for the novel process. Likewise, it is possible and advantageous to use synthetic promoters, either additionally or alone, especially when they mediate a seed-specific expression, such as, for example, as described in WO 99/16890. In a particular embodiment, seed-specific promoters are utilized to enhance the production of the desired PUFA or LCPUFA.

**[0032]** The term “operatively linked” as used herein means that the expression control sequence and the nucleic acid of interest are linked so that the expression of the said nucleic acid of interest can be governed by the said expression control sequence, i.e. the expression control sequence shall be functionally linked to the said nucleic acid sequence to be expressed. Accordingly, the expression control sequence and, the nucleic acid sequence to be expressed may be physically linked to each other, e.g., by inserting the expression control sequence at the 5' end of the nucleic acid sequence to be expressed. Alternatively, the expression control sequence and the nucleic acid to be expressed may be merely in physical proximity so that the expression control sequence is capable of governing the expression of at least one nucleic acid sequence of interest. The expression control sequence and the nucleic acid to be expressed are, preferably, separated by not more than 500 bp, 300 bp, 100 bp, 80 bp, 60 bp, 40 bp, 20 bp, 10 bp or 5 bp.

**[0033]** In a further preferred embodiment of the polynucleotide of the present invention, said polynucleotide further comprises a terminator sequence operatively linked to the nucleic acid sequence.

**[0034]** The term “terminator” as used herein refers to a nucleic acid sequence which is capable of terminating transcription. These sequences will cause dissociation of the transcription machinery from the nucleic acid sequence to be transcribed. Preferably, the terminator shall be active in plants and, in particular, in plant seeds. Suitable terminators are known in the art and, preferably, include polyadenylation signals such as the SV40-poly-A site or the tk-poly-A site or one of the plant specific signals indicated in Loke et al.

(Loke 2005, Plant Physiol 138, pp. 1457-1468), downstream of the nucleic acid sequence to be expressed.

**[0035]** The present invention also relates to a vector comprising the polynucleotide of the present invention.

**[0036]** The term “vector”, preferably, encompasses phage, plasmid, viral vectors as well as artificial chromosomes, such as bacterial or yeast artificial chromosomes. Moreover, the term also relates to targeting constructs which allow for random or site-directed integration of the targeting construct into genomic DNA. Such target constructs, preferably, comprise DNA of sufficient length for either homologous or heterologous recombination as described in detail below. The vector encompassing the polynucleotide of the present invention, preferably, further comprises selectable markers for propagation and/or selection in a host. The vector may be incorporated into a host cell by various techniques well known in the art. If introduced into a host cell, the vector may reside in the cytoplasm or may be incorporated into the genome. In the latter case, it is to be understood that the vector may further comprise nucleic acid sequences which allow for homologous recombination or heterologous insertion. Vectors can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. The terms “transformation” and “transfection”, conjugation and transduction, as used in the present context, are intended to comprise a multiplicity of prior-art processes for introducing foreign nucleic acid (for example DNA) into a host cell, including calcium phosphate, rubidium chloride or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, natural competence, carbon-based clusters, chemically mediated transfer, electroporation or particle bombardment. Suitable methods for the transformation or transfection of host cells, including plant cells, can be found in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) and other laboratory manuals, such as Methods in Molecular Biology, 1995, Vol. 44, *Agrobacterium* protocols, Ed.: Gartland and Davey, Humana Press, Totowa, N.J. Alternatively, a plasmid vector may be introduced by heat shock or electroporation techniques. Should the vector be a virus, it may be packaged in vitro using an appropriate packaging cell line prior to application to host cells.

**[0037]** Preferably, the vector referred to herein is suitable as a cloning vector, i.e. replicable in microbial systems. Such vectors ensure efficient cloning in bacteria and, preferably, yeasts or fungi and make possible the stable transformation of plants. Those which must be mentioned are, in particular, various binary and co-integrated vector systems which are suitable for the T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they contain at least the vir genes, which are required for the *Agrobacterium*-mediated transformation, and the sequences which delimit the T-DNA (T-DNA border). These vector systems, preferably, also comprise further cis-regulatory regions such as promoters and terminators and/or selection markers with which suitable transformed host cells or organisms can be identified. While co-integrated vector systems have vir genes and T-DNA sequences arranged on the same vector, binary systems are based on at least two vectors, one of which bears vir genes, but no T-DNA, while a second one bears T-DNA, but no vir gene. As a consequence, the last-mentioned vectors are relatively small, easy to manipulate and can be replicated both in *E. coli* and in *Agrobac-*

*terium*. These binary vectors include vectors from the pBIB-HYG, pPZP, pBecks, pGreen series. Preferably used in accordance with the invention are Bin19, pBI101, pBinAR, pGPTV and pCambia. An overview of binary vectors and their use can be found in Hellens et al, Trends in Plant Science (2000) 5, 446-451. Furthermore, by using appropriate cloning vectors, the polynucleotides can be introduced into host cells or organisms such as plants or animals and, thus, be used in the transformation of plants, such as those which are published, and cited, in: Plant Molecular Biology and Biotechnology (CRC Press, Boca Raton, Fla.), chapter 6/7, pp. 71-119 (1993); F. F. White, Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, 15-38; B. Jené et al., Techniques for Gene Transfer, in: Transgenic Plants, vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press (1993), 128-143; Potrykus 1991, Annu. Rev. Plant Physiol. Plant Molec. Biol. 42, 205-225.

**[0038]** More preferably, the vector of the present invention is an expression vector. In such an expression vector, i.e. a vector which comprises the polynucleotide of the invention having the nucleic acid sequence operatively linked to an expression control sequence (also called "expression cassette") allowing expression in prokaryotic or eukaryotic cells or isolated fractions thereof. Suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (Invitrogen) or pSPORT1 (GIBCO BRL). Further examples of typical fusion expression vectors are pGEX (Pharmacia Biotech Inc; Smith 1988, Gene 67:31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.), where glutathione S-transferase (GST), maltose E-binding protein and protein A, respectively, are fused with the recombinant target protein. Examples of suitable inducible nonfusion *E. coli* expression vectors are, inter alia, pTrc (Amann 1988, Gene 69:301-315) and pET 11d (Studier 1990, Methods in Enzymology 185, 60-89). The target gene expression of the pTrc vector is based on the transcription from a hybrid trp-lac fusion promoter by host RNA polymerase. The target gene expression from the pET 11d vector is based on the transcription of a T7-gn10-lac fusion promoter, which is mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is provided by the host strains BL21 (DE3) or HMS174 (DE3) from a resident  $\lambda$ -prophage which harbors a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter. The skilled worker is familiar with other vectors which are suitable in prokaryotic organisms; these vectors are, for example, in *E. coli*, pLG338, pACYC184, the pBR series such as pBR322, the pUC series such as pUC18 or pUC19, the M113mp series, pKC30, pRep4, pHS1, pHS2, pPLc236, pMBL24, pLG200, pUR290, pIN-III113-B1,  $\lambda$ gt11 or pBdCl, in *Streptomyces* pIJ101, pIJ364, pIJ702 or pIJ361, in *Bacillus* pUB110, pC194 or pBD214, in *Corynebacterium* pSA77 or pAJ667. Examples of vectors for expression in the yeast *S. cerevisiae* comprise pYep Sec1 (Baldari 1987, Embo J. 6:229-234), pMFa (Kurjan 1982, Cell 30:933-943), pJRY88 (Schultz 1987, Gene 54:113-123) and pYES2 (Invitrogen Corporation, San Diego, Calif.). Vectors and processes for the construction of vectors which are suitable for use in other fungi, such as the filamentous fungi, comprise those which are described in detail in: van den Hondel, C. A. M. J. J., & Punt, P. J. (1991) "Gene

transfer systems and vector development for filamentous fungi, in: Applied Molecular Genetics of fungi, J. F. Peberdy et al., Ed., pp. 1-28, Cambridge University Press: Cambridge, or in: More Gene Manipulations in Fungi (J. W. Bennett & L. L. Lasure, Ed., pp. 396-428: Academic Press: San Diego). Further suitable yeast vectors are, for example, pAG-1, YEpl6, YEpl13 or pEMBLYe23. As an alternative, the polynucleotides of the present invention can be also expressed in insect cells using baculovirus expression vectors. Baculovirus vectors which are available for the expression of proteins in cultured insect cells (for example Sf9 cells) comprise the pAc series (Smith 1983, Mol. Cell Biol. 3:2156-2165) and the pVL series (Lucklow 1989, Virology 170:31-39).

**[0039]** The polynucleotide of the present invention can be expressed in single-cell plant cells (such as algae), see Falcitatore 1999, Marine Biotechnology 1 (3):239-251 and the references cited therein, and plant cells from higher plants (for example Spermatophytes, such as arable crops) by using plant expression vectors. Examples of plant expression vectors comprise those which are described in detail in: Becker 1992, Plant Mol. Biol. 20:1195-1197; Bevan 1984, Nucl. Acids Res. 12:8711-8721; Vectors for Gene Transfer in Higher Plants; in: Transgenic Plants, Vol. 1, Engineering and Utilization, Ed.: Kung and R. Wu, Academic Press, 1993, p. 15-38. A plant expression cassette, preferably, comprises regulatory sequences which are capable of controlling the gene expression in plant cells and which are functionally linked so that each sequence can fulfill its function, such as transcriptional termination, for example polyadenylation signals. Preferred polyadenylation signals are those which are derived from *Agrobacterium tumefaciens* T-DNA, such as the gene 3 of the Ti plasmid pTiACH5, which is known as octopine synthase (Gielen 1984, EMBO J. 3, 835) or functional equivalents of these, but all other terminators which are functionally active in plants are also suitable. Since plant gene expression is very often not limited to transcriptional levels, a plant expression cassette preferably comprises other functionally linked sequences such as translation enhancers, for example the overdrive sequence, which comprises the 5'-untranslated tobacco mosaic virus leader sequence, which increases the protein/RNA ratio (Gallie 1987, Nucl. Acids Research 15:8693-8711). As described above, plant gene expression must be functionally linked to a suitable promoter which performs the expression of the gene in a timely, cell-specific or tissue-specific manner. Promoters which can be used are constitutive promoters (Benfey 1989, EMBO J. 8:2195-2202) such as those which are derived from plant viruses such as 35S CAMV (Franck 1980, Cell 21:285-294), 19S CaMV (see U.S. Pat. No. 5,352,605 and WO 84/02913) or plant promoters such as the promoter of the Rubisco small subunit, which is described in U.S. Pat. No. 4,962,028. Other preferred sequences for the use in functional linkage in plant gene expression cassettes are targeting sequences which are required for targeting the gene product into its relevant cell compartment (for a review, see Kermode 1996, Crit. Rev. Plant Sci. 15, 4: 285-423 and references cited therein), for example into the vacuole, the nucleus, all types of plastids, such as amyloplasts, chloroplasts, chromoplasts, the extracellular space, the mitochondria, the endoplasmic reticulum, oil bodies, peroxisomes and other compartments of plant cells. As described above, plant gene expression can also be facilitated via a chemically inducible promoter (for

a review, see Gatz 1997, *Annu. Rev. Plant Physiol. Plant Mol. Biol.*, 48:89-108). Chemically inducible promoters are particularly suitable if it is desired that genes are expressed in a time-specific manner. Examples of such promoters are a salicylic-acid-inducible promoter (WO 95/19443), a tetracyclin-inducible promoter (Gatz 1992, *Plant J.* 2, 397-404) and an ethanol-inducible promoter. Promoters which respond to biotic or abiotic stress conditions are also suitable promoters, for example the pathogen-induced PRP1-gene promoter (Ward 1993, *Plant Mol. Biol.* 22:361-366), the heat-inducible hsp80 promoter from tomato (U.S. Pat. No. 5,187,267), the cold-inducible alpha-amylase promoter from potato (WO 96/12814) or the wound-inducible pinII promoter (EP 0 375 091 A). The promoters which are especially preferred are those which bring about the expression of genes in tissues and organs in which fatty acid, lipid and oil biosynthesis takes place, in seed cells such as the cells of endosperm and of the developing embryo. Suitable promoters are the napin gene promoter from oilseed rape (U.S. Pat. No. 5,608,152), the USP promoter from *Vicia faba* (Baeumlein 1991, *Mol. Gen. Genet.* 225 (3):459-67), the oleosin promoter from *Arabidopsis* (WO 98/45461), the phaseolin promoter from *Phaseolus vulgaris* (U.S. Pat. No. 5,504,200), the Bce4 promoter from *Brassica* (WO 91/13980) or the legumin B4 promoter (LeB4; Baeumlein 1992, *Plant Journal*, 2 (2):233-9), and promoters which bring about the seed-specific expression in monocotyledonous plants such as maize, barley, wheat, rye, rice and the like. Suitable promoters to be taken into consideration are the Ipt2 or Ipt1 gene promoter from barley (WO 95/15389 and WO 95/23230) or those which are described in WO 99/16890 (promoters from the barley hordein gene, the rice glutelin gene, the rice oryzin gene, the rice prolamin gene, the wheat gliadin gene, wheat glutelin gene, the maize zein gene, the oat glutelin gene, the *sorghum* kasirin gene, the rye secalin gene). Likewise, especially suitable are promoters which bring about the plastid-specific expression since plastids are the compartment in which the precursors and some end products of lipid biosynthesis are synthesized. Suitable promoters such as the viral RNA-polymerase promoter, are described in WO 95/16783 and WO 97/06250, and the clpP promoter from *Arabidopsis*, described in WO 99/46394.

**[0040]** The abovementioned vectors are only a small overview of vectors to be used in accordance with the present invention. Further vectors are known to the skilled worker and are described, for example, in: *Cloning Vectors* (Ed., Pouwels, P. N., et al., Elsevier, Amsterdam-New York-Oxford, 1985, ISBN 0 444 904018). For further suitable expression systems for prokaryotic and eukaryotic cells see the chapters 16 and 17 of Sambrook, loc cit.

**[0041]** It follows from the above that, preferably, said vector is an expression vector. More preferably, the said polynucleotide of the present invention is under the control of a seed-specific promoter in the vector of the present invention. A preferred seed-specific promoter as meant herein is selected from the group consisting of Conlinin 1, Conlinin 2, napin, LuFad3, USP, LeB4, Arc, Fae, ACP, LuPXR, and SBP. For details, see, e.g., US 2003-0159174.

**[0042]** Moreover, the present invention relates to a host cell comprising the polynucleotide or the vector of the present invention.

**[0043]** Preferably, said host cell is a plant cell and, more preferably, a plant cell obtained from an oilseed crop. More preferably, said oilseed crop is selected from the group

consisting of flax (*Linum* sp.), rapeseed (*Brassica* sp.), soybean (*Glycine* sp.), sunflower (*Helianthus* sp.), cotton (*Gossypium* sp.), corn (*Zea mays*), olive (*Olea* sp.), safflower (*Carthamus* sp.), cocoa (*Theobroma* cocoa), peanut (*Arachis* sp.), hemp, camelina, *crambe*, oil palm, coconuts, groundnuts, sesame seed, castor bean, *lesquerella*, tallow tree, sheanuts, tungnuts, kapok fruit, poppy seed, jojoba seeds and *perilla*.

**[0044]** Also preferably, said host cell is a microorganism. More preferably, said microorganism is a bacterium, a fungus or algae. More preferably, it is selected from the group consisting of *Candida*, *Cryptococcus*, *Lipomyces*, *Rhodospiridium*, *Yarrowia* and *Schizochytrium*.

**[0045]** Moreover, a host cell according to the present invention may also be an animal cell. Preferably, said animal host cell is a host cell of a fish or a cell line obtained therefrom. More preferably, the fish host cell is from herring, salmon, sardine, redfish, eel, carp, trout, halibut, mackerel, zander or tuna.

**[0046]** Generally, the controlling steps in the production of LCPUFAs, i.e., the long chain unsaturated fatty acid biosynthetic pathway, are catalyzed by membrane-associated fatty acid desaturases and elongases. Plants and most other eukaryotic organisms have specialized desaturase and elongase systems for the introduction of double bonds and the extension of fatty acids beyond C18 atoms. The elongase reactions have several important features in common with the fatty acid synthase complex (FAS). However, the elongase complex is different from the FAS complex as the complex is localized in the cytosol and membrane bound, ACP is not involved and the elongase 3-keto-acyl-CoA-synthase catalyzes the condensation of malonyl-CoA with an acyl primer. The elongase complex consists of four components with different catalytic functions, the keto-acyl-synthase (condensation reaction of malonyl-CoA to acyl-CoA, creation of a 2 C atom longer keto-acyl-CoA fatty acid), the keto-acyl-reductase (reduction of the 3-keto group to a 3-hydroxy-group), the dehydratase (dehydration results in a 3-enoyl-acyl-CoA fatty acid) and the enoyl-CoA-reductase (reduction of the double bond at position 3, release from the complex). For the production of LCPUFAs including ARA, EPA and/or DHA the elongation reactions, beside the desaturation reactions, are essential. Higher plants do not have the necessary enzyme set to produce LCPUFAs (4 or more double bonds, 20 or more C atoms). Therefore the catalytic activities have to be conferred to the plants or plant cells. The polynucleotides of the present invention catalyze the desaturation and elongation activities necessary for the formation of ARA, EPA and/or DHA. By delivering the novel desaturases and elongases increased levels of PUFAs and LCPUFAs are produced.

**[0047]** However, person skilled in the art knows that dependent on the host cell, further, enzymatic activities may be conferred to the host cells, e.g., by recombinant technologies. Accordingly, the present invention, preferably, envisages a host cell which in addition to the polynucleotide of the present invention comprises polynucleotides encoding such desaturases and/or elongases as required depending on the selected host cell. Preferred desaturases and/or elongases which shall be present in the host cell are at least one enzyme selected from the group consisting of:  $\Delta$ -4-desaturase,  $\Delta$ -5-desaturase,  $\Delta$ -5-elongase,  $\Delta$ -6-desaturase, 2-desaturase, 5-desaturase,  $\omega$ 3-desaturase and  $\Delta$ -6-elongase. Especially preferred are the bifunctional d12d15-Desaturases

d12d15Des(Ac) from *Acanthamoeba castellanii* (WO2007042510), d12d15Des(Cp) from *Claviceps purpurea* (WO2008006202) and d12d15Des(Lg)1 from *Lottia gigantea* (WO2009016202), the d12-Desaturases d12Des(Co) from *Calendula officinalis* (WO200185968), d12Des(Lb) from *Laccaria bicolor* (WO2009016202), d12Des(Mb) from *Monosiga brevicollis* (WO2009016202), d12Des(Mg) from *Mycosphaerella graminicola* (WO2009016202), d12Des(Nh) from *Nectria haematococca* (WO2009016202), d12Des(OI) from *Ostreococcus lucimarinus* (WO2008040787), d12Des(Pb) from *Phycomyces blakesleeanus* (WO2009016202), d12Des(Ps) from *Phytophthora sojae* (WO2006100241) and d12Des(Tp) from *Thalassiosira pseudonana* (WO2006069710), the d15-Desaturases d15Des(Hr) from *Helobdella robusta* (WO2009016202), d15Des(Mc) from *Microcoleus chthonoplastes* (WO2009016202), d15Des(Mf) from *Mycosphaerella fijiensis* (WO2009016202), d15Des(Mg) from *Mycosphaerella graminicola* (WO2009016202) and d15Des(Nh)2 from *Nectria haematococca* (WO2009016202), the d4-Desaturases d4Des(Eg) from *Euglena gracilis* (WO2004090123), d4Des(Tc) from *Thraustochytrium* sp. (WO2002026946) and d4Des(Tp) from *Thalassiosira pseudonana* (WO2006069710), the d5-Desaturases d5Des(OI)2 from *Ostreococcus lucimarinus* (WO2008040787), d5Des(Pp) from *Physcomitrella patens* (WO2004057001), d5Des(Pt) from *Phaeodactylum tricornutum* (WO2002057465), d5Des(Tc) from *Thraustochytrium* sp. (WO2002026946), d5Des(Tp) from *Thalassiosira pseudonana* (WO2006069710) and the d6-Desaturases d6Des(Cp) from *Ceratodon purpureus* (WO2000075341), d6Des(OI) from *Ostreococcus lucimarinus* (WO2008040787), d6Des(Ot) from *Ostreococcus tauri* (WO2006069710), d6Des(Pf) from *Primula farinosa* (WO2003072784), d6Des(Pir)\_BO from *Pythium irregulare* (WO2002026946), d6Des(Pir) from *Pythium irregulare* (WO2002026946), d6Des(Plu) from *Primula luteola* (WO2003072784), d6Des(Pp) from *Physcomitrella patens* (WO200102591), d6Des(Pt) from *Phaeodactylum tricornutum* (WO2002057465), d6Des(Pv) from *Primula vialii* (WO2003072784) and d6Des(Tp) from *Thalassiosira pseudonana* (WO2006069710), the d8-Desaturases d8Des(Ac) from *Acanthamoeba castellanii* (EP1790731), d8Des(Eg) from *Euglena gracilis* (WO200034439) and d8Des(Pm) from *Perkinsus marinus* (WO2007093776), the o3-Desaturases o3Des(Pi) from *Phytophthora infestans* (WO2005083053), o3Des(Pir) from *Pythium irregulare* (WO2008022963), o3Des(Pir)2 from *Pythium irregulare* (WO2008022963) and o3Des(Ps) from *Phytophthora sojae* (WO2006100241), the bifunctional d5d6-elongases d5d6Elo(Om)2 from *Oncorhynchus mykiss* (WO2005012316), d5d6Elo(Ta) from *Thraustochytrium aureum* (WO2005012316) and d5d6Elo(Tc) from *Thraustochytrium* sp. (WO2005012316), the d5-elongases d5Elo(At) from *Arabidopsis thaliana* (WO2005012316), d5Elo(At)2 from *Arabidopsis thaliana* (WO2005012316), d5Elo(Ci) from *Ciona intestinalis* (WO2005012316), d5Elo(OI) from *Ostreococcus lucimarinus* (WO2008040787), d5Elo(Ot) from *Ostreococcus tauri* (WO2005012316), d5Elo(Tp) from *Thalassiosira pseudonana* (WO2005012316) and d5Elo(XI) from *Xenopus laevis* (WO2005012316), the d6-elongases d6Elo(OI) from *Ostreococcus lucimarinus* (WO2008040787), d6Elo(Ot) from *Ostreococcus tauri* (WO2005012316), d6Elo(Pi) from *Phytophthora infestans* (WO2003064638), d6Elo(Pir) from *Pythium irregulare*

(WO2009016208), d6Elo(Pp) from *Physcomitrella patens* (WO2001059128), d6Elo(Ps) from *Phytophthora sojae* (WO2006100241), d6Elo(Ps)2 from *Phytophthora sojae* (WO2006100241), d6Elo(Ps)3 from *Phytophthora sojae* (WO2006100241), d6Elo(Pt) from *Phaeodactylum tricornutum* (WO2005012316), d6Elo(Tc) from *Thraustochytrium* sp. (WO2005012316) and d6Elo(Tp) from *Thalassiosira pseudonana* (WO2005012316), the d9-elongases d9Elo(Ig) from *Isochrysis galbana* (WO2002077213), d9Elo(Pm) from *Perkinsus marinus* (WO2007093776) and d9Elo(Ro) from *Rhizopus oryzae* (WO2009016208). Particularly, if the manufacture of ARA is envisaged in higher plants, the enzymes recited in Table 3, below (i.e. additionally a d6-desaturase, d6-elongase, d5-elongase, d5-desaturase, d12-desaturase, and d6-elongase) or enzymes having essentially the same activity may be combined in a host cell. If the manufacture of EPA is envisaged in higher plants, the enzymes recited in Table 4, below (i.e. additionally a d6-desaturase, d6-elongase, d5-desaturase, d12-desaturase, d6-elongase, omega 3-desaturase and d15-desaturase), or enzymes having essentially the same activity may be combined in a host cell. If the manufacture of DHA is envisaged in higher plants, the enzymes recited in Table 5, below (i.e. additionally a d6-desaturase, d6-elongase, d5-desaturase, d12-desaturase, d6-elongase, omega 3-desaturase, d15-desaturase, d5-elongase, and d4-desaturase), or enzymes having essentially the same activity may be combined in a host cell.

**[0048]** The present invention also relates to a cell, preferably a host cell as specified above or a cell of a non-human organism specified elsewhere herein, said cell comprising a polynucleotide which is obtained from the polynucleotide of the present invention by a point mutation, a truncation, an inversion, a deletion, an addition, a substitution and homologous recombination. How to carry out such modifications to a polynucleotide is well known to the skilled artisan and has been described elsewhere in this specification in detail.

**[0049]** The present invention furthermore pertains to a method for the manufacture of a polypeptide encoded by a polynucleotide of any the present invention comprising

**[0050]** a) cultivating the host cell of the invention under conditions which allow for the production of the said polypeptide; and

**[0051]** b) obtaining the polypeptide from the host cell of step a).

**[0052]** Suitable conditions which allow for expression of the polynucleotide of the invention comprised by the host cell depend on the host cell as well as the expression control sequence used for governing expression of the said polynucleotide. These conditions and how to select them are very well known to those skilled in the art. The expressed polypeptide may be obtained, for example, by all conventional purification techniques including affinity chromatography, size exclusion chromatography, high pressure liquid chromatography (HPLC) and precipitation techniques including antibody precipitation. It is to be understood that the method may—although preferred—not necessarily yield an essentially pure preparation of the polypeptide. It is to be understood that depending on the host cell which is used for the aforementioned method, the polypeptides produced thereby may become posttranslationally modified or processed otherwise.

**[0053]** The present invention also encompasses a polypeptide encoded by the polynucleotide of the present invention or which is obtainable by the aforementioned method.

**[0054]** The term “polypeptide” as used herein encompasses essentially purified polypeptides or polypeptide preparations comprising other proteins in addition. Further, the term also relates to the fusion proteins or polypeptide fragments being at least partially encoded by the polynucleotide of the present invention referred to above. Moreover, it includes chemically modified polypeptides. Such modifications may be artificial modifications or naturally occurring modifications such as phosphorylation, glycosylation, myristylation and the like (Review in Mann 2003, Nat. Biotechnol. 21, 255-261, review with focus on plants in Huber 2004, Curr. Opin. Plant Biol. 7, 318-322). Currently, more than 300 posttranslational modifications are known (see full ABFRC Delta mass list at [abrf.org/index.cfm/dm.home](http://abrf.org/index.cfm/dm.home)). The polypeptides of the present invention shall exhibit the desaturase or elongase activity referred to above.

**[0055]** Encompassed by the present invention is, furthermore, an antibody or fragments thereof which specifically recognizes the polypeptide of the invention.

**[0056]** Antibodies against the polypeptides of the invention can be prepared by well known methods using a purified polypeptide according to the invention or a suitable fragment derived therefrom as an antigen. A fragment which is suitable as an antigen may be identified by antigenicity determining algorithms well known in the art. Such fragments may be obtained either from the polypeptide of the invention by proteolytic digestion or may be a synthetic peptide. Preferably, the antibody of the present invention is a monoclonal antibody, a polyclonal antibody, a single chain antibody, a chimerized antibody or a fragment of any of these antibodies, such as Fab, Fv or scFv fragments etc. Also comprised as antibodies by the present invention are bispecific antibodies, synthetic antibodies or chemically modified derivatives of any of the aforementioned antibodies. The antibody of the present invention shall specifically bind (i.e. does significantly not cross react with other polypeptides or peptides) to the polypeptide of the invention. Specific binding can be tested by various well known techniques. Antibodies or fragments thereof can be obtained by using methods which are described, e.g., in Harlow and Lane “Antibodies, A Laboratory Manual”, CSH Press, Cold Spring Harbor, 1988. Monoclonal antibodies can be prepared by the techniques originally described in Köhler 1975, Nature 256, 495, and Galfré 1981, Meth. Enzymol. 73, 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals. The antibodies can be used, for example, for the immunoprecipitation, immunolocalization or purification (e.g., by affinity chromatography) of the polypeptides of the invention as well as for the monitoring of the presence of said variant polypeptides, for example, in recombinant organisms, and for the identification of proteins or compounds interacting with the proteins according to the invention.

**[0057]** Moreover, the present invention contemplates a non-human transgenic organism comprising the polynucleotide or the vector of the present invention.

**[0058]** Preferably, the non-human transgenic organism is a plant, plant part, or plant seed. Preferred plants to be used for introducing the polynucleotide or the vector of the invention are plants which are capable of synthesizing fatty acids, such as all dicotyledonous or monocotyledonous plants, algae or

mosses. It is to be understood that host cells derived from a plant may also be used for producing a plant according to the present invention. Preferred plants are selected from the group of the plant families Adelotheceaceae, Anacardiaceae, Asteraceae, Apiaceae, Betulaceae, Boraginaceae, Brassicaceae, Bromeliaceae, Caricaceae, Cannabaceae, Convolvulaceae, Chenopodiaceae, Crypthecodiniaceae, Cucurbitaceae, Ditrachaceae, Elaeagnaceae, Ericaceae, Euphorbiaceae, Fabaceae, Geraniaceae, Gramineae, Juglandaceae, Lauraceae, Leguminosae, Linaceae, Prasinophyceae or vegetable plants or ornamentals such as *Tagetes*. Examples which may be mentioned are the following plants selected from the group consisting of: Adelotheceaceae such as the genera *Physcomitrella*, such as the genus and species *Physcomitrella patens*, Anacardiaceae such as the genera *Pistacia*, *Mangifera*, *Anacardium*, for example the genus and species *Pistacia vera* [pistachio], *Mangifera indica* [mango] or *Anacardium occidentale* [cashew], Asteraceae, such as the genera *Calendula*, *Carthamus*, *Centaurea*, *Cichorium*, *Cynara*, *Helianthus*, *Lactuca*, *Locusta*, *Tagetes*, *Valeriana*, for example the genus and species *Calendula officinalis* [common marigold], *Carthamus tinctorius* [safflower], *Centaurea cyanus* [cornflower], *Cichorium intybus* [chicory], *Cynara scolymus* [artichoke], *Helianthus annuus* [sunflower], *Lactuca sativa*, *Lactuca crisa*, *Lactuca esculenta*, *Lactuca scariola* L. ssp. *sativa*, *Lactuca scariola* L. var. *integrata*, *Lactuca scariola* L. var. *integrifolia*, *Lactuca sativa* subsp. *romana*, *Locusta communis*, *Valeriana locusta* [salad vegetables], *Tagetes lucida*, *Tagetes erecta* or *Tagetes tenuifolia* [african or french marigold], Apiaceae, such as the genus *Daucus*, for example the genus and species *Daucus carota* [carrot], Betulaceae, such as the genus *Corylus*, for example the genera and species *Corylus avellana* or *Corylus colurna* [hazelnut], Boraginaceae, such as the genus *Borago*, for example the genus and species *Borago officinalis* [borage], Brassicaceae, such as the genera *Brassica*, *Melanosinapis*, *Sinapis*, *Arabadopsis*, for example the genera and species *Brassica napus*, *Brassica rapa* ssp. [oilseed rape], *Sinapis arvensis* *Brassica juncea*, *Brassica juncea* var. *juncea*, *Brassica juncea* var. *crispifolia*, *Brassica juncea* var. *foliosa*, *Brassica nigra*, *Brassica sinapioides*, *Melanosinapis communis* [mustard], *Brassica oleracea* [fodder beet] or *Arabidopsis thaliana*, Bromeliaceae, such as the genera *Anana*, *Bromelia* (pineapple), for example the genera and species *Anana comosus*, *Ananas ananas* or *Bromelia comosa* [pineapple], Caricaceae, such as the genus *Carica*, such as the genus and species *Carica papaya* [pawpaw], Cannabaceae, such as the genus *Cannabis*, such as the genus and species *Cannabis sativa* [hemp], Convolvulaceae, such as the genera *Ipomea*, *Convolvulus*, for example the genera and species *Ipomea batatas*, *Ipomea pandurata*, *Convolvulus batatas*, *Convolvulus tiliaceus*, *Ipomea fastigiata*, *Ipomea tiliacea*, *Ipomea triloba* or *Convolvulus panduratus* [sweet potato, batate], Chenopodiaceae, such as the genus *Beta*, such as the genera and species *Beta vulgaris*, *Beta vulgaris* var. *altissima*, *Beta vulgaris* var. *Vulgaris*, *Beta maritima*, *Beta vulgaris* var. *perennis*, *Beta vulgaris* var. *conditiva* or *Beta vulgaris* var. *esculents* [sugarbeet], Crypthecodiniaceae, such as the genus *Crypthecodinium*, for example the genus and species *Crypthecodinium cohnii*, Cucurbitaceae, such as the genus *Cucurbita*, for example the genera and species *Cucurbita maxima*, *Cucurbita mixta*, *Cucurbita pepo* or *Cucurbita moschata* [pumpkin/squash], Cymbellaceae such as the genera *Amphora*, *Cymbella*,



*Okenia*, *Phaeodactylum*, *Reimeria*, for example the genus and species *Phaeodactylum tricornutum*, Ditrichaceae such as the genera *Ditrichaceae*, *Astomiopsis*, *Ceratodon*, *Chrysoblastella*, *Ditrichum*, *Distichium*, *Eccremidium*, *Lophidion*, *Philibertiella*, *Pleuridium*, *Saelania*, *Trichodon*, *Skottsbergia*, for example the genera and species *Ceratodon antarcticus*, *Ceratodon columbiae*, *Ceratodon heterophyllus*, *Ceratodon purpureus*, *Ceratodon purpureus*, *Ceratodon purpureus* ssp. *convolutus*, *Ceratodon*, *purpureus* spp. *stenocarpus*, *Ceratodon purpureus* var. *rotundifolius*, *Ceratodon ratodon*, *Ceratodon stenocarpus*, *Chrysoblastella chilensis*, *Ditrichum ambiguum*, *Ditrichum brevisetum*, *Ditrichum crispatisimum*, *Ditrichum difficile*, *Ditrichum falcifolium*, *Ditrichum flexicaule*, *Ditrichum giganteum*, *Ditrichum heteromallum*, *Ditrichum lineare*, *Ditrichum lineare*, *Ditrichum montanum*, *Ditrichum montanum*, *Ditrichum pallidum*, *Ditrichum punctulatum*, *Ditrichum pusillum*, *Ditrichum pusillum* var. *tortile*, *Ditrichum rhynchostegium*, *Ditrichum schimperii*, *Ditrichum tortile*, *Distichium capillaceum*, *Distichium hagenii*, *Distichium inclinatum*, *Distichium macounii*, *Eccremidium floridanum*, *Eccremidium whiteleggei*, *Lophidion strictus*, *Pleuridium acuminatum*, *Pleuridium alternifolium*, *Pleuridium holdridgei*, *Pleuridium mexicanum*, *Pleuridium ravenelii*, *Pleuridium subulatum*, *Saelania glaucescens*, *Trichodon borealis*, *Trichodon cylindricus* or *Trichodon cylindricus* var. *oblongus*, Elaeagnaceae such as the genus *Elaeagnus*, for example the genus and species *Olea europaea* [olive], Ericaceae such as the genus *Kalmia*, for example the genera and species *Kalmia latifolia*, *Kalmia angustifolia*, *Kalmia microphylla*, *Kalmia polifolia*, *Kalmia occidentalis*, *Cistus chamaerhodendros* or *Kalmia lucida* [mountain laurel], Euphorbiaceae such as the genera *Manihot*, *Janipha*, *Jatropha*, *Ricinus*, for example the genera and species *Manihot utilissima*, *Janipha manihot*, *Jatropha manihot*, *Manihot aipil*, *Manihot dulcis*, *Manihot manihot*, *Manihot melanobasis*, *Manihot esculenta* [manihot] or *Ricinus communis* [castor-oil plant], Fabaceae such as the genera *Pisum*, *Albizia*, *Cathormion*, *Feuillea*, *Inga*, *Pithecolobium*, *Acacia*, *Mimosa*, *Medicago*, *Glycine*, *Dolichos*, *Phaseolus*, *Soja*, for example the genera and species *Pisum sativum*, *Pisum arvense*, *Pisum humile* [pea], *Albizia berteriana*, *Albizia julibrissin*, *Albizia lebbek*, *Acacia berteriana*, *Acacia littoralis*, *Albizia berteriana*, *Albizia berteriana*, *Cathormion berteriana*, *Feuillea berteriana*, *Inga fragrans*, *Pithecellobium berterianum*, *Pithecellobium fragrans*, *Pithecolobium berterianum*, *Pseudalbizia berteriana*, *Acacia julibrissin*, *Acacia nemu*, *Albizia nemu*, *Feuillea julibrissin*, *Mimosa julibrissin*, *Mimosa speciosa*, *Sericanrda julibrissin*, *Acacia lebbek*, *Acacia macrophylla*, *Albizia lebbek*, *Feuillea lebbek*, *Mimosa lebbek*, *Mimosa speciosa* [silk tree], *Medicago sativa*, *Medicago falcata*, *Medicago varia* [alfalfa], *Glycine max* *Dolichos soja*, *Glycine gracilis*, *Glycine hispida*, *Phaseolus max*, *Soja hispida* or *Soja max* [soybean], Funariaceae such as the genera *Aphanorrhagma*, *Entosthodon*, *Funaria*, *Physcomitrella*, *Physcomitrium*, for example the genera and species *Aphanorrhagma serratum*, *Entosthodon attenuatus*, *Entosthodon bolanderi*, *Entosthodon bonplandii*, *Entosthodon californicus*, *Entosthodon drummondii*, *Entosthodon jamesonii*, *Entosthodon leibergii*, *Entosthodon neoscoticus*, *Entosthodon rubrissetus*, *Entosthodon spathulifolius*, *Entosthodon tucsoni*, *Funaria americana*, *Funaria bolanderi*, *Funaria calcarea*, *Funaria californica*, *Funaria calvescens*, *Funaria convoluta*, *Funaria*

*flavicans*, *Funaria groutiana*, *Funaria hygrometrica*, *Funaria hygrometrica* var. *arctica*, *Funaria hygrometrica* var. *calvescens*, *Funaria hygrometrica* var. *convoluta*, *Funaria hygrometrica* var. *muralis*, *Funaria hygrometrica* var. *utahensis*, *Funaria microstoma*, *Funaria microstoma* var. *obtusifolia*, *Funaria muhlenbergii*, *Funaria orcuttii*, *Funaria plano-convexa*, *Funaria polaris*, *Funaria ravenelii*, *Funaria rubriseta*, *Funaria serrata*, *Funaria sonora*, *Funaria sublimbatus*, *Funaria tucsoni*, *Physcomitrella californica*, *Physcomitrella patens*, *Physcomitrella readeri*, *Physcomitrium australe*, *Physcomitrium californicum*, *Physcomitrium collenchymatum*, *Physcomitrium coloradense*, *Physcomitrium cupuliferum*, *Physcomitrium drummondii*, *Physcomitrium euryostomum*, *Physcomitrium flexifolium*, *Physcomitrium hookeri*, *Physcomitrium hookeri* var. *serratum*, *Physcomitrium immersum*, *Physcomitrium kellermanii*, *Physcomitrium megalocarpum*, *Physcomitrium pyriforme*, *Physcomitrium pyriforme* var. *serratum*, *Physcomitrium rufipes*, *Physcomitrium sandbergii*, *Physcomitrium subsphaericum*, *Physcomitrium washingtoniense*, Geraniaceae, such as the genera *Pelargonium*, *Cocos*, *Oleum*, for example the genera and species *Cocos nucifera*, *Pelargonium grossularioides* or *Oleum cocois* [coconut], Gramineae, such as the genus *Saccharum*, for example the genus and species *Saccharum officinarum*, Juglandaceae, such as the genera *Juglans*, *Wallia*, for example the genera and species *Juglans regia*, *Juglans ailanthifolia*, *Juglans sieboldiana*, *Juglans cinerea*, *Wallia cinerea*, *Juglans bixbyi*, *Juglans californica*, *Juglans hindsii*, *Juglans intermedia*, *Juglans jamaicensis*, *Juglans major*, *Juglans microcarpa*, *Juglans nigra* or *Wallia nigra* [walnut], Lauraceae, such as the genera *Persea*, *Laurus*, for example the genera and species *Laurus nobilis* [bay], *Persea americana*, *Persea gratissima* or *Persea persea* [avocado], Leguminosae, such as the genus *Arachis*, for example the genus and species *Arachis hypogaea* [peanut], Linaceae, such as the genera *Linum*, *Adenolinum*, for example the genera and species *Linum usitatissimum*, *Linum humile*, *Linum austriacum*, *Linum bienne*, *Linum angustifolium*, *Linum catharticum*, *Linum flavum*, *Linum grandiflorum*, *Adenolinum grandiflorum*, *Linum lewisii*, *Linum narbonense*, *Linum perenne*, *Linum perenne* var. *lewisii*, *Linum pratense* or *Linum trigynum* [linseed], Lythraeae, such as the genus *Punica*, for example the genus and species *Punica granatum* [pomegranate], Malvaceae, such as the genus *Gossypium*, for example the genera and species *Gossypium hirsutum*, *Gossypium arboreum*, *Gossypium barbadense*, *Gossypium herbaceum* or *Gossypium thurberi* [cotton], Marchantiaceae, such as the genus *Marchantia*, for example the genera and species *Marchantia berteriana*, *Marchantia foliacea*, *Marchantia macropora*, Musaceae, such as the genus *Musa*, for example the genera and species *Musa nana*, *Musa acuminata*, *Musa paradisiaca*, *Musa* spp. [banana], Onagraceae, such as the genera *Camissonia*, *Oenothera*, for example the genera and species *Oenothera biennis* or *Camissonia brevipes* [evening primrose], Palmae, such as the genus *Elais*, for example the genus and species *Elais guineensis* [oil palm], Papaveraceae, such as the genus *Papaver*, for example the genera and species *Papaver orientale*, *Papaver rhoeas*, *Papaver dubium* [poppy], Pedaliaceae, such as the genus *Sesamum*, for example the genus and species *Sesamum indicum* [sesame], Piperaceae, such as the genera *Piper*, *Artanthe*, *Peperomia*, *Steffensia*, for example the genera and species *Piper aduncum*, *Piper amalago*, *Piper angustifolium*, *Piper auritum*, *Piper betel*,



*Piper cubeba*, *Piper longum*, *Piper nigrum*, *Piper retrofractum*, *Artanthe adunca*, *Artanthe elongata*, *Peperomia elongata*, *Piper elongatum*, *Steffensia elongata* [cayenne pepper], Poaceae, such as the genera *Hordeum*, *Secale*, *Avena*, *Sorghum*, *Andropogon*, *Holcus*, *Panicum*, *Oryza*, *Zea* (maize), *Triticum*, for example the genera and species *Hordeum vulgare*, *Hordeum jubatum*, *Hordeum murinum*, *Hordeum secalinum*, *Hordeum distichon*, *Hordeum aegiceris*, *Hordeum hexastichon*, *Hordeum hexastichum*, *Hordeum irregulare*, *Hordeum sativum*, *Hordeum secalinum* [barley], *Secale cereale* [rye], *Avena sativa*, *Avena fatua*, *Avena byzantina*, *Avena fatua* var. *sativa*, *Avena hybrida* [oats], *Sorghum bicolor*, *Sorghum halepense*, *Sorghum saccharatum*, *Sorghum vulgare*, *Andropogon drummondii*, *Holcus bicolor*, *Holcus sorghum*, *Sorghum aethiopicum*, *Sorghum arundinaceum*, *Sorghum caffrorum*, *Sorghum cernuum*, *Sorghum dochna*, *Sorghum drummondii*, *Sorghum durra*, *Sorghum guineense*, *Sorghum lanceolatum*, *Sorghum nervosum*, *Sorghum saccharatum*, *Sorghum subglabrescens*, *Sorghum verticilliflorum*, *Sorghum vulgare*, *Holcus halepensis*, *Sorghum miliaceum*, *Panicum militaceum* [millet], *Oryza sativa*, *Oryza latifolia* [rice], *Zea mays* [maize], *Triticum aestivum*, *Triticum durum*, *Triticum turgidum*, *Triticum hybernum*, *Triticum macha*, *Triticum sativum* or *Triticum vulgare* [wheat], Porphyridiaceae, such as the genera *Chrootheca*, *Flintiella*, *Petrovanella*, *Porphyridium*, *Rhodella*, *Rhodorus*, *Vanhoeffenia*, for example the genus and species *Porphyridium cruentum*, Proteaceae, such as the genus *Macadamia*, for example the genus and species *Macadamia intergrifolia* [macadamia], Prasinophyceae such as the genera *Nephroselmis*, *Prasinococcus*, *Scherffelia*, *Tetraselmis*, *Mantoniella*, *Ostreococcus*, for example the genera and species *Nephroselmis olivacea*, *Prasinococcus capsulatus*, *Scherffelia dubia*, *Tetraselmis chui*, *Tetraselmis suecica*, *Mantoniella squamata*, *Ostreococcus tauri*, Rubiaceae such as the genus *Coffea*, for example the genera and species *Coffea* spp., *Coffea arabica*, *Coffea canephora* or *Coffea liberica* [coffee], Scrophulariaceae such as the genus *Verbascum*, for example the genera and species *Verbascum blattaria*, *Verbascum chaixii*, *Verbascum densiflorum*, *Verbascum lagurus*, *Verbascum longifolium*, *Verbascum lychnitis*, *Verbascum nigrum*, *Verbascum olympicum*, *Verbascum phlomoides*, *Verbascum phoenicum*, *Verbascum pulverulentum* or *Verbascum thapsus* [mullein], Solanaceae such as the genera *Capsicum*, *Nicotiana*, *Solanum*, *Lycopersicon*, for example the genera and species *Capsicum annuum*, *Capsicum annuum* var. *glabriusculum*, *Capsicum frutescens* [pepper], *Capsicum annuum* [paprika], *Nicotiana tabacum*, *Nicotiana alata*, *Nicotiana attenuata*, *Nicotiana glauca*, *Nicotiana langsdorffii*, *Nicotiana obtusifolia*, *Nicotiana quadrivalvis*, *Nicotiana repanda*, *Nicotiana rustica*, *Nicotiana sylvestris* [tobacco], *Solanum tuberosum* [potato], *Solanum melongena* [eggplant], *Lycopersicon esculentum*, *Lycopersicon lycopersicum*, *Lycopersicon pyriforme*, *Solanum integrifolium* or *Solanum lycopersicum* [tomato], Sterculiaceae, such as the genus *Theobroma*, for example the genus and species *Theobroma cacao* [cacao] or Theaceae, such as the genus *Camellia*, for example the genus and species *Camellia sinensis* [tea]. In particular preferred plants to be used as transgenic plants in accordance with the present invention are oil fruit crops which comprise large amounts of lipid compounds, such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, sesame, *Calendula*, *Punica*, evening primrose, mullein,

thistle, wild roses, hazelnut, almond, *macadamia*, avocado, bay, pumpkin/squash, linseed, soybean, pistachios, borage, trees (oil palm, coconut, walnut) or crops such as maize, wheat, rye, oats, triticale, rice, barley, cotton, cassava, pepper, *Tagetes*, Solanaceae plants such as potato, tobacco, eggplant and tomato, *Vicia* species, pea, alfalfa or bushy plants (coffee, cacao, tea), *Salix* species, and perennial grasses and fodder crops. Preferred plants according to the invention are oil crop plants such as peanut, oilseed rape, canola, sunflower, safflower, poppy, mustard, hemp, castor-oil plant, olive, *Calendula*, *Punica*, evening primrose, pumpkin/squash, linseed, soybean, borage, trees (oil palm, coconut). Especially preferred are sunflower, safflower, tobacco, mullein, sesame, cotton, pumpkin/squash, poppy, evening primrose, walnut, linseed, hemp, thistle or safflower. Very especially preferred plants are plants such as safflower, sunflower, poppy, evening primrose, walnut, linseed, or hemp.

[0059] Preferred mosses are *Physcomitrella* or *Ceratodon*. Preferred algae are *Isochrysis*, *Mantoniella*, *Ostreococcus* or *Cryptocodinium*, and algae/diatoms such as *Phaeodactylum* or *Thraustochytrium*. More preferably, said algae or mosses are selected from the group consisting of: *Emiliana*, *Shewanella*, *Physcomitrella*, *Thraustochytrium*, *Fusarium*, *Phytophthora*, *Ceratodon*, *Isochrysis*, *Aleurita*, *Muscarioides*, *Mortierella*, *Phaeodactylum*, *Cryptothecodinium*, specifically from the genera and species *Thallasiosira pseudonona*, *Euglena gracilis*, *Physcomitrella patens*, *Phytophthora infestans*, *Fusarium gramineum*, *Cryptocodinium cohnii*, *Ceratodon purpureus*, *Isochrysis galbana*, *Aleurita farinosa*, *Thraustochytrium* sp., *Muscarioides viallii*, *Mortierella alpine*, *Phaeodactylum tricornutum* or *Caenorhabditis elegans* or especially advantageously *Phytophthora infestans*, *Thallasiosira pseudonona* and *Cryptocodinium cohnii*.

[0060] Transgenic plants may be obtained by transformation techniques as elsewhere in this specification. Preferably, transgenic plants can be obtained by T-DNA-mediated transformation. Such vector systems are, as a rule, characterized in that they contain at least the vir genes, which are required for the *Agrobacterium*-mediated transformation, and the sequences which delimit the T-DNA (T-DNA border). Suitable vectors are described elsewhere in the specification in detail.

[0061] Also encompassed are transgenic non-human animals comprising the vector or polynucleotide of the present invention. Preferred non-human transgenic animals envisaged by the present invention are fish, such as herring, salmon, sardine, redfish, eel, carp, trout, halibut, mackerel, zander or tuna.

[0062] However, it will be understood that dependent on the non-human transgenic organism specified above, further, enzymatic activities may be conferred to the said organism, e.g., by recombinant technologies. Accordingly, the present invention, preferably, envisages a non-human transgenic organism specified above which in addition to the polynucleotide of the present invention comprises polynucleotides encoding such desaturases and/or elongases as required depending on the selected host cell. Preferred desaturases and/or elongases which shall be present in the organism are at least one enzyme selected from the group of desaturases and/or elongases or the combinations specifically recited elsewhere in this specification (see above and Tables 3, 4 and 5).

**[0063]** Furthermore, the present invention encompasses a method for the manufacture of polyunsaturated fatty acids comprising:

**[0064]** a) cultivating the host cell of the invention under conditions which allow for the production of polyunsaturated fatty acids in said host cell;

**[0065]** b) obtaining said polyunsaturated fatty acids from the said host cell.

**[0066]** The term "polyunsaturated fatty acids (PUFA)" as used herein refers to fatty acids comprising at least two, preferably, three, four, five or six, double bonds. Moreover, it is to be understood that such fatty acids comprise, preferably from 18 to 24 carbon atoms in the fatty acid chain. More preferably, the term relates to long chain PUFA (LCPUFA) having from 20 to 24 carbon atoms in the fatty acid chain. Preferred unsaturated fatty acids in the sense of the present invention are selected from the group consisting of DGLA 20:3 (8,11,14), ARA 20:4 (5,8,11,14), iARA 20:4(8,11,14,17), EPA 20:5 (5,8,11,14,17), DPA 22:5 (4,7,10,13,16), DHA 22:6 (4,7,10,13,16,19), 20:4 (8,11,14,17), more preferably, arachidonic acid (ARA) 20:4 (5,8,11,14), eicosapentaenoic acid (EPA) 20:5 (5,8,11,14,17), and docosahexaenoic acid (DHA) 22:6 (4,7,10,13,16,19). Thus, it will be understood that most preferably, the methods provided by the present invention pertaining to the manufacture of ARA, EPA or DHA. Moreover, also encompassed are the intermediates of LCPUFA which occur during synthesis. Such intermediates are, preferably, formed from substrates by the desaturase or elongase activity of the polypeptides of the present invention. Preferably, substrates encompass LA 18:2 (9,12), ALA 18:3(9,12,15), Eicosadienoic acid 20:2 (11,14), Eicosatrienoic acid 20:3 (11,14,17), DGLA 20:3 (8,11,14), ARA 20:4 (5,8,11,14), eicosatetraenoic acid 20:4 (8,11,14,17), Eicosapentaenoic acid 20:5 (5,8,11,14,17), Docosahexapentaenoic acid 22:5 (7,10,13,16,19).

**[0067]** The term "cultivating" as used herein refers maintaining and growing the host cells under culture conditions which allow the cells to produce the said polyunsaturated fatty acid, i.e. the PUFA and/or LCPUFA referred to above. This implies that the polynucleotide of the present invention is expressed in the host cell so that the desaturase and/or elongase activity is present. Suitable culture conditions for cultivating the host cell are described in more detail below.

**[0068]** The term "obtaining" as used herein encompasses the provision of the cell culture including the host cells and the culture medium as well as the provision of purified or partially purified preparations thereof comprising the polyunsaturated fatty acids, preferably, ARA, EPA, DHA, in free or in -CoA bound form, as membrane phospholipids or as triacylglyceride esters. More preferably, the PUFA and LCPUFA are to be obtained as triglyceride esters, e.g., in form of an oil. More details on purification techniques can be found elsewhere herein below.

**[0069]** The host cells to be used in the method of the invention are grown or cultured in the manner with which the skilled worker is familiar, depending on the host organism.

**[0070]** Usually, host cells are grown in a liquid medium comprising a carbon source, usually in the form of sugars, a nitrogen source, usually in the form of organic nitrogen sources such as yeast extract or salts such as ammonium sulfate, trace elements such as salts of iron, manganese and magnesium and, if appropriate, vitamins, at temperatures of between 0° C. and 100° C., preferably between 10° C. and

60° C. under oxygen or anaerobic atmosphere dependent on the type of organism. The pH of the liquid medium can either be kept constant, that is to say regulated during the culturing period, or not. The cultures can be grown batchwise, semibatchwise or continuously. Nutrients can be provided at the beginning of the fermentation or administered semicontinuously or continuously: The produced PUFA or LCPUFA can be isolated from the host cells as described above by processes known to the skilled worker, e.g., by extraction, distillation, crystallization, if appropriate precipitation with salt, and/or chromatography. It might be required to disrupt the host cells prior to purification. To this end, the host cells can be disrupted beforehand. The culture medium to be used must suitably meet the requirements of the host cells in question. Descriptions of culture media for various microorganisms which can be used as host cells according to the present invention can be found in the textbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Culture media can also be obtained from various commercial suppliers. All media components are sterilized, either by heat or by filter sterilization. All media components may be present at the start of the cultivation or added continuously or batchwise, as desired. If the polynucleotide or vector of the invention which has been introduced in the host cell further comprises an expressible selection marker, such as an antibiotic resistance gene, it might be necessary to add a selection agent to the culture, such as an antibiotic in order to maintain the stability of the introduced polynucleotide. The culture is continued until formation of the desired product is at a maximum. This is normally achieved within 10 to 160 hours. The fermentation broths can be used directly or can be processed further. The biomass may, according to requirement, be removed completely or partially from the fermentation broth by separation methods such as, for example, centrifugation, filtration, decanting or a combination of these methods or be left completely in said broth. The fatty acid preparations obtained by the method of the invention, e.g., oils, comprising the desired PUFA or LCPUFA as triglyceride esters are also suitable as starting material for the chemical synthesis of further products of interest. For example, they can be used in combination with one another or alone for the preparation of pharmaceutical or cosmetic compositions, foodstuffs, or animal feeds. Chemically pure triglycerides comprising the desired PUFA or LCPUFA can also be manufactured by the methods described above. To this end, the fatty acid preparations are further purified by extraction, distillation, crystallization, chromatography or combinations of these methods. In order to release the fatty acid moieties from the triglycerides, hydrolysis may be also required. The said chemically pure triglycerides or free fatty acids are, in particular, suitable for applications in the food industry or for cosmetic and pharmacological compositions.

**[0071]** Moreover, the present invention relates to a method for the manufacture of poly-unsaturated fatty acids comprising:

**[0072]** a) cultivating the non-human transgenic organism of the invention under conditions which allow for the production of poly-unsaturated fatty acids in said non-human transgenic organism; and

**[0073]** b) obtaining said poly-unsaturated fatty acids from the said non-human transgenic organism.

**[0074]** Further, it follows from the above that a method for the manufacture of an oil, lipid or fatty acid composition is

also envisaged by the present invention comprising the steps of any one of the aforementioned methods and the further step of formulating PUFA or LCPUFA as oil, lipid or fatty acid composition. Preferably, said oil, lipid or fatty acid composition is to be used for feed, foodstuffs, cosmetics or medicaments. Accordingly, the formulation of the PUFA or LCPUFA shall be carried out according to the GMP standards for the individual envisaged products. For example, an oil may be obtained from plant seeds by an oil mill. However, for product safety reasons, sterilization may be required under the applicable GMP standard. Similar standards will apply for lipid or fatty acid compositions to be applied in cosmetic or pharmaceutical compositions. All these measures for formulating oil, lipid or fatty acid compositions as products are comprised by the aforementioned manufacture.

**[0075]** The term "oil" refers to a fatty acid mixture comprising unsaturated and/or saturated fatty acids which are esterified to triglycerides. Preferably, the triglycerides in the oil of the invention comprise PUFA or LCPUFA as referred to above. The amount of esterified PUFA and/or LCPUFA is, preferably, approximately 30%, a content of 50% is more preferred, a content of 60%, 70%, 80% or more is even more preferred. The oil may further comprise free fatty acids, preferably, the PUFA and LCPUFA referred to above. For the analysis, the fatty acid content can be, e.g., determined by GC analysis after converting the fatty acids into the methyl esters by transesterification. The content of the various fatty acids in the oil or fat can vary, in particular depending on the source. The oil, however, shall have a non-naturally occurring composition with respect to the PUFA and/or LCPUFA composition and content. It will be understood that such a unique oil composition and the unique esterification pattern of PUFA and LCPUFA in the triglycerides of the oil shall only be obtainable by applying the methods of the present invention specified above. Moreover, the oil of the invention may comprise other molecular species as well. Specifically, it may comprise minor impurities of the polynucleotide or vector of the invention. Such impurities, however, can be detected only by highly sensitive techniques such as PCR.

**[0076]** The contents of all references cited throughout this application are herewith incorporated by reference in general and with respect to their specific disclosure content referred to above.

## FIGURES

**[0077]** FIG. 1 shows a schematical figure of the different enzymatic activities leading to the production of ARA, EPA and DHA.

**[0078]** FIG. 2 shows a yeast expression experiment with feeding of 22:5n-3 in the presence (A) and absence (B) of d4Des(Eh)

**[0079]** FIG. 3 shows a yeast expression experiment with feeding of 20:3n-3 in the presence (A) and absence (B) of d8Des(Eh)

**[0080]** FIG. 4 shows a yeast expression experiment with feeding of 18:3n-3 in the presence (A) and absence (B) of d9Elo(Eh)

**[0081]** FIG. 5 shows a yeast expression experiment with feeding of 18:3n-6 (GLA) and 18:4n-3 (SDA) in the presence (A) and absence (B) of d5Elo(Eh)

**[0082]** FIG. 6 shows a yeast expression experiment with feeding of 20:4n-6 (ARA) and 20:5n-3 (EPA) in the presence (A) and absence (B) of d5Elo(Eh)

**[0083]** FIG. 7 shows the expression of d9Elo(Eh) in seeds of two *Arabidopsis* events. As control seeds not expressing d9Elo(Eh) are shown (WT).

**[0084]** FIG. 8 shows the Acyl-CoA analysis of mature *Arabidopsis* seeds from both events expressing the d9Elo(Eh) in comparison to seeds not expressing d9Elo(Eh) (Col0).

**[0085]** FIG. 9 shows the expression of d9Elo(Eh), d8Des(Eh) and d5Des(Eh) in seeds of various *Arabidopsis* events.

**[0086]** FIG. 10 shows gas chromatographic analysis of mature *Arabidopsis* seeds transformed with the construct OstELO5EmD4. Peaks were quantified and listed in the table below. The products of d5Elo(Eh) and d4Des(Eh) activity are 22:6n-3 (DHA).

**[0087]** FIG. 11 is a comparison between two d4-desaturases (Tc and Eh) showing that d4Des(Eh) is different from known d4-desaturases in producing a high ratio of DHA: DPA.

**[0088]** FIG. 12 shows the expression of d5Elo(Eh) in seeds of various *Arabidopsis* events.

**[0089]** FIG. 13 is a comparison between three different d6-desaturases and the substrate specificity of d5Des(Eh).

**[0090]** This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application, as well as the figures, are incorporated herein by reference.

## EXAMPLES

### Example 1

#### Organism and Culture Conditions

**[0091]** *Emiliana huxleyi* was grown as described in Sciana et al. (2003) Marine Ecology Progress Series 261:111-122 with following conditions:

**[0092]** Growth in 50 ml in conical flasks using K/2 medium (Keller et al. (1987) Journal of Phycology 23:633-638). The flasks were placed in a growth chamber at a temperature of  $17 \pm 0.1^\circ \text{C}$ . under 14L:10D irradiance. Light was provided by fluorescent lamps giving a photon flux density (400 to 700 nm) of  $170 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ .

### Example 2

#### Cloning of Novel Desaturase and Elongase Sequences

**[0093]** RNA from cells grown as described under Example 1 was extracted using the RNA-extraction Kit from Qiagen, a RACE-library was generated using the RACE-Kit from Clontech. From the RACE-library sequences for desaturase and elongases were amplified with PCR using following primer pairs and PCR conditions.

**[0094]** PCR reaction (50  $\mu\text{L}$ ):

**[0095]** 5.00  $\mu\text{L}$  Template cDNA

**[0096]** 5.00  $\mu\text{L}$  10 $\times$  Puffer (Advantage-Polymerase)+25 mM  $\text{MgCl}_2$

**[0097]** 5.00  $\mu\text{L}$  2 mM dNTP

**[0098]** 1.25  $\mu\text{L}$  je Primer (10 pmol/ $\mu\text{L}$ )

**[0099]** 0.50  $\mu\text{L}$  Advantage-Polymerase

- [0100] The Advantage polymerase mix from Clontech was used.  
 [0101] Reaction conditions of the PCR:  
 [0102] Annealing: 1 min 55° C.  
 [0103] Denaturation: 1 min 94° C.  
 [0104] Elongation: 2 min 72° C.  
 [0105] Cycles: 35  
 [0106] Primer pairs used in PCR:

Name	Primer pair (5' orientation)	SEQ ID NO.
Eh4ff	CCATGGGAGGCGCGCGCGAG	11
Eh4rv	CTAGTCCGCCTTGAGTTCTC	12
Eh5ff	ACCATGTGCAAGGCGAGCGGCCT	13
Eh5rv	TCACCAATCATGAGGAAGGT	14
Eh8ff	CCATGGGCAAGGGCGGCAACGC	15
Eh8rv	GGGCAGAGATGCCGCACTAG	16
Eh9ff	ACCATGCTCGATCGCGCCTCGTC	17
Eh9rv	TCACAGCGCCTTGCGGGTAGC	18

- [0107] The PCR reactions resulted in following polynucleotide sequences:

Gene	Activity	Length in bp	SEQ ID NO.
D4Des(Eh)	D4-desaturase	1280	5
D8Des(Eh)	D8-desaturase	1256	1
D9Elo(Eh)	D9-elongase	804	3
D5Elo(Eh)	Multi-elongase	921	7

- [0108] A list of identified full-length coding sequences is shown in Table 1.

TABLE 1

List of full-length coding sequences and deduced amino acid sequences			
SEQ ID NO:	Gene	Coding sequence in bp	Amino acid sequence
1	D8Des(Eh)	1254	417
3	D9Elo(Eh)	801	266
5	D4Des(Eh)	1278	425
7	D5Elo(Eh)	918	305

- [0109] Open reading frames as shown in Table 1 were cloned into the pESC(Leu) vector from Stratagene according to manufactures reaction conditions. Reactions were transformed into *E. coli* DH5 $\alpha$  and plasmid DNA was isolated. The plasmids pESC-d4Des(Eh), pESC-d8Des(Eh), pESC-d9Elo(Eh), pESC-d5Elo(Eh) were then used for yeast transformation.

## Example 3

## Yeast Transformation and Growth Conditions

- [0110] *S. cerevisiae* strain INVSC from Invitrogen was transformed with the constructs pESC-d4Des(Eh), pESC-d8Des(Eh), pESC-d9Elo(Eh), pESC-d5Elo(Eh) and pESC

using the S. C. EasyComp Transformation Kit (Invitrogen, Carlsbad, Calif.) with selection on leucine-deficient medium.

- [0111] Yeast were grown after transformation in complete medium containing all amino acids and nucleotides. Then yeast were plated on different medium containing either the complete medium (SD) or the complete medium lacking leucine (SD-Leu). Only yeast containing pESC-d4Des(Eh), pESC-d8Des(Eh), pESC-d9Elo(Eh), pESC-d5Elo(Eh) or pESC vector can grow on this medium.

## Example 4

## Functional Expression of Desaturases and Elongases in Yeast and Gas Chromatographic Analysis

- [0112] Yeast cells containing the respective pESC plasmids as prepared above were incubated 12 h in liquid DOB-U medium at 28° C., 200 rpm inkubiert and than additional 12 h in induction medium (DOB-U+2% (w/v) galactose+2% (w/v) raffinose). To the induction medium 250  $\mu$ M of the respective fatty acids were added to check for enzyme activity and specificity.

- [0113] Yeast cells were analyzed as following:

- [0114] Yeast cells from induction medium were harvested by centrifugation (100 $\times$ g, 5 min, 20° C.) and washed with 100 mM NaHCO<sub>3</sub>, pH 8.0, to remove residual fatty acids. From the yeast pellet a total extract of fatty acid methylsters (FAME) was generated by adding 2 ml 1 N methanolic sulfuric acid and 2% (v/v) Dimethoxypropan for 1 h at 80° C. FAME were extracted two times with Petrolether (PE). Not derivased fatty acids were removed by washing with 2 ml 100 mM NaHCO<sub>3</sub>, pH 8.0 and 2 ml Aqua dest. The PE-phases were dried with Na<sub>2</sub>SO<sub>4</sub> and eluted in 100  $\mu$ l PE. The samples were then separated with a DB-23-column (30 m, 0.25 mm, 0.25  $\mu$ m, Agilent) in a Hewlett-Packard 6850-machine with FID using following conditions: oven temperature 50° C. to 250° C. with a rate of 5° C./min and finally 10 min at 250° C.

- [0115] The identification of the fatty acids was done using the retention times of known fatty acid standards (Sigma). The method is described e.g. in Napier and Michaelson, 2001, Lipids. 36(8):761-766; Sayanova et al., 2001, Journal of Experimental Botany. 52(360):1581-1585, Sperling et al., 2001, Arch. Biochem. Biophys. 388(2):293-298 and Michaelson et al., 1998, FEBS Letters. 439(3):215-218.

## Example 5

## Functional Characterization of d4Des(Eh)

- [0116] As described above d4Des(Eh) was functionally characterized in yeast. The result of the analysis is shown in FIG. 2. Yeast transformed with pESC-d4Des(Eh) was compared to yeast transformed with pESC (control) while feeding the yeast cells with the fatty acid DPA 22:5n-3. Based on this comparison pESC-d4Des(Eh) exhibits d4-desaturase activity as in the control no 22:6 is observed. Therefore d4Des(Eh) is a functional d4-desaturase.

## Example 6

## Functional Characterization of d8Des(Eh)

- [0117] As described above d8Des(Eh) was functionally characterized in yeast. The result of the analysis is shown in

FIG. 3. Yeast transformed with pESC-d8Des(Eh) was compared to yeast transformed with pESC (control) while feeding the fatty acid 20:3n-3. Based on this comparison a new fatty acid was formed compared to the control, which is 20:4n-3. The formation of this fatty acid proves that d8Des (Eh) was functionally expressed and has d8-desaturase activity. The conversion rate of 20:3n-3 to 20:4n-3 was 5%.

#### Example 7

##### Functional Characterization of d9Elo(Eh)

[0118] As described above d9Elo(Eh) was functionally characterized in yeast. The result of the analysis is shown in FIG. 4. Yeast transformed with pESC-d9Elo(Eh) was compared to yeast transformed with pESC (control) while feeding the fatty acids 18:3n-3 (ALA) or 18:2 (LA). Based on this comparison a new fatty acid was formed compared to the control, which is 20:3n-3 or 20:2n-6, respectively. The formation of these fatty acids proves that d9Elo(Eh) was functionally expressed and has d9-elongase activity. The conversion rate of 18:3n-3 to 20:3n-3 was 17%, the conversion rate of 18:2n-6 to 20:2n-6 was 49%.

#### Example 8

##### Functional Characterization of d5Elo(Eh)

[0119] As described above d5Elo(Eh) was functionally characterized in yeast. The result of the analysis is shown in FIGS. 5 and 6. Yeast transformed with pESC-d5Elo(Eh) was compared to yeast transformed with pESC (control) while feeding the fatty acids 18:3n-6 (GLA), 18:4 (SDA) or 20:4n-6 (ARA), 20:5n-3 (EPA), respectively. Based on this comparison new fatty acids formation was observed when compared to the control, which is 20:3n-6 or 20:4n-3 when fed GLA or SDA and 22:4n-6 or 22:5n-3 when fed ARA or EPA, respectively. The formation of these fatty acids proves that d5Elo(Eh) was functionally expressed and has d5-elongase activity. The conversion rate of GLA was 13%, the conversion rate of 18:4n-3 was 30%, the conversion rate of ARA was 38% and the conversion rate of EPA was 30%. Surprisingly the elongase used a wide variety of substrates of elongation. The specification indicates a multifunctional elongase activity with higher specificities for omega3 fatty acids.

#### Example 9

##### Expression of Novel Elongases from *Emilana huxleyi* in Plants

[0120] The novel desaturases and elongases were cloned into a plant transformation vector as described in WO2003/093482, WO2005/083093 or WO2007/093776. Exemplary suitable combinations of genes are described in Table 2, 3 and 4.

TABLE 2

Gene combinations for the production of ARA.		
Gene	Activity	SEQ ID NO:
D6Des(Ot)	$\Delta 6$ -Desaturase	19
D6Elo(Pp)	$\Delta 6$ -Elongase	21
D5Des(Eh)	$\Delta 5$ -Desaturase	9

TABLE 2-continued

Gene combinations for the production of ARA.		
Gene	Activity	SEQ ID NO:
D12Des(Ps)	$\Delta 12$ -Desaturase	23
D6Elo(Tp)	$\Delta 6$ -Elongase	25
D8Des(Eh)	$\Delta 8$ -Desaturase	1
D9Elo(Eh)	$\Delta 9$ -Elongase	3

TABLE 3

Gene combinations for the production of EPA.		
Gene	Activity	SEQ ID NO:
D6Des(Ot)	$\Delta 6$ -Desaturase	19
D5Elo(Eh)	$\Delta 5$ -Elongase	7
D5Des(Eh)	$\Delta 5$ -Desaturase	9
D12Des(Ps)	$\Delta 12$ -Desaturase	23
D6Elo(Tp)	$\Delta 6$ -Elongase	25
$\omega 3$ -Des(Pi)	Omega 3-Desaturase	27
D15Des(Cp)	$\Delta 15$ -Desaturase	29
D8Des(Eh)	$\Delta 8$ -Desaturase	1
D9Elo(Eh)	$\Delta 9$ -Elongase	3

TABLE 4

Gene combinations for the production of DHA.		
Gene	Activity	SEQ ID NO:
D6Des(Ot)	$\Delta 6$ -Desaturase	19
D5Elo(Eh)	$\Delta 5$ -Elongase	7
D5Des(Eh)	$\Delta 5$ -Desaturase	9
D12Des(Ps)	$\Delta 12$ -Desaturase	23
D6Elo(Tp)	$\Delta 6$ -Elongase	25
$\omega 3$ -Des(Pi)	Omega 3-Desaturase	27
D15Des(Cp)	$\Delta 15$ -Desaturase	29
D5Elo(Ot)	$\Delta 5$ -elongase	31
D4Des(Eh)	$\Delta 4$ -desaturase	5
D8Des(Eh)	$\Delta 8$ -Desaturase	1
D9Elo(Eh)	$\Delta 9$ -Elongase	3

[0121] Based on the gene combinations as described in Table 2, Table 3 or Table 4 following combinations were designed

[0122] AP2: LuCnI-d5Des(Eh)\_LuCnI-d8Des8Eh)\_Napin- $\omega 3$ Des(PO\_Napin-d12Des(Ps)\_LuCnI-d9Elo(Eh)

[0123] OstELO5EmD4: VfUSP-d6Elo(Pp)\_LuCnI-d5Des8Tc)\_VfSBP-d6Des(Ot)\_Napin- $\omega 3$ Des(PO\_Napin-d12Des(Ps)\_LuCnI-d5Elo(Ot)\_LuCnI-d4Des(Eh)

[0124] OstELO5TcD4: VfUSP-d6Elo(Pp)\_LuCnI-d5Des8Tc)\_VfSBP-d6Des(Ot)\_Napin- $\omega 3$ Des(PO\_Napin-d12Des(Ps)\_LuCnI-d5Elo(Ot)\_LuCnI-d4Des(Tc)

[0125] Transgenic rapeseed lines were generated as described in Deblaere et al, 1984, Nucl. Acids. Res. 13, 4777-4788 and seeds of transgenic rapeseed plants are analyzed as described in Qiu et al. 2001, J. Biol. Chem. 276, 31561-31566.

[0126] Transgenic *Arabidopsis* plants were generated as described in Bechtholdt et al. 1993 *C. R. Acad. Sci. Ser. III Sci. Vie.*, 316, 1194-1199. Seeds of transgenic *Arabidopsis* plants expressing d9Elo(Eh) by using the seed-specific promoter Glycinin from soybean (Lelievre et al. (1992) Plant Physiol 98:387-391) were analyzed by gas chromatography (FIG. 7). Compared to non-transgenic control plants (WT)

there are changes in the fatty acid profile, proving that d9Elo(Eh) was functionally expressed in seeds. The major shifts in the fatty acid profile is directed to a 10fold increase in the fatty acid 20:2n-6 and 20:3n-3 (FIG. 7). Therefore d9Elo(Eh) exhibits a  $\Delta 9$ -elongase activity, which is consistent with the yeast characterization. Further, the levels of 18:2 and ALA in the transgenic events expressing d9Elo(Eh) are lowered compared to WT, as these fatty acids are direct substrates for the d9Elo(Eh). Further, the endogenous elongation system in the plant is unchanged as levels of 20:1 and 22:1 are similar between transgenic plants expression d9Elo(Eh) and WT control. This indicates that the expression of d9Elo(Eh) does not disturb endogenous elongation process, but delivers additional activity.

[0127] To further prove the activity of d9Elo(Eh) expressed in seeds of *Arabidopsis thaliana* AcylCoA-measurements were done. Substrates and products of the d9Elo(Eh) elongation reaction are AcylCoA-esters, which are then further incorporated into triacylglycerides (oil). The analysis of the acylCoA-pool reveals the formation and flux of the elongation reaction.

[0128] FIG. 8 summarizes the AcylCoA measurements for *Arabidopsis* event expressing d9Elo(Eh) in comparison to controls not expressing d9Elo(Eh) (Col0). The change in the chromatogram is indicated by a star. At this position a massive amount of 20:2n-6 is detected, which is much lower in the control. The conditions for separation of the fatty acid CoA-esters does not allow the detection of 20:3n-3 as this CoA ester is not separated from 18:3CoA.

[0129] The massive occurrence of 20:2n-6-CoA proves the expression of d9Elo(Eh) as this is the direct product of its enzymatic activity.

[0130] Further, transgenic *Arabidopsis* lines have been generated to validate the activity of d8Des(Eh) and d5Des(Eh). Vector AP2 has been constructed according to standard molecular biology steps as described in WO2003/093482, WO2005/083093, WO2007/093776 or WO2009/016202 and transformed into *Arabidopsis thaliana* as described above. Analysis of transgenic seeds is shown in FIG. 9. The products of d9Elo(Eh) are 20:2 and 20:3n-3.

[0131] Further, transgenic *Arabidopsis* lines have been generated to validate the activity of d4Des(Eh). Construct OstELO5EmD4 was transformed into *Arabidopsis* as described above and seeds of a number of individual lines have been analyzed by gas chromatography (FIG. 10). The activity of d4Des(Eh) is demonstrated by the formation of DHA 22:6 (last column). All lines show the production of DHA with levels of up to 4.7%. Of special interest is the ratio of DHA to DPA. Surprisingly the ratio of d4Des(Eh) is much higher than in d4-desaturases known in the art. A comparison against the d4-desaturase from *Thraustochytrium* ssp. of WO2002/026946 is shown in FIG. 11. The enzyme from *Thraustochytrium* ssp. showed so far highest levels of DHA (WO2005/083093), but with an unfavorable ratio of DPA to DHA. A high ratio of DHA:DPA is for the commercial use of such oils of importance.

[0132] Further, transgenic *Arabidopsis* lines have been generated to validate the activity of d5Elo(Eh). Construct EmELO5TcD4 was transformed into *Arabidopsis* as described above and seeds of a number of individual lines have been analyzed by gas chromatography (FIG. 12). The activity of d5Elo(Eh) is demonstrated by the formation of

DPA 22:5 and DHA 22:6. Most lines show the production of these two fatty acids, proving that d5Elo(Em) is functionally expressed in the seeds.

[0133] Further, transgenic *Arabidopsis* lines have been generated to validate the activity and substrate specificity of d5Des(Eh). For this purpose two  $\Delta 6$ -desaturases were selected based on their different substrate specificity. The borage $\Delta 6$  is expected to use phosphatidylcholin-18:2 as substrate (WO96/21022), whereas the *Ostreococcus*  $\Delta 6$  (Ostr $\Delta 6$ ) uses Acyl-CoA ester (WO2005/012316). In combination with the d6-elongase from *Physcomitrella patens* (WO2001/059128) both d6-desaturases produce DGLA or 20:4n-3, respectively. The ratio of ARA to EPA is for the borage $\Delta 6$  2.9, for the Ostr $\Delta 6$  2.3. It is noted that the use of Ostr $\Delta 6$  results in 3-4 times higher levels of products compared to the borage $\Delta 6$ . The further combination of the d5Des(Eh) resulted in the production of ARA and EPA, demonstrating the functionality of the d5Des(Eh). The conversion of d5Des(Eh) of DGLA to ARA is 29% (borage $\Delta 6$ ) or 47% (Ostr $\Delta 6$ ). For 20:4n-3 to EPA it is 33% (borage $\Delta 6$ ) or 26% (Ostr $\Delta 6$ ).

[0134] Based on these results it is concluded that for Acyl-CoA substrates d5Des(Eh) is specific for the omega6 fatty acid DGLA. This is a novel substrate specificity not observed in the state of the art d5-desaturases.

#### REFERENCE LIST

- [0135] Arondel, V., Lemieux, B., Hwang, I., Gibson, S., Goodman, H. M., and Somerville, C. R. (1992). Map-based cloning of a gene controlling omega-3 fatty acid desaturation in *Arabidopsis*. *Science* 258, 1353-1355.
- [0136] Broadwater, J. A., Whittle, E., and Shanklin, J. (2002). Desaturation and hydroxylation. Residues 148 and 324 of *Arabidopsis* FAD2, in addition to substrate chain length, exert a major influence in partitioning of catalytic specificity. *J. Biol. Chem.* 277, 15613-15620.
- [0137] Broun, P., Shanklin, J., Whittle, E., and Somerville, C. (1998b). Catalytic plasticity of fatty acid modification enzymes underlying chemical diversity of plant lipids. *Science* 282, 1315-1317.
- [0138] Calvo, A. M., Gardner, H. W., and Keller, N. P. (2001). Genetic connection between fatty acid metabolism and sporulation in *Aspergillus nidulans*. *J. Biol. Chem.* 276, 25766-25774.
- [0139] Knutzon, D. S., Thurmond, J. M., Huang, Y. S., Chaudhary, S., Bobik, E. G., Jr., Chan, G. M., Kirchner, S. J., and Mukerji, P. (1998). Identification of Delta5-dehydratase from *Mortierella* alpine by heterologous expression in Bakers' yeast and canola. *J. Biol. Chem.* 273, 29360-29366.
- [0140] Mantle, P. G. and Nisbet, L. J. (1976). Differentiation of *Claviceps purpurea* in axenic culture. *J. Gen. Microbiol.* 93, 321-334.
- [0141] Mey, G., Oeser, B., Lebrun, M. H., and Tudzynski, P. (2002). The biotrophic, non-appressorium-forming grass pathogen *Claviceps purpurea* needs a Fus3/Pmk1 homologous mitogen-activated protein kinase for colonization of rye ovarian tissue. *Mol. Plant Microbe Interact.* 15, 303-312.
- [0142] Okuley, J., Lightner, J., Feldmann, K., Yadav, N., Lark, E., and Browse, J. (1994). *Arabidopsis* FAD2 gene encodes the enzyme that is essential for polyunsaturated lipid synthesis. *Plant Cell* 6, 147-158.

- [0143] Qi, B., Fraser, T., Mugford, S., Dobson, G., Sayanova, O., Butler, J., Napier, J. A., Stobart, A. K., and Lazarus, C. M. (2004). Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants. *Nat. Biotechnol.* 22, 739-745.
- [0144] Qiu, X., Hong, H., and McKenzie, S.L. (2001). Identification of a Delta 4 fatty acid desaturase from *Thraustochytrium* sp. involved in the biosynthesis of docosahexanoic acid by heterologous expression in *Saccharomyces cerevisiae* and *Brassica juncea*. *J Biol Chem* 276, 31561-6.
- [0145] Shanklin, J. and Cahoon, E. B. (1998). DESATURATION AND RELATED MODIFICATIONS OF FATTY ACIDS1. *Annu. Rev. Plant Physiol Plant Mol. Biol.* 49, 611-641.
- [0146] Tudzynski, P., Correia, T., and Keller, U. (2001). Biotechnology and genetics of ergot alkaloids. *Appl. Microbiol. Biotechnol.* 57, 593-605.
- [0147] All references cited in this specification are hereby incorporated by reference with respect to their entire disclosure content and the disclosure content specifically mentioned in this specification.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 32

<210> SEQ ID NO 1

<211> LENGTH: 1256

<212> TYPE: DNA

<213> ORGANISM: *Emiliana huxleyi*

<400> SEQUENCE: 1

```
ccatgggcaa gggcggaac gcgaacccgc gggagctcaa aggcggcaag gccgagcagc    60
tgacagtcta cctgtatggc aaggctgtcg acgtctcgaa gtctcggaag ctgcacccgg    120
gaggcgccaa ggcgtgtcgc atcttcaaca accgcgacgc caccgagcag ttcgagatgt    180
accactcgcc cgccgcccac aagatgatgc gtgcgatgtc gaagagcgcg ccggaggccc    240
cgaggggagag cgaggtcgcg acgtcggtcg ttgggacgga ctctcgccaag ctgacgcaga    300
cgctgcacga cgctggatgc ttcgacccgc actaccctga cgaggccttc aagctcggcc    360
tcacgtgtgt gcccggtatc ctgcgcttct acctgtgtcg gagcggcatg ccggcgctcg    420
gataccttct gatcgctttc tcgtactaca tgctggggtg gacctccac gattacttgc    480
accacggctg cctcaagggc ggccaaaagc agctggtgca ctggaacaac gccgtcggct    540
acgcaatcgg cgcttgccag ggctacgcgg tcggctggtg gcgagcgcgc cacaacacgc    600
accacctcgt cactaacgaa gaaggcaacg accccgacat catgaccgcg cccgtgctca    660
tcttcgtgcg caacagcccc gtgatgcgcg ctgccctcaa cgcggcgag cggtggcagc    720
agtactacta cgtgcccgcg atgagcctca tggacatgta ctggcgcttc gactcgatgc    780
agtacctggc cgcgcgaccc ttcaacaagg tgtgggcctc gtggcgctc ctgcgctgctc    840
actactcctt tgctggctac atgttccacg gacagtacca gtggctgctg ctgacgatgc    900
tggtgcgcgg ctctctcacg ggcatcgtcg tcttctcgac gcattatggc gaggaggtca    960
tcccgggcga ccacggcatg aactcgtcg agcagacggc gctcactct cgcaacatca    1020
ccggcgggta cctcgtcaac ctgctcacgg gctacatctc gctgcagacg gagcaccacc    1080
tctggccgat gatgccacc gcgcgcctcg aggcggcgca gccctacgcg cgcgccttct    1140
tcaagaagca cggcttcgtc taccgcgagt cgaacctcgt cgagtgcgtc aagtacaaca    1200
tcgccgccct cgacatccgc acgcgcaacg gcgagtgggc agagatgccg cactag      1256
```

<210> SEQ ID NO 2

<211> LENGTH: 417

<212> TYPE: PRT

<213> ORGANISM: *Emiliana huxleyi*

<400> SEQUENCE: 2

Met Gly Lys Gly Gly Asn Ala Asn Pro Arg Glu Leu Lys Gly Gly Lys

-continued

1	5	10	15
Ala Glu Gln Leu Thr Val Tyr Leu Tyr Gly Lys Ala Val Asp Val Ser	20	25	30
Lys Phe Ala Lys Leu His Pro Gly Gly Ala Lys Ala Leu Arg Ile Phe	35	40	45
Asn Asn Arg Asp Ala Thr Glu Gln Phe Glu Met Tyr His Ser Pro Ala	50	55	60
Ala His Lys Met Met Arg Ala Met Ser Lys Ser Ala Pro Glu Ala Pro	65	70	80
Arg Glu Ser Glu Val Ala Thr Ser Val Val Gly Thr Asp Phe Ala Lys	85	90	95
Leu Thr Gln Thr Leu His Asp Val Gly Cys Phe Asp Pro His Tyr Pro	100	105	110
Asp Glu Ala Phe Lys Leu Gly Leu Thr Leu Leu Pro Gly Phe Leu Gly	115	120	125
Phe Tyr Leu Leu Arg Ser Gly Met Pro Ala Leu Gly Ser Phe Leu Ile	130	135	140
Ala Phe Ser Tyr Tyr Met Ser Gly Trp Thr Ser His Asp Tyr Leu His	145	150	155
His Gly Cys Leu Lys Gly Gly Gln Lys Gln Leu Val His Trp Asn Asn	165	170	175
Ala Val Gly Tyr Ala Ile Gly Ala Trp Gln Gly Tyr Ala Val Gly Trp	180	185	190
Trp Arg Ala Arg His Asn Thr His His Leu Val Thr Asn Glu Glu Gly	195	200	205
Asn Asp Pro Asp Ile Met Thr Ala Pro Val Leu Ile Phe Val Arg Asn	210	215	220
Ser Pro Val Ile Ala Ala Leu Asn Ala Ala Gln Arg Trp Gln Gln	225	230	235
Tyr Tyr Tyr Val Pro Ala Met Ser Leu Met Asp Met Tyr Trp Arg Phe	245	250	255
Glu Ser Met Gln Tyr Leu Ala Ala Arg Pro Phe Asn Lys Val Trp Ala	260	265	270
Ser Trp Ala Leu Leu Ala Leu His Tyr Ser Phe Val Gly Tyr Met Phe	275	280	285
His Gly Gln Tyr Gln Trp Leu Leu Leu Thr Met Leu Val Arg Gly Phe	290	295	300
Leu Thr Gly Ile Val Val Phe Ser Thr His Tyr Gly Glu Glu Val Ile	305	310	315
Pro Gly Asp His Gly Met Thr Leu Val Glu Gln Thr Ala Leu Thr Ser	325	330	335
Arg Asn Ile Thr Gly Gly Tyr Leu Val Asn Leu Leu Thr Gly Tyr Ile	340	345	350
Ser Leu Gln Thr Glu His His Leu Trp Pro Met Met Pro Thr Ala Arg	355	360	365
Leu Glu Ala Ala Gln Pro Tyr Ala Arg Ala Phe Phe Lys Lys His Gly	370	375	380
Phe Val Tyr Arg Glu Ser Asn Leu Val Glu Cys Val Lys Tyr Asn Ile	385	390	395
Ala Ala Leu Asp Ile Arg Thr Arg Asn Gly Glu Trp Ala Glu Met Pro	405	410	415



---

-continued

---

His

<210> SEQ ID NO 3  
<211> LENGTH: 804  
<212> TYPE: DNA  
<213> ORGANISM: Emiliana huxleyi

&lt;400&gt; SEQUENCE: 3

```
accatgctcg atcgcgccctc gtccgacgcg gccatctggt ctgcggtgtc cgatccggaa    60
atcctgatcg gcactttctc ctacctgctg ctcaagccgc tgcacgcaa ctcagggtctc    120
gtggacgagc ggaaaggcgc ctaccggacc tcgatgatct ggtacaacgt ggtgctcgcg    180
ctcttctccg cgacgagctt ctacgtgact gcgaccgcgc tcgggtggga caagggcacc    240
ggcgagtggc tccgcagtct cacggggcgc agcccgcagc agctgtggca atgcccgctc    300
agggtatggg actccaagct gttcctgtgg acggccaagg ccttctacta ctcaaagtac    360
gtggagtacc tcgacacggc gtggctcgtc ctcaagggga agaaggtctc cttcctgcag    420
ggcttcacc actttggcgc gccgtgggac gtgtacctgg gcattcggct gaagaacgag    480
ggcgtgtgga tcttcagtgt cttcaactcg ttcateccaca cggtcagtga cagctactac    540
ggcctcaccg ccgcgggcta caagatccgc ggcaagccga tcaccaccgc gatgcaaata    600
agccagttcg tcggcggttt tgtcctagtg tgggactaca tcaacgtgcc gtgcttcacc    660
gccgacgccg ggcaggtctt cagctgggtc tttaactatg cttacgtcgg ctccgtcttt    720
ctgctctttt gccacttctt ctacatggac aacatcgcca aggccaaaggc caagaaggcc    780
gtcgctaccc gcaaggcgct gtga                                           804
```

<210> SEQ ID NO 4  
<211> LENGTH: 266  
<212> TYPE: PRT  
<213> ORGANISM: Emiliana huxleyi

&lt;400&gt; SEQUENCE: 4

```
Met Leu Asp Arg Ala Ser Ser Asp Ala Ala Ile Trp Ser Ala Val Ser      1
      5              10              15
Asp Pro Glu Ile Leu Ile Gly Thr Phe Ser Tyr Leu Leu Leu Lys Pro      20
      25              30
Leu Leu Arg Asn Ser Gly Leu Val Asp Glu Arg Lys Gly Ala Tyr Arg      35
      40              45
Thr Ser Met Ile Trp Tyr Asn Val Val Leu Ala Leu Phe Ser Ala Thr      50
      55              60
Ser Phe Tyr Val Thr Ala Thr Ala Leu Gly Trp Asp Lys Gly Thr Gly      65
      70              75              80
Glu Trp Leu Arg Ser Leu Thr Gly Asp Ser Pro Gln Gln Leu Trp Gln      85
      90              95
Cys Pro Ser Arg Val Trp Asp Ser Lys Leu Phe Leu Trp Thr Ala Lys     100
      105              110
Ala Phe Tyr Tyr Ser Lys Tyr Val Glu Tyr Leu Asp Thr Ala Trp Leu     115
      120              125
Val Leu Lys Gly Lys Lys Val Ser Phe Leu Gln Gly Phe His His Phe     130
      135              140
Gly Ala Pro Trp Asp Val Tyr Leu Gly Ile Arg Leu Lys Asn Glu Gly     145
      150              155              160
```

-continued

---

Val Trp Ile Phe Met Phe Phe Asn Ser Phe Ile His Thr Val Met Tyr  
                   165                  170                  175

Thr Tyr Tyr Gly Leu Thr Ala Ala Gly Tyr Lys Ile Arg Gly Lys Pro  
                   180                  185                  190

Ile Ile Thr Ala Met Gln Ile Ser Gln Phe Val Gly Gly Phe Val Leu  
                   195                  200                  205

Val Trp Asp Tyr Ile Asn Val Pro Cys Phe His Ala Asp Ala Gly Gln  
                   210                  215                  220

Val Phe Ser Trp Val Phe Asn Tyr Ala Tyr Val Gly Ser Val Phe Leu  
                   225                  230                  235                  240

Leu Phe Cys His Phe Phe Tyr Met Asp Asn Ile Ala Lys Ala Lys Ala  
                   245                  250                  255

Lys Lys Ala Val Ala Thr Arg Lys Ala Leu  
                   260                  265

<210> SEQ ID NO 5  
 <211> LENGTH: 1280  
 <212> TYPE: DNA  
 <213> ORGANISM: Emiliana huxleyi

<400> SEQUENCE: 5

```
ccatgggagg cgccggcgcg agcgaggctg aacggcccaa gtggaccacg atccacgggc      60
ggcacgtcga tgtgtcaaaag ttccgccacc cgggtgggaa catcatcgag ctcttctatg      120
gcattgactc gacgagcgcg ttcgagcagt tccacggcca ccacaagggc gcgtggaaga      180
tgctcaaggc gctgccgacc aaggaggtcg accccgccga cgtgcccgag cagccgcagg      240
agcacgttgc cgagatgacg cggctgatga cgtcgtggcg cgagcgcggc ctctttaagc      300
cgcgccccgt cgctcgggac atctacggtc tcgcccgtcg cgtgccatc gtcgctgca      360
tcgctcgcg cgccgcacgc cgggtgctga gcgggatcgg gctcggcagc tgctgggcgc      420
agtgcggctt cctgcagcac atggggcggc accgcgagtg ggggggtcgg tactccttcc      480
tcctgcagca ctctctcgag ggcctcctca agggcgggtc cgcctcgtgg tggcgcaacc      540
gccacaacaa gcatcacgca aagactaacg tgctcggcga ggacggcgac ctgcggacga      600
ctcccttctt cgcctgggac ccgacgctcg ccaagaaggt tccagactgg tcgctcaaga      660
cgcaggcctt cactctctcc ccgcccctcg gacgctacgt ctttgtcttt gccttcacga      720
tccgcaagta tgcgctcgtc aagaagctct ggacagagct cgcactcatg atcgcgact      780
acgcgatggt ctactacgcg ctgcagctcg ccggtgcgtc gctcggcagc ggctcgcct      840
tttactgcac cggctacgcc tggcaaggca tctacctcgg cttcttcttc ggctgtccc      900
acttcgcggc cgagcgagtc cctccacccg ccacctggct cgagtcgtcc atgateggca      960
ccgtcgactg gggaggctcc tccgctttt gcggctacgt ctccggcttc ctcaacatcc     1020
agatcgagca ccacatggcg ccgcagatgc cgatggagaa cctgcgccag atccgcgcgc     1080
actgcaaggc gagcgcgagg aagctcgggc ttccctatcg cgagctctcc ttgcgcggcg     1140
cggtaagct gatgatggtc ggcctctggc gcacggggag ggacgagctg cagctgcgct     1200
ccgacaggcg caagtactcg cgcacccagg cctacatggc ggccgcctcg gcggtggtgg     1260
agaacctcaa ggcggactag                                     1280
```

<210> SEQ ID NO 6

-continued

&lt;211&gt; LENGTH: 425

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Emiliana huxleyi

&lt;400&gt; SEQUENCE: 6

```

Met Gly Gly Ala Gly Ala Ser Glu Ala Glu Arg Pro Lys Trp Thr Thr
1          5          10          15
Ile His Gly Arg His Val Asp Val Ser Lys Phe Arg His Pro Gly Gly
20          25          30
Asn Ile Ile Glu Leu Phe Tyr Gly Met Asp Ser Thr Ser Ala Phe Glu
35          40          45
Gln Phe His Gly His His Lys Gly Ala Trp Lys Met Leu Lys Ala Leu
50          55          60
Pro Thr Lys Glu Val Asp Pro Ala Asp Val Pro Gln Gln Pro Gln Glu
65          70          75          80
His Val Ala Glu Met Thr Arg Leu Met Thr Ser Trp Arg Glu Arg Gly
85          90          95
Leu Phe Lys Pro Arg Pro Val Ala Ser Gly Ile Tyr Gly Leu Ala Val
100         105         110
Val Ala Ala Ile Val Ala Cys Ile Ala Cys Ala Pro His Ala Pro Val
115         120         125
Leu Ser Gly Ile Gly Leu Gly Ser Cys Trp Ala Gln Cys Gly Phe Leu
130         135         140
Gln His Met Gly Gly His Arg Glu Trp Gly Val Arg Tyr Ser Phe Leu
145         150         155         160
Leu Gln His Phe Phe Glu Gly Leu Leu Lys Gly Gly Ser Ala Ser Trp
165         170         175
Trp Arg Asn Arg His Asn Lys His His Ala Lys Thr Asn Val Leu Gly
180         185         190
Glu Asp Gly Asp Leu Arg Thr Thr Pro Phe Phe Ala Trp Asp Pro Thr
195         200         205
Leu Ala Lys Lys Val Pro Asp Trp Ser Leu Lys Thr Gln Ala Phe Thr
210         215         220
Phe Leu Pro Ala Leu Gly Ala Tyr Val Phe Val Phe Ala Phe Thr Ile
225         230         235         240
Arg Lys Tyr Ala Val Val Lys Lys Leu Trp His Glu Leu Ala Leu Met
245         250         255
Ile Ala His Tyr Ala Met Phe Tyr Tyr Ala Leu Gln Leu Ala Gly Ala
260         265         270
Ser Leu Gly Ser Gly Leu Ala Phe Tyr Cys Thr Gly Tyr Ala Trp Gln
275         280         285
Gly Ile Tyr Leu Gly Phe Phe Phe Gly Leu Ser His Phe Ala Val Glu
290         295         300
Arg Val Pro Ser Thr Ala Thr Trp Leu Glu Ser Ser Met Ile Gly Thr
305         310         315         320
Val Asp Trp Gly Gly Ser Ser Ala Phe Cys Gly Tyr Val Ser Gly Phe
325         330         335
Leu Asn Ile Gln Ile Glu His His Met Ala Pro Gln Met Pro Met Glu
340         345         350
Asn Leu Arg Gln Ile Arg Ala Asp Cys Lys Ala Ser Ala Glu Lys Leu
355         360         365
Gly Leu Pro Tyr Arg Glu Leu Ser Phe Ala Gly Ala Val Lys Leu Met

```

-continued

---

370	375	380	
Met Val Gly Leu Trp Arg Thr Gly Arg Asp Glu Leu Gln Leu Arg Ser			
385	390	395	400
Asp Arg Arg Lys Tyr Ser Arg Thr Gln Ala Tyr Met Ala Ala Ala Ser			
	405	410	415
Ala Val Val Glu Asn Leu Lys Ala Asp			
	420	425	

<210> SEQ ID NO 7  
 <211> LENGTH: 921  
 <212> TYPE: DNA  
 <213> ORGANISM: Emiliana huxleyi

<400> SEQUENCE: 7

accatgtgta aggcctcttg tcttgcttca ggtgctaaac ctgctgctgc ttcaactatt	60
gatcagctcg ctggacttgg aagagtgtgt gttattgttg gatctttcac tgctgctatg	120
tggtatgctc ttcaacctct tgattcacct ggtactatct atcatgattc agctgttatg	180
ggtgctcttt tgtcttggcc aatgggttac attgctcctc ttgcttacgt ttgtgctgtt	240
atggctggat gtacacttat gtcacaaaga gcttctatta agccattttt gaaacaatac	300
gttcagcctg ttacaatgt ttcccaaatt gttatgtgtt cttacatggt ttgggggttg	360
gctcctaaag ttgatgttct tggacttaac cctttcgcta tgaatacaga aagagataaa	420
aagactgagt ggtttatgtt cggttcattac ctttctaaat tcgttgattg gacagatact	480
ttcttgatga ttggatctaa atcttttaga cagggttcat tcttgcaagt ttttcatcat	540
gctacagttg gtatgatatt gggtgctttg ttgagaaagg gatggggttg aggtacttgt	600
gtttggggag cttttattaa ctctgttaca catgttctta tgtatacaca ttacttggtt	660
acatctcttg gtcttcataa cctctttaag tctcaactta ctaattttca acttgctcaa	720
ttcgcttcat gtgttttgca tgctgctttg gtttttgctt cagagacagt tcttcctgct	780
agacttgctt atattcaatt gggttaccat cctactcttt tgtttctttt cggttttcag	840
atgaagtggg ttccttcttg gatcactgga caaacaatca ctggtagaga gtcagaggct	900
cctgaaaaga aagttgcttg a	921

<210> SEQ ID NO 8  
 <211> LENGTH: 305  
 <212> TYPE: PRT  
 <213> ORGANISM: Emiliana huxleyi

<400> SEQUENCE: 8

Met Cys Lys Ala Ser Gly Leu Ala Ser Gly Ala Lys Pro Ala Ala Ala	
1	15
Ser Thr Ile Asp Gln Ser Ala Gly Leu Gly Arg Val Ala Val Ile Val	
20	30
Gly Ser Phe Thr Ala Ala Met Cys Tyr Ala Leu Gln Pro Leu Asp Ser	
35	45
Pro Gly Thr Ile Tyr His Asp Ser Ala Val Met Gly Ala Leu Leu Ser	
50	60
Trp Pro Met Val Tyr Ile Ala Pro Leu Ala Tyr Val Cys Ala Val Met	
65	80
Ala Gly Cys Arg Leu Met Ser Gln Arg Ala Ser Ile Lys Pro Phe Leu	
85	95

-continued

---

Lys Gln Tyr Val Gln Pro Val Tyr Asn Val Phe Gln Ile Val Met Cys  
                   100                  105                  110  
 Ser Tyr Met Val Trp Gly Leu Ala Pro Lys Val Asp Val Leu Gly Leu  
                   115                  120                  125  
 Asn Pro Phe Ala Met Asn Thr Glu Arg Asp Lys Lys Thr Glu Trp Phe  
                   130                  135                  140  
 Met Phe Val His Tyr Leu Ser Lys Phe Val Asp Trp Thr Asp Thr Phe  
                   145                  150                  155                  160  
 Leu Met Ile Gly Ser Lys Ser Phe Arg Gln Val Ser Phe Leu Gln Val  
                   165                  170                  175  
 Phe His His Ala Thr Val Gly Met Ile Trp Gly Ala Leu Leu Arg Lys  
                   180                  185                  190  
 Gly Trp Gly Gly Gly Thr Cys Val Trp Gly Ala Phe Ile Asn Ser Val  
                   195                  200                  205  
 Thr His Val Leu Met Tyr Thr His Tyr Leu Val Thr Ser Leu Gly Leu  
                   210                  215                  220  
 His Asn Pro Leu Lys Ser Gln Leu Thr Asn Phe Gln Leu Ala Gln Phe  
                   225                  230                  235                  240  
 Ala Ser Cys Val Leu His Ala Ala Leu Val Phe Ala Ser Glu Thr Val  
                   245                  250                  255  
 Leu Pro Ala Arg Leu Ala Tyr Ile Gln Leu Val Tyr His Pro Thr Leu  
                   260                  265                  270  
 Leu Phe Leu Phe Gly Phe Gln Met Lys Trp Val Pro Ser Trp Ile Thr  
                   275                  280                  285  
 Gly Gln Thr Ile Thr Gly Arg Glu Ser Glu Ala Pro Glu Lys Lys Val  
                   290                  295                  300  
 Ala  
 305

<210> SEQ ID NO 9  
 <211> LENGTH: 1368  
 <212> TYPE: DNA  
 <213> ORGANISM: Emiliana huxleyi

<400> SEQUENCE: 9

atgtcattgg ctgctaaga tgcagcctcg gccactcat ccgtcttgga ccctaagtat	60
cacggagcta caaataagtc aagaactgat gcagcagacc ttacagttag ttctatcgac	120
acttctaagg agatgatcat aaggggtcgt gtgtatgatg tctctgattt tattaagg	180
cacccgggag gaagcattat taaactctcc ttaggttctg atgcaacaga cgcttataac	240
aacttcata ttaggtctaa aaaagcggat aaaatgttga gagctttgcc aagtaggcca	300
gtagcggatg gattcgctag agacgctttg tctgcagact tcgaggccct gagagcccaa	360
ctcgaggccg aaggttactt cgaaccgaat ctgtggcatg tagcttatcg agttgaggaa	420
gtcgttgcta tgtactgggc ggggtattaga cttatctggg cgggttattg gtttttagga	480
gccattgtag caggaataga tcaggggaga tgcggttggc ttcagcatga ggggtgtcat	540
tattcgctca caggaatat taaacttgat cgacacatgc aaatgattat ctatggatta	600
ggttgcggaa tgtccggttg ttattggaga aaccaacata acaagcacca tgcgacaccg	660
caaaagtgg gtgcagatcc agacctcaa acaatgcctc tggttgcgtt ccatggactc	720
atcggtgcta aggctagggg agcaggaaag tcgtggctag catggcaagc tccacttttc	780

-continued

---

```

tttggaggcg ttatcacaac cctggatatct tttggttggc agttcgtcca acatccaaag      840
cacgcattga gagtaggaaa ccaactcgaa ttaggctata tggctttacg atatgcttta      900
tggtagtcag cattcgggtca tcttgggctt ggtggtgctt tcagattgta cgctttttat      960
gtggcagtcg gaggtacata tatcttcacg aactttgcgg tgtctcacac acataaggat     1020
gttgttccac acgataagca tatttcttgg accttgtatt ctgcaaacca taccactaat     1080
caatctaaca cacctctagt caattggtgg atggcctatc tgaattttca aattgaacat     1140
caccttttcc ctagcatgcc acaatataac catcctaaaa tctgcggaag agtgaacaaa     1200
ttgtttgaaa aacatggcgt agagtacgat gtcagaactt acgcgaagtc aatgcgtgat     1260
acatacgtga atctcttggc tgtgggaaat gcatttcatt cccttcacga gagaaacgag     1320
ggattaacga ctaggagatc tgcggctggt agagttacag gtcattga                    1368

```

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 455

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Emiliana huxleyi

&lt;400&gt; SEQUENCE: 10

```

Met Ser Leu Ala Ala Lys Asp Ala Ala Ser Ala His Ser Ser Val Leu
1           5           10           15
Asp Pro Lys Tyr His Gly Ala Thr Asn Lys Ser Arg Thr Asp Ala Ala
20          25          30
Asp Leu Thr Val Ser Ser Ile Asp Thr Ser Lys Glu Met Ile Ile Arg
35          40          45
Gly Arg Val Tyr Asp Val Ser Asp Phe Ile Lys Arg His Pro Gly Gly
50          55          60
Ser Ile Ile Lys Leu Ser Leu Gly Ser Asp Ala Thr Asp Ala Tyr Asn
65          70          75          80
Asn Phe His Ile Arg Ser Lys Lys Ala Asp Lys Met Leu Arg Ala Leu
85          90          95
Pro Ser Arg Pro Val Ala Asp Gly Phe Ala Arg Asp Ala Leu Ser Ala
100         105         110
Asp Phe Glu Ala Leu Arg Ala Gln Leu Glu Ala Glu Gly Tyr Phe Glu
115         120         125
Pro Asn Leu Trp His Val Ala Tyr Arg Val Ala Glu Val Val Ala Met
130         135         140
Tyr Trp Ala Gly Ile Arg Leu Ile Trp Ala Gly Tyr Trp Phe Leu Gly
145         150         155         160
Ala Ile Val Ala Gly Ile Ala Gln Gly Arg Cys Gly Trp Leu Gln His
165         170         175
Glu Gly Gly His Tyr Ser Leu Thr Gly Asn Ile Lys Leu Asp Arg His
180         185         190
Met Gln Met Ile Ile Tyr Gly Leu Gly Cys Gly Met Ser Gly Cys Tyr
195         200         205
Trp Arg Asn Gln His Asn Lys His His Ala Thr Pro Gln Lys Leu Gly
210         215         220
Ala Asp Pro Asp Leu Gln Thr Met Pro Leu Val Ala Phe His Gly Leu
225         230         235         240
Ile Gly Ala Lys Ala Arg Gly Ala Gly Lys Ser Trp Leu Ala Trp Gln
245         250         255

```

-continued

---

Ala Pro Leu Phe Phe Gly Gly Val Ile Thr Thr Leu Val Ser Phe Gly  
260 265 270

Trp Gln Phe Val Gln His Pro Lys His Ala Leu Arg Val Gly Asn Gln  
275 280 285

Leu Glu Leu Gly Tyr Met Ala Leu Arg Tyr Ala Leu Trp Tyr Ala Ala  
290 295 300

Phe Gly His Leu Gly Leu Gly Gly Ala Phe Arg Leu Tyr Ala Phe Tyr  
305 310 315 320

Val Ala Val Gly Gly Thr Tyr Ile Phe Thr Asn Phe Ala Val Ser His  
325 330 335

Thr His Lys Asp Val Val Pro His Asp Lys His Ile Ser Trp Thr Leu  
340 345 350

Tyr Ser Ala Asn His Thr Thr Asn Gln Ser Asn Thr Pro Leu Val Asn  
355 360 365

Trp Trp Met Ala Tyr Leu Asn Phe Gln Ile Glu His His Leu Phe Pro  
370 375 380

Ser Met Pro Gln Tyr Asn His Pro Lys Ile Cys Gly Arg Val Lys Gln  
385 390 395 400

Leu Phe Glu Lys His Gly Val Glu Tyr Asp Val Arg Thr Tyr Ala Lys  
405 410 415

Ser Met Arg Asp Thr Tyr Val Asn Leu Leu Ala Val Gly Asn Ala Ser  
420 425 430

His Ser Leu His Gln Arg Asn Glu Gly Leu Thr Thr Arg Glu Ser Ala  
435 440 445

Ala Val Arg Val Thr Gly His  
450 455

<210> SEQ ID NO 11  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 11

ccatgggagg cgccggcgcg ag

22

<210> SEQ ID NO 12  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 12

ctagtccgcc ttgaggttct c

21

<210> SEQ ID NO 13  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 13

accatgtgca aggcgagcgg cct

23

---

-continued

---

<210> SEQ ID NO 14  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 14

tcaccaatca tgaggaaggt 20

<210> SEQ ID NO 15  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 15

ccatgggcaa gggcggcaac gc 22

<210> SEQ ID NO 16  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 16

gggcagagat gccgcactag 20

<210> SEQ ID NO 17  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 17

accatgctcg atcgccctc gtc 23

<210> SEQ ID NO 18  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic

<400> SEQUENCE: 18

tcacagcgcc ttgcgggtag c 21

<210> SEQ ID NO 19  
<211> LENGTH: 1371  
<212> TYPE: DNA  
<213> ORGANISM: *Ostreococcus tauri*

<400> SEQUENCE: 19

atgtgtgttg agaccgagaa caacgatgga atccctactg tggagatcgc ttctgatgga 60

gagagagaaa gagctgaggg taacgtgaag ttgtctgctg agaagatgga acctgctgct 120

ttggctaaga ccttcgctag aagatacgtg gttatcgagg gagttgagta cgatgtgacc 180

gatttcaaac atcctggagg aacctgtgatt ttctacgctc tctctaacac tggagctgat 240



-continued

---

```

gctactgagg ctttcaagga gttccaccac agatctagaa aggctaggaa ggctttggct 300
gctttgcctt ctagacctgc taagaccgct aaagtggatg atgctgagat gctccaggat 360
ttcgctaagt ggagaaagga gttggagagg gacggattct tcaagccttc tctgtctcat 420
gttgcttaca gattcgctga gttggctgct atgtacgctt tgggaacctc cttgatgtac 480
gctagatacg ttgtgtcctc tgtgttggtt tacgcttgct tcttcggagc tagatgtgga 540
tgggttcaac atgagggagg acattcttct ttgaccggaa acatctggtg ggataagaga 600
atccaagctt tcactgctgg attcggattg gctggatctg gagatatgtg gaactccatg 660
cacaacaagc accatgctac tctcaaaaa gtgaggcacg atatggattt ggataccact 720
cctgctgttg ctttcttcaa caccgctgtg gaggataata gacctagggg attctctaag 780
tactggctca gattgcaagc ttggaccttc attcctgtga cttctggatt ggtgttgctc 840
ttctggatgt tcttctcca tcttctaag gctttgaagg gaggaagta cgaggagctt 900
gtgtggatgt tggctgctca tgtgattaga acctggacca ttaaggctgt tactggattc 960
accgctatgc aatcctacgg actcttcttg gctacttctt gggtttccgg atgctacttg 1020
ttcgctcact tctctacttc tcacacccat ttggatgttg ttctgctga tgagcatttg 1080
tcttgggtta ggtacgctgt ggatcacacc attgatatcg atccttctca gggatgggtt 1140
aactggttga tgggatactt gaactgcaa gtgattcatc acctcttccc ttctatgctt 1200
caattcagac aacctgaggt gtccagaaga ttcgttgctt tcgctaagaa gtggaacctc 1260
aactacaagg tgatgactta tgctggagct tggaaggcta ctttgggaaa cctcgataat 1320
gtgggaaagc actactacgt gcacggacaa cattctggaa agaccgcttg a 1371

```

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 456

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Ostreococcus tauri*

&lt;400&gt; SEQUENCE: 20

```

Met Cys Val Glu Thr Glu Asn Asn Asp Gly Ile Pro Thr Val Glu Ile
1           5           10          15

Ala Phe Asp Gly Glu Arg Glu Arg Ala Glu Ala Asn Val Lys Leu Ser
20          25          30

Ala Glu Lys Met Glu Pro Ala Ala Leu Ala Lys Thr Phe Ala Arg Arg
35          40          45

Tyr Val Val Ile Glu Gly Val Glu Tyr Asp Val Thr Asp Phe Lys His
50          55          60

Pro Gly Gly Thr Val Ile Phe Tyr Ala Leu Ser Asn Thr Gly Ala Asp
65          70          75          80

Ala Thr Glu Ala Phe Lys Glu Phe His His Arg Ser Arg Lys Ala Arg
85          90          95

Lys Ala Leu Ala Ala Leu Pro Ser Arg Pro Ala Lys Thr Ala Lys Val
100         105         110

Asp Asp Ala Glu Met Leu Gln Asp Phe Ala Lys Trp Arg Lys Glu Leu
115         120         125

Glu Arg Asp Gly Phe Phe Lys Pro Ser Pro Ala His Val Ala Tyr Arg
130         135         140

Phe Ala Glu Leu Ala Ala Met Tyr Ala Leu Gly Thr Tyr Leu Met Tyr
145         150         155         160

```

-continued

Ala	Arg	Tyr	Val	Val	Ser	Ser	Val	Leu	Val	Tyr	Ala	Cys	Phe	Phe	Gly	
				165					170						175	
Ala	Arg	Cys	Gly	Trp	Val	Gln	His	Glu	Gly	Gly	His	Ser	Ser	Leu	Thr	
			180						185					190		
Gly	Asn	Ile	Trp	Trp	Asp	Lys	Arg	Ile	Gln	Ala	Phe	Thr	Ala	Gly	Phe	
		195					200						205			
Gly	Leu	Ala	Gly	Ser	Gly	Asp	Met	Trp	Asn	Ser	Met	His	Asn	Lys	His	
	210					215					220					
His	Ala	Thr	Pro	Gln	Lys	Val	Arg	His	Asp	Met	Asp	Leu	Asp	Thr	Thr	
	225				230					235					240	
Pro	Ala	Val	Ala	Phe	Phe	Asn	Thr	Ala	Val	Glu	Asp	Asn	Arg	Pro	Arg	
			245						250					255		
Gly	Phe	Ser	Lys	Tyr	Trp	Leu	Arg	Leu	Gln	Ala	Trp	Thr	Phe	Ile	Pro	
		260					265						270			
Val	Thr	Ser	Gly	Leu	Val	Leu	Leu	Phe	Trp	Met	Phe	Phe	Leu	His	Pro	
		275					280					285				
Ser	Lys	Ala	Leu	Lys	Gly	Gly	Lys	Tyr	Glu	Glu	Leu	Val	Trp	Met	Leu	
	290						295				300					
Ala	Ala	His	Val	Ile	Arg	Thr	Trp	Thr	Ile	Lys	Ala	Val	Thr	Gly	Phe	
	305				310					315					320	
Thr	Ala	Met	Gln	Ser	Tyr	Gly	Leu	Phe	Leu	Ala	Thr	Ser	Trp	Val	Ser	
			325						330					335		
Gly	Cys	Tyr	Leu	Phe	Ala	His	Phe	Ser	Thr	Ser	His	Thr	His	Leu	Asp	
		340						345					350			
Val	Val	Pro	Ala	Asp	Glu	His	Leu	Ser	Trp	Val	Arg	Tyr	Ala	Val	Asp	
		355					360					365				
His	Thr	Ile	Asp	Ile	Asp	Pro	Ser	Gln	Gly	Trp	Val	Asn	Trp	Leu	Met	
	370					375					380					
Gly	Tyr	Leu	Asn	Cys	Gln	Val	Ile	His	His	Leu	Phe	Pro	Ser	Met	Pro	
	385				390					395					400	
Gln	Phe	Arg	Gln	Pro	Glu	Val	Ser	Arg	Arg	Phe	Val	Ala	Phe	Ala	Lys	
			405						410					415		
Lys	Trp	Asn	Leu	Asn	Tyr	Lys	Val	Met	Thr	Tyr	Ala	Gly	Ala	Trp	Lys	
		420						425				430				
Ala	Thr	Leu	Gly	Asn	Leu	Asp	Asn	Val	Gly	Lys	His	Tyr	Tyr	Val	His	
		435				440					445					
Gly	Gln	His	Ser	Gly	Lys	Thr	Ala									
	450					455										

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 873

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Physcomitrella patens

&lt;400&gt; SEQUENCE: 21

atggaagtgtg	ttgagagggt	ctacggagag	ttggatggaa	aggtttccca	aggagtgaac	60
gctttgttgg	gatctttcgg	agttgagttg	actgataccc	caactactaa	gggattgcca	120
ctcgttgatt	ctccaactcc	aattgtgttg	ggagtgtctg	tttacttgac	catcgtgatc	180
ggaggattgc	tttggatcaa	ggctagagat	ctcaagccaa	gagcttctga	gccattcttg	240
ttgcaagctt	tggtgttggt	gcacaacttg	ttctgcttcg	ctttgtctct	ttacatgtgc	300
gtgggtatcg	cttaccaagc	tatcacctgg	agatattcct	tgtggggaaa	cgcttataac	360

-continued

```

ccaaagcaca aggagatggc tatcctcggt tacctcttct acatgtccaa gtacgtggag 420
ttcatggata ccgatgcat gatcctcaag agatccacca gacagatttc ttctctccac 480
gtgtaccacc attcttctat ctcccttate tgggtgggcta ttgtcatca tgctccagga 540
ggagaggcctt attggagtgc tgctctcaac tctggagtgc atgtgttgat gtacgcttac 600
tacttcttgg ctgcttgctt gagatcttcc ccaaagctca agaacaagta cctcttctgg 660
ggaagatacc tcaccaatt ccagatgttc cagttcatgc tcaacttggg gcaagcttac 720
tacgatatga aaaccaacgc tccatatcca caatggctca tcaagatcct cttctactac 780
atgatctccc tcttgcttct cttcggaaac ttctacgtgc aaaagtacat caagccatcc 840
gatggaaagc aaaaggagc taagaccgag tga 873

```

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 290

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Physcomitrella patens*

&lt;400&gt; SEQUENCE: 22

```

Met Glu Val Val Glu Arg Phe Tyr Gly Glu Leu Asp Gly Lys Val Ser
1           5           10          15
Gln Gly Val Asn Ala Leu Leu Gly Ser Phe Gly Val Glu Leu Thr Asp
20          25          30
Thr Pro Thr Thr Lys Gly Leu Pro Leu Val Asp Ser Pro Thr Pro Ile
35          40          45
Val Leu Gly Val Ser Val Tyr Leu Thr Ile Val Ile Gly Gly Leu Leu
50          55          60
Trp Ile Lys Ala Arg Asp Leu Lys Pro Arg Ala Ser Glu Pro Phe Leu
65          70          75          80
Leu Gln Ala Leu Val Leu Val His Asn Leu Phe Cys Phe Ala Leu Ser
85          90          95
Leu Tyr Met Cys Val Gly Ile Ala Tyr Gln Ala Ile Thr Trp Arg Tyr
100         105         110
Ser Leu Trp Gly Asn Ala Tyr Asn Pro Lys His Lys Glu Met Ala Ile
115        120        125
Leu Val Tyr Leu Phe Tyr Met Ser Lys Tyr Val Glu Phe Met Asp Thr
130        135        140
Val Ile Met Ile Leu Lys Arg Ser Thr Arg Gln Ile Ser Phe Leu His
145        150        155        160
Val Tyr His His Ser Ser Ile Ser Leu Ile Trp Trp Ala Ile Ala His
165        170        175
His Ala Pro Gly Gly Glu Ala Tyr Trp Ser Ala Ala Leu Asn Ser Gly
180        185        190
Val His Val Leu Met Tyr Ala Tyr Tyr Phe Leu Ala Ala Cys Leu Arg
195        200        205
Ser Ser Pro Lys Leu Lys Asn Lys Tyr Leu Phe Trp Gly Arg Tyr Leu
210        215        220
Thr Gln Phe Gln Met Phe Gln Phe Met Leu Asn Leu Val Gln Ala Tyr
225        230        235        240
Tyr Asp Met Lys Thr Asn Ala Pro Tyr Pro Gln Trp Leu Ile Lys Ile
245        250        255
Leu Phe Tyr Tyr Met Ile Ser Leu Leu Phe Leu Phe Gly Asn Phe Tyr

```

-continued

260	265	270	
Val Gln Lys Tyr Ile Lys Pro Ser Asp Gly Lys Gln Lys Gly Ala Lys			
275	280	285	
Thr Glu			
290			
 <210> SEQ ID NO 23			
<211> LENGTH: 1197			
<212> TYPE: DNA			
<213> ORGANISM: Phytophthora sojae			
 <400> SEQUENCE: 23			
atggctat	ttt tgaaccctga	ggctgattct	gctgctaacc tcgctactga ttctgaggct 60
aagcaaagac	aattggctga	ggctggatac	actcatgttg aggggtgctcc tgetcctttg 120
cctttggagt	tgcctcattt	ctctctcaga	gatctcagag ctgctattcc taagcactgc 180
ttcgagagat	ctttcgtgac	ctccacctac	tacatgatca agaacgtgtt gacttgcgct 240
gctttgttct	acgctgctac	cttcattgat	agagctggag ctgctgctta tgttttgtgg 300
cctgtgtact	ggttcttcca	gggatcttac	ttgactggag tgtgggttat cgctcatgag 360
tgtggacatc	aggcttattg	ctctctcgag	gtgggtgaaca acttgattgg actcgtgttg 420
cattctgctt	tgttggtgcc	ttaccactct	tggagaatct ctcacagaaa gcaccattcc 480
aacactggat	cttgcgagaa	cgatgagggt	ttcggttcctg tgaccagatc tgtgttggtc 540
tcttcttgga	acgagacctt	ggaggattct	cctctctacc aactctaccg tatcgtgtac 600
atgttggttg	ttggatggat	gcctggatac	ctcttcttca acgctactgg acctactaag 660
tactggggaa	agtcctaggtc	tcacttcaac	ccttactccg ctatctatgc tgataggag 720
agatggatga	tcgtgctctc	cgatattttc	ttgggtggcta tgttggtgtt tttggtgct 780
ttggtgcaca	ctttctcctt	caacaccatg	gtgaagtctt acgtgggtgcc ttacttcatt 840
gtgaacgctt	acttggtgtt	gattacctac	ctccaacaca ccgataccta catccctcat 900
ttcagagagg	gagagtggaa	ttggttgaga	ggagctttgt gcactgtgga tagatcattt 960
gggtccattcc	tcgattctgt	gggtgcataga	atcgtggata cccatgtttg ccaccacatc 1020
ttctccaaga	tgcccttcta	tcattgctgag	gaggctacca acgctattaa gcctctcctc 1080
ggaaagtctc	acttgaagga	taccactcct	gttcctgttg ctctctggag atcttacacc 1140
cattgcaagt	tcgttgagga	tgatggaaag	gtggtgttct acaagaacaa gctctag 1197

<210> SEQ ID NO 24  
 <211> LENGTH: 398  
 <212> TYPE: PRT  
 <213> ORGANISM: Phytophthora sojae

<400> SEQUENCE: 24

Met Ala Ile Leu Asn Pro Glu Ala Asp Ser Ala Ala Asn Leu Ala Thr	
1 5 10 15	
Asp Ser Glu Ala Lys Gln Arg Gln Leu Ala Glu Ala Gly Tyr Thr His	
20 25 30	
Val Glu Gly Ala Pro Ala Pro Leu Pro Leu Glu Leu Pro His Phe Ser	
35 40 45	
Leu Arg Asp Leu Arg Ala Ala Ile Pro Lys His Cys Phe Glu Arg Ser	
50 55 60	

-continued

Phe	Val	Thr	Ser	Thr	Tyr	Tyr	Met	Ile	Lys	Asn	Val	Leu	Thr	Cys	Ala	
65					70					75					80	
Ala	Leu	Phe	Tyr	Ala	Ala	Thr	Phe	Ile	Asp	Arg	Ala	Gly	Ala	Ala	Ala	
				85					90					95		
Tyr	Val	Leu	Trp	Pro	Val	Tyr	Trp	Phe	Phe	Gln	Gly	Ser	Tyr	Leu	Thr	
			100					105					110			
Gly	Val	Trp	Val	Ile	Ala	His	Glu	Cys	Gly	His	Gln	Ala	Tyr	Cys	Ser	
		115					120					125				
Ser	Glu	Val	Val	Asn	Asn	Leu	Ile	Gly	Leu	Val	Leu	His	Ser	Ala	Leu	
		130				135						140				
Leu	Val	Pro	Tyr	His	Ser	Trp	Arg	Ile	Ser	His	Arg	Lys	His	His	Ser	
145					150					155					160	
Asn	Thr	Gly	Ser	Cys	Glu	Asn	Asp	Glu	Val	Phe	Val	Pro	Val	Thr	Arg	
			165						170					175		
Ser	Val	Leu	Ala	Ser	Ser	Trp	Asn	Glu	Thr	Leu	Glu	Asp	Ser	Pro	Leu	
		180						185					190			
Tyr	Gln	Leu	Tyr	Arg	Ile	Val	Tyr	Met	Leu	Val	Val	Gly	Trp	Met	Pro	
		195					200					205				
Gly	Tyr	Leu	Phe	Phe	Asn	Ala	Thr	Gly	Pro	Thr	Lys	Tyr	Trp	Gly	Lys	
		210				215					220					
Ser	Arg	Ser	His	Phe	Asn	Pro	Tyr	Ser	Ala	Ile	Tyr	Ala	Asp	Arg	Glu	
225					230					235					240	
Arg	Trp	Met	Ile	Val	Leu	Ser	Asp	Ile	Phe	Leu	Val	Ala	Met	Leu	Ala	
			245						250					255		
Val	Leu	Ala	Ala	Leu	Val	His	Thr	Phe	Ser	Phe	Asn	Thr	Met	Val	Lys	
			260					265					270			
Phe	Tyr	Val	Val	Pro	Tyr	Phe	Ile	Val	Asn	Ala	Tyr	Leu	Val	Leu	Ile	
		275					280					285				
Thr	Tyr	Leu	Gln	His	Thr	Asp	Thr	Tyr	Ile	Pro	His	Phe	Arg	Glu	Gly	
		290				295					300					
Glu	Trp	Asn	Trp	Leu	Arg	Gly	Ala	Leu	Cys	Thr	Val	Asp	Arg	Ser	Phe	
305					310					315					320	
Gly	Pro	Phe	Leu	Asp	Ser	Val	Val	His	Arg	Ile	Val	Asp	Thr	His	Val	
			325						330					335		
Cys	His	His	Ile	Phe	Ser	Lys	Met	Pro	Phe	Tyr	His	Cys	Glu	Glu	Ala	
			340					345					350			
Thr	Asn	Ala	Ile	Lys	Pro	Leu	Leu	Gly	Lys	Phe	Tyr	Leu	Lys	Asp	Thr	
		355				360						365				
Thr	Pro	Val	Pro	Val	Ala	Leu	Trp	Arg	Ser	Tyr	Thr	His	Cys	Lys	Phe	
		370				375					380					
Val	Glu	Asp	Asp	Gly	Lys	Val	Val	Phe	Tyr	Lys	Asn	Lys	Leu			
385					390					395						

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 819

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Thalassiosira pseudonana

&lt;400&gt; SEQUENCE: 25

atggatgctt ataacgctgc tatggataag attggagctg ctatcatcga ttggagtgat 60

ccagatggaa agttcagagc tgatagggag gattggtggt tgtgcgattt cagatccgct 120

atcaccattg ctctcatcta catcgctttc gtgatcttgg gatctgctgt gatgcaatct 180

-continued

---

```

ctcccagcta tggaccata cccatcaag ttcctctaca acgtgtctca aatcttctc 240
tgcgcttaca tgactgttga ggctggattc ctgcttata ggaacggata caccgttatg 300
ccatgcaacc acttcaacgt gaacgatcca ccagttgcta acttgccttg gctcttctac 360
atctccaaag tgtgggattt ctgggatacc atcttcattg tgctcggaag gaagtggaga 420
caactctctt tcttgacagt gtaccatcat accaccatct tctcttctca ctggttgaac 480
gctaacgtgc tctacgatgg agatatcttc ttgaccatcc tctcaacgg attcattcac 540
accgtgatgt acacctacta cttcatctgc atgcacacca aggattctaa gaccggaaag 600
tctttgccaa tctggtgga gtcactcttg accgctttcc aactcttgca attcaccatc 660
atgatgtccc aagctaccta cttgggtttc cacggatgcg ataaggttcc cctcagaatc 720
accatcgtgt acttcgtgta cattctctcc cttttcttcc tcttcgctca gttcttcgtg 780
caatcctaca tggctccaaa gaagaagaag tccgcttga 819

```

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 272

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Thalassiosira pseudonana*

&lt;400&gt; SEQUENCE: 26

```

Met Asp Ala Tyr Asn Ala Ala Met Asp Lys Ile Gly Ala Ala Ile Ile
1           5           10          15
Asp Trp Ser Asp Pro Asp Gly Lys Phe Arg Ala Asp Arg Glu Asp Trp
20          25          30
Trp Leu Cys Asp Phe Arg Ser Ala Ile Thr Ile Ala Leu Ile Tyr Ile
35          40          45
Ala Phe Val Ile Leu Gly Ser Ala Val Met Gln Ser Leu Pro Ala Met
50          55          60
Asp Pro Tyr Pro Ile Lys Phe Leu Tyr Asn Val Ser Gln Ile Phe Leu
65          70          75          80
Cys Ala Tyr Met Thr Val Glu Ala Gly Phe Leu Ala Tyr Arg Asn Gly
85          90          95
Tyr Thr Val Met Pro Cys Asn His Phe Asn Val Asn Asp Pro Pro Val
100         105         110
Ala Asn Leu Leu Trp Leu Phe Tyr Ile Ser Lys Val Trp Asp Phe Trp
115         120         125
Asp Thr Ile Phe Ile Val Leu Gly Lys Lys Trp Arg Gln Leu Ser Phe
130         135         140
Leu His Val Tyr His His Thr Thr Ile Phe Leu Phe Tyr Trp Leu Asn
145         150         155         160
Ala Asn Val Leu Tyr Asp Gly Asp Ile Phe Leu Thr Ile Leu Leu Asn
165         170         175
Gly Phe Ile His Thr Val Met Tyr Thr Tyr Tyr Phe Ile Cys Met His
180         185         190
Thr Lys Asp Ser Lys Thr Gly Lys Ser Leu Pro Ile Trp Trp Lys Ser
195         200         205
Ser Leu Thr Ala Phe Gln Leu Leu Gln Phe Thr Ile Met Met Ser Gln
210         215         220
Ala Thr Tyr Leu Val Phe His Gly Cys Asp Lys Val Ser Leu Arg Ile
225         230         235         240

```

-continued

---

Thr Ile Val Tyr Phe Val Tyr Ile Leu Ser Leu Phe Phe Leu Phe Ala  
                   245                  250                  255  
 Gln Phe Phe Val Gln Ser Tyr Met Ala Pro Lys Lys Lys Lys Ser Ala  
                   260                  265                  270

<210> SEQ ID NO 27  
 <211> LENGTH: 1086  
 <212> TYPE: DNA  
 <213> ORGANISM: Phytophthora infestans

<400> SEQUENCE: 27

```

atggcgacga aggaggcgta tgtgttcccc actctgacgg agatcaagcg gtcgtacac 60
aaagactgtt tcgaggtctt ggtgcctctg tcgctctact acaccgtgcg ttgtctggtg 120
atcgcggtgg ctetaacctt cgggtctaac tacgtctcgcg ctctgcccga ggtcgagagc 180
ttctgggctc tggacgccgc actctgcacg ggctacatct tgctgcaggg catcgtgttc 240
tgggggcttct tcacggtggg ccacgatgcc ggccacggcg ccttctcgcg ctaccacctg 300
cttaacttcg tggtagggcac ttctatgcac tcgctcatcc tcacgccctt cgagtcgtgg 360
aagctcacgc accgtcacca ccacaagaac acgggcaaca ttgacctga cgaggtcttc 420
taccgcgaac gcaaggccga cgaccaccgc ctgtctcgca acctgattct ggcgtcggg 480
gcagcgtggc tcgcctattt ggtcgagggc ttccctcctc gtaaggtcaa ccacttcaac 540
ccgttcgagc ctctgttctg gcgtcagggt tcagctgtgg taatctctct tctcgccac 600
ttcttcgtgg ccggactctc catctatctg agcctccagc tgggcttaa gacgatggca 660
atctactact atggacctgt ttttgtgttc ggcagcatgc tggtcattac caccttcta 720
caccacaatg atgaggagac cccatggtac gccgactcgg agtggacgta cgtcaagggc 780
aacctctcgt ccgtggaccg atcgtacggc gcgtcattg acaacctgag ccacaacatc 840
ggcacgcacc agatccacca ccttttcctc atcattccgc actacaaact caagaaagcc 900
actgcggcct tccaccaggc tttccctgag ctggtgcgca agagcgacga gccaatatc 960
aaggctttct tccgggttgg acgtctctac gcaaactacg gcgttgtgga ccaggaggcg 1020
aagctcttca cgctaaagga agccaaggcg gcgaccgagg cggcggccaa gaccaagtcc 1080
acgtaa 1086

```

<210> SEQ ID NO 28  
 <211> LENGTH: 361  
 <212> TYPE: PRT  
 <213> ORGANISM: Phytophthora infestans

<400> SEQUENCE: 28

```

Met Ala Thr Lys Glu Ala Tyr Val Phe Pro Thr Leu Thr Glu Ile Lys
1           5           10          15
Arg Ser Leu Pro Lys Asp Cys Phe Glu Ala Ser Val Pro Leu Ser Leu
20          25          30
Tyr Tyr Thr Val Arg Cys Leu Val Ile Ala Val Ala Leu Thr Phe Gly
35          40          45
Leu Asn Tyr Ala Arg Ala Leu Pro Glu Val Glu Ser Phe Trp Ala Leu
50          55          60
Asp Ala Ala Leu Cys Thr Gly Tyr Ile Leu Leu Gln Gly Ile Val Phe
65          70          75          80
Trp Gly Phe Phe Thr Val Gly His Asp Ala Gly His Gly Ala Phe Ser

```

-continued

85								90				95			
Arg	Tyr	His	Leu	Leu	Asn	Phe	Val	Val	Gly	Thr	Phe	Met	His	Ser	Leu
			100						105				110		
Ile	Leu	Thr	Pro	Phe	Glu	Ser	Trp	Lys	Leu	Thr	His	Arg	His	His	His
			115				120					125			
Lys	Asn	Thr	Gly	Asn	Ile	Asp	Arg	Asp	Glu	Val	Phe	Tyr	Pro	Gln	Arg
	130					135					140				
Lys	Ala	Asp	Asp	His	Pro	Leu	Ser	Arg	Asn	Leu	Ile	Leu	Ala	Leu	Gly
145					150					155					160
Ala	Ala	Trp	Leu	Ala	Tyr	Leu	Val	Glu	Gly	Phe	Pro	Pro	Arg	Lys	Val
				165					170					175	
Asn	His	Phe	Asn	Pro	Phe	Glu	Pro	Leu	Phe	Val	Arg	Gln	Val	Ser	Ala
			180					185					190		
Val	Val	Ile	Ser	Leu	Leu	Ala	His	Phe	Phe	Val	Ala	Gly	Leu	Ser	Ile
		195					200					205			
Tyr	Leu	Ser	Leu	Gln	Leu	Gly	Leu	Lys	Thr	Met	Ala	Ile	Tyr	Tyr	Tyr
	210					215					220				
Gly	Pro	Val	Phe	Val	Phe	Gly	Ser	Met	Leu	Val	Ile	Thr	Thr	Phe	Leu
225					230					235					240
His	His	Asn	Asp	Glu	Glu	Thr	Pro	Trp	Tyr	Ala	Asp	Ser	Glu	Trp	Thr
				245					250					255	
Tyr	Val	Lys	Gly	Asn	Leu	Ser	Ser	Val	Asp	Arg	Ser	Tyr	Gly	Ala	Leu
			260					265					270		
Ile	Asp	Asn	Leu	Ser	His	Asn	Ile	Gly	Thr	His	Gln	Ile	His	His	Leu
	275						280					285			
Phe	Pro	Ile	Ile	Pro	His	Tyr	Lys	Leu	Lys	Lys	Ala	Thr	Ala	Ala	Phe
	290					295					300				
His	Gln	Ala	Phe	Pro	Glu	Leu	Val	Arg	Lys	Ser	Asp	Glu	Pro	Ile	Ile
305					310					315					320
Lys	Ala	Phe	Phe	Arg	Val	Gly	Arg	Leu	Tyr	Ala	Asn	Tyr	Gly	Val	Val
				325					330					335	
Asp	Gln	Glu	Ala	Lys	Leu	Phe	Thr	Leu	Lys	Glu	Ala	Lys	Ala	Ala	Thr
		340						345					350		
Glu	Ala	Ala	Ala	Lys	Thr	Lys	Ser	Thr							
	355						360								

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 1434

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Claviceps purpurea

&lt;400&gt; SEQUENCE: 29

```

atggctgcta ctacctctgc tatgagcaag gatgctgttc ttagaagaac tgctgctgct    60
actactgcta tcgatcacga aagctctacc tctgcttctc cagctgattc tcttagactc    120
tctgcttctt ctacctctct ctcttctctc agctctctcg acgctaagga taaggatgat    180
gagtacgctg gacttcttga tacttacgga aacgctttca cccctcctga tttcactatc    240
aaggatatca gagatgctat cctaagcac tgcttcgagc gttctgctat caagggatac    300
gcttatatcc tcagagatgt ggcttgccct tctaccactt tctacctctt ccacaacttc    360
gttaccctcg agaagcttcc ttacaccctt cttagagttt tctctggggg agttttacac    420
gctcttcagg gacttttcgg aactggactc tggattatcg ctcacgagtg tggacacggt    480

```



-continued

```

gctttctctc cttctaccct cactaacgat cttactggat gggttctcca ctctgctctt 540
ctcgtgcctt acttctcttg gaagttctct cactctgctc accacaaggg aaccggaaat 600
atggaaaggg atatggcttt cctccctaga actagggtct aatacgctac cagattcgga 660
agagctatgg atcagcttgg agatctttgc gaggaaaccc ctatctacac tgctggatct 720
cttgttttcc agcagcttct tggatggcct tcttacttga tcgctaacgt tactggacac 780
gatcttcacg agagacagag agaggggaaga ggaaaggga agaagaacgg attcggagga 840
actgttaacc acttcgacct tcgttctcct atcttcgatg acaagcacgc taagtttacc 900
gttctcagcg atatcggaat tggacttgct atcgctgctc ttgtttacct cggaacaga 960
ttcggatggg ctaacgttgc tggttggtac ttcgttcctt acctctgggt taaccactgg 1020
atcgttgcta tcactttcct tcagcacact gatcctactc ttctcacta cactgctgag 1080
gaatggaact tcgttcgtgg agctgctgct acaatcgata gagagatggg atttatcggt 1140
agacacctct tccacggaat cggtgagact cactgctctc accactacgt ttcttcaatc 1200
cctttctaca acgctgatga ggcttctgag gctatcaagc ctgttatggg aaagcactac 1260
cggtctgaga ctaaggatgg acctatgggt ttatcaggg ctttgtggaa aactgctaga 1320
tggtgtcaat gggttgagcc ttctgctgat gctcaagtg ctggtgaagg tgttctcttc 1380
ttcaggaaca gaaacggact tggaactaag cctatctcta tgaggacca gtga 1434

```

&lt;210&gt; SEQ ID NO 30

&lt;211&gt; LENGTH: 477

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Claviceps purpurea*

&lt;400&gt; SEQUENCE: 30

```

Met Ala Ala Thr Thr Ser Ala Met Ser Lys Asp Ala Val Leu Arg Arg
1           5           10          15
Thr Ala Ala Ala Thr Thr Ala Ile Asp His Glu Ser Ser Thr Ser Ala
20          25          30
Ser Pro Ala Asp Ser Pro Arg Leu Ser Ala Ser Ser Thr Ser Leu Ser
35          40          45
Ser Leu Ser Ser Leu Asp Ala Lys Asp Lys Asp Asp Glu Tyr Ala Gly
50          55          60
Leu Leu Asp Thr Tyr Gly Asn Ala Phe Thr Pro Pro Asp Phe Thr Ile
65          70          75          80
Lys Asp Ile Arg Asp Ala Ile Pro Lys His Cys Phe Glu Arg Ser Ala
85          90          95
Ile Lys Gly Tyr Ala Tyr Ile Leu Arg Asp Val Ala Cys Leu Ser Thr
100         105         110
Thr Phe Tyr Leu Phe His Asn Phe Val Thr Pro Glu Asn Val Pro Tyr
115         120         125
Thr Pro Leu Arg Val Phe Leu Trp Gly Val Tyr Thr Ala Leu Gln Gly
130         135         140
Leu Phe Gly Thr Gly Leu Trp Ile Ile Ala His Glu Cys Gly His Gly
145         150         155         160
Ala Phe Ser Pro Ser Thr Leu Thr Asn Asp Leu Thr Gly Trp Val Leu
165         170         175
His Ser Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Phe Ser His Ser
180         185         190

```

-continued

---

Ala	His	His	Lys	Gly	Thr	Gly	Asn	Met	Glu	Arg	Asp	Met	Ala	Phe	Leu
	195						200					205			
Pro	Arg	Thr	Arg	Ala	Gln	Tyr	Ala	Thr	Arg	Phe	Gly	Arg	Ala	Met	Asp
	210					215					220				
Gln	Leu	Gly	Asp	Leu	Cys	Glu	Glu	Thr	Pro	Ile	Tyr	Thr	Ala	Gly	Phe
225					230					235				240	
Leu	Val	Phe	Gln	Gln	Leu	Leu	Gly	Trp	Pro	Ser	Tyr	Leu	Ile	Ala	Asn
			245						250					255	
Val	Thr	Gly	His	Asp	Leu	His	Glu	Arg	Gln	Arg	Glu	Gly	Arg	Gly	Lys
			260					265						270	
Gly	Lys	Lys	Asn	Gly	Phe	Gly	Gly	Thr	Val	Asn	His	Phe	Asp	Pro	Arg
			275					280					285		
Ser	Pro	Ile	Phe	Asp	Asp	Lys	His	Ala	Lys	Phe	Ile	Val	Leu	Ser	Asp
	290					295					300				
Ile	Gly	Leu	Gly	Leu	Ala	Ile	Ala	Ala	Leu	Val	Tyr	Leu	Gly	Asn	Arg
305					310					315				320	
Phe	Gly	Trp	Ala	Asn	Val	Ala	Val	Trp	Tyr	Phe	Val	Pro	Tyr	Leu	Trp
				325					330					335	
Val	Asn	His	Trp	Ile	Val	Ala	Ile	Thr	Phe	Leu	Gln	His	Thr	Asp	Pro
			340					345						350	
Thr	Leu	Pro	His	Tyr	Thr	Ala	Glu	Glu	Trp	Asn	Phe	Val	Arg	Gly	Ala
		355					360					365			
Ala	Ala	Thr	Ile	Asp	Arg	Glu	Met	Gly	Phe	Ile	Gly	Arg	His	Leu	Phe
	370					375					380				
His	Gly	Ile	Val	Glu	Thr	His	Val	Leu	His	His	Tyr	Val	Ser	Ser	Ile
385				390						395					400
Pro	Phe	Tyr	Asn	Ala	Asp	Glu	Ala	Ser	Glu	Ala	Ile	Lys	Pro	Val	Met
			405					410						415	
Gly	Lys	His	Tyr	Arg	Ser	Glu	Thr	Lys	Asp	Gly	Pro	Met	Gly	Phe	Ile
			420					425					430		
Arg	Ala	Leu	Trp	Lys	Thr	Ala	Arg	Trp	Cys	Gln	Trp	Val	Glu	Pro	Ser
	435					440						445			
Ala	Asp	Ala	Gln	Gly	Ala	Gly	Glu	Gly	Val	Leu	Phe	Phe	Arg	Asn	Arg
	450				455						460				
Asn	Gly	Leu	Gly	Thr	Lys	Pro	Ile	Ser	Met	Arg	Thr	Gln			
465					470					475					

&lt;210&gt; SEQ ID NO 31

&lt;211&gt; LENGTH: 903

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Ostreococcus tauri*

&lt;400&gt; SEQUENCE: 31

atgtctgctt	ctggagcttt	gttgctgct	attgctttcg	ctgcttacgc	ttacgetacc	60
tacgcttatg	ctttcgagtg	gtctcatgct	aacggaatcg	ataacgtgga	tgctagagag	120
tggattggag	ctttgtcttt	gagactccct	gcaattgcta	ccaccatgta	cctcttgctt	180
tgccttgctg	gacctagatt	gatggctaag	agggaggctt	ttgacctaag	gggattcatg	240
ctcgttaca	acgcttacca	aaccgctttc	aacgttgctg	tgctcggaat	gttcgctaga	300
gagatctctg	gattgggaca	acctgtttgg	ggatctacta	tgccttgag	cgataggaag	360
tccttcaaga	ttttgtggg	agtggtgctc	cattacaaca	ataagtaact	cgagttgttg	420

-continued

---

```

gatactgtgt tcattgtggc taggaaaaag accaagcagc tctctttctt gcatgtgtac 480
catcatgctt tgttgatttg ggcttgggtg cttgtttgtc atctcatggc taccaacgat 540
tgcacgatg cttatttcgg agctgcttgc aactctttca tccacatcgt gatgtactcc 600
tactacctca tgtctgcttt gggaattaga tgcccttgga agagatatat caccaggct 660
cagatgttgc aattcgtgat cgtgttcgct catgctgttt tcgtgctcag acaaaagcac 720
tgccctgtta ctttgccttg ggcacaaatg ttcgtgatga caaatatggt ggtgctcttc 780
ggaaacttct acctcaaggc ttactctaac aagtctaggg gagatggagc ttcttctgtt 840
aagcctgctg agactactag agcaccttct gtgagaagaa ccaggctccag gaagatcgat 900
tga 903

```

```

<210> SEQ ID NO 32
<211> LENGTH: 300
<212> TYPE: PRT
<213> ORGANISM: Ostreococcus tauri

```

```

<400> SEQUENCE: 32

```

```

Met Ser Ala Ser Gly Ala Leu Leu Pro Ala Ile Ala Phe Ala Ala Tyr
1      5      10      15
Ala Tyr Ala Thr Tyr Ala Tyr Ala Phe Glu Trp Ser His Ala Asn Gly
20     25     30
Ile Asp Asn Val Asp Ala Arg Glu Trp Ile Gly Ala Leu Ser Leu Arg
35     40     45
Leu Pro Ala Ile Ala Thr Thr Met Tyr Leu Leu Phe Cys Leu Val Gly
50     55     60
Pro Arg Leu Met Ala Lys Arg Glu Ala Phe Asp Pro Lys Gly Phe Met
65     70     75     80
Leu Ala Tyr Asn Ala Tyr Gln Thr Ala Phe Asn Val Val Val Leu Gly
85     90     95
Met Phe Ala Arg Glu Ile Ser Gly Leu Gly Gln Pro Val Trp Gly Ser
100    105    110
Thr Met Pro Trp Ser Asp Arg Lys Ser Phe Lys Ile Leu Leu Gly Val
115    120    125
Trp Leu His Tyr Asn Asn Lys Tyr Leu Glu Leu Leu Asp Thr Val Phe
130    135    140
Met Val Ala Arg Lys Lys Thr Lys Gln Leu Ser Phe Leu His Val Tyr
145    150    155    160
His His Ala Leu Leu Ile Trp Ala Trp Trp Leu Val Cys His Leu Met
165    170    175
Ala Thr Asn Asp Cys Ile Asp Ala Tyr Phe Gly Ala Ala Cys Asn Ser
180    185    190
Phe Ile His Ile Val Met Tyr Ser Tyr Tyr Leu Met Ser Ala Leu Gly
195    200    205
Ile Arg Cys Pro Trp Lys Arg Tyr Ile Thr Gln Ala Gln Met Leu Gln
210    215    220
Phe Val Ile Val Phe Ala His Ala Val Phe Val Leu Arg Gln Lys His
225    230    235    240
Cys Pro Val Thr Leu Pro Trp Ala Gln Met Phe Val Met Thr Asn Met
245    250    255
Leu Val Leu Phe Gly Asn Phe Tyr Leu Lys Ala Tyr Ser Asn Lys Ser

```

260							265					270				
Arg	Gly	Asp	Gly	Ala	Ser	Ser	Val	Lys	Pro	Ala	Glu	Thr	Thr	Arg	Ala	
		275					280					285				
Pro	Ser	Val	Arg	Arg	Thr	Arg	Ser	Arg	Lys	Ile	Asp					
	290					295					300					

- a) the nucleic acid sequence of SEQ ID NO: 1, 3, 5, 7 or 9;
- b) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, 4, 6, 8 or 10;
- c) a nucleic acid sequence having at least 70% sequence identity to the nucleic acid sequence of a) or b), wherein said nucleic acid sequence encodes a polypeptide having desaturase or elongase activity;

d) a nucleic acid sequence encoding a polypeptide having desaturase or elongase activity and comprising an amino acid sequence which has at least 82% sequence identity to an amino acid sequence encoded by any one of the nucleic acid sequences of a) to c); and

e) a nucleic acid sequence which is capable of hybridizing under stringent conditions to any one of the nucleic acid sequences of a) to d), wherein said nucleic acid sequence encodes a polypeptide having desaturase or elongase activity,

wherein said expression control sequence is heterologous to said nucleic acid sequence.

\* \* \* \* \*